

Table 216-1.--Walleye pollock harvest (round lb) in groundfish fisheries in Southeast Alaska management areas (Central Southeast Outside (CSEO), Southern Southeast Outside (SSEO), East Yakutat (EYKT), Northern Southeast Inside (NSEI) and Southern Southeast Inside (SSEI)). Pelagic and non-pelagic trawl, longline and mechanical jig gears are combined here. Prior to 1998, the bulk of the harvest is from pelagic trawl gear; after 1998 a trawl ban was enacted in federal waters of the Eastern Gulf of Alaska (CSEO, SSEO, EYKT). Gear types are combined to minimize confidential data.

year	CSEO	EYKT	NSEI	NSEO	SSEI	SSEO
1985	-	-	360	-	confidential	-
1986	-	-	4,115	-	766	-
1987	-	-	3,933	-	4,913	confidential
1988	1,002	8,543	3,405	-	34	confidential
1989	19,139	14,726	confidential	confidential	confidential	2,500
1990	92,162	28,582	11,620	confidential	7,162	16,501
1991	10,015	47,085	7,457	confidential	103	confidential
1992	9,554	4,104	2,283	confidential	confidential	-
1993	-	-	4,370	-	1,505	-
1994	-	-	confidential	-	-	-
1995	-	-	2,647	-	confidential	confidential
1996	-	confidential	2,157	-	-	confidential
1997	confidential	-	30,412	-	-	confidential
1998	-	-	24,728	126	confidential	-
1999	confidential	-	17,049	-	-	-
2000	confidential	-	4,347	-	confidential	-
2001	-	-	confidential	-	-	-
2002	-	-	26	-	-	-
2003	-	-	-	-	-	-
2004	-	-	-	confidential	-	-
2005	-	confidential	-	confidential	-	-
2007	-	-	169	confidential	-	-
2008	-	confidential	1,128	confidential	-	-
2009	-	-	183	confidential	-	-
2010	-	confidential	4,567	confidential	-	-
2011	-	-	confidential	-	-	-
2012	-	-	confidential	-	-	-
2013	-	-	confidential	-	-	-
2014	-	-	confidential	-	-	-
2015	-	134,401	confidential	confidential	confidential	-




Figure 216-1. Groundfish management areas in Southeast Alaska. NMFS Areas 650 includes EYKT, NSEO, CSEO, and SSEO, while Area 659 encompasses portions of NSEI and SSEI.