Annual Report to the Alaska Board of Game on Intensive Management for Moose with Wolf, Black Bear, and Grizzly Bear Predation Control in Game Management Unit 19D East

Prepared by the Division of Wildlife Conservation February 2019



- 1) Description of IM Program¹
 - A) This report is an annual evaluation for a predation control program authorized by the Alaska Board of Game (Board) under 5 AAC 92.123
 - B) Month this report was submitted by the Department to the Board:

Annual Report, February 2019

- C) Program name: Unit 19D East wolf and bear predation control program (Fig. 1)
- D) Existing program has an associated Operational Plan
- E) Game Management Unit fully or partly included in IM program area: Unit 19D (East)
- F) IM objective is a moose population size of 6,000–8,000 with a harvest of 400–600.
- G) Month and year the current predation control program was originally authorized by the Board: Fall 1995. Indicate date(s) if renewed: January 2000, March 2003, January 2006, May 2006, March 2009, February 2014.
- H) Predation control is currently active in this IM area.
- I) If active, month and year the current predation control program began: December 2003
- J) A habitat management program funded by the Department or from other sources is currently active in this IM area: No
- **K) Size of IM program area (square miles) and geographic description:** Unit 19D East is 8,513 mi²
- L) Size and geographic description of area for assessing ungulate abundance: Wolf Control Focus Area (WCFA) is 4,484 mi²; Bear Control Focus Area (BCFA) is 528 mi².
- M) Size and geographic description of area for ungulate harvest reporting: WCFA is 4.484 mi².
- N) Size and geographic description of area for assessing predator abundance: WCFA is 4,484 mi²; BCFA is 528 mi².
- O) Size and geographic description of predation control area: WCFA is 4,484 mi²; BCFA is 528 mi².
- P) Criteria for evaluating progress toward IM objectives: Moose abundance and harvest.

¹ For purpose and context of this report format, see *Intensive Management Protocol, section on Tools for Program Implementation and Assessment*

- **Q)** Criteria for success with this program: BCFA abundance=2.0 moose/mi² (~1,100 moose) moose and WCFA harvest=180 moose.
- **R)** Department recommendation for IM program in this reporting period: Continue program (details provided in section 6).




Figure 1. Unit 19D (East) intensive management area.

2) Prey data

Date(s) and method of most recent abundance assessment for moose: November 2017 Geospatial Population Estimator (GSPE) in a 1,118 mi² area surrounding the BCFA.

Compared to IM area, was a similar trend and magnitude of difference in abundance observed in nearby non-treatment area(s) since program inception: Non-treatment area not established.

Date(s) of most recent age and sex composition survey: November 2017 Geospatial Population Estimator in a 1,118 mi² area surrounding the BCFA.

Compared to IM area, was a similar composition trend and magnitude of difference in composition observed in nearby non-treatment area(s) since program inception: Non-treatment area not established.

Table 1. Moose abundance, age and sex composition in a 1,118 mi² area surrounding the BCFA since program implementation in year 1. Regulatory year is 1 July to 30 June (e.g.,

Regulatory Year 2001 is 1 July 2001 to 30 June 2002).

			Composition (number per 100 Cow					
Period	Regu- latory Year	Abundance ^a (90% CI)	Calves (90% CI)	Yearling Bulls (90% CI)	Bulls (90% CI)	Total n		
Year 1	2001	868(<u>+</u> 147)	36(±10)	8(<u>+</u> 3)	21(<u>+</u> 6)	455		
Year 2	2002							
Year 3	2003							
Year 4	2004	1,192(<u>+</u> 228)	66(<u>+</u> 18)	8(<u>+</u> 4)	18(<u>+</u> 6)	578		
Year 5	2005							
Year 6	2006	1,308(<u>+</u> 174)	55(<u>+</u> 10)	12(<u>+3</u>)	30(<u>+</u> 8)	762		
Year 7	2007	1,720(<u>+</u> 306)	53(<u>+</u> 14)	15(<u>+</u> 4)	36(<u>+</u> 10)	844		
Year 8	2008	1,718(<u>+</u> 352)	44(<u>+</u> 12)	14(<u>+</u> 5)	40(<u>+</u> 11)	678		
Year 9	2009	$1,820(\pm 323)$	38(±10)	11(±4)	40(±11)	711		
Year 10	2010	1,796(±312)	43(±11)	16(±5)	49(±13)	712		
Year 11	2011	$1,647(\pm 296)$	42(±11)	10(±3)	$33(\pm 10)$	639		
Year 12	2012	$1,337(\pm 199)$	35(±11)	7(±2)	38(±5)	650		
Year 13	2013	-			-			
Year 14	2014	-			-			
Year 15	2015	2,014(±398)	41(±12)	14(±5)	36(±11)	811		
Year 16	2016							
Year 17	2017	2,389 (±372)	42 (±3)	N/A	31 (±6)	1089		
Year 18	2018							

^aEstimate with sightability correction factor applied

Describe trend in abundance or composition: Results of the trend analysis for regulatory years 2001–2009 indicate a statistically significant increasing linear trend in abundance within a 1,118 mi² area surrounding the BCFA (115 moose/year, SE=19.2, P=0.004). The most recent survey in 2017 indicated the highest estimate in the program to date.

Table 2. Moose harvest from a 1,118 mi² area surrounding the BCFA since program implementation in year 1. Moose harvest from WCFA (4,484 mi²) since year 13. Regulatory year is 1 July to 30 June (e.g., Regulatory Year 2001 is 1 July 2001 to 30 June 2002).

	Regulatory	Reported		Other	
Period	Year	Male	Female	mortality ^a	Total
Year 1	2001	29	0	_b	29
Year 2	2002	23	0	_b	23
Year 3	2003	32	0	_b	32
Year 4	2004	7	0	_b	7
Year 5	2005	14	0	_b	14
Year 6	2006	12	0	3	15
Year 7	2007	25	0	1	26
Year 8	2008	61	0	1	62
Year 9	2009	56	0	2	58
Year 10	2010	50	0	2	52
Year 11	2011	100	0	1	101
Year 12	2012	73	0	1	74
Year 13	2013	94	1	2	97
Year 14	2014	111	0	3	114
Year 15	2015	162	1	2	164
Year 16	2016	143	0	2	145
Year 17	2017	138	0	2	140
Year 18	2018	107	0	2	109

^a Mortuary harvest

Describe trend in harvest: Harvest reached a peak in Regulatory Year 2015. The lower harvest in Regulatory Year 2018 was likely due in part to poor weather.

Describe any other harvest related trend if appropriate: None.

3) Predator data

Wolves

Date(s) and method of most recent spring abundance assessment for wolves in the WCFA: March 2018, pilot interviews.

Date(s) and method of most recent fall abundance assessment for wolves in the WCFA: Calculated for fall 2017 by adding total removal from WCFA to spring 2018 abundance estimate.

Other research or evidence of trend or abundance status in wolves:

Gardner, C. L., and N. J. Pamperin. 2014. Intensive aerial wolf survey operations manual for Interior Alaska. Alaska Department of Fish and Game, Wildlife Special Publication ADF&G/DWC/WSP-2014-01, Juneau.

^b Records destroyed by fire

^c Preliminary data

- Keech, M. A., M. S. Lindberg, R. D. Boertje, P. Valkenburg, B. D. Taras, T. A. Boudreau, K. B. Beckmen. 2011. Effects of Predator Treatments, Individual Traits, and Environment on Moose Survival in Alaska. The Journal of Wildlife Management 75(6):1361–1380.
- Keech, M. A. 2012. Response of moose and their predators to wolf reduction and short-term bear removal in a portion of Unit 19D. Alaska Department of Fish and Game, Federal Aid in Wildlife Restoration, Final Wildlife Research Report ADF&G/DWC/WRR-2012-#, Grants W-33-4 through W-33-10, Project 1.62, Juneau, Alaska

Table 3. Wolf abundance objectives and removal in Wolf Control Focus Area (WCFA) since program implementation in year 1. Removal objectives are to reduce wolf numbers as low as possible in the WCFA and to maintain a minimum of 40 wolves in all of Unit 19D East to ensure wolves persist in the unit. The current WCFA was established in RY 2009. Prior to RY 2009, control was conducted in various different geographic areas. All values listed are for the current WCFA. Regulatory year (RY) is 1 July to 30 June (e.g, RY 2001 is 1 July 2001 to 30 June 2002).

	Regu- latory	Fall	Harvest removal		Dept.	Public control	Total	Spring
Period	Year	abundance ^a	Trap	Hunt	removal	removal ^b	removal	abundance
Year 1	2001	89	19	3	0	N/A	22	67 ^c
Year 2	2002		28	5	0	N/A	33	
Year 3	2003		9	1	0	17	27	
Year 4	2004		12	2	0	12	26	
Year 5	2005	26	9	1	0	3	13	13°
Year 6	2006	29	13	1	0	2	16	13°
Year 7	2007		6	2	0	19	27	
Year 8	2008		4	3	0	19	26	
Year 9	2009	37	7	4	0	4	15	22 ^c
Year 10	2010		4	2	0	13	19	
Year 11	2011	55–57	11	0	0	22	33	22-24 ^d
Year 12	2012	33	5	0	0	8	13	20 ^d
Year 13	2013	27	9	0	0	9	18	9 ^d
Year 14	2014	42	13	0	0	10	23	19 ^d
Year 15	2015		18	1	0	12	31	
Year 16	2016	105	14	0	0	3	17	88°
Year 17	2017	75	22	1_	0	21	44	31 ^d
Year 18	2018 ^e		3	2	0	7	12	

^aCalculated by adding total removal to WCFA spring abundance during each regulatory year.

^bPublic control removal began in regulatory year 2003

[°]Calculated by extrapolating density within a 3,210 mi² aerial reconnaissance survey area within the WCFA to the entire WCFA

^dAbundance based on private pilot and department biologist observations.

^ePreliminary

Black Bears

Date(s) and method of most recent spring abundance assessment for black bears in the BCFA: May 2014, mark-recapture estimator

Date(s) and method of most recent fall abundance assessment for black bears in the BCFA: August 2014, calculated for fall 2014 by subtracting total removal in RY13 from May 2014 abundance estimate

Other research or evidence of trend or abundance status in black bears:

Keech, M. A., M. S. Lindberg, R. D. Boertje, P. Valkenburg, B. D. Taras, T. A. Boudreau, K. B. Beckmen. 2011. Effects of Predator Treatments, Individual Traits, and Environment on Moose Survival in Alaska. The Journal of Wildlife Management 75(6):1361–1380

Keech, M. A. 2012. Response of moose and their predators to wolf reduction and short-term bear removal in a portion of Unit 19D. Alaska Department of Fish and Game, Federal Aid in Wildlife Restoration, Final Wildlife Research Report ADF&G/DWC/WRR-2012-#, Grants W-33-4 through W-33-10, Project 1.62, Juneau, Alaska.

Table 4. Black bear abundance and removal in Bear Control Focus Area (BCFA) since program implementation in year 1. Public bear control ended in RY14. When active the removal objective is to reduce bear numbers as low as possible within the BCFA. The May 2004 estimated black bear population for all of Unit 19D (East) was approximately 1,700. The regulatory year is 1 July to 30 June (e.g, RY 2001 is 1 July 2001 to 30 June 2002).

					De	pt.	Pub	lic		
	Regu-	Spring	Har	vest	con	trol	cont	rol		Fall
	latory	abundance ^a	rem		rem		remo		Total	abundance
Period	Year	(95% CI)	FAb	SPc	FAb	SPc	FAb	SPc	removal	a,d
Year 1	2001		1	0	0	0	0	0	1	
Year 2	2002	96(<u>+</u> 13) ^e	4	0	0	67 ^f	0	0	73	
Year 3	2003	$30(\pm 9)^{e}$	1	5	0	26 ^f	0	0	32	23
Year 4	2004		0	1	0	0	0	0	1	Near 0
Year 5	2005		1	5	0	0	0	0	6	
Year 6	2006	$70(\pm 14)^{g}$	0	0	0	0	0	0	0	
Year 7	2007		1	7	0	0	0	0	8	70
Year 8	2008		1	5	0	0	0	0	9	
Year 9	2009	$123(96-162)^g$	4	0	0	0	0	6	10	
Year 10	2010		1	3	0	0	4	13	21	113
Year 11	2011		7	1	0	0	1	2	11	
Year 12	2012		0	0	0	0	0	0	0	
Year 13	2013	113(89–149) ^g	1	1	0	0	4	0	6	107
Year 14	2014		13	2	0	0	0	0	13	
Year 15	2015		1	2	0	0	0	0	3	
Year 16	2016		0	5	0	0	0	0	5	
Year 17	2017		5	3	0	0	0	0	8	
Year 18	2018 ^h		2		0	0	0	0	2^{h}	

^aDoes not include cubs of the year

Brown Bears

Date(s) and method of most recent spring abundance assessment for brown bears in the BCFA: May 2004, Estimated by extrapolation from BCFA.

Date(s) and method of most recent fall abundance assessment for brown bears in the BCFA: November 2003, calculated by subtracting total removal from May 2004 abundance estimate.

^bFall

^cSpring

^dCalculated by subtracting total removal from spring abundance estimate in the previous RY

^eRemoval estimator

fNon-lethal removal

gMark-recapture estimator

^hPreliminary

Other research or evidence of trend or abundance status in brown bears:

Keech, M. A., M. S. Lindberg, R. D. Boertje, P. Valkenburg, B. D. Taras, T. A. Boudreau, K. B. Beckmen. 2011. Effects of Predator Treatments, Individual Traits, and Environment on Moose Survival in Alaska. The Journal of Wildlife Management 75(6):1361–1380

Keech, M. A. 2012. Response of moose and their predators to wolf reduction and short-term bear removal in a portion of Unit 19D. Alaska Department of Fish and Game, Federal Aid in Wildlife Restoration, Final Wildlife Research Report ADF&G/DWC/WRR-2012-#, Grants W-33-4 through W-33-10, Project 1.62, Juneau, Alaska.

Table 5. Brown bear abundance and removal in Bear Control Focus Area (BCFA) since program implementation in year 1. When active the removal objective is to reduce bear numbers as low as possible within the BCFA. Public bear control ended in RY14. The May 2004 estimated brown bear population for all of Unit 19D (East) was approximately 128. The regulatory year is 1 July to 30 June (e.g, Regulatory Year 2001 is 1 July 2001 to 30 June 2002).

					De	pt.	Pul	blic		
	Regu-		Har	vest	con	trol	control			
	latory	Spring	rem	oval	reme	oval	rem	oval	Total	Fall
Period	Year	abundance ^a	FAb	SP ^c	FAb	SP ^c	FAb	SP ^c	removal	abundance ^{a,d}
Year 1	2001	-	0	0	0	0	0	0	0	
Year 2	2002	12e	0	0	0	6 ^f	0	0	6	
Year 3	2003	-	0	0	0	0	0	0	0	6
Year 4	2004	-	0	0	0	0	0	0	0	
Year 5	2005	-	0	0	0	0	0	0	0	
Year 6	2006		0	2	0	0	0	0	2	
Year 7	2007		0	2	0	0	0	0	2	
Year 8	2008		0	0	0	0	0	0	0	
Year 9	2009		2	0	0	0	0	0	2	
Year 10	2010		0	0	0	0	0	0	0	
Year 11	2011		0	0	0	0	0	0	0	
Year 12	2012		0	0	0	0	0	0	0	
Year 13	2013		0	0	0	0	0	0	0	
Year 14	2014		1	1	0	0	0	0	2	
Year 15	2015		0	0	0	0	0	0	0	
Year 16	2016		2	0	0	0	0	0	2	
Year 17	2017		0	0	0	0	0	0	0	
Year 18	2018 ^g		0		0	0	0	0		

^aDoes not include cubs

^bFall

Spring

^dCalculated by subtracting total removal from spring abundance estimate in the previous regulatory year

^eEstimated by using density extrapolated from other areas of Interior Alaska with comparable habitat

fNon-lethal removal

^gPreliminary

4) Habitat data and nutritional condition of prey species

Where active habitat enhancement is occurring or was recommended in the Operational Plan, describe progress toward objectives: No active habitat enhancement occurring.

Table 6. Nutritional indicators for moose in a 1,118 mi² area surrounding the BCFA since program implementation in year 1. A regulatory year is 1 July to 30 June (e.g, Regulatory Year 2001 is 1 July 2001 to 30 June 2002).

	Regulatory	Twinning rate for radiocollared cows	Twinning rate uncollared cows
Period	year	>2 yrs (n)	(n)
Year 1	2001	59% (22)	39% (46)
Year 2	2002	24% (25)	36% (39)
Year 3	2003	32% (31)	39% (31)
Year 4	2004	44% (45)	50% (40)
Year 5	2005	40% (60)	35% (29)
Year 6	2006	52% (56)	50% (30)
Year 7	2007	55% (51)	-
Year 8	2008	33% (43)	26% (87)
Year 9	2009	33% (40)	29% (45)
Year 10	2010		37% (38)
Year 11	2011		34% (47)
Year 12	2012		21% (51)
Year 13	2013		
Year 14	2014		49% (45)
Year 15	2015		47% (36)
Year 16	2016		27% (44)
Year 17	2017		13% (39)

5) Costs specific to implementing Intensive Management

Table 7. Unit 19D East program cost (\$1,000 = 1.0) of agency salary based on estimate of proportional time of field level staff and cost of operations for intensive management activities (e.g., predator control or habitat enhancement beyond normal Survey and Inventory work) performed by personnel in the Department or work by other state agencies (e.g., Division of Forestry) or contractors in Unit 19D (East) during years 10-15. Fiscal year (FY) is also 1 July to 30 June but the year is one greater than the comparable regulatory year (e.g, FY 2011 is 1 July 2010 to 30 June 2011).

Predation controla Other IM activities **Total IM** Research Fiscal Time^b Year Costc Time Cost costd **Period** cost Year 10 2011 0.4 3.5 0.4 5.0 8.5 56.0 Year 11 2012 1.2 7.3 43.6 50.9 39.0 4.0 Year 12 2013 8.0 2.0 44.2 52.2 119.3 1.3 Year 13 2014 1.0 11.3 0.4 5.0 16.3 256 Year 14 2015 1.4 11.5 0.4 5.0 16.5 0 Year 15 2016 9.5 5.0 14.5 242.2 1.4 0.4 2017 1.4 9.5 5.0 242.2 Year 16 0.4 14.5 2018 1.2 7.9 Year 17 0.1 139.4 140.6 190.3

6) Evaluation (February 2018) for program renewal following Year 17 and Department recommendations for Unit 19D (East)

Has progress toward defined criteria been achieved? Yes. Moose population and harvest have increased compared to precontrol.

Has achievement of success criteria occurred? Population objective has been achieved, but harvest objective has not been achieved.

Recommendation for IM program: Continue program

Rationale for recommendation on overall program: Population objectives have been achieved, and progress towards harvest objectives has occurred. The program was modified during the February 2014 Board of Game meeting by eliminating the public bear control program due to insufficient bear removal; providing an option for department bear control; continuing public wolf control; establishing population criteria of 2.0 moose/mi² within the BCFA; establishing harvest criteria of 180 moose from within the WCFA; and evaluating harvest from within the WCFA. This predation control program expires June 30, 2020 and will be reevaluated at the Region III Board of Game meeting in February, 2020.

^aState or private funds only.

^bPerson-months (22 days per month)

^cSalary plus operations

^dSeparate from implementing IM program but beneficial for understanding of ecological or human response to management treatment (scientific approach that is not unique to IM).