Interim Report to the Alaska Board of Game on Intensive Management for Moose with Wolf Predation Control in Unit 15A

Prepared by the Division of Wildlife Conservation August 2015



- 1) Description of IM Program and Department recommendation for reporting period.
- A) This report is an annual evaluation for a predation control program authorized by the Alaska Board of Game (Board) under 5 AAC 92.118
- B) Month this report was submitted by the Department to the Board: August 2015
- C) Program name: GMU 15A Moose
- **D**) Existing program has an associated Operational Plan: Operational Plan for Intensive Management of Moose in Game Management Unit 15A During Regulatory years 2012-2017.
- E) Game Management Unit(s) fully or partly included in IM program area: GMU 15A
- F) IM objectives for Moose: Population size 3,000-3,500. Harvest 180-350.
- **G)** Month and year the current predation control program was originally authorized by the Board: January 2012, revised at the March 2013 Alaska Board of Game (BOG) meeting.
- H) Predation control is currently active in this IM area: Yes
- I) The current predation control program began: November 2013.
- J) A habitat management program funded by the Department or from other sources is currently active in this IM area: Yes
- **K**) Size of IM program area (square miles) and geographic description: 1,314 mi², GMU 15A
- L) Size and geographic description of area for assessing ungulate abundance: 1,314 mi², GMU 15A
- **M**) Size and geographic description of area for ungulate harvest reporting: 1,314mi², GMU 15A
- **N) Size and geographic description of area for assessing predator abundance:** 1,314 mi², GMU 15A
- **O)** Size and geographic description of predation control area: Approx. 49 mi², includes portions of Salamatof and Kenai Native Associations lands in GMU 15A
- **P)** Criteria for evaluating progress toward IM objectives: An increase in calf:cow ratio, increased survival of calves, no further population decline.
- **Q)** Criteria for success with this program: Increased calf survival and recruitment. The overall program will be successful when we attain IM population and harvest objectives in GMU 15A.

R) Department recommendation for IM program in this reporting period: Continue

Refer to one or more scaled maps in the Operational Plan for areas described in this section

2) Prey data

Date(s) and method of most recent abundance assessment for Moose:

Please refer to Figure 3 on page 5 of the Operational Plan for Intensive Management of Moose in Unit 15A. Prey data for RY 2012 include a November 2012 composition survey and a February 2013 GSPE population estimate. Data for RY 2013 is from a November/December composition survey (Table 1). Surveys during 2014 were not completed due to inadequate snow cover.

Compared to IM area, was a similar trend and magnitude of difference in abundance observed in nearby non-treatment area(s) **since program inception:** NO, we are in the first year of the program and it is premature to detect a difference.

Date(s) of most recent age and sex composition survey: November 25 – December 3, 2013.

Compared to IM area, was a similar composition trend and magnitude of difference in composition observed in nearby non-treatment area(s) since program inception: NO, we are in the first year of the program and it is premature to detect a difference.

Table 1. Moose abundance, age and sex composition in assessment area (L) since program implementation in year 1 to reauthorization review in year 2017 in GMU 15A. Regulatory year is 1 July to 30 June (e.g, RY 2012 is 1 July 2012 to 30 June 2013).

			Composition (number per 100 females)					
Period	RY	Abundance (variation)	Calves	Yearlings	Males	Total <i>n</i>		
	2012		25		30	372		
	2012	1569 (±13.4%; 95% C.I.)						
Year 1	2013		25		29	332		
Year 2	2014 ^a		NA		NA	NA		
Year 3	2015							
Year 4	2016							
Year 5	2017							

^a Composition flights were not completed due to inadequate snow cover.

Describe trend in abundance or composition: No noticeable difference in composition data between RY 2012 and RY 2013. Composition flights were not completed during RY 2014 due to lack of adequate snow cover. There are no census data for RY 2013 and RY 2014.

Period	RY	Reported		Estimated		Total harvest	Other mortality ^a	Total
		Male	Female	Unreported	Illegal	-		
	2012	6	0			6	107	113
Year 1	2013	30	0			30	87	117
Year 2	2014	41	0		10	51	68	119
Year 3	2015							
Year 4	2016							
Year 5	2017							

Table 2. Moose harvest in assessment area (M). Methods for estimating unreported harvest are described in Survey and Inventory reports.

^a vehicle mortality, and mortuary.

Describe trend in harvest: Please refer to Figure 4 on page 6 of the Operational Plan for Intensive Management of Moose in Unit 15A. The increased harvest from 2012 to 2013 was primarily attributed to liberalized antler configuration for a legal bull. In 2012 a legal bull had to have an antler spread of at least 50 inches or at least 4 brow tines on at least one side. In 2013 a bull with no more than a spike on at least one side was added to the 2012 definition for a legal bull to harvest.

Describe any other harvest related trend if appropriate: During 2012, 293 individuals reported hunting in GMU 15A and the reported legal harvest was 6 bulls (2% success rate). During 2013, 351 individuals reported hunting in GMU 15A and the reported legal harvest was 30 bulls (9% success rate). During 2014, 540 individuals reported hunting in GMU 15A and reported harvest was 41 bulls (8% success rate).

3) Predator data: Wolves

Dates and method of most recent spring abundance assessment for wolves:

Survey data for RY 2010 were collected in March 2011, for RY 2011 in November 2011, and for RY 2012 in February 2013. All of GMU 15A was flown and the total numbers are based on the number of wolves observed and an assessment of tracks observed. A partial survey was conducted during December 2013, but no surveys were flown during 2014 due to inadequate snow cover. It appears that wolf numbers have remained relatively constant since 2010 (Table 3.). The spring abundance is our best estimate of what remained post-harvest. For this report hunting mortality is included under trapping because it is difficult to distinguish between the two. Only a few wolves are taken under the hunting regulation. We believe 10-20 wolves spend at least some of their time in the areas open to wolf control.

Dates and method of most recent fall abundance assessment for wolves:

Fall abundance was estimated by adding the estimated number of wolves removed prior to the date the wolf survey was flown to the number of wolves counted during the survey.

Other research or evidence of trend or abundance status in wolves: Interviews with trappers and information from trapper survey reflect concurrence with estimated abundance.

Table 3. Population estimates and human caused mortalities for wolves in GMU 15A. Removal objective is 100% of pre-control fall abundance from control area (49 mi²) in year 1 of wolf predation control program, so estimated or confirmed number remaining by spring (30 April) each RY in all of GMU 15A (1,314 mi²) must be at least 15.

by spring (so repri) cach Kr in an or Sivie Tori (1,514 in) must be at least 10.								
Period	RY	Fall	Harvest		Dept.	Public	Total removal	Spring
		abundance	removal		control	control	from area N	abundance
			from GMU		removal	removal		
			15A		from	from		
			Trap	Hunt	-15A	15A		
	2010	53-63	15				15 ^a	38-48
	2011	60-62	10				10	50-52
Year	2012	60-65 ^a	20				20 ^a	45-50
Year 1	2013	45-60 ^b	2			3	5	45-50
Year 2	2014	45-60 ^d	8			8		45-50
Year 3	2015							

^a Given the date of the survey we estimated that 5 wolves were harvested by trappers/hunters from the date of the survey until the close of the season.

^b This is estimate based on a partial survey of GMU 15A and other reported sightings.

^c Harvest data was obtained from the State Winfonet database.

^d No surveys were flown, however there is no indication abundance changed from the previous year.

Black Bear

The latest estimate for black bear abundance occurred in the mid-1980's. Extrapolating data from that time period resulted in a current estimate of 700-900 black bears in GMU 15A. It is not known if these data accurately portray current black bear numbers in this area, but black bears do occur throughout the unit.

Brown Bear

The Kenai National Wildlife Refuge completed a study in 2013 estimating the brown bear population on the Kenai Peninsula. Using their density calculation $(42/1000 \text{ km}^2)$ there were approximately 142 brown bears in GMU 15A during June 2010 (the time they completed gathering field data). There have been no brown bear censuses since that time.

The most significant action effecting brown bear mortality in GMU 15A is the recent liberalization of hunting seasons and bag limits. Prior to Fall 2012 the hunting season was managed through a limited drawing permit season with a 1 bear/4 years bag limit. In January 2012 the BOG liberalized hunting opportunity for Kenai brown bears by adding a fall registration hunt with unlimited number of permits and season dates of October 1 – November 30. The BOG furthered liberalized brown bear hunting opportunity in March 2013 including expanded season dates of September 1 – May 31, a bag limit of 1 bear/regulatory year, and maintained the unlimited number of registration permits. The BOG also set a cap (to begin in calendar year 2014) on human caused brown bear mortalities of 70 human caused mortalities annually based on at least a 3 year running average. During calendar year 2013 the recorded human caused brown bear mortalities reached 71. For calendar year 2014, total human caused brown bear mortalities in GMU 15A was 13 and 4 in 2015 as of August 28, 2015.

There are no identified Intensive Management control efforts for black or brown bears.

4) Habitat data and nutritional condition of prey species:

Where active habitat enhancement is occurring or was recommended in the Operational Plan, describe progress toward objectives

Objective(s): Increase available moose browse through mechanical treatment and work with other agencies to develop a long term habitat management strategy. The department received funding to expand this effort in the future.

Area treated and method: Timber (mainly spruce, aspen, and some birch) was harvested on about 85 acres during 2013 in GMU 15A. Portions of that area were scarified and approximately 1,000 birch seedlings were planted.

Observation on treatment: Initial visits to the site indicate good survival of the planted seedlings and regeneration of early successional species has started. Since it is early in the project, future updates will provide better insights regarding the success of this project. While the 85 acres treated is not expected to impact GMU 15A on a population scale, this project laid the foundation for future habitat enhancement efforts.

Evidence of progress toward objectives: Department staff will continue to work with other government and private companies or organizations to develop a long term habitat management strategy. The department did receive a significant allocation (\$1,000,000) from the legislature for habitat enhancement projects and there are plans to use the majority of these funds in GMU 15A. Staff have been working with various agencies regarding to determine the best use of these funds, but efforts are still in the planning stage.

Nutritional indicators for moose in assessment area (L) of the GMU 15A Intensive management area: Current research efforts addressing moose productivity and body condition are in the early stages and data are not summarized at this time. Preliminary data indicate that adult cow moose are in relatively poor condition in GMU 15A compared to adult cows in GMU 15C, based on body condition indices.

5) Costs specific to implementing Intensive Management

Table 4. Cost (\$1000 = 1.0) of agency salary based on estimate of proportional time of field level staff and cost of operations for intensive management activities (e.g., predator control or habitat enhancement beyond normal Survey and Inventory work) performed by personnel in the Department or work by other state agencies (e.g., Division of Forestry) or contractors in GMU 15A Intensive management area. Fiscal year (FY) is also 1 July to 30 June but the year is one greater than the comparable RY (e.g, FY 2010 is 1 July 2009 to 30 June 2010).

		Predation control ^a		Other IM	activities	Total IM	Research
Period	FY	Time ^b	Cost ^c	Time	Cost ^c	Cost ^c	Cost ^d
	2013			12	35.5	35.5	150.0
	2014			13	136.3	136.3	250.0
Year 1	2015	5.0	34	11	30.9	64.9	150.0
Year 2	2016	1.0	6.8	1	6.8		150.0
Year 3	2017						
Year 4	2018						
Year 5	2019						

^aState or private funds only.

^bPerson-months (22 days per month)

^cSalary plus operations

^dSeparate from implementing IM program but beneficial for understanding of ecological or human response to management treatment (scientific approach that is not unique to IM).

6) Department recommendations for interim

Has progress toward defined criteria been achieved? Yes, we completed initial habitat improvements (85 acres) and are continuing discussions with private organizations and government agency to develop a long term habitat improvement plan. We also issued permits for the wolf control portion of the program, and the public took 3 wolves by aerial shooting in December 2013 and March 2014. A private contractor was hired to attempt ground based trapping efforts within the control area. Lack of adequate snow cover hampered initial efforts for ground based methods. During RY 2014 no wolves were taken under predator control efforts due to lack of adequate snow cover.

Has achievement of success criteria occurred? No surveys have been completed. There is no indication this moose population has grown.

Recommendation for IM practice: continue habitat improvement efforts. We completed two years of active wolf control in this area and have taken 3 wolves total. Given the small areas where control efforts can be conducted and the limited take, these efforts have not had any impact on the moose population in GMU 15A. However, we recommend continuing predator control efforts at least through spring 2016.

We recommend continuation of habitat enhancement efforts. As stated in the Operational Plan, we consider lack of habitat as the primary factor for low moose numbers (and consequently moose harvest) in GMU 15A. The lack of adequate snow cover did impact the ability to effectively pursue wolves for control efforts, but should have favored moose. While body

condition indices did improve in GMU 15A, they were still below the condition indices measured for GMU 15C moose. This provides additional evidence that habitat conditions are the primary factor limiting moose population growth in this area.