

**SALMON SPAWNING GROUND SURVEYS  
IN THE BRISTOL BAY AREA, ALASKA, 2000**



By

**Tim Sands, James Browning, Keith A. Weiland,  
and Slim Morstad.**

**Regional Information Report<sup>1</sup> No. 2A01-08**

**Alaska Department of Fish and Game  
Division of Commercial Fisheries  
333 Raspberry Road  
Anchorage, Alaska 99518-1599**

**May 2001**

---

<sup>1</sup> The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished divisional reports. These reports frequently serve diverse ad hoc informational purposes or archive basic un-interpreted data. To accommodate timely reporting of recently collected information, reports in this series undergo only limited internal review and may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Division of Commercial Fisheries.

## **AUTHORS**

Tim Sands is the Togiak Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, P.O. Box 230, Dillingham, AK 99576-0230.

Keith A. Weiland is the Egegik and Ugashik Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, 333 Raspberry Road, Anchorage, AK 99518.

James Browning is the Nushagak Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, P.O. Box 230, Dillingham, AK 99576-0230.

Slim Morstad is the Naknek/Kvichak Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, P.O. Box 37, King Salmon, AK 99613.

## **ACKNOWLEDGMENTS**

We would like to thank the U.S. Fish and Wildlife Service and the University of Washington, Fisheries Research Institute for equipment, personnel and funding they provided to help gather escapement data in 2000.

"The Alaska Department of Fish and Game administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972. If you believe you have been discriminated against in any program, activity, or facility, or if you desire further information please write to ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; U.S. Fish and Wildlife Service, 4040 N. Fairfield Drive, Suite 300, Arlington, VA 22203; or O.E.O., U.S. Department of the Interior, Washington DC 20240. For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-4120, (TDD) 907-465-3646, or (FAX) 907-465-2440."

## TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES .....	iv
LIST OF FIGURES .....	v
LIST OF APPENDICES .....	vi
INTRODUCTION .....	1
Naknek/Kvichak District .....	1
Egegik District .....	1
Ugashik District .....	2
Nushagak District .....	2
Togiak District .....	4
METHODS .....	4
Naknek/Kvichak District .....	5
Egegik District .....	5
Ugashik District .....	5
Nushagak District .....	5
Togiak District .....	6
RESULTS AND DISCUSSION .....	6
Naknek-Kvichak District .....	6
Egegik District .....	7
Ugashik District .....	8
Nushagak District .....	9
Togiak District .....	9
LITERATURE CITED .....	11
TABLES .....	13
FIGURES .....	29
APPENDICES .....	39

**LIST OF TABLES**

<u>Table</u>	<u>Page</u>
1. Aerial survey counts of sockeye salmon, Alagnak River system, 2000 .....	13
2. Aerial survey counts of chinook, chum, pink, and coho salmon, Naknek-Kvichak District, 2000.....	14
3. Aerial survey peak counts of chinook salmon escapement, Egegik District, 2000 .....	15
4. Aerial survey peak counts of chum salmon escapement, Egegik District, 2000 .....	16
5. Aerial survey peak counts of coho salmon escapement, Egegik District, 2000 .....	17
6. Aerial survey peak counts of sockeye salmon escapement, King Salmon and Dog Salmon Rivers, Ugashik District, 2000 .....	18
7. Peak survey counts of chinook salmon escapement, Ugashik District, 2000 .....	19
8. Peak survey counts of chum salmon escapement, Ugashik District, 2000 .....	20
9. Aerial survey peak counts of coho salmon escapement, Ugashik District, 2000 .....	21
10. Peak aerial counts of live sockeye salmon and total escapement estimates, Wood River system, 2000 .....	22
11. Peak aerial counts of live sockeye salmon and total escapement estimates, Togiak District, 2000 .....	25
12. Peak aerial counts of live chinook salmon and total escapement estimates, Togiak District, 2000 .....	26
13. Peak aerial counts of live chum salmon and total escapement estimates, Togiak District, 2000 .....	27
14. Peak aerial counts of live coho salmon and total escapement estimates, Togiak District, 2000 .....	28

**LIST OF FIGURES**

<u>Figure</u>		<u>Page</u>
1.	Bristol Bay management area, Alaska .....	29
2.	Alagnak River drainage, Bristol Bay, Alaska .....	30
3.	Egegik River drainage, Bristol Bay, Alaska .....	31
4.	Ugashik River drainage, Bristol Bay, Alaska .....	32
5.	Wood River Lakes system, Bristol Bay, Alaska .....	33
6.	Lake Nunavaugaluk system, Bristol Bay, Alaska .....	34
7.	Nushagak-Mulchatna River system, Bristol Bay, Alaska .....	35
8.	Tikchik Lakes system, Bristol Bay, Alaska .....	36
9.	Togiak River system, Bristol Bay, Alaska .....	37
10.	Kulukak River system, Bristol Bay, Alaska .....	38

## LIST OF APPENDICES

<u>Table</u>	<u>Page</u>
1. Sockeye salmon total escapement estimates, Naknek-Kvichak District, 1955-2000 .....	39
2. Aerial survey counts of chinook salmon escapements, Naknek River Drainage, 1971-2000 .....	41
3. Chinook salmon escapement survey history, mainstem Naknek River, 1929-2000 .....	42
4. Chinook salmon escapement survey history, Big Creek, Naknek River Drainage, 1963-2000 .....	45
5. Chinook salmon escapement survey history, King Salmon Creek, Naknek River drainage, 1964-2000 .....	48
6. Chinook salmon escapement survey history, Paul's Creek, Naknek River Drainage, 1971-2000 .....	50
7. Chinook salmon escapement survey history, Alagnak River, 1963-2000 .....	52
8. Chinook salmon escapement survey history, Kvichak River, 1932-2000 .....	54
9. Chinook salmon escapement data, Naknek-Kvichak District, 1970-2000 .....	55
10. Chum salmon escapement survey history, Alagnak River, 1961-2000 .....	56
11. Pink salmon escapement survey history, Alagnak River, 1968-2000 .....	58
12. Pink salmon escapement survey history, Kvichak River, 1966-2000 .....	59
13. Pink salmon escapement survey history, Naknek River, 1974-2000 .....	60
14. Aerial survey counts of chinook salmon escapement, Egegik District, 1981-2000 .....	61
15. Aerial survey counts of chum salmon escapement, Egegik District, 1982-2000 .....	62

**LIST OF APPENDICES (Continued)**

<u>Table</u>	<u>Page</u>
16. Aerial survey counts of pink salmon escapement, Egegik District, 1974-2000 .....	63
17. Aerial survey counts of coho salmon escapement, Egegik District, 1981-2000 .....	64
18. Aerial survey counts of chinook salmon escapement, Ugashik District, 1980-2000 .....	65
19. Aerial survey counts of chum salmon escapement, Ugashik District, 1980-2000 .....	66
20. Aerial survey counts of pink salmon escapement, Ugashik District, 1980-2000 .....	67
21. Aerial survey counts of coho salmon escapement, Ugashik District, 1981-2000.....	68
22. Spawner distribution and total escapement estimates of sockeye salmon, Wood River system, 1959-2000 .....	69
23. Total escapement estimates of pink salmon, Nushagak and Togiak Districts, 1962-2000 .....	70
24. Aerial estimates of sockeye salmon escapements, Togiak District 1980-2000 .....	71
25. Peak aerial counts of live sockeye salmon, Togiak River drainage, 1980-2000 .....	72
26. Peak aerial counts of live sockeye salmon, Togiak District, 1980-2000 .....	73
27. Peak aerial counts of live chinook salmon, Togiak River drainage, 1980-2000 .....	74
28. Peak aerial counts of live chinook salmon, Togiak District, 1980-2000 .....	75
29. Peak aerial counts of live chum salmon, Togiak River drainage, 1980-2000 .....	76
30. Peak aerial counts of live chum salmon, Togiak District, 1980-2000 .....	77
31. Peak aerial counts of live coho salmon, Togiak River drainage, 1980-2000 .....	78
32. Peak aerial counts of live coho salmon, Togiak District, 1980-2000 .....	79

## INTRODUCTION

Aerial surveys of salmon spawning streams have been conducted in the Bristol Bay area of Alaska (Figure 1) for many years. Surveys provide biologists with information regarding the abundance and distribution of sockeye salmon *Oncorhynchus nerka*, chinook salmon *O. tshawytscha*, chum salmon *O. keta*, pink salmon *O. gorbuscha*, and coho salmon *O. kisutch* escapements. This information is important to fishery managers for several reasons. It supplements data gathered at counting towers on the mainstem rivers, provides data from rivers where counting towers are not utilized, and provides data for time periods and species not covered by counting tower operations. Collected information is used to: (1) evaluate escapement goals and escapement/return relationships, (2) forecast future returns, (3) identify possible management problems relating to escapements, and (4) contribute to strategies designed to alleviate escapement problems. This report summarizes the 2000 salmon spawning ground surveys conducted in the Bristol Bay area.

### *Naknek/Kvichak District*

The Naknek-Kvichak District is comprised of three major rivers: (1) the Kvichak River, issuing from Iliamna Lake and its tributaries, (2) the Alagnak or Branch River flowing from Kukaklek and Nonvianuk Lakes, and (3) the Naknek River emanating from Naknek Lake and its tributaries (Figure 2). All of these systems flow into Kvichak Bay.

Since 1955, Kvichak River sockeye salmon escapements have been estimated using counting towers located on the Kvichak's mainstem, approximately one-quarter mile downstream of Lake Iliamna's outlet. From 1957 to 1976, Alagnak River sockeye escapements were estimated using a counting tower located near the upper extent of tidal influence. Since 1977, all Alagnak sockeye escapements have been estimated using aerial surveys. From 1950 to 1957, sockeye escapements to the Naknek River system were counted using a weir on the mainstem of the river just upstream of the tidal influence. From 1958 to the present, escapements have been estimated using counting towers near the Naknek River 'Rapids' downstream of the outlet of Naknek Lake. Escapements of other salmon species into Naknek-Kvichak District drainages have been estimated using aerial surveys.

### *Egegik District*

The Egegik River system contains two major watersheds: (1) the Egegik River, emanating from Becharof Lake and nearby coastal lowlands, and (2) the King Salmon River, issuing from runoff from the Kejulik Mountains and southern portions of Katmai National Park (Figure 3). Both rivers flow into Egegik Bay near Egegik village.

From 1952 through 1956, a weir was used in the Egegik River to count sockeye salmon escapements. The weir was located near the bottom of the Egegik River 'rapids'. From

1957 to the present, counting towers situated between the outlet of Becharof Lake and Egegik Lagoon have been used to estimate sockeye escapements. Escapements for other salmon species have been estimated using aerial surveys.

### *Ugashik District*

The Ugashik River system is comprised of four major watersheds: (1) the Ugashik River, flowing from Lower Ugashik Lake and nearby coastal lowlands, (2) the Dog Salmon River, emanating from glacial melt and runoff from peaks in the Aleutian Range, (3) the King Salmon River, issuing from Mother Goose Lake and three major runoff tributaries, and (4) Dago Creek, emitting from a large lowland coastal area (Figure 4). All of these systems flow into the intertidal reaches of Ugashik River and Ugashik Bay.

From 1949 to 1956, a weir located downstream from the outlet of Lower Ugashik Lake was used to count sockeye salmon escapements. From 1957 to the present, sockeye escapements have been estimated using counting towers located between the outlet of Lower Ugashik Lake and Ugashik Lagoon. Escapements for other salmon species have been estimated using aerial surveys.

### *Nushagak District*

The Nushagak watershed is comprised of four major rivers: (1) the Wood River, draining Grant, Kulik, Beverley, Nerka, and Aleknagik Lakes, (2) the Nushagak River, draining Tikchik Lakes and the Nuyakuk, upper Nushagak, and Mulchatna Rivers, (3) the Igushik River, draining Ualik and Amanka Lakes, and (4) the Snake River, draining Lake Nunavaugaluk (Figures 5 through 8). All of these systems empty into Nushagak Bay.

Abundance and age composition of sockeye salmon escapements in the Wood River Lake system has been estimated annually from counting towers at the outlet of Lake Aleknagik since 1953.

Sockeye salmon distribution in the Wood River Lake system is an important element in establishing escapement goals and measuring success in achieving escapement goals for this system. Interconnecting rivers between the large lakes in the system are primarily used by three-ocean sockeye for spawning, while the lake beaches and tributary streams are used more by two-ocean sockeye. Knowledge of the age composition of returning sockeye gives managers the ability to use a variable escapement goal policy to minimize overcrowding of spawners in the interconnecting rivers while taking advantage of the extensive beach spawning areas and numerous tributary streams.

Periodically, ADF&G personnel conduct aerial surveys to assess sockeye spawner distribution within the Wood River Lake system. Personnel from the University of Washington, Fisheries Research Institute also conducted ground surveys on major creeks and some rivers of the system. Surveys of the actual spawning distribution within the

creeks, rivers, and beaches of the system provide a measure of management success in obtaining the desired spawning distribution.

Salmon escapement in the Nushagak River is estimated by a sonar project, located on the Nushagak River below Portage Creek, approximately 32 km (20 miles) upstream from the river mouth. The Nushagak River sonar project has been used since 1980 to estimate annual escapements for all salmon species in the entire Nushagak drainage (Miller 1997). Prior to the advent of the sonar project, sockeye escapement was estimated by a counting tower project on the Nuyakuk River (1959-1988). Aerial surveys of the Nushagak-Mulchatna system were conducted annually beginning in 1966. Initial surveys provided escapement estimates for chinook and chum salmon, and surveys in the Nushagak and Mulchatna systems since 1977 were used to estimate sockeye abundance in that system. Together, the combined estimates from counting towers and aerial surveys were used by fishery managers as estimates of the Nushagak River drainage sockeye escapement.

ADF&G staff continued to survey the upper Nushagak and Mulchatna areas after the development of the sonar project to provide a comparison with sonar estimates and document spawner distribution for all species except coho salmon. Chum salmon surveys were discontinued in the Nushagak District in 1980, and surveys of the Nushagak-Mulchatna Rivers for all other species were discontinued in 1991 due to the success of the sonar project and limited funding. The Nuyakuk tower project was halted after the 1988 season, but was reinitiated for the 1995 season and has been operated since that time. Aerial surveys of the Nushagak and Mulchatna systems have been conducted sporadically since 1991 providing infrequent information on spawning sockeye distribution in the Nushagak River.

Aerial surveys were conducted sporadically in the Tikchik Lakes system from 1954 to 1987 to assess spawner distribution of sockeye salmon. Surveys of the Tikchik Lakes were conducted infrequently since 1990 to document an apparent change in spawner distribution, evidenced by changes observed in the age composition of Nushagak River sockeye escapement, and supported by reports of low numbers of spawners in the Tikchik Lake system. These surveys have documented lower than expected numbers of spawners in the Tikchik Lakes system, based on sonar estimates in the lower Nushagak River and historical distribution patterns (Browning et. al. 1998). However, few corresponding surveys were conducted in the Nushagak and Mulchatna drainages to completely assess distribution.

Sockeye escapement is measured in the Igushik Lakes system at a counting tower located at the outlet of Amanka Lake. Spawner distribution has not been documented annually, and surveys have not been conducted on the Igushik system for sockeye salmon and other species since 1991 (Russell, et. al. 1992). Spawning escapement and distribution of sockeye salmon in the Snake Lake system was estimated annually prior to 1998 by aerial surveys, but with the closure of the Snake River section and funding shortages in recent years, these surveys have not been continued.

## *Togiak District*

Two major river drainages flow into the Togiak District: (1) the Togiak River, draining Togiak, Gechiak, Pungokebuk, and Ongivinuk Lakes and Nayorurun and Kemuk Rivers (Figure 9), and (2) the Kulukak River, draining Kulukak Lake (Figure 10). Various smaller systems within the district include the Kanik River draining Tithe Creek Ponds and the Quigmy, Matogak, Osviak, Slug, Negukthlik, and Ungalikthluk Rivers. Kulukak River and the Kanik River flow into Kulukak Bay, located in the eastern portion of the district. The Togiak and Quigmy Rivers flow into Togiak Bay, located in the middle of the district, and the Matogak, Osviak, and Slug Rivers flow into Hagemeister Straits and coastal waters in the western portion of the district (Figure 1).

Sockeye salmon escapement is estimated for the Togiak Lake system from counting towers operated at the outlet of Togiak Lake. Abundance and distribution of spawning populations of sockeye salmon in the Togiak River and tributaries below the counting towers, as well as other systems within the Togiak District, are estimated by aerial surveys. Abundance and distribution of chinook, chum, pink, and coho salmon spawning in Togiak District watersheds are also estimated entirely from aerial surveys.

Since 1991, the operational budget has not had sufficient funds to conduct spawning ground aerial surveys in the Togiak District. The U.S. Fish and Wildlife Service Togiak National Wildlife Refuge (USFWS/TNWR) has provided funding for aircraft charters for aerial surveys, and has assisted with aerial surveys in the Togiak District to monitor salmon populations within drainages on the refuge.

## **METHODS**

All survey flights were conducted from small fixed-wing, high-wing, wheeled aircraft (Super Cub, Cessna 180, Cessna 185, or Cessna 206) or helicopter (Robinson R-22) chartered from local air charter companies and flown by experienced survey pilots. Alaska Department of Fish and Game (ADF&G) or USFWS biologists familiar with the streams and target species counted salmon. USFWS pilots and aircraft flew several of the surveys in the Togiak National Wildlife Refuge. Counts were made from low altitudes (200 to 400 feet) at air speeds of 50 to 90 mph. Polarized sunglasses and aircraft positioning were used to minimize effects of glare off the water. Surveys were scheduled to coincide as closely as possible to the historic peak of spawning for the target species, taking into account weather, water conditions, and aircraft availability. Peak of spawning was defined as that point when the greatest number of spawning salmon are occupying redds. Counts were registered on a hand tally counter or on a tape player. This information was transferred to survey data forms either sometime during the survey or upon returning to the office.

Aerial surveys account for only a portion of the known spawning populations (Evzerof, 1975; Nielson and Green, 1981; Rogers, 1984). At the time of each survey, some of the salmon have yet to reach the spawning grounds, some have already spawned and died,

some are still schooled, and some are either misidentified or not seen. Methods used to interpret aerial survey counts are described below for each commercial fishing district.

### ***Naknek/Kvichak District***

Aerial surveys were flown during late summer and fall to assess escapements of sockeye, chinook and chum salmon in portions of the Naknek/Kvichak District. Salmon counts for these drainages are indices of the total number of each species present in the spawning area at the time of the survey. Three surveys were flown, August 7, 15 and September 11, to provide estimates of the Alagnak River Drainage sockeye, chinook and chum escapements. Additionally, all major chinook spawning areas in the Naknek River Drainage were surveyed on August 1, and August 7; the Kvichak River survey was flown on August 7. These survey counts were not expanded to provide instantaneous population estimates, although expansions have been made in some earlier years based on subjective criteria. Counting towers were used to estimate total sockeye escapement to the Kvichak and Naknek Rivers. ADF&G, Commercial Fisheries Division staff made all aerial survey counts in the district.

### ***Egegik District***

No system-wide aerial surveys were flown for sockeye salmon in 2000. An aerial survey of known chinook and chum salmon spawning areas in both the Egegik and King Salmon Rivers was flown on July 31. An aerial survey was flown on September 24, with funding provided by the USFWS, to estimate coho salmon escapements. All aerial survey counts in the Egegik drainage reflect only the actual numbers of salmon sighted and should be considered a minimum indication of abundance.

### ***Ugashik District***

Salmon counts in the Ugashik District reflect the actual numbers of salmon sighted on the spawning grounds for 2000. Aerial surveys of known chinook and chum salmon spawning areas in the Ugashik drainage were flown on August 8. An aerial survey, funded by the ADF&G, was flown on October 12 to estimate coho salmon escapement. The survey was flown at the preferred time. Aerial survey counts should be considered a minimum indication of total abundance.

### ***Nushagak District***

Due to poor weather, lack of suitable aircraft with qualified pilots, no aerial spawning ground surveys were flown in the Nushagak District during the 2000 season.

### *Togiak District*

Survey and data analysis methods used in the Togiak District were similar to those described by Nelson (1979), Bucher (1981), and Russell, et. al. (1990). This year surveys were flown on August 14, 16, 22, 24, 26 and 27. ADF&G staff surveyed the Togiak and Kulukak River drainages for sockeye and chinook salmon and the Quigmy, Negukthlik and Ungalikthluk Rivers for sockeye salmon. USFWS/TNWR staff conducted surveys of the Quigmy, Negukthlik and Ungalikthluk Rivers for chinook salmon and the Slug, Osviak, and Matogak Rivers for chum and sockeye salmon.

Total escapement was estimated for sockeye salmon in systems without counting towers (i.e. Kulukak River, mainstem and tributaries of the Togiak River below the towers) by multiplying peak aerial counts by an expansion factor between 1.5 and 3.0 depending on survey and water conditions (Table 11 lists expansion factors by stream). Since 1980, total escapement, for chinook salmon, in the Togiak District has been calculated by aerial counts using a multiplier of 2.5 if the survey was timed properly relative to the spawning peak and visibility conditions were average. Since 1968, escapement for chum salmon has been calculated by aerial counts using a multiplier of 2.0 (Nelson 1968). Since 1978, total pink salmon escapements have also been estimated by multiplying aerial counts by 2.0. An expansion factor of 3.0 has been used for coho salmon in all areas of the Togiak District since the initiation of coho surveys in 1980. Expansion factors have been subjectively adjusted based on weather conditions, visibility, and survey timing with respect to the peak spawning activity.

## **RESULTS AND DISCUSSION**

### *Naknek/Kvichak District*

Aerial surveys of sockeye salmon escapement into the Alagnak River and its tributaries were conducted on August 15 and September 11. The sockeye salmon escapement index count totaled 451,300 for the system (Table 1). This is above the 1955-1998 average aerial count of 234,150 (Appendix Table 1), and was more than twice the escapement point goal of 185,000.

Aerial surveys of chinook salmon escapements into the Naknek River drainage were flown on August 1 and August 7. Chinook salmon escapement counts were made in each of the four main spawning areas: mainstem Naknek River, Big Creek, King Salmon Creek and Paul's Creek. In 2000, the mainstem Naknek River was flown earlier than normal, due to the poor weather and lateness of the 1999 survey. A total of 3,233 chinook were observed in the Naknek drainage. Over the period from 1971-2000 there have been 21 years in which chinook salmon escapement indices have been obtained from all four main spawning areas (Appendix Tables 2-6). The chinook escapement index for these 21 years has ranged from a low of 2,691 in 1992 to a high of 11,730 in 1988.

Alagnak River drainage chinook salmon escapement was surveyed on August 7, estimating a total of 2,220 (Table 2). From 1970-2000, Alagnak chinook salmon counts have ranged from a low of 824 in 1973 to a high of 15,210 in 1997 (Appendix Table 7). An aerial survey of chinook salmon escapement into the Kvichak River was conducted on August 7 estimating only 6 chinook in the Kaskanak Flats area (Appendix Tables 8 and 9).

Chum salmon were counted only during the August 7 Alagnak and Kvichak River survey. Alagnak River has been the principle chum salmon producing system in the Naknek/Kvichak District (Appendix Table 10). A total of 12,600 chum salmon were observed during the survey.

### *Egegik District*

The 2000 Egegik River sockeye escapement past the counting towers totaled 1,032,138 fish, or 6% below the mid-range objective of 1.1 million. The BEG range for Becharof Lake is 800 thousand to 1.4 million. Aerial survey counts of known chinook salmon spawning areas in the Egegik drainage yielded a total count of 560 chinook salmon. No additional chinook salmon were counted at the Egegik River counting towers (Table 3). This total was 51% below the average count of 1,146 (Appendix Table 14). The commercial chinook harvest in the Egegik District totaled approximately 1,061 fish, or 57% below the 1980 to 1999 average harvest of 2,483. Fishing time was reduced to three days per week between June 1 and June 16, compared to a four day per week schedule in past seasons. Using gillnets with larger than 5.5-inch mesh in the commercial fishery from June 1 to July 1 was prohibited by regulation. All of these factors probably contributed to the reduced commercial harvest of chinook salmon, but in general chinook salmon runs to Egegik have been low in recent years. Given the catch and escapement figures above, the Egegik chinook salmon removal rate for 2000 was probably less than 65%.

The chum salmon escapement index was 1,189 fish (Table 4). The 2000 index was well below the 1982-1999 average of 8,214 fish (Appendix Table 15). The 2000 commercial chum harvest from the Egegik District totaled approximately 38,000 fish, or 63% below the 1980 to 1999 average catch of 103,000. Escapement indices of less than 10,000 chum salmon have been recorded since 1989, but aerial surveys for chum salmon are not reliable indicators and it is believed that chum escapement indices documented over the last several years have probably greatly underestimated chum salmon escapements (Browning et.al. 1998). In 1999, a comparison of the Gertrude Creek Weir count of 16,000 and an aerial survey count on August 6 shows that the aerial count revealed only about 2% of the weir count.

No pink salmon were noted during the July 31 aerial survey. A total of 32 pink salmon were reported from the commercial catch. The 1974 to 2000 pink salmon escapement indices are listed in Appendix Table 16.

The coho salmon escapement was documented with an aerial survey conducted on September 24 (Table 5). The USFWS in King Salmon provided funding for this survey. A total of 4,870 coho salmon were counted in the Egegik River and in numerous tributaries of Becharof Lake. A little over half of this total was counted in tributaries of Becharof Lake. The aerial counts were focused on main coho salmon producing areas, which are listed in Table 5. Compared to the last five years, the 2000 index count of 4,870 was about average. The commercial harvest totaled approximately 13,200 fish, which was 66% below the 20-year (1980-1999) average of 39,200. Deliveries occurred through August 28, though the fishery remained open until September 30. Historical survey counts are listed in Appendix Table 17.

### *Ugashik District*

The 2000 Ugashik sockeye salmon escapement tower count was approximately 620,000 fish, or 27% under the mid-range objective of 850,000. However, tower counts were 59,000 and 68,000 sockeye salmon on the first two days of counting, July 3 and 4, and these were the highest counts ever recorded for these dates. It is likely that the total escapement was substantially greater than the final tower count. No system-wide aerial surveys were conducted due to a lack of funding, however; an additional 5,480 and 12,900 sockeye salmon were counted in the Dog Salmon and King Salmon Rivers, respectively, during a chinook and chum salmon survey (Table 6).

Chinook salmon escapement surveys of Dog Salmon, King Salmon, and Ugashik Rivers were flown on August 8 and yielded a count of 1,324 fish. Additionally, 26 chinook salmon were counted past the counting towers bringing the cumulative escapement count to about 1,350 (Table 7). The Figure Eight Creek count of 378 was the largest escapement component for the system, but no count was made for Ugashik River and Grassy Creek. The 2000 escapement count was 70% below the 1980 to 1999 average count of 4,513 chinook salmon (Appendix Table 18), and like Egegik, chinook salmon runs have been low in recent years. The Ugashik District's commercial catch of approximately 893 chinook was 72% below the 20-year average harvest of 3,226, and about 46% below the recent 10-year average of 1,655.

Aerial surveys of Dog Salmon, King Salmon, and Ugashik Rivers on August 8, yielded a count of 15,810 chum salmon (Table 8). The survey was considered to be near the peak of spawning though a fair number of chum carcasses were observed. The 2000 aerial count was well below the 1980 to 1999 average of 30,568 (Appendix Table 19). The District's commercial chum salmon harvest of approximately 38,000 fish, was also well below average.

The Ugashik pink salmon returns have historically been very small. This year a total of 4 pink salmon were reported in the commercial catch and 46 pink salmon were counted past the Ugashik counting towers before that project ceased operation on July 28 (Appendix Table 20).

An aerial survey for coho salmon was again made this year in the Ugashik drainage (Table 9). A total of 12,070 coho salmon were observed the October 12 flight most of which were seen in Painter Creek a tributary of the King Salmon River. For the Ugashik lakes a count of 4,400 was 83% above the 1996, 1997 and 1999 average count of 2,400. The coho harvest of approximately 1,270 fish was about one twentieth of the recent 10-year average. Historical coho salmon escapement data is recorded in Appendix Table 21.

### *Nushagak District*

There were no spawning ground surveys flown this year, for any species. The sonar project at Portage Creek produced apportioned estimates of 56,374 chinook salmon, 403,500 sockeye salmon, 172,846 coho salmon, 135,285 pink salmon and 141,323 chum salmon in the Nushagak River for 2000. A counting tower was operated on the Nuyakuk River again this year; 129,468 sockeye salmon were counted past this site in 2000.

Spawning escapement of sockeye salmon in the Wood River system was estimated to be 1.3 million fish, 8% over the upper end of the biological escapement goal range (700,000 – 1.2 million). Two-ocean sockeye comprised approximately 61% of the Wood River escapement while three-ocean sockeye contributed the other 39% of the escapement. Once again the Wood to Nushagak sockeye ratio was projected to exceed 3:1, triggering the OEG minimum escapement goal of 235,000 sockeye salmon for the Nushagak River. The management staff had to balance a strong sockeye return to the Wood River with an average return to the Nushagak.

### *Togiak District*

Peak aerial counts and total population estimates were derived from aerial surveys for sockeye salmon in major river systems of the Togiak District in 2000 (Table 11). The expanded aerial survey estimate of 40,275 sockeye salmon for the Togiak River and its tributaries below the counting tower was 47% above the 1980-1999 average of 27,270 fish (Appendix Table 24). Escapement past the counting tower was 311,970 sockeye, exceeding the upper end of the BEG range (100,000 – 200,000) by 56%. The spawning escapement of sockeye salmon in Kulukak Section, including Kulukak River, Kulukak Lake, and Tithe Creek Ponds, was estimated at 22,350 fish, 72% of the 1980-1999 average of 31,105. Peak aerial sockeye salmon live counts in the Kemuk, Pungokebuk and Ongivunuck Rivers were below the 1980-1999 average (Appendix Table 25); while counts for the Gechiak and Nayorurun tributaries to the Togiak River were above average. Total sockeye salmon escapement for Togiak District was 390,080. In the tributaries to the Togiak River and the smaller systems in the western portion of the district where conditions were average, the standard factor of 2.5 was used. The expanded escapement estimate for chinook salmon in the Togiak District was 16,896 fish. The 2000 district-wide chinook escapement was 33% above the 5-year average of 12,668 fish. The aerial peak live count for the Togiak River and tributaries was 5% above the 1980-1999 average, and aerial counts for all chinook systems in the Togiak District combined were 5% above the 20-year average (Appendix Tables 27 and 28). The

escapement goal of 10,000 chinook salmon into the Togiak River was slightly exceeded with an expanded spawning estimate of 11,813 fish. Chinook peak aerial counts for Kulukak, Matogak, Osviak and Slug Rivers were all below average, while Quigmy, Negukthlik and Ungalikthluk Rivers were at or above average. The Kulukak River escapement estimate (340 chinook) was poor and comprised only 44% of the 1980-1999 Kulukak River average.

Chum salmon counts were conducted coincidentally with the chinook salmon surveys. Chum salmon escapements were mixed throughout the Togiak District; the Slug, Osviak and Matogak River escapements exceeded the 20-year averages. The Quigmy, Kulukak, Negukthlik and Ungalikthluk River escapements were below the 20-year averages. There was no estimate of chum salmon escapement for the Togiak River drainage due to poor survey conditions and therefore no estimate of the total chum salmon escapement for the district is available (Table 13, Appendix Tables 29 and 30).

Pink salmon aerial escapement surveys are flown in even years. In 2000 no surveys were done due to poor weather and lack of a survey plane. Historical counts (for even years only) may be perused in Appendix Table 23.

Aerial surveys for coho salmon were not done in 2000; the lack of a suitable survey plane and poor weather prevented these surveys. Total coho escapement for Togiak River and tributaries was, therefore, not estimated. There was little effort to harvest coho in 2000 because no market was available after August 15. Anecdotal information from a lodge on the Togiak River did indicate a strong return of coho salmon with good success for sport fishermen.

## LITERATURE CITED

- ADF&G (Alaska Department of Fish and Game). Annual Management Report, 1999, Bristol Bay Area. Alaska Department of Fish and Game, Commercial Fisheries Division. Anchorage.
- Brookover T.E., J.B. Browning, J.R. Regnart, R.B.Russell, and K.A. Weiland. 1996. Salmon spawning ground surveys in the Bristol Bay area, 1995. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2A96-31, Anchorage.
- Browning J.B., J.R. Regnart, and K.A. Weiland. 1998. Salmon spawning ground surveys in the Bristol Bay area, 1997. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2A98-34, Anchorage.
- Bucher, W.A. 1981. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1980. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 81, Anchorage.
- Evzerov, A.V. 1975. An evaluation of the errors occurring in salmon census by aerial survey. In Canadian Translation Fisheries and Aquatic Sciences. No. 4714, 1981. [Transl. from Russian] (Otsenka pogreshnogtei aerovizual' nogo metoda ucheta Lososei), p. 82-84. From Lososevye Dal'Nego Vostoka (CVI:1975).
- Miller, J.X. 1997. Sonar enumeration of pacific salmon escapement into the Nushagak River, 1997. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Regional Information Report No. 2A96-08, Anchorage.
- Nielson, J.D. and G.H. Green, 1981. Enumeration of spawning salmon from spawner residence time and aerial counts. Transactions of the American Fisheries Society 110:554-556.
- Nelson, M.L. 1967. Red salmon spawning ground surveys in the Nushagak and Togiak Districts, Bristol Bay, 1966. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet 96.
- Nelson, M.L. 1968. Spawning ground surveys in the Nushagak and Togiak Districts, Bristol Bay, 1968. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 5.
- Nelson, M.L. 1973. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1973. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 46.
- Nelson, M.L. 1979. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1977-1979. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 73, Anchorage.

## LITERATURE CITED (Continued)

- Regnart, J.R. 1996. Kvichak River sockeye salmon spawning ground surveys, 1955-1996. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report No. 2A95-48, Anchorage.
- Rogers, D.E. 1984. Aerial survey estimates of Bristol Bay sockeye salmon escapements. Proceedings of the Workshop on Stream Indexing for Salmon Escapement Estimation. Canadian Technical Report of Fisheries and Aquatic Sciences. 1326:197-208.
- Russell, R.B., D.L. Bill, and W.A. Bucher. 1990. Salmon spawning ground surveys in the Bristol Bay area, 1989. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2K89-15, Anchorage.
- Russell, R.B., J.R. Regnart, and T.E. Brookover. 1992. Salmon spawning ground surveys in the Bristol Bay area, 1991. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2A92-01, Anchorage.

**Table 1.** Aerial survey counts of sockeye salmon, Alagnak River system, 2000<sup>a</sup>

Location	Number of Fish	Percent of Total
Nonvianuk River	0	0.0
Nonvianuk Lake	0	0.0
Kulik River	109,600	24.3
Kulik Lake	400	0.1
Alagnak River	0	0.0
Kukaklek Lake	45,000	10.0
Nanuktuk Creek	82,500	18.3
Battle River	77,500	17.2
Battle Lake	10,400	2.3
Spectacle Creek	74,300	16.5
Funnel Creek	51,600	11.4
Total	451,300	100.0

<sup>a</sup> Aerial surveys were conducted with fixed-wing aircraft.

**Table 2.** Aerial survey counts of chinook, chum, pink, and coho salmon, Naknek-Kvichak District, 2000.<sup>a</sup>

Location	Survey Date	Number of Salmon			
		Chinook	Chum	Pink	Coho
Kvichak River	7-Aug	6	2,500 <sup>b</sup>	7,000	400
Alagnak River	7-Aug	2,220	10,120 <sup>b</sup>	30,000	NC
Naknek River :					
Paul's Creek	1-Aug	43	NC	NC	-
King Salmon Creek	1-Aug	178	NC	NC	-
Big Creek	7-Aug	1,112	NC	NC	NC
Mainstem Naknek River	7-Aug	1,900	0	10,000	-
Total		5,459	12,620	47,000	400

<sup>a</sup> Aerial surveys were conducted with fixed-wing aircraft.

<sup>b</sup> Incidental observation.

**Table 3.** Aerial survey peak counts of chinook salmon escapement, Egegik District, 2000.

Location	Survey Date	Number of Chinook Salmon Counted
Egegik River	July 17 <sup>a</sup>	0
Shosky Creek	July 31	4
Whale Mountain Creek	July 31	0
Mossy Creek	July 31	16
Mink Creek	July 31	0
Gertrude Creek	July 31	85
Kaye's Creek	July 31	41
Takayoto Creek	July 31	73
Angle Creek	July 31	<sup>b</sup>
Contact Creek	July 31	341
Mainstem King Salmon River	July 31	<sup>b</sup>
Total		560

<sup>a</sup> Tower count.

<sup>b</sup> No Count.

**Table 4.** Aerial survey peak counts of chum salmon escapement, Egegik District, 2000.

Location	Survey Date	Number of Chum Salmon Counted
Egegik River	17-Jul <sup>a</sup>	0
Shosky Creek	31-Jul	<sup>b</sup>
Whale Mountain Creek	31-Jul	492
Mossy Creek	31-Jul	<sup>b</sup>
Mink Creek	31-Jul	4
Gertrude Creek	31-Jul	475
Kaye's Creek	31-Jul	32
Takayoto Creek	31-Jul	6
Angle Creek	31-Jul	<sup>b</sup>
Contact Creek	31-Jul	180
Mainstem King Salmon River	31-Jul	<sup>b</sup>
Total		1,189

<sup>a</sup> Tower count.

<sup>b</sup> No Count.

**Table 5.** Aerial survey counts of coho salmon escapement, Egegik District, 2000.

Location	Survey Date	Number of Coho Salmon Counted	Comments
<u>Egegik River Drainage<sup>1</sup></u>			
Egegik River Rapids	September 24	2,300	Most in the rapids above Myer's Creek
Stream 115.8 (Featherly Creek)	September 24	500	Most were schooled up at the mouth.
Stream 107.6 (Burl's Creek)	September 24	190	Most were schooled up at the mouth.
Stream 90.3 (Salmon Creek)	September 24	730	Most were schooled up at the mouth.
Stream 89.8	September 24	300	Most were schooled up at the mouth.
Stream 87.0 (Bear Creek)	September 24	450	Most were schooled up at the mouth.
Stream 73.5 (Becharof Creek)	September 24	180	Most were schooled up at the mouth.
Stream 48.1 (Kejulik River)	September 24	220	Includes Margaret & Albert Creeks, poor visibility.
Total		4,870	

<sup>1</sup> Streams tributary to Becharof Lake are designated by the number of miles between their mouth and the outlet of Becharof Lake (Egegik River) as one travels around the lake in a clockwise fashion from the Becharof lake outlet. This is the same system of designation used for years by previous investigators. Due to logistic problems only one survey was flow this season.

**Table 6.** Aerial survey peak counts of sockeye salmon escapement, King Salmon and Dog Salmon River, Ugashik District, 2000.

Location	Survey Date	Number of Sockeye Salmon Counted
<u>King Salmon River System:</u>		
Goose Lake	Aug. 8	5,700
Needle Lake	Aug. 8	<sup>a</sup>
Volcano Creek	Aug. 8	<sup>a</sup>
Painter Creek	Aug. 8	7,200
Indecision Creek	Aug. 8	<sup>a</sup>
Sub-total		12,900
<u>Dog Salmon River System:</u>		
Figure-Eight Creek	Aug. 8	4,200
Goblet Creek	Aug. 8	0
Oldham Creek	Aug. 8	1,000
Wandering Creek	Aug. 8	280
Mainstem Dog Salmon River	Aug. 8	<sup>a</sup>
Sub-total		5,480
 Total		 18,380

<sup>a</sup> No Count.

**Table 7.** Peak survey counts of chinook salmon escapement, Ugashik District, 2000.

Location	Survey Date	Number of Chinook Salmon Counted
<u>King Salmon River System</u>		
Old Creek	Aug. 8	246
Pumice Creek	Aug. 8	339
Painter Creek	Aug. 8	314
Mainstem King Salmon River	Aug. 8	<sup>a</sup>
Indecision Creek	Aug. 8	<sup>a</sup>
Volcano Creek	Aug. 8	<sup>a</sup>
Sub-total		899
<u>Dog Salmon River System</u>		
Figure-Eight Creek	Aug. 8	378
Goblet Creek	Aug. 8	37
Oldham Creek	Aug. 8	<sup>a</sup>
Wandering Creek	Aug. 8	10
Mainstem Dog Salmon River	Aug. 8	<sup>a</sup>
Sub-total		425
<u>Ugashik River System</u>		
Mainstem Ugashik River	28-Jul	26 <sup>b</sup>
Grassy Creek	Aug. 8	<sup>a</sup>
Sub-total		26
Total		1,350

<sup>a</sup> No Count.

<sup>b</sup> Tower count only.

**Table 8.** Peak survey counts of chum salmon escapement, Ugashik District, 2000.

Location	Survey Date	Number of Chum Salmon Counted
<u>King Salmon River System</u>		
Old Creek	Aug. 8	5,850 <sup>a</sup>
Pumice Creek	Aug. 8	7,300 <sup>a</sup>
Painter Creek	Aug. 8	2,150 <sup>a</sup>
Mainstem King Salmon River	Aug. 8	<sup>b</sup>
Needle Lake	Aug. 8	<sup>b</sup>
Indecision Creek	Aug. 8	<sup>b</sup>
Volcano Creek	Aug. 8	<sup>b</sup>
Sub-total		15,300
<u>Dog Salmon River System</u>		
Figure-Eight Creek	Aug. 8	140
Goblet Creek	Aug. 8	40
Oldham Creek	Aug. 8	150
Wandering Creek	Aug. 8	180
Mainstem Dog Salmon River	Aug. 8	<sup>b</sup>
Sub-total		510
<u>Ugashik River System</u>		
Mainstem Ugashik River	July 28	7 <sup>c</sup>
Grassy Creek	Aug. 8	<sup>b</sup>
Sub-total		7
Total		15,817

<sup>a</sup> Includes carcasses.

<sup>b</sup> No Count.

<sup>c</sup> Includes tower and below tower counts.

**Table 9.** Aerial survey counts of coho salmon escapement, Ugashik District, 2000. <sup>a</sup>

Location	Survey Date	Number of Coho Salmon Counted	Comments
<u>Ugashik Drainage</u>			
<u>Upper Ugashik Lake</u>			
Crooked Creek	October 12	950	Most fish in small schools.
Deer Creek	October 12	150	Most fish in small schools.
<u>Lower Ugashik Lake</u>			
Black Creek to Cabin	October 12	150	Up Cabin Creek.
Black Creek to Elizabeth Lake	October 12	3,010	Most fish were up the creeks, but some still holding off the mouths.
Ugashik Outlet	October 12	140	
<u>King Salmon River Tributaries</u>			
Pumice Creek	October 12	710	
Old Creek	October 12	310	
Painter Creek	October 12	6,500	Most fish at the mouth, few were on redds.
<u>Dog Salmon River Tributaries</u>			
Figure Eight Creek	October 12	150	
District Total		12,070	

<sup>a</sup> Only one survey was flown.

**Table 10.** Peak aerial live counts and total escapement estimates of sockeye salmon in the Wood River system, 2000.

Area	Date	Aerial Count	Population Estimate	Distribution %
<b>Wood River</b>				
<b>Lake Aleknagik</b>		<b>17,822</b>		
Eagle Creek <sup>a</sup>	8/12/00	73		
Hansen Creek <sup>a</sup>	8/1/00	75		
Happy Creek <sup>a</sup>	8/5/00	3,385		
Bear Creek <sup>a</sup>	8/8/00	4,158		
Yako Creek <sup>a</sup>	8/4/00	2,624		
Whitefish Creek <sup>a</sup>	8/16/00	687		
Ice Creek <sup>a</sup>	8/9/00	6,743		
Mission Creek <sup>a</sup>	8/15/00	77		
Sunshine Creek				
Youth Creek				
Northshore Beaches				
Southshore Beaches				
Yako Beaches				
<b>Agulowok River &amp; lower River Bay</b>				
<b>Lake Nerka</b>		<b>23,149</b>		
Fenno Creek <sup>a</sup>	8/7/00	8,089		
Pike Creek <sup>a</sup>	8/19/00	893		
Stovall Creek <sup>a</sup>	8/25/00	413		
Bear Creek				
Teal Creek <sup>a</sup>	8/19/00	355		
Pick Creek <sup>a</sup>	8/18/00	4,915		
Elva Creek <sup>a</sup>	8/27/00	321		
Kema Creek <sup>a</sup>	8/26/00	181		
Hidden Lake Creek <sup>a</sup>	8/18/00	3,165		
Lynx Creek <sup>a</sup>	8/25/00	4,817		
Upper River Bay Beaches, NW				
Upper River Bay Beaches, SE				
Allan Cr. - Ross Cr. Beaches				
N6 - River Bay Beach				

(Continued)

**Table 10.** (Continued)

Area	Date	Aerial Count	Population Estimate	Distribution %
Pick Creek Beach				
Elva Creek Beach				
Amakuk Arm Beaches				
Amakuk Arm - Ott's Bay Beach				
Ott's Bay Beach				
Anvil Bay Beaches				
Anvil Bay - Elbow Pt. Beach				
Elbow Pt. - Lynx Cr. Beach				
Lynx Cr. - Teal Cr. Beach				
Kema Lake Beaches				
Hidden Lake Beaches				
Lynx Lake Beaches				
<b>Little Togiak River<sup>a</sup></b>				
<b>Little Togiak Lake</b>		<b>0</b>		
Northshore Beaches				
Southshore Beaches				
D Slough Beaches				
<b>Agulukpak River</b>				
<b>Lake Beverley</b>		<b>1,261</b>		
Tsun Creek				
Moose Creek <sup>a</sup>	8/21/00	1,261		
Hope Creek				
Hardluck Bay Beaches				
Sam's Beach				
Golden Horn Beaches				
Silver Horn Beaches				
B12 & B9 Beaches				
Hope Lake Beach				

(Continued)

**Table 10.** (Continued)

Area	Date	Aerial Count	Population Estimate	Distribution %
<b>Peace River</b>				
<b>Lake Mikchalk</b>		<b>0</b>		
Narrows				
Northshore Beaches				
Southshore Beaches				
<b>Wind River</b>				
<b>Lake Kulik</b>		<b>0</b>		
K1 & K2 Creeks				
K5 Creek - Grant River Beaches				
Grant River - K2 Creek Beaches				
Southshore Beaches				
<b>Grant River</b>				
<b>Total</b>		42,232	0	0.0%

<sup>a</sup> Ground survey counts conducted by FRI, University of Washington.

<sup>b</sup> No aerial surveys were done in 2000. Information in this table is based solely on stream surveys conducted by FRI personnel. No population estimates or distribution information was calculated.

**Table 11.** Peak aerial counts of live sockeye salmon and total escapement estimates, Togiak District, 2000.

Stream	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor <sup>a</sup>	Number
<u>Togiak Section</u>				
Togiak Tower				311,970
Togiak River mainstem	16-Aug	12,600	2.0	25,200
Gechiak Lake System	16-Aug	8,100	1.5	12,150
Pungokepuk Lake	16-Aug	925	1.5	1,388
Nayorurun River	16-Aug	150	1.5	225
Kemuk River	16-Aug	100	1.5	150
Ongivinuk Lake System	16-Aug	775	1.5	1,163
Subtotal		22,650		40,275
<u>Kulukak Section</u>				
Kulukak River	16-Aug	1,825	2.0	3,650
Kulukak Lake	16-Aug	4,275	2.0	8,550
Tithe Creek Ponds	16-Aug	5,075	2.0	10,150
Subtotal		11,175		22,350
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River <sup>b</sup>	14-Aug	526	2.5	1,315
Osviak River <sup>b</sup>	14-Aug	1,512	2.5	3,780
Slug River <sup>b</sup>	14-Aug	1,201	2.0	2,402
Subtotal		3,239		7,497
<u>Other</u>				
Quigmy River <sup>b</sup>	22-Aug	125	2.5	313
Negukthlik River	22-Aug	2,175	3.0	6,525
Ungalikthluk River	22-Aug	575	2.0	1,150
Subtotal		2,875		7,988
Total		39,939		390,080

<sup>a</sup> Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

<sup>b</sup> USFWS estimate.

**Table 12.** Peak aerial counts of live chinook salmon and total escapement estimates, Togiak District, 2000.

River	Aerial Counts		Total Escapement Estimates	
	Date	Number	Factor <sup>a</sup>	Number
<u>Togiak Section</u>				
Togiak River mainstem				
A	16-Aug	75	3.0	225
B	16-Aug	50	3.0	150
C	16-Aug	500	3.0	1,500
D	16-Aug	400	3.0	1,200
E	16-Aug	850	3.0	2,550
F	16-Aug	1,450	3.0	4,350
Subtotal		3,325		9,975
Gechiak River	16-Aug	350	2.5	875
Pungokepuk River	16-Aug	85	2.5	213
Nayorurun River	16-Aug	125	2.5	313
Kemuk River	16-Aug	100	2.5	250
Ongivinuk River	16-Aug	75	2.5	188
Subtotal		735		1,838
Togiak River Drainage Total		4,060		11,813
<u>Kulukak Section</u>				
Kulukak River	16-Aug	340	2.5	850
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River <sup>b</sup>	27-Jul	65	2.5	163
Osviak River <sup>b</sup>	26-Jul	42	3.0	126
Slug River <sup>b</sup>	26-Jul	6	2.5	15
Subtotal		113		304
<u>Other</u>				
Quigmy River <sup>b</sup>	27-Jul	26	2.5	65
Negukthlik River <sup>b</sup>	27-Jul	1,100	3.0	3,300
Ungalikthluk River <sup>b</sup>	27-Jul	226	2.5	565
Subtotal		1,352		3,930
Total		5,865		16,896

a Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

b USFWS estimate.

**Table 13.** Peak aerial counts of live chum salmon and total escapement estimates, Togiak District, 2000.

River	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor <sup>a</sup>	Estimate
<u>Togiak Section</u>				
Togiak River mainstem				
A			2.5	0
B			2.5	0
C			2.5	0
D			2.5	0
E			2.5	0
F			2.5	0
Subtotal		0		0
Gechiak River			2.0	0
Pungokepuk River			2.0	0
Nayorurun River			2.0	0
Kemuk River			2.0	0
Ongivinuk River			2.0	0
Subtotal		0		0
Togiak River Drainage Total		0		0
<u>Kulukak Section</u>				
Kulukak River	16-Aug	4,950	2.0	9,900
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River <sup>b</sup>	27-Jul	9,090	2.0	18,180
Osviak River <sup>b</sup>	26-Jul	10,880	2.5	27,200
Slug River <sup>b</sup>	26-Jul	4,150	2.0	8,300
Subtotal		24,120		53,680
<u>Other</u>				
Quigmy River <sup>b</sup>	27-Jul	2,870	2.0	5,740
Negukthlik River <sup>b</sup>	27-Jul	200	2.5	500
Ungalikthluk River <sup>b</sup>	27-Jul	5,520	2.0	11,040
Subtotal		8,590		17,280
Total		37,660		80,860

a Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

b U.S. Fish and Wildlife Service estimate.

**Table 14.** Peak aerial counts of live coho salmon and total escapement estimates, Togiak District, 2000.

Stream	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor <sup>a</sup>	Number
<u>Togiak Section</u>				
Togiak River mainstem				
A				0
B				0
C	No Surveys Done			0
D				0
E				0
F				0
Subtotal		0		0
Gechiak River				0
Pungokebuk River	No Surveys Done			0
Nayorurun River				0
Kemuk River				0
Ongivinuk River				0
Subtotal		0		0
Togiak River Drainage		0		0
<u>Kulukak Section</u>				
Kulukak River				0
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River <sup>b</sup>				0
Osviak River <sup>b</sup>	No Surveys Done			0
Slug River <sup>b</sup>				0
Subtotal		0		0
<u>Other</u>				
Quigmy River	No Surveys Done			0
Negukthlik River				0
Ungalikthluk River				0
Subtotal		0		0
Total	No Surveys Done			0

<sup>a</sup> Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

<sup>b</sup> U.S.F.W.S. survey.

<sup>c</sup> Survey precluded by muddy water.

Figure 1. Bristol Bay management area, Alaska.




Figure 2. Alagnak River drainage, Bristol Bay, Alaska.





Figure 3. Egegik river drainage, Bristol Bay, Alaska.






Figure 4. Ugashik River drainage, Bristol Bay, Alaska.




Figure 5. Wood River Lakes system, Bristol Bay, Alaska.




Figure 6. Lake Nunavaugaluk system, Bristol Bay, Alaska.




Figure 7. Nushagak-Mulchatna River system, Bristol Bay, Alaska.




Figure 8. Tikchik Lakes system, Bristol Bay, Alaska.




Figure 9. Togiak River system, Bristol Bay, Alaska.




Figure 10. Kulukak River system, Bristol Bay, Alaska.

**Appendix Table 1.** Sockeye salmon total escapement estimates, Naknek-Kvichak District, 1955-2000.  
 Estimates based on visual counts from towers unless otherwise noted.

Year	Kvichak	Naknek	Alagnak	Total	Alagnak Percent of Total
1955	250,546	278,500 <sup>b</sup>	171,500 <sup>a</sup>	700,546	24
1956	9,443,318	1,772,595 <sup>b</sup>	784,000 <sup>a</sup>	11,999,913	7
1957	2,842,810	634,645 <sup>b</sup>	126,595	3,604,050	4
1958	534,785	278,118	94,650	907,553	10
1959	680,000	2,231,807	825,431	3,737,238	22
1960	14,630,000	828,381	1,240,530	16,698,911	7
1961	3,705,849	351,078	90,036	4,146,963	2
1962	2,580,884	723,066	90,630	3,394,580	3
1963	338,760	905,358	203,304	1,447,422	14
1964	957,120	1,349,604	248,700	2,555,424	10
1965	24,325,926	717,798	175,020	25,218,744	1
1966	3,775,184	1,016,445	174,336	4,965,965	4
1967	3,216,208	755,640	202,626	4,174,474	5
1968	2,557,440	1,023,222	193,872	3,774,534	5
1969	8,394,204	1,331,202	122,490	9,847,896	1
1970	13,935,306	732,502	177,060	14,844,868	1
1971	2,387,392	935,754	187,302	3,510,448	5
1972	1,009,962	586,518	151,188	1,747,668	9
1973	226,554	356,676	35,280	618,510	6
1974	4,433,844	1,241,058	214,848	5,889,750	4
1975	13,140,450	2,026,686	100,480	15,267,616	1
1976	1,965,282	1,320,750	81,822	3,367,854	2
1977	1,341,144	1,085,856	100,000 <sup>a</sup>	2,527,000	4
1978	4,149,288	813,378	229,400 <sup>a</sup>	5,192,066	4
1979	11,218,434	925,362	294,200 <sup>a</sup>	12,437,996	2
1980	22,505,268	2,644,698	297,900 <sup>a</sup>	25,447,866	1
1981	1,754,358	1,796,220	82,210 <sup>a</sup>	3,632,788	2
1982	1,134,840	1,155,552	239,300 <sup>a</sup>	2,529,692	9
1983	3,569,982	888,294	96,220 <sup>a</sup>	4,554,496	2
1984	10,490,670	1,242,474	215,370 <sup>a</sup>	11,948,514	2
1985	7,211,046	1,849,938	118,030 <sup>a</sup>	9,179,014	1
1986	1,179,322	1,977,645	230,180 <sup>a</sup>	3,387,147	7
1987	6,065,880	1,061,806	154,210 <sup>a</sup>	7,281,896	2
1988	4,065,216	1,037,862	194,630 <sup>a</sup>	5,297,708	4
1989	8,317,500	1,161,984	196,760 <sup>a</sup>	9,676,244	2
1990	6,970,020	2,092,578	168,760 <sup>a</sup>	9,231,358	2
1991	4,222,788	3,578,508	277,589 <sup>a</sup>	8,078,885	3
1992	4,725,864	1,606,650	226,643 <sup>a</sup>	6,559,157	3
1993	4,025,166	1,535,658	347,975 <sup>a</sup>	5,908,799	6

(Continued)

**Appendix Table 1.** (page 2 of 2)

---

Year	Kvichak	Naknek	Alagnak	Total	Alagnak Percent of Total
1994	8,337,840	990,810	242,595 <sup>a</sup>	9,571,245	3
1995	10,038,720	1,111,140	215,713 <sup>a</sup>	11,365,573	2
1996	1,450,578	1,078,098	306,750 <sup>a</sup>	2,835,426	11
1997	1,503,732	1,025,664	218,115 <sup>a</sup>	2,747,511	8
1998	2,296,074	1,202,172	252,200 <sup>a</sup>	3,750,446	7
1999	6,196,914	1,625,364	463,600 <sup>a</sup>	8,285,878	6
2000	1,827,780	1,375,488	451,300 <sup>a</sup>	3,654,568	12
Mean	5,433,266	1,223,057	234,152 <sup>c</sup>	6,902,222	5

---

<sup>a</sup> Aerial survey counts.

<sup>b</sup> Weir counts.

<sup>c</sup> Mean of counts from 1977 to present.

**Appendix Table 2.** Aerial survey counts of chinook salmon escapements, Naknek River drainage, 1971-2000.

Year	Mainstem Naknek River	Paul's Creek	King Salmon Creek	Big Creek	Total
1971	1,639	52	704	490	2,885
1972	351	156	1,224	1,060	2,791
1973	1,315		115	1,106	2,536
1974		91	495	860	1,446
1975	2,250	144	279	779	3,452
1976	5,950	31	180	970	7,131
1977	4,830		1,860		6,690
1978					<sup>a</sup>
1979					<sup>a</sup>
1980	300	17		30	347
1981	2,890		591	790	4,271
1982	5,360	340	980	1,930	8,610
1983	2,860	290	460	4,220	7,830
1984	790	400	385	3,420	4,995
1985	590				590
1986	2,200	73	102	1,542	3,917
1987	2,800	7	290	1,353	4,450
1988	7,380	150	600	3,600	11,730
1989	1,700	50	100	860	2,710
1990	4,500	150	350	2,000	7,000
1991	1,655	121	275	2,340	4,391
1992	1,550	88	158	895	2,691
1993	5,520	86	700	1,710	8,016
1994	5,970	203	974	2,531	9,678
1995	2,790	26	239	1,905	4,960
1996	2,965	157	312	1,576	5,010
1997	7,520	248	902	1,783	10,453
1998	2,150	210	1,060	2,085	5,505
1999		223	847	2,250	3,320
2000	1,900	43	178	1,112	3,233
Mean	3,066	140	552	1,661	5,420 <sup>b</sup>
Percent	57	3	10	31	100

<sup>a</sup> Counts unavailable.

<sup>b</sup> The sum of mean indices.

**Appendix Table 3.** Chinook salmon escapement survey history, mainstem Naknek River, 1929-2000.

Year	Count Dates	Surveyors	Actual Weir Count <sup>a</sup>	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>b</sup>	Comments
1929	7/03-7/31		1,498			Chinook count peaked 7/27.
1930	6/20-8/09		1,999			Chinook count peaked 8/09.
1931	6/17-8/09		896			Chinook count peaked 8/07.
1932	6/23-8/10		1,869			
1950	7/08-8/20		3,097			Chinook count peaked 8/09.
1951	6/28-8/07		1,876			Chinook count peaked 8/04.
1952	6/25-8/10		633			Chinook count peaked 8/06.
1953	6/24-8/10		2,074			Chinook count peaked 7/26.
1954	6/20-8/11		3,474			Chinook count peaked 8/10.
1955	6/13-8/17		4,188			Chinook count peaked 8/16.
1956	6/22-8/28		7,378			Chinook count peaked 8/18.
1957	6/28-8/04		8,504			Chinook count peaked 8/03.
1966		Redick				300 were counted 8/26 from a skiff in the Rapids.
1967	Mid-Aug.	Paddock			800	
1968					1,200	Conservative estimate.
1969					1,200	
1970	7/31	Whitehead		845		
	8/03	Siedelman		3,060		Visibility very good. Super-cub.
	8/22	Siedelman		1,540	1,750	Water high & murky. Spawning pre-peak.
	8/22	Whitehead		1,310		
	8/25	Whitehead		2,225		Counting conditions optimal.
	8/25	Siedelman		2,536	2,500	Conditions good. Spawning pre-peak.

(Continued)

Appendix Table 3. (page 2 of 3)

Year	Count Dates	Surveyors	Actual Weir Count <sup>a</sup>	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>b</sup>	Comments
1971	8/26	Cunningham		1,639		Fish concentrated near Rapids Camp. Few dead.
1972	8/23	Cunningham & McCurdy		351		Poor counting conditions. Post-peak.
1973	8/19	Russell		1,315		Counting conditions good. Peak near at hand.
1974	8/19	Russell			450	Count accuracy questionable. Many fish were deep.
1975	8/17	Russell		2,250		Good viewing, peak near. Still fish spawning 9/08.
1976	8/13	Bill		2,615		Spawning near peak. Very few dead.
	8/16	Russell		5,950	7,250	Pre-peak. Still lots fish holding in large groups.
1977	8/22	Russell		4,830	5,750	Pre-peak. Few dead. Some still holding deep.
1978	8/09	Gwartney			4,000	Near peak.
1983	8/14	Bill		2,860	3,000	Pre-peak. Still fish holding in large groups.
1984	8/14	Bill		790	2,370	
1985	8/06	Bill			600	Pre-peak.
	8/27	Bill		590	700	
1986	8/18	Russell		1,990		Spawning pre-peak. Still many fish holding.
	8/19	Meyer		2,200		Peak of spawning drawing near.
1987	8/19	Meyer		2,800		Pre-peak. Fish still in large groups. Few redds.
	8/28	Bill		2,655	2,855	
1988	8/09	Minard		7,380	7,400	Approaching peak. Most fish on redds.
1989	8/14	Minard		1,700		Fish actively spawning. Few carcasses observed.
1990	8/06	Minard		4,500		
1991	8/20	Russell		1,655		Pre-peak. Still many fish schooled & waiting.

(Continued)

Appendix Table 3. (page 3 of 3)

Year	Count Dates	Surveyors	Actual Weir Count <sup>a</sup>	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>b</sup>	Comments
1992	8/21	Regnart		877		Water clarity poor in deeper pools.
	8/27	Regnart		1,550		At Peak...all fish on redds.
1993	8/23	Regnart		5,520		Near peak. Still some fish schooled.
1994	8/24	Regnart		5,970		Near peak. Most on redds.
1995	8/21	Regnart		2,790		Near peak. Most on redds.
1996	8/21	Regnart		2,965		At Peak...all fish on redds.
1997	8/16	Regnart		7,520		Near peak. Most on redds.
1998	8/18	Regnart		2,150		At Peak...all fish on redds.
1999	no survey					Survey flown September 8 no count available.
2000	8/07	Morstad		1,900		Early still fish schooled
Mean			3,124	2,752		

<sup>a</sup> Weir count did not account for estimated 15-20% of chinook that spawn downstream of weir site. Also does not account for fish that migrated upstream past the weir site before and after weir operation.

<sup>b</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 4.** Chinook salmon escapement survey history, Big Creek, Naknek River Drainage, 1963-2000.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1963	8/01	Paddock		362		Covered only half stream length. Helicopter.
	8/13	Paddock		1,345	2,690	Spawning near peak. Good survey.
1964	7/31	Paddock		484		Survey too early.
	8/15	Siedelman & Williamson		636		Survey fair to good. Near peak. Helicopter.
	8/15-8/18	Siedelman & Williamson	1,130			Peak of spawning over.
1965	8/05-8/08	Andrews	578			Fair survey. Began below Index Area No. 1.
1966	8/13-8/16	Redick	979			Spawning at peak. Included Index Area No. 1. Count affected by rain/turbid water in lower areas.
1967	8/10-8/14	Whitehead & Bury	1,129			Upstream redds occupied while those in the lower stream area were abandoned.
1968	8/10-8/14	Meyers & Preyer	3,827			Counting conditions fair to poor.
1969	8/12-8/14	Parkinson & Faro	1,012			High murky waters hampered float count.
	Mid-Aug.	??			5,000	Flown due to poor count conditions during float.
1970	7/19	Whitehead		825		
	8/15-8/17	Parkinson & Brooks	1,601			High murky waters in lower 2/3 of stream.

(Continued)

Appendix Table 4. (page 2 of 3)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1971	8/13	Cunningham		490	1,200	Only upper 1/3 of stream surveyed due to murky water in lower 2/3.
	8/28	Siedelman		277		Past peak. Survey affected by winds of 30+ mph.
1972	8/08	Cunningham		695		Pre-peak.
	8/18	Siedelman		1,060		Post-peak.
1973	8/17	Russell		1,106		At peak of spawning. Many fish beaten up (fungus).
1974	8/01	Russell		520	850	Pre-peak. No dead chinook. Lots dead chums.
	8/11	Russell		860	1,250	Didn't survey lower 8 miles of creek 8/11. Could add 150 fish to survey as Russell saw that many in the unsurveyed portion from skiff 8/10. Near peak.
1975	8/09	Russell		779		Survey pre-peak.
1976	8/13	Bill		970	1,400	Not total stream coverage due to winds & low fuel.
1983	8/14	Bill		4,220	9,000	
1984	8/08	Bill		3,420	8,800	At peak of spawning.
1985	8/06	Bill			2,900	Survey conditions..high water & gusty winds.
1986	8/08	Meyer		1,542	6,000	Excellent conditions. Fish at spawning peak.
1987	8/21	Meyer		1,353	2,500	
1988	8/09	Minard		3,600		
1989	8/14	Minard		860		
1990	8/06	Minard		2,000		
1991	8/12	Regnart		2,340		At spawning peak..all fish on redds, only 20 dead.
1992	8/18	Regnart		895		Est. 5-6 days post-peak. Count includes 125 dead.

(Continued)

**Appendix Table 5.** Chinook salmon escapement survey history, King Salmon Creek, Naknek River drainage, 1964-2000.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1964	7/31	Paddock		378		Survey conditions fair. Helicopter.
	8/11	Paddock		55		Visibility poor. Helicopter.
	8/11-8/14	Paddock & Siedelman	104			Peak of spawning long past. Poor survey (turbid).
1966	7/31-8/03	Redick	633			Spawning at or near peak.
1967	7/24-7/26	Paddock	289			Poor visibility. Estimated 600 fish present.
1968	7/17	Whitehead		282		Pre-peak. Helicopter.
	7/17	Meyers		242		Pre-peak. Helicopter.
	7/20	Whitehead		868		Optimum conditions. Count from H-21 Helicopter.
	7/20	Meyers		575		Optimum conditions. Count from H-21 Helicopter.
	7/20-7/23	Whitehead & Meyers	2,204			Counting conditions optimum.
1969	7/23-7/25	Parkinson & Berry	2,722			Pre-peak. Count fair-to-poor last 2 days (weather).
1970	7/19	Whitehead		260		Counting conditions poor. Pre-peak.
1971	7/28	Cunningham		704		Visibility was good.
1972	7/29	Siedelman		1,224		Peak of spawning.
1973	8/01	Siedelman		115		Visibility only fair. Survey possibly post-peak.
1974	7/15	Russell		164	350	Pre-peak. Many fish holding in pools.
	7/28	Russell		495	625	At or near peak. Only one carcass obsd. Good vis.
1975	7/28	Russell		279	375	Survey pre-peak. Good viewing conditions.
	8/10	Russell	67			Floated only lower 12 miles of creek.
	8/17	Russell		0		Excellent viewing conditions. Spawning is done.
1976	8/03	Bill		180	400	Peak within next 3 days.
1977	7/29	Russell		1,860	2,350	At peak of spawning.

(Continued)

Appendix Table 5. (page 2 of 2)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1978	8/09	Gwartney			350	Past peak. Viewing good. Most fish dead or spent.
1979	??	Gwartney			1,750	
1980	8/08	Bill				Creek too high & muddy to census.
1981	7/30	Russell		591	1,500	Peak of spawning in progress. Vis = fair-to-poor.
1982	8/07	Bill		980	3,920	Good visibility.
1983	8/14	Bill		460	1,400	Poor visibility. Muddy. 30% spawners dead already.
1984	8/08	Bill		385	1,155	
1988	8/08	Minard		600		At peak.
1989	8/14	Minard		100		Past peak.
1990	8/06	Minard		350		
1991	7/30	Russell		100		Pre-peak and water clarity only "Fair".
	8/05	Russell		275		Est. at spawning peak, most fish on redds, 2 dead.
1992	8/09	Russell		158		Post-peak as 47 dead counted & aband. redds numerous.
1993	7/31	Russell		700	900	Slightly pre-peak. Most fish on redds. Water clear.
1994	7/29	Russell		974		Slightly pre-peak. Most fish on redds. Only 6 carcasses.
1995	8/05	Russell		239		A little past peak. Several singles on redds. Vis. only
1996	8/05	Regnard		312		Slightly post peak. 26 dead counted.
1997	7/18	Regnard		902		Pre-peak and water clarity "Good".
1998	8/18	Regnard		1,060		Estimate is at peak of spawn.
1999	8/02	Morstad		847		Estimate near peak of spawn, fair to good conditions.
2000	8/01	Morstad		178		Estimate near peak, survey conditions fair
Mean			1,190	497		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for conditions, river areacounting at time of survey.

**Appendix Table 6.** Chinook salmon escapement survey history, Paul's Creek, Naknek River drainage, 1971-2000.

Year	Count Dates	Surveyors	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1971	7/28	Cunningham	52		
1972	7/28	Siedelman	156		Prior to peak.
1973	8/01	Siedelman			Too murky to survey.
1974	7/15	Russell	2		
	7/26	Russell	91	250	Prior to spawning peak.
1975	7/28	Russell	144	225	Prior to peak. Good conditions.
1976	8/03	Bill	31	100	Poor conditions. Fish paired & spawning.
1977					No count.
1978	8/09	Gwartney		300	Past peak. 75% of fish dead.
1979					No count.
1980	8/08	Bill	17		All carcasses. Creek high & muddy.
1981					No count.
1982	8/07	Bill	340	1,020	Good visibility. Spawning near peak.
1983	8/14	Bill	290	800	Poor visibility.
1984	8/08	Bill	400	800	Fair visibility. About 25% dead already.
1985	8/06	Bill		170	Pre-peak.
1986	8/08	Meyer	73	236	Approximately 30% dead already.
1987	8/13	Russell	7		Poor survey conditions. Past peak.
	??	Meyer		400	Estimat 400 present based on jet boat surveys.
1988	8/08	Minard	150		At peak.
1989	8/14	Minard	50		Past peak. Excellent visibility.
1990	8/06	Minard	150		Excellent survey conditions.

(Continued)

Appendix Table 6. (page 2 of 2)

Year	Count Dates	Surveyors	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1991	7/30	Russell	121		Slightly pre-peak. Only 1 carcass noted.
1992	8/01	Russell	88		Slightly pre-peak. Stream clarity only "Fair".
1993	7/31	Russell	86	140	Slightly pre-peak. Overflow approx 60% of stream.
1994	7/29	Russell	203	300	Pre-peak...but many fish on redds.
1995	8/05	Russell	26		Water clarity poor. 5 carcasses noted
1996	8/05	Regnart	157		Peak of spawning. 12 dead counted.
1997	7/18	Regnart	248		Pre-peak. Excellent visibility
1998	8/18	Regnart	210		
1999	8/02	Morstad	223		Pre spawning, 10% on redds and two carcasses
2000	8/02	Morstad	43		Pre spawning, 10% on redds no carcasses
Mean			134		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 7.** Chinook salmon escapement survey history, Alagnak River, 1963-2000.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1963	8/12	Siedelman		551		Excellent conditions. No side channels flown.
1966	8/06	Redick		13		Poor conditions.
	8/06-8/10	Redick	238			Nonvianuk & mainstem portions only (not Kukaklek).
	8/11	Redick		1,465		Pre-peak. Still many fish upmigrating.
1967	8/16	VanValin		1,250		
1968	8/18	Siedelman		6,717	8,500	Fairly good survey.
1969	8/19	Siedelman		4,781	6,000	Marginal survey conditions, (20kn NW winds).
1970	8/22	Siedelman		5,250	5,000	Peak of spawning. Visibility good
	8/22	Whitehead		4,590		Peak of spawning. Visibility good
1971	8/25	Siedelman		1,420	1,500	Water high, but count okay.
	8/25	Cunningham		1,475		
1972	8/23	Cunningham		2,256	2,400	Past peak. Many dead. Many unoccupied redds.
1973	8/16	Russell		824	1,250	Near peak of spawning. No dead though.
1974	8/13	Russell		1,411	1,700	Pre-peak.
	8/19	Russell		1,596	1,900	Spawning near peak.
1975	8/17	Russell		6,620	7,250	About a week pre-peak. Some large groups holding.
1976	8/16	Bill		7,593	8,750	Pre-peak. Not many dead yet.
1977	8/18	Bill		3,634	12,000	Pre-peak. Didn't count river below Pfaff Pond.
	8/18	Sanders		9,425		Pre-peak. Didn't count river below Pfaff Pond.
1978	8/24	Bill		11,650	25,100	
1979						No survey.
1980	8/08	Bill		2,020	5,090	Pre-peak. Fog over lower river.
	8/21	Bill		2,930	5,860	
1981	8/26	Bill		2,430	8,540	

(Continued)

Appendix Table 7. (page 2 of 2)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1982	8/09	Bill		3,400	4,700	At least a weak too early.
	8/19	Bill		3,350	5,480	Peak survey.
1983	8/15	Bill		2,980	3,500	At peak of spawning.
1984	8/14	Bill		6,090	9,135	
1985	8/17	Bill		3,920	9,518	About peak for chinook spawning. 30% dead already.
1986	8/11	Bill		3,090	7,200	Peak of spawning.
1987	8/22	Bill		2,420		
1988	8/12	Bill		4,600		
1989	8/15	Bill		3,650		
1990	8/08	Bill		1,720		
1991	8/09	Regnart		2,023		Pre-peak. Most fish schooled yet. Few on redds.
	8/19	Regnart		2,531		Near peak. Most fish on redds.
1992	8/10	Regnart		3,042		Pre-peak. Most fish still schooled.
	8/21	Regnart		2,275		Near peak...but water clarity worse than earlier.
1993	8/09	Regnart		10,170		Near peak. Most on redds.
1994	8/08	Regnart		8,480		About half the fish on redds. Others schooled.
1995	8/10	Regnart		6,860		About 2/3 of chinook noted on redds.
1996	8/12	Regnart		9,885		Near peak. Most on redds.
1997	8/7	Regnart		15,210		Peak. Excellent visibility
1998	8/12	Anderson		4,148		About 1/3 of braids poor light; most on redds.
1999	8/10	Morstad		2,178		Peak of spawning, good survey conditions
2000	8/7	Morstad		2,220		Peak of spawning, good survey conditions
Mean			238	4,185		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 4.** (page 3 of 3)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1993	8/17	Regnart		1,710		Estimated survey 3-4 days past peak.
1994	8/16	Regnart		2,531		Est. 2-3 days post-peak. Count includes 159 dead.
1995	8/15	Regnart		1,905		Estimate survey was several days past peak.
1996	8/12	Regnart		1,576		At spawning peak....38 dead observed
1997	8/7	Regnart		1,783		At spawning peak....48 dead observed
1998	8/18	Regnart		2,085		At spawning peak....no carcasses present
1999		Morstad		2,250		At spawning peak....no carcasses present
2000	8/7	Morstad		1,112		At spawning peak....4 dead observed
Mean			1,465	1,471		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 8.** Chinook salmon escapement survey history, Kvichak River, 1932-2000.

Year	Count Dates	Surveyors	Weir Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1932	6/28-8/5		5,753			Peak count was on 7/05 (1,168 fish).
1976	8/16	Bill		35	45	Survey timed to count pink salmon.
1980 <sup>b</sup>	8/08	Bill		900	1,000	Chinook actively spawning.
1984	8/14	Bill		200		
1988	8/13	Bill		190	570	Nearly all on redds.
1989	8/16	Bill		100	260	
1990	8/19	Bill		170	510	
1992	8/13	Regnart		264		All fish on redds in Kaskanak Flats.
1993	8/16	Regnart		115		All fish on redds in Kaskanak Flats.
1994	8/12	Regnart		306		
1995	8/14	Regnart		96		
1996	8/18	Regnart		132		
1997	8/15	Regnart		103		
1998	8/14	Anderson		187		All fish on redds in Kaskanak Flats
1999	8/10	Morstad		1,200		
2000	8/07	Morstad		6		
Mean			5,753	267		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

<sup>b</sup> Pecks Creek, a Kvichak River tributary, was float surveyed 7/30-8/03, 1980 by R. Russell and 99 spawning chinook salmon were counted.

**Appendix Table 9.** Chinook salmon escapement data, Naknek-Kvichak District, 1970-2000.

Non-expanded Escapement Indices by Drainage <sup>a</sup>				
Year	Naknek	Alagnak	Kvichak	Total
1970	4,145 <sup>b</sup>	5,250		9,395
1971	2,885	1,420		4,305
1972	2,791	2,256		5,047
1973	2,536 <sup>b</sup>	824		3,360
1974	1,446 <sup>c</sup>	1,596		3,042
1975	3,452	6,620		10,072
1976	7,131	7,593	35	14,759
1977	6,690 <sup>b</sup>	3,634		10,324
1978	<sup>d</sup>	11,650		11,650
1979	<sup>d</sup>	<sup>d</sup>		0
1980	347 <sup>e</sup>	2,930	900	4,177
1981	4,271 <sup>b</sup>	2,430		6,701
1982	8,610	3,400		12,010
1983	7,830	2,980		10,810
1984	4,995	6,090	200	11,285
1985	590 <sup>f</sup>	3,920		4,510
1986	3,917	3,090		7,007
1987	4,450	2,420		6,870
1988	11,730	4,600	190	16,520
1989	2,710	3,650	100	6,460
1990	7,000	1,720	170	8,890
1991	4,391	2,531		6,922
1992	2,691	3,042	264	5,997
1993	8,016	10,170	115	18,301
1994	9,678	8,480	306	18,464
1995	4,960	6,860	96	11,916
1996	5,010	9,885	132	15,027
1997	10,453	15,210	103	25,766
1998	5,505	4,148	187	9,840
1999	3,320 <sup>c</sup>	2,178	1,200	6,698
2000	3,233	2,220	6	5,459
Mean <sup>g</sup>	4,670	4,760	267	9,697

<sup>a</sup> Includes aerial indices from all streams surveyed in drainage.<sup>b</sup> No index count for Paul's Creek.<sup>c</sup> No index count for Naknek River.<sup>d</sup> No non-expanded index counts exist for this year.<sup>e</sup> Includes only index counts for mainstem Naknek River, Paul's Creek, & Big Creek.<sup>f</sup> Naknek River mainstem only.<sup>g</sup> Sum of mean indices.

**Appendix Table 10.** Chum salmon escapement survey history, Alagnak River, 1961-2000.

Year	Count Dates	Surveyors	Tower Counts	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1961			18,906			
1962			3,846			
1963	8/12	Siedelman	20,124	4,120		
1964			2,562			
1965			132			
1966						
1967			9,990			
1968			72			
1969			210			
1970			5,790			
1971			402			
1972			48			
1973						
1974						
1975						
1976	8/16	Bill		2,125	5,250	
1977	8/18	Bill		35,000		
1978	8/24	Bill		9,900		
1979						
1980	8/21	Bill		7,300	14,600	
1981	8/26	Bill		75,000	75,000	
1982	8/09	Bill		14,000	42,000	
	8/19	Bill		12,000	30,000	
1983	8/15	Bill		8,800		Pre-peak.

(Continued)

Appendix Table 10. (page 2 of 2)

Year	Count Dates	Surveyors	Tower Counts	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1984	8/14	Bill		48,000	87,500	
1990	8/08	Bill		8,500	30,000	Pre-peak.
	8/18	Bill		48,800		Close to peak of spawning.
1991	8/09	Regnart		43,000		Pre-peak.
	8/19	Regnart		64,300		Peak of spawning.
1992	8/10	Regnart		114,000		Near Peak.
1993	8/09	Regnart		4,600		Near Peak.
1994	8/08	Regnart		62,900		Near Peak.
1995	8/10	Regnart		132,000		Near Peak.
1996	8/12	Regnart		145,000		Near Peak
1997	8/07	Regnart		37,800		Near Peak
1998	8/12	Anderson		3,150		Poor survey conditions
1999	8/10	Morstad		11,800		Near Peak
2000	8/07	Morstad		10,120		Near Peak
Mean			3,575	39,227		

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of chum salmon spawners in the river at time of aerial survey, based on survey conditions, coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for river area counting at time of survey.

**Appendix Table 11.** Pink salmon escapement survey history, Alagnak River, 1968-2000.

Year	Count Dates	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1968	8/27	Siedelman	97,000	125,000	
1970					No survey.
1972					No survey.
1974	8/14	Bill	20,600		Big schools. Pre-peak.
1976	8/16	Bill	6,375	13,000	Pre-peak.
1978	8/24	Bill	330,300	736,000	Just starting to spawn. Many still in lower river.
1980	8/21	Bill	121,000	242,000	
1982	8/09	Bill	21,300	63,900	
	8/19	Bill	24,800	43,000	Pre-peak.
1984	8/14	Bill	296,500	567,100	Survey too early for peak. Most fish schooled.
1986	8/11	Bill	48,600	145,800	
1988	8/12	Bill	415,000	620,000	Pre-peak.
1990	8/08	Bill	45,100		
	8/18	Bill	240,500		Estimated to be about 1 week pre-peak.
1992	8/10	Regnart	15,000		Pre-peak.
1994	8/08	Regnart			No pinks noted.
1996	8/12	Regnart			No pinks noted.
1998	8/12	Anderson	3,200		High water poor light conditions
2000	8/7	Morstad	30,000		
Mean			114,352	212,983	

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting a time of survey.

**Appendix Table 12.** Pink salmon escapement survey history, Kvichak River, 1966-2000.

Year	Count Dates	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1966		Robertson		67,500	
1968	8/26	Siedelman		88,000	
1970					No survey.
1972					No survey.
1974	8/14	Bill		30,560	
1976	8/16	Bill		16,100	Most still schooled.
1978	8/28	Bill	88,000	440,000	Still numerous fish migrating & some schooled.
1980	8/08	Bill	7,000	25,000	Still schooled.
1982					No Survey.
1984	8/14	Bill	111,000	165,000	
1986					No survey.
1988	8/13	Bill	94,000		
1990	8/19	Bill	25,300	47,000	
1992					No survey.
1994					No survey.
1996					No Survey
1998					No Survey.
2000	8/07	Morstad	7,000		Still schooled.
Mean			55,383	109,895	

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 13.** Pink salmon escapement survey history, Naknek River, 1974-2000.

Year	Count Dates	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate <sup>a</sup>	Comments
1974	8/14	Bill	161,800	362,000	
1976	8/13	Bill	94,600	110,000	Just pre-peak. Many still schooled.
1978	8/24	Bill	312,000	780,000	
1980	8/08	Bill	80,000	160,000	Pre-peak.
1982	8/19	Bill	33,600	34,000	Pre-peak.
1984	8/14	Bill	27,000	125,000	
1986	8/18	Russell	286,000	375,000	Most fish still schooled and holding. Pre-peak.
1988	8/24	Russell	187,000		
1990	8/18	Bill		65,000	
1992					No survey.
1994					No survey.
1996					No Survey.
1998					No survey.
2000	8/07	Morstad	10,000		
Mean			132,444	251,375	

<sup>a</sup> Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

**Appendix Table 14.** Aerial survey counts of chinook salmon escapement, Egegik District, 1981-2000. <sup>a</sup>

Year	Egegik River	Shosky Creek	Whale Mountain Creek	Mossy Creek	Mink Creek	Gertrude Creek	Kaye's Creek	Takayoto Creek	Angle Creek <sup>f</sup>	Contact Creek	King Salmon River	Total
1981						515						515
1982	300					900				300		1,500
1983						860		380		375		1,615
1984	40	300				600		350		110		1,400
1985	75	80	0	15	10	260	230	315		95		1,080
1986	65	150	48	0	0	150	46	40		18	15	532 <sup>b</sup>
1987	15	174	2	74	0	408	284	232	2	88		1,279
1988	50	151	0	12		248	120	177		110		868
1989	14	90	13	43	7	310	120	300		100		997
1990	24 <sup>c</sup>	85	7	35	2	260	175	175		205		968
1991	0 <sup>c</sup>	62	60	30	33	83	117	95		73		553
1992 <sup>d</sup>	15	143	52	54	22	416	320	190		296		1,508
1993	80	58	6	38	6	350	170	200		235		1,143
1994 <sup>d</sup>	66 <sup>c</sup>	48	32	118	77	840	214	230		705		2,330
1995 <sup>d</sup>	60 <sup>c</sup>	32	10	53	103	456	248	130		275		1,367
1996	42 <sup>c</sup>	102	8	38	20	230	74	123	6	203	<sup>g</sup>	846
1997	30 <sup>c</sup>	39	2	18	10	260	173	374	<sup>g</sup>	740	<sup>g</sup>	1,646
1998	0 <sup>c</sup>	29	45	55	<sup>g</sup>	320	165	120	<sup>g</sup>	329	<sup>g</sup>	1,063
1999	6 <sup>c</sup>	75	10	51	<sup>g</sup>	165	6	115	<sup>g</sup>	145	<sup>g</sup>	573
Average	52	101	20	42	24	402	164	209	4	245	15	1,146
2000	0 <sup>c</sup>	4	0	16	0	85	41	73	<sup>g</sup>	341	<sup>g</sup>	560
Deviation <sup>e</sup>		-96%	-100%	-62%		-79%	-75%	-65%		39%		-51%

<sup>a</sup> Peak aerial counts unless otherwise noted. Data not expanded.

<sup>b</sup> Survey 10-14 days later than normal.

<sup>c</sup> Tower count.

<sup>d</sup> Helicopter surveys.

<sup>e</sup> 2000 deviation from 1981-1999 average.

<sup>f</sup> Angle Creek is usually too turbid to survey.

<sup>g</sup> No Count.

**Appendix Table 15.** Aerial survey counts of chum salmon escapement, Egegik District, 1982-2000. <sup>a</sup>

Year	Egegik River	Shosky Creek	Whale Mountain Creek	Mossy Creek	Mink Creek	Gertrude Creek	Kaye's Creek	Takayoto Creek	Angle Creek <sup>b</sup>	Contact Creek	King Salmon River	Total
1982						12,000				2,000		14,000
1983	6 <sup>c</sup>					5,000		3,500		6,000		14,500
1984	800	200				13,000		2,400		10,000		26,400
1985	400	0	600	200	35	2,600	800	0		500	50	5,185
1986	0	0	6,025			140	3	5	0	15	25	6,213 <sup>d</sup>
1987	150	0	19,000	16	1,000	3,770	2,780	0		2,850		29,566
1988	500	50	4,400	100	50	5,200	1,600	0		3,200		15,100
1989	0	10	3,200	25	100	1,100	0	0		200	14	4,649
1990	72 <sup>c</sup>	0	2,000	0	150	1,675	80	0		750		4,727
1991	0 <sup>c</sup>	0	1,500	70	100	990	280	0		480		3,420
1992 <sup>e</sup>	50	0	680	15	25	4,500	400	0		3,630	200	9,500
1993	100	0	1,020	8	1	1,075	0	0		100		2,304
1994 <sup>e</sup>	42 <sup>c</sup>	0	1,700	5	7	760	175	30		260		2,979
1995 <sup>e</sup>	144 <sup>c</sup>	2	395	15	30	560	162	5		600		1,913
1996	12	<sup>f</sup>	438	4	20	530	<sup>f</sup>	24	<sup>f</sup>	633	<sup>f</sup>	1,661
1997	0 <sup>c</sup>	<sup>f</sup>	220	8	10	495	290	60	<sup>f</sup>	640	<sup>f</sup>	1,723
1998	17 <sup>c</sup>	8	1,480	4	<sup>f</sup>	920	4	4	<sup>f</sup>	140	<sup>f</sup>	2,577
1999	6 <sup>c</sup>	<sup>f</sup>	1,040	4	<sup>f</sup>	243	<sup>f</sup>	4	<sup>f</sup>	140	<sup>f</sup>	1,437
Average	143	21	2,913	34	127	3,031	506	355	0	1,785	72	8,214
2000	0 <sup>c</sup>	<sup>f</sup>	492	<sup>f</sup>	4	475	32	6	<sup>f</sup>	180	<sup>f</sup>	1,189

<sup>a</sup> Peak aerial counts unless otherwise noted. Data not expanded.

<sup>b</sup> Angle Creek is usually too turbid to survey.

<sup>c</sup> Tower count.

<sup>d</sup> Survey 10-14 days later than normal.

<sup>e</sup> Helicopter surveys.

<sup>f</sup> No Count.

**Appendix Table 16.** Aerial survey counts of pink salmon escapement, Egegik District, 1974-2000.<sup>a</sup>

Year	Egegik River	Whale Mountain Creek	Gertrude Creek	Contact Creek	Other	Total
1974	3,912 <sup>b</sup>					3,912
1976	0 <sup>b</sup>					0
1977	84 <sup>b</sup>					84
1980	0 <sup>b</sup>					0
1982	15,000					15,000
1983			58 <sup>c</sup>			58
1984	17,000					17,000
1985						
1986	2,500					2,500
1987						
1988	23,000					23,000
1989	300					300
1990	17,000		40 <sup>c</sup>			17,040
1991		88 <sup>d</sup>	24 <sup>d</sup>	36 <sup>d</sup>		148
1992 <sup>e</sup>	6 <sup>b</sup>	10			3	13
1993	50					50
1994	21,282 <sup>b</sup>					21,282
1995	24 <sup>b</sup>					24
1996	103,116 <sup>b</sup>					103,116
1997	0 <sup>b</sup>		1,290 <sup>f</sup>			1,290
1998	2 <sup>b</sup>		2,487 <sup>f</sup>			2,489
1999	6 <sup>b</sup>		1,125 <sup>f</sup>			1,131
Average	13,550	49	837	36	3	10,422
2000 <sup>g</sup>	0 <sup>b</sup>					

<sup>a</sup> Non-expanded aerial peak counts unless otherwise noted.

<sup>b</sup> Tower counts.

<sup>c</sup> Float count.

<sup>d</sup> Foot survey (USFWS).

<sup>e</sup> Helicopter surveys.

<sup>f</sup> Gertrude Creek Weir count.

<sup>g</sup> No Counts.

**Appendix Table 17.** Aerial survey counts of coho salmon escapement, Egegik District, 1981-2000.

Year	Number of Surveys	Coho Salmon Count	Comments
1981	1 <sup>a</sup>	4,000	Only Becharof tributaries surveyed.
1982	1	20,000	Surveyed on August 20.
1983	0	0	No surveys done.
1984	3	43,225	40,000 counted in Egegik Lagoon on August 15.
1985	3	5,260	Peak surveys on August 26.
1986	1	12,575	Surveyed August 19.
1987	6	6,930	Included King Salmon River & tributaries.
1988	6	13,715	Included King Salmon River & tributaries.
1989	9	4,485	Included Gertrude & Whale Mountain Creeks.
1990	7	13,400	Peak survey on August 17.
1991	0	220	Incidental observation made August 6.
1992	<sup>b</sup> 0	200	Incidental observation in Egegik River August 6.
1993	0	1,130	Incidental observation from Egegik River August 16.
1994	<sup>b,c</sup> 2	7,412	Included King Salmon River & tributaries.
1995	<sup>d</sup> 2	5,258	Included King Salmon River & tributaries.
1996	<sup>e</sup> 2	9,043	Included King Salmon River & tributaries.
1997	3	4,106	Gertrude Weir Count & selected Becharof Lake tributaries.
1998	1	6,075	Gertrude Weir Count & selected Becharof Lake tributaries.
1999	1	4,353	Gertrude Weir Count & selected Becharof Lake tributaries.
2000	1	4,870	Selected Becharof Lake tributaries

<sup>a</sup> Survey done by USFWS personnel.

<sup>b</sup> Helicopter surveys.

<sup>c</sup> The Egegik River Tower was maintained through September 11 and approximately 10,140 coho salmon were counted.

<sup>d</sup> The Egegik River Tower was maintained through August 30 and approximately 7,470 coho salmon were counted.

<sup>e</sup> The Egegik River Tower was maintained August 7 to September 11 and approximately 24,918 coho salmon were counted.

**Appendix Table 18.** Aerial survey counts of chinook salmon escapement, Ugashik District, 1980-2000.

Year	Ugashik River	Dog <sup>g</sup> Salmon River	King Salmon	Painter Creed	Pumice Creek	Old Creek	Total
1980	0 <sup>a</sup>		900	1,000			1,900
1981	18 <sup>a</sup>		50	300			368
1982	0 <sup>a</sup>		700	700			1,400
1983	50 <sup>a</sup>	1,635	525	635	1,800	660	5,305
1984	108 <sup>a</sup>	836	4,100	1,875	1,100	880	8,899
1985	150 <sup>b</sup>	560	4,600	410	930	410	7,060
1986	66 <sup>b</sup>	252	1,777	646	705	739	4,185
1987	54 <sup>a</sup>	751	981	1,051	1,602	1,155	5,594
1988	249 <sup>c</sup>	900	5,820	1,170	1,025	660	9,824
1989	226 <sup>bc</sup>	848	1,670	1,030	510	520	4,804
1990	67 <sup>ac</sup>	540	1,500	590	450	610	3,757
1991	131 <sup>ac</sup>	449	700	365	375	420	2,440
1992 <sup>d</sup>	260 <sup>ac</sup>	821	1,260	855	750	815	4,761
1993	188 <sup>ac</sup>	579	1,970	865	450	635	4,687
1994 <sup>d</sup>	233 <sup>ac</sup>	1,741	2,225	1,005	2,530	1,490	9,224
1995	149 <sup>ac</sup>	882	440	366	501	505	2,843
1996	76 <sup>ac</sup>	1,079	1,200	403	<sup>f</sup>	30 <sup>f</sup>	2,788
1997	839 <sup>ac</sup>	906	802	525	536	558	4,166
1998	458 <sup>ac</sup>	1,411	883	1,230	352	438	4,772
1999	237 <sup>ac</sup>	535	<sup>f</sup>	166	340	213	1,491
Average	178	866	1,690	759	872	632	4,513
2000	26 <sup>a</sup>	425	<sup>f</sup>	314	339	246	1,350
Deviation <sup>e</sup>		-51%		-59%	-61%	-61%	-70%

<sup>a</sup> Tower counts

<sup>b</sup> Tower count plus later aerial survey counts of main river.

<sup>c</sup> Survey included Grassy Creek (tributary downstream of Ugashik Lagoon).

<sup>d</sup> Helicopter surveys.

<sup>e</sup> 2000 deviation from 1980-2000 average.

<sup>f</sup> No Count.

<sup>g</sup> Includes Figure-Eight, Goblet, Oldham, and Wandering Creeks.

**Appendix Table 19.** Aerial survey counts of chum salmon escapement, Ugashik District, 1980-2000.

Year	Ugashik River	Dog <sup>i</sup> Salmon River	King Salmon River	Painter Creek	Pumice Creek	Old Creek	Other	Total
1980	18 <sup>a</sup>		7,000	3,000				10,018
1981	0 <sup>a</sup>		200					200
1982	12 <sup>a</sup>		19,000	35,000			650	54,662
1983	0 <sup>a</sup>	1,650	2,700	4,000	20,000 <sup>b</sup>	3,300		31,650
1984	132 <sup>a</sup>	750	119,000	16,000	16,000	14,500	2,500	168,882
1985	42 <sup>c</sup>	350	20,000	1,925	6,000	670	300	29,287
1986	0 <sup>c</sup>	120	8,650	1,200	2,000	630	125	12,725
1987	130 <sup>c</sup>	340	9,750	2,290	10,340	2,090	40	24,980
1988	752 <sup>c,d</sup>	2,290	25,000	10,500	11,650	5,800	950	56,942
1989	600 <sup>c,d</sup>	1,005	7,500	3,700	2,200	2,010	625	17,640
1990	312 <sup>c,d</sup>	170	6,200	1,150	1,630	410	10	9,882
1991	315 <sup>c,d</sup>	240	7,400	750	2,550	2,525	130	13,910
1992 <sup>e</sup>	510 <sup>a,c,d</sup>	1,210	8,525	4,000	14,000	15,000	0	43,245
1993	93 <sup>c,d</sup>	105	7,000	720	2,040	1,025	8	10,991
1994 <sup>e</sup>	66 <sup>a,c</sup>	851	9,150	1,625	12,750	6,975	150	31,567
1995	6 <sup>a,c</sup>	160	3,900	1,370	2,600	1,800	0	9,836
1996	138 <sup>a</sup>	85	16,500	700	7,400	2,500	0	27,323
1997	100 <sup>a,c</sup>	450	10,500	4,200	5,300	9,480	115	30,145
1998	607 <sup>a,c</sup>	840	10,600	3,800	2,000	4,350	224	22,421
1999	278 <sup>a,c</sup>	400	<sup>h</sup>	650	1,660	2,020	50	5,058
Average	206	648	15,714	5,083	7,066	4,417	346	30,568 <sup>f</sup>
2000	7 <sup>a</sup>	510	<sup>h</sup>	2,150	7,300	5,850		15,817
Deviation <sup>g</sup>	-97%	-21%		-58%	3%	32%		-48%

<sup>a</sup> Tower counts

<sup>b</sup> Float count done from a raft.

<sup>c</sup> Survey included Grassy Creek (tributary downstream of Ugashik Lagoon).

<sup>d</sup> Included tower count plus later aerial survey count.

<sup>e</sup> Helicopter surveys.

<sup>f</sup> Average of the sums of indices for all locations.

<sup>g</sup> 2000 deviation from 1980-1999 average.

<sup>h</sup> No Count.

<sup>i</sup> Includes Figure-Eight, Goblet, Oldham, and Wandering Creeks.

**Appendix Table 20.** Aerial survey counts of pink salmon escapement, Ugashik District, 1980-2000.

Year	Number of Surveys <sup>a</sup>	Pink Salmon Count	Comments
1980	1	2,000	
1982	1	6,000	4,000 in King Salmon River, 2,000 in Painter Creek.
1983	2	803	Survey of Dog Salmon River conducted by USFWS.
1984	3	656	650 counted in King Salmon River during September 21 float trip.
1985	3	0	
1986	1	350	Observed in King Salmon River on August 19.
1987	2	1	
1988	7	2,800	Peak count on August 23: 2,000 in King Salmon River.
1989	8	50	Observed in Ugashik River on August 9.
1990	5	2,000	Peak count on August 13.
1991	0	660	Ugashik River tower count.
1992 <sup>b</sup>	0	1,728	Ugashik River tower count.
1993	0	0	
1994 <sup>b</sup>	0	425	Observed near Ugashik Lake Outlet on August 11.
1995	0	36	Ugashik River tower count.
1996	0	550	Observed in King Salmon River on August 12.
1997	0	0	
1998	0	57	Ugashik River tower count.
1999	0	6	Ugashik River tower count.
2000	0	46	Ugashik River tower count.

<sup>a</sup> Zero indicates no surveys designated to look for pink salmon and any observations recorded would be incidental to surveying for other species.

<sup>b</sup> Helicopter survey.

**Appendix Table 21.** Aerial survey counts of coho salmon escapement, Ugashik District, 1981-2000.

Year	Number of Surveys	Coho Salmon Counts	Comments
1981	1	13,300	Surveyed on September 7.
1982	1	10,000	Surveyed on August 26.
1983	0		
1984	1	6,100	Surveyed on August 31.
1985	2	18,880	16,500 in King Salmon River on September 12.
1986	2	8,455	Surveyed on August 19 and 25.
1987	2	17,000	16,700 in King Salmon River on August 23.
1988	7	28,280	12,900 in King Salmon River on September 7.
1989	4	11,515	7,615 observed on August 14.
1990	5	12,610	
1991	0	400	Incidental observation made August 12.
1992 <sup>a</sup>	0	790	Incidental observation made August 11.
1993	0	705	Incidental observation made August 16.
1994 <sup>a</sup>	0	760	Incidental observation made August 11.
1995	0		
1996 <sup>b</sup>	1	8,275	Surveyed on September 27 and 28.
1997 <sup>b</sup>	2	9,400	Surveyed on September 30 and October 17.
1998 <sup>b</sup>	1	1,459	Surveyed on November 19.
1999 <sup>b</sup>	1	10,210	Surveyed on October 14.
2000 <sup>b</sup>	1	12,070	Surveyed on October 12.

<sup>a</sup> Helicopter survey.

<sup>b</sup> Surveys are of selected areas in the Ugashik Lakes, King Salmon and Dog Salmon River drainages.

**Appendix Table 22.** Spawner distribution and total escapement estimates of sockeye salmon, Wood River system, 1959-2000.

Year	Spawner Distribution (%)			Total Escapement <sup>a</sup>
	Creeks	Beaches	Rivers	
1959	32.8	50.3	16.9	2,209,300
1960	27.4	55.5	17.1	1,016,100
1961	11.4	32.3	56.3	460,700
1962	24.0	65.2	10.8	873,900
1963	12.1	68.5	19.4	721,400
1964	18.9	64.0	17.1	1,076,100
1965	40.6	11.1	48.3	675,100
1966	16.4	54.9	28.7	1,208,700
1967	9.3	66.2	24.5	515,800
1968	9.9	50.8	39.3	649,300
1969	8.6	42.4	49.0	604,300
1970	14.0	52.4	33.6	1,162,000
1971	11.2	56.8	32.0	851,200
1972	17.4	45.1	37.5	430,600
1973	11.5	23.9	64.6	330,500
1974	14.1	63.9	22.0	1,708,800
1975	14.5	34.4	51.1	1,270,100
1976	12.7	33.5	53.8	817,000
1977	11.3	39.5	49.2	561,800
1978	14.2	51.3	34.5	2,267,200
1979	7.3	60.4	32.3	1,706,400
1980	20.8	24.5	54.7	2,969,000
1981	23.0	20.7	56.3	1,233,000
1982	14.0	17.2	68.8	976,400
1983	14.3	60.9	24.8	1,361,000
1984	11.4	27.6	61.0	1,002,800
1985	18.6	22.2	59.1	939,000
1986	16.1	23.3	60.6	819,000
1987	27.6	56.1	16.3	1,337,000
1988	31.0	44.4	24.6	866,800
1989	19.6	28.9	51.5	1,186,400
1990				1,069,400
1991			19.0	1,159,900
1992	24.9	56.7	18.4	1,286,300
1993	40.9	34.1	25.0	1,176,100
1994	25.5	36.4	38.1	1,471,900
1995	33.5	52.9	13.6	1,482,200
1996	25.8	39.3	34.9	1,649,600
1997	15.6	60.8	23.6	1,512,400
1998	20.0	66.2	13.8	1,755,800
1999				1,512,400
Mean	19.0	44.6	36.0	1,167,871
2000				1,300,000

<sup>a</sup> Estimated from Wood River tower counts. Rounded to the nearest hundred.

**Appendix Table 23.** Total escapement estimates of pink salmon, Nushagak and Togiak Districts, 1962-2000.<sup>a</sup>

Year	Nushagak District <sup>b</sup>	Togiak District <sup>c</sup>
1962	543,000	
1964	910,560	
1974	585,520	8,620 <sup>f</sup>
1976	863,430	37,570
1978	9,386,480	150,000 <sup>f</sup>
1980	2,785,200	102,820
1982	1,656,660	44,300
1984	2,926,450	269,950
1986	72,190 <sup>d</sup>	80,000 <sup>f</sup>
1988	494,610 <sup>d</sup>	142,500 <sup>f</sup>
1990	801,730 <sup>d</sup>	207,000
1992	<sup>e</sup>	235,000 <sup>f</sup>
1994	192,780 <sup>d</sup>	88,000 <sup>f</sup>
1996	821,312 <sup>d</sup>	<sup>e</sup>
1998	132,400 <sup>d</sup>	134,780 <sup>f</sup>
2000	135,285 <sup>d</sup>	<sup>e</sup>
Mean	1,487,174	125,045
2000	135,285 <sup>d</sup>	<sup>e</sup>

<sup>a</sup> Only those years of comprehensive aerial coverage are included: even years only; all counts rounded to the nearest 10 fish.

<sup>b</sup> Includes Wood, Igushik, Snake, Nushagak, and Nuyakuk Rivers, and Ice, Youth, and Sunshine Creeks, unless otherwise noted.

<sup>c</sup> Includes Togiak, Matogak and Osviak Rivers; 1982, 1990 and 1998 also Include Slug River.

<sup>d</sup> Sonar estimate of Nushagak-Mulchatna Rivers only.

<sup>e</sup> No escapement estimate.

<sup>f</sup> Togiak River estimate only.

**Appendix Table 24.** Aerial estimates of sockeye salmon escapements, Togiak District, 1980 - 2000.<sup>a</sup>

Year	Togiak River & Tributaries <sup>b</sup>	Kulukak Systems <sup>c</sup>
1980	50,700	45,700
1981	39,700	58,800
1982	25,300	52,800
1983	13,200	27,000
1984	30,900	49,800
1985	8,800	36,600
1986	35,000	42,800
1987	28,600	37,800
1988	32,400	31,700
1989	19,800	10,800
1990	47,100	49,600
1991	23,700	23,900
1992	16,500	26,400
1993	15,900	31,800
1994	19,400	29,700
1995	25,500	14,600
1996	30,200	19,000
1997	20,600	8,000
1998	21,900	13,000
1999	40,200	12,300
1980-99 Mean (20-Year)	27,270	31,105
1980-89 Mean (10-Year)	28,440	39,380
1990-99 Mean (10-Year)	26,100	22,830
2000	40,275	22,350

<sup>a</sup> All counts are rounded to the nearest hundred.

<sup>b</sup> Estimates do not include fish spawning above the counting tower (Togiak Lake outlet); estimates for Ungalikthluk, Osviak,. Matogak and Slug Rivers are not included in the 1977-94 data as reported in Bristol Bay Data Reports 73 and 81.

<sup>c</sup> Includes Kulukak River, Kulukak Lake, and Tithe Creek Ponds.

**Appendix Table 25.** Peak aerial counts of live sockeye salmon, Togiak River drainage, 1980 - 2000.

Year	Togiak Mainstem	Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuk River	Total
1980	18,600	3,200	2,500	500	3,200	2,000	30,000
1981	14,100	2,700	3,150			3,400	23,350
1982	2,300	3,600	2,500	0	100	4,800	13,300
1983	4,800	1,100	700	0	0	1,200	7,800
1984	10,550	2,800	2,450	0	0	2,300	18,100
1985	1,800	400	500	0	0	1,700	4,400
1986	13,500						13,500
1987	5,200	3,600	600	0	0	4,900	14,300
1988	9,400	2,000	1,100	0	0	3,700	16,200
1989	7,600	1,500	630			150	9,880
1990	8,770	5,720	5,980	0	2,550	1,190	24,210
1991	7,990	1,640	1,220			1,010	11,860
1992	3,030	1,280	1,400			2,200	7,910
1993	2,300	1,270	540			2,950	7,060
1994	3,100	560	1,870			3,900	9,430
1995	3,260	1,745	1,000		4,200	2,330	12,535
1996	9,160	2,270	150	100	240	3,190	15,110
1997	8,200	1,600	450	50	650	2,800	13,750
1998	4,890	3,100	150	10	0	2,800	10,950
1999	5,400	11,275	1,475	100	75	6,700	25,025
Mean 1980-1999	7,198	2,703	1,493	63	847	2,801	15,105 <sup>a</sup>
%	47.6%	17.9%	9.9%	0.4%	5.6%	18.5%	100.0%
2000	12,600	8,100	925	150	100	775	22,650

<sup>a</sup> Sum of means for all streams.

**Appendix Table 26.** Peak aerial counts of live sockeye salmon, Togiak District, 1980-2000.

Year	Togiak River <sup>a</sup>	Kulukak River <sup>b</sup>	Tithe Creek Ponds	Quigmy River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1980	30,000	12,200	14,200		500	200	1,900			63,500
1981	23,350	15,700	18,250		700	6,400	5,900	3,900	12,800 <sup>c</sup>	74,200
1982	13,300	11,900	19,300		0	1,000	5,500	300	2,400	53,700
1983	7,800	8,430	2,720		80	20	2,000	230	940	22,220
1984	18,100	7,400	14,000		200	6,800		100	5,200	51,800
1985	4,400	6,700	11,600		0	200	2,300	260	1,310	26,770
1986	13,500	10,900	14,000							38,400
1987	14,300	10,500	8,400							33,200
1988	16,200	12,600	3,250	250	100	380	5,880	200	2,700	41,560
1989	9,880	2,920	2,500					5,000		20,300
1990	24,210	10,600	14,200	100	400	2,200	3,540	9,700	3,800	68,750
1991	11,860	8,650	3,320	35	860	2,530	560	3,400	2,650	33,865
1992	7,910	7,530	4,950	40	300	3,340	1,460	3,600	3,760	32,890
1993	7,060	9,600	6,300					3,100	5,680	31,740
1994	9,430	10,270	4,600	580	990	1,750	6,070	2,230	3,240	39,160
1995	12,535	3,000	4,310	200	610	1,470	2,820	390	1,720	27,055
1996	15,110	2,490	7,000 <sup>d</sup>		360	780	1,045	1,000 <sup>d</sup>		27,785 <sup>d</sup>
1997	13,750	2,300	3,000		360	780	1,045	1,000		22,235
1998	10,950	2,175	4,300	20	900	2,600	5,010	2,300	240	28,495
1999	25,025	3,250	3,200	1,100	2,400	750	1,400	1,625	625	39,375
Mean	14,434	7,956	8,170	258	548	1,950	3,095	2,255	2,448	41,113 <sup>e</sup>
%	35.1%	19.4%	19.9%	0.6%	1.3%	4.7%	7.5%	5.5%	6.0%	100.0%
2000	22,650	6,100	5,075	125	526	1,512	1,201	2,175	575	39,939

<sup>a</sup> Includes all surveyed sections of Togiak River proper and all tributaries to the Togiak River.

<sup>b</sup> Includes surveys of Kulukak Lake. Counts prior to 1977 include Kulukak Lake only and are not included in the mean.

<sup>c</sup> Includes a combined count for the Negukthlik and Ungalikthluk of 4,500 fish.

<sup>d</sup> Complete count not available

<sup>e</sup> Sum of means for all streams.

Appendix Table 27. Peak aerial counts of live chinook salmon, Togiak River drainage, 1980-2000.

Year	Togiak River Section <sup>a</sup>						Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuk River	Total
	A	B	C	D	E	F						
1980	180	120	340	230	120	140	910	200	510	170	190	3,110
1981	420	390	500	200	300	740	980	310	370	390	290	4,890
1982					80	320	470	170	190	130	470	1,830
1983	120	220	370	290	360	850	820	240	340	430	350	4,390
1984	250	560	900	560	820	1,920	760	580	270	580	430	7,630
1985	270	320	640	340	470	970	470	250	290	310	460	4,790
1986	150	80	160	30	110	350						880
1987	20	70	170	120	200	480	610	180	100	120	320	2,390
1988	70	70	160	160	170	710	390	180	60	70	90	2,130
1989	10	30	370			940	190	80			40	1,660
1990	230	170	680	365	805	1,085	370	125	75	400	10	4,315
1991	505	165	475	225	520	455	460	105	90	100	150	3,250
1992	150	250	440	225	450	690	250	160	70	175	105	2,965
1993	170	120	220	160		1,810 <sup>b</sup>	595	240	130	65	440	3,950
1994				215	815	1,580	420	215	225	570	380	4,420
1995	120	220	750	255	800	800	715	140	425	520	295	5,040
1996	75	150	160	100	255	625	335	120	120	235	325	2,500
1997	100	350	1,300	600	820	1,000	275	180	150	275	100	5,150
1998	10	20	250	50	400	1,200	400	150	275	140	275	3,170
1999	150	210	540	510	225	480	365	90	240	305	270	3,385
Mean 1980-1999	167	195	468	258	429	857	515	196	218	277	263	3,842 <sup>c</sup>
%	4.3%	5.1%	12.2%	6.7%	11.2%	22.3%	13.4%	5.1%	5.7%	7.2%	6.8%	100.0%
2000	75	50	500	400	850	1450	350	85	125	100	75	4,060

<sup>a</sup> Section A; Togiak Bay - Gechiak River  
Section B; Gechiak River - Pungokepuk River  
Section C; Pungokepuk River - Nayorurun River  
Section D; Nayorurun River - Kemuk River  
Section E; Kemuk River - Ongivinuck River  
Section F; Ongivinuck River - Togiak Lake

<sup>b</sup> Includes count for Section E.

<sup>c</sup> Sum of means for all streams.

**Appendix Table 28.** Peak aerial counts of live chinook salmon, Togiak District, 1980-2000.

Year	Togiak River <sup>a</sup>	Quigmy River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1980	3,110	0	700	70	40		260	160	4,340
1981	4,890	0	1,290	470	1,730	350	1,460	180	10,370
1982	1,830	90	1,690	290	320		1,600	280	6,100
1983	4,390	40	2,460	190	120		1,080	260	8,540
1984	7,630	30	1,190	150	360		680	20	10,060
1985	4,790	0	540	100	50		80	90	5,650
1986	880								880
1987	2,390		300	30	40		660	80	3,500
1988	2,130	10	490	0	40	0	650	170	3,490
1989	1,660		740				560		2,960
1990	4,315	30	635	75	60	0	930	25	6,070
1991	3,250	25	285	75	100		1,175	55	4,965
1992	2,965	15	485	40	105	30	490	35	4,165
1993	3,950		1,140	80	110	100	830	70	6,280
1994	4,420	20	835	40	60	10	540	190	6,115
1995	5,040	35	430	65	135	50	740	80	6,575
1996	2,500	35	698	35	71	30	402	<sup>b</sup>	3,771
1997	5,150	10	310	50	65	33		10	5,628
1998	3,170	45	375	92	58	39	75	25	3,879
1999	3,385	10	240	105	40	150	345	130	4,405
Mean 1980-1999	3,592	25	781	109	195	72	698	103	5,574 <sup>c</sup>
%	64.4%	0.4%	14.0%	2.0%	3.5%	1.3%	12.5%	1.9%	100.0%
2000	4,060	26	340	65	42	6	1,100	226	5,865

<sup>a</sup> Includes all surveyed sections of Togiak River proper and all tributaries to the Togiak River.

<sup>b</sup> Complete count not available.

<sup>c</sup> Sum of means for all streams.

**Appendix Table 29.** Peak aerial counts of live chum salmon, Togiak River drainage, 1980-2000.

Year	Togiak River Section <sup>a</sup>						Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuk River	Total
	A	B	C	D	E	F						
1980	41,300	11,000	9,200	900	6,000	3,100	10,200	2,700	10,100	800	3,500	98,800
1981	11,800	4,500	2,400	1,000	3,000	6,000	3,100	500	4,300	1,700	4,200	42,500
1982				200	1,200	2,500	500	400	1,300	100	1,000	7,200
1983	8,160	3,050	3,780	1,100	2,780	6,070	150	140	5,560	570	3,790	35,150
1984	3,900	6,300	800	0	2,600	6,400	3,700	2,000	4,200	700	3,500	34,100
1985	8,300	6,500	3,200	900	6,700	10,200	4,100	600	9,600	1,800	8,300	60,200
1986 <sup>b</sup>												
1987	12,000	9,400	2,700	500	13,200	33,000	2,600	1,200	4,100	700	13,100	92,500
1988	10,000				4,900	3,800	3,700	5,000	3,500	200	3,800	34,900
1989		2,600	2,100		5,000	8,100	290	700			1,200	19,990
1990	2,200	1,275	1,350	400	650	4,200	3,150	1,150	3,400	250	125	18,150
1991	10,200	3,900	2,800	600	5,500	6,000	2,300	500	3,500	800	3,480	39,580
1992 <sup>c</sup>	1,800	1,800	300	100	1,200	1,500	2,000	500	1,800	900	800	22,700 <sup>d</sup>
1993	6,500	3,500	2,300	60		4,400	1,950 <sup>e</sup>	450	4,380	620	3,500	27,660
1994				1,300	5,200	10,400	900	2,400	7,100	900	5,700	33,900
1995	15,700	7,100	4,700	1,800	6,800	5,900	4,800	1,900	9,700	2,700	8,200	69,300
1996	3,700	10,250	5,500	1,300	5,750	8,250	2,600	750	900	550	3,400	42,950
1997	3,900	3,100	3,800	2,750	7,100	4,550	3,200	800	4,750	1,800	3,900	39,650
1998	2,300	1,400	2,750	1,300	4,300	8,950	3,600	1,050	3,000	250	1,650	30,550
1999	3,975	1,950	2,375	1,300	1,725	2,200	1,840	440	4,230	480	2,540	23,055
Mean	9,596	5,055	3,317	963	4,847	7,446	0	2,927	1,260	4,919	878	45,367 <sup>f</sup>
%	21.2%	11.1%	7.3%	2.1%	10.7%	16.4%		6.5%	2.8%	10.8%	1.9%	100.0%
2000 <sup>b</sup>												

<sup>a</sup> Section A; Togiak Bay - Gechiak River  
 Section B; Gechiak River - Pungokepuk River  
 Section C; Pungokepuk River - Nayorurun River  
 Section D; Nayorurun River - Kemuk River  
 Section E; Kemuk River - Ongivinuck River  
 Section F; Ongivinuck River - Togiak Lake

<sup>b</sup> No aerial surveys conducted.  
<sup>c</sup> Counts by section are not representative due to post-peak survey, and are not included in the mean.  
<sup>d</sup> Preferred total estimate; management survey count conducted 7/15/92.  
<sup>e</sup> Includes count for Section E.  
<sup>f</sup> Sum of means for all streams.

**Appendix Table 30.** Peak aerial counts of live chum salmon, Togiak District, 1980-2000.

Year	Togiak River <sup>a</sup>	Quigmy River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1980	98,800	2,700	27,300	5,700	29,500	6,700	18,500	15,000	204,200
1981	42,500	10,800	11,200	21,700	53,000	3,900	3,800	14,600	161,500
1982	7,200	1,300	8,300	3,100	5,500	2,400	160	1,270	29,230
1983	35,150	4,900	12,960	7,600	11,900	1,210	300	7,360	81,380
1984	34,100	6,300	8,500	10,200	18,400		2,100	3,000	82,600
1985	60,200	1,800	7,800	2,860	5,460	8,800	130	14,650	101,700
1986 <sup>b</sup>									
1987	92,500	1,500	22,000	2,300	2,160				120,460
1988	34,900	10,800	35,000	12,000	17,400	7,600	400	11,300	129,400
1989	19,990	2,820	5,580	7,450	4,900		560		41,300
1990	18,150	555	5,550	1,475	2,300	3,650	750	1,300	33,730
1991	39,580	4,420	9,540	4,730	8,700		120	3,020	70,110
1992	22,700 <sup>c</sup>	600	4,800 <sup>c</sup>	4,400	7,100	1,700	100	4,000	45,400
1993	27,660		6,950	1,970	1,360	3,060	20	4,020	45,040
1994	33,900	890	10,700	1,630	2,000	4,360	230	1,090	54,800
1995	138,600	2,200	7,600	5,200	13,920	6,440	1,000	7,200	182,160
1996	42,950	960	7,560	560	810	2,670	40	<sup>d</sup>	55,550
1997	39,650	1,700	4,550	3,000	2,500	1,890	<sup>b</sup>	<sup>b</sup>	53,290
1998	30,550	2,630	2,700	4,980	3,870	1,060	150	1,300	47,240
1999	23,055	1,340	3,430	5,700	3,650	4,750	410	11,360	53,695
Mean 1980-1999	44,323	3,234	10,633	5,608	10,233	4,013	1,598	5,910	85,552 <sup>e</sup>
%	51.8%	3.8%	12.4%	6.6%	12.0%	4.7%	1.9%	6.9%	100.0%
2000	<sup>b</sup>	2,870	4,950	9,090	10,880	4,150	200	5,520	37,660

<sup>a</sup> Includes all surveyed sections of Togiak River proper and tributaries to the Togiak River.

<sup>b</sup> No aerial surveys conducted.

<sup>c</sup> Preferred estimate from a management survey due to post-peak spawning ground survey.

<sup>d</sup> Complete count not available.

<sup>e</sup> Sum of means for all streams.

Appendix Table 31. Peak aerial counts of live coho salmon, Togiak River drainage, 1980-2000.

Year	Togiak River Section <sup>a</sup>						Mainstem Total	Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuk River	Total
	A	B	C	D	E	F							
1980	3,620	1,010	1,740	1,270	5,080	1,860	14,580	3,460	760	1,310	860	740	21,710
1981	9,280	580	100	800	370	750	11,880	520	360	230	210	1,300	14,500
1982	2,200	1,500	150	100	1,400	1,700	7,050	1,930	1,740	510	200	11,870	23,300
1983 <sup>b</sup>													
1984	1,440	1,190	200	120	620	1,480	5,050	4,750	2,240	990	1,110	6,140	20,280
1985	800 <sup>c</sup>	660 <sup>c</sup>	110 <sup>c</sup>	70 <sup>c</sup>	150	820	2,610	1,340	750	40	80	6,250	11,070
1986			60	400	100	400	960					2,560	3,520
1987	340	500	250	200	240	530	2,060	1,020	70			1,060	4,210
1988	950	370		140	210	360	2,030	1,530				4,100	7,660
1989 <sup>b</sup>													
1990	1,650	390	400	0	540	660	3,640	920	450	260	130	1,730	7,130
1991 <sup>d</sup>	4,900	400	700	600	1,680	140	8,020					100	140 <sup>d</sup>
1992	4,420	1,120	1,180	540	2,940	3,080	13,280	5,240	1,440	780	1,500	4,460	26,700
1993 <sup>b</sup>													
1994 <sup>b</sup>								1,290 <sup>d</sup>	220 <sup>d</sup>	120 <sup>d</sup>	95 <sup>d</sup>	1,930	3,655
1995 <sup>b</sup>								1,450			200	1,180	2,830
1996	2,550	1,090	150	250	1,600	5,020	10,910	2,080	1,170	575	725	6,450	21,910
1997	600	200	400	100	400	1,800	3,500	1,000	650	350	475	900	6,875
1998	460	625	100	100	310	1,075	2,670	2,550	575	400	500	1,750	8,445
1999	250	75	50	25	100	75	575	275	35	100	25	175	1,185
Mean	2,390	694	399	314	1,049	1,317	6,163	1,957	805	472	470	3,100	12,967 <sup>e</sup>
%	18.43%	5.35%	3.08%	2.42%	8.09%	10.15%	47.53%	15.09%	6.21%	3.64%	3.62%	23.91%	100.00%

2000<sup>b</sup>

- a Section A; Togiak Bay - Gechiak River  
 Section B; Gechiak River - Pungokepuk River  
 Section C; Pungokepuk River - Nayorurun River  
 Section D; Nayorurun River - Kemuk River  
 Section E; Kemuk River - Ongivinuck River

- b No aerial surveys conducted.  
 c Proportional estimates based on 1984 data.  
 d Timing of aerial surveys did not coincide with the period of peak spawning activity, and therefore, counts were not included in the mean or percent.  
 e Sum of means for all streams.

Appendix Table 32. Peak aerial counts of live coho salmon, Togiak District, 1980-2000.

Year	Togiak River <sup>a</sup>	Quigmy River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1980	21,710		10,300						32,010
1981	14,500		3,790				100	840	19,230
1982	23,300		3,380						26,680
1983 <sup>b</sup>									0
1984	20,280		10,750	1,850	1,080	670			34,630
1985	9,430	200	7,790	610	420				18,450
1986	3,520								3,520
1987	4,210	30	910	440	120			130	5,840
1988	8,590	460	1,840	310	490	470	370	3,170	15,700
1989 <sup>b</sup>									0
1990	7,130	1,029	5,195	2,675	1,491	810		4,153	22,483
1991 <sup>c</sup>	140		4,200						4,340
1992	26,700		12,640						39,340
1993 <sup>b</sup>									0
1994 <sup>b</sup>									0
1995		855	1,185	1,392	1,080	1,149		5,196 <sup>d</sup>	10,857
1996	21,660	1,211	10,290	3,062	2,805	1,944	851	5,917	47,740
1997	6,875	325	1,675	150	1,046	1,397		1,690	13,158
1998	8,445	390	3,650	1,785	2,001	523		2,770	19,564
1999	1,185	169	500	220	213	117	95	450	2,949 <sup>e</sup>
Mean	12,681	519	5,278	1,249	1,075	885	354	2,702	17,178
%	73.8%	3.0%	30.7%	7.3%	6.3%	5.2%	2.1%	15.7%	100.0%
2000 <sup>b</sup>									0

<sup>a</sup> Includes all surveyed sections of Togiak River proper and tributaries to the Togiak River.

<sup>b</sup> No aerial surveys conducted.

<sup>c</sup> Timing of aerial surveys did not coincide with the period of peak spawning activity, and therefore, counts were not included in the mean or percent.

<sup>d</sup> Negukthlik and Ungalikthluk Rivers combined.

<sup>e</sup> Sum of means for all streams.

The Alaska Department of Fish and Game administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility, or if you desire further information please write to ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; U.S. Fish and Wildlife Service, 4040 N. Fairfax Drive, Suite 300 Webb, Arlington, VA 22203 or O.E.O., U.S. Department of the Interior, Washington DC 20240.

For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-6077, (TDD) 907-465-3646, or (FAX) 907-465-6078.