

**Draft Report for the 2020 Upper Cook Inlet Board of
Fish Meeting: Northern Cook Inlet Chinook Salmon
Marine Harvest Stock Composition, 2016–2017**

by

Adam St. Saviour

Andrew W. Barclay

and

Nick Logelin

February 2020

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code		all standard mathematical signs, symbols and abbreviations	
deciliter	dL		AAC		
gram	g	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H _A
hectare	ha			base of natural logarithm	<i>e</i>
kilogram	kg	all commonly accepted		catch per unit effort	CPUE
kilometer	km	professional titles	e.g., Dr., Ph.D., R.N., etc.	coefficient of variation	CV
liter	L			common test statistics	(F, t, χ^2 , etc.)
meter	m	at	@	confidence interval	CI
milliliter	mL	compass directions:		correlation coefficient (multiple)	R
millimeter	mm	east	E		
		north	N	correlation coefficient (simple)	r
Weights and measures (English)		south	S		cov
cubic feet per second	ft ³ /s	west	W	covariance	°
foot	ft			degree (angular)	df
gallon	gal	copyright	©	degrees of freedom	<i>E</i>
inch	in	corporate suffixes:		expected value	>
mile	mi	Company	Co.	greater than	≥
nautical mile	nmi	Corporation	Corp.	greater than or equal to	HPUE
ounce	oz	Incorporated	Inc.	harvest per unit effort	<
pound	lb	Limited	Ltd.	less than	≤
quart	qt	District of Columbia	D.C.	less than or equal to	ln
yard	yd	et alii (and others)	et al.	logarithm (natural)	log
		et cetera (and so forth)	etc.	logarithm (base 10)	log ₂ , etc.
Time and temperature		exempli gratia		logarithm (specify base)	'
day	d	(for example)	e.g.	minute (angular)	NS
degrees Celsius	°C	Federal Information Code	FIC	not significant	H ₀
degrees Fahrenheit	°F			null hypothesis	%
degrees kelvin	K	id est (that is)	i.e.	percent	P
hour	h	latitude or longitude	lat or long	probability	
minute	min	monetary symbols		probability of a type I error (rejection of the null hypothesis when true)	α
second	s	(U.S.)	\$, ¢	probability of a type II error (acceptance of the null hypothesis when false)	β
		months (tables and figures): first three letters	Jan.,...,Dec	second (angular)	"
Physics and chemistry		registered trademark	®	standard deviation	SD
all atomic symbols		trademark	™	standard error	SE
alternating current	AC	United States		variance	
ampere	A	(adjective)	U.S.	population	Var
calorie	cal	United States of America (noun)	USA	sample	var
direct current	DC	U.S.C.	United States Code		
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH	U.S. state	use two-letter abbreviations (e.g., AK, WA)		
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

REGIONAL INFORMATION REPORT 2A20-03

DRAFT REPORT FOR THE 2020 UPPER COOK INLET BOARD OF FISH MEETING: NORTHERN COOK INLET CHINOOK SALMON MARINE HARVEST STOCK COMPOSITION 2016–2017

by

Adam St. Saviour
Alaska Department of Fish and Game, Division of Sport Fish, Palmer

Andrew W. Barclay
Alaska Department of Fish and Game, Division of Commercial Fisheries, Gene Conservation Laboratory,
Anchorage

and

Nick Logelin
Alaska Department of Fish and Game, Division of Sport Fish, Palmer

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1565

February 2020

This investigation was financed by the Alaska Sustainable Salmon Fund project 44908 and by Pacific States Marine Fishery Commission (PSMFC) (grant no. 16-108G) and (1037B.15).

The Regional Information Report Series was established in 1987 and was redefined in 2007 to meet the Division of Commercial Fisheries regional need for publishing and archiving information such as project operational plans, area management plans, budgetary information, staff comments and opinions to Board of Fisheries proposals, interim or preliminary data and grant agency reports, special meeting or minor workshop results and other regional information not generally reported elsewhere. Reports in this series may contain raw data and preliminary results. Reports in this series receive varying degrees of regional, biometric and editorial review; information in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author or the Division of Commercial Fisheries if in doubt of the level of review or preliminary nature of the data reported. Regional Information Reports are available through the Alaska State Library and on the Internet at: <http://www.adfg.alaska.gov/sf/publications/>

*Adam St. Saviour,
Alaska Department of Fish and Game, Division of Sport Fish,
1800 Glenn Hwy, Palmer, AK 99645-6736, USA*

*Andrew W. Barclay
Alaska Department of Fish and Game, Division of Commercial Fisheries, Gene Conservation Laboratory
333 Raspberry Rd, Anchorage AK 99518-1599, USA*

and

*Nick Logelin,
Alaska Department of Fish and Game, Division of Sport Fish,
1800 Glenn Hwy, Palmer, AK 99645-6736, USA*

This document should be cited as follows:

St. Saviour, A., A. W. Barclay, and N. Logelin. 2020. Draft Report for the 2020 Upper Cook Inlet Board of Fish Meeting: Northern Cook Inlet Chinook salmon marine harvest stock composition, 2016–2017. Alaska Department of Fish and Game, Regional Information Report 2A20-03, Anchorage.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	i
LIST OF FIGURES	ii
LIST OF APPENDICES	ii
ABSTRACT	1
INTRODUCTION.....	1
Objectives.....	3
Primary Objectives.....	3
Secondary Objectives	3
METHODS	3
Sample Collection.....	3
Tissue Sampling.....	3
Age, Sex, and Length Sampling.....	4
Laboratory Analysis	4
Assaying Genotypes.....	4
Laboratory Failure Rates and Quality Control.....	5
Statistical Analysis	5
Data Retrieval and Quality Control.....	5
Stratification and Subsampling for Genetic Mixed-Stock Analysis	6
Genetic Baseline	6
Mixed-Stock Analysis Reporting Groups	6
Genetic Mixed-Stock Analysis.....	7
RESULTS	8
Sample Collection.....	8
Tissue Sampling.....	8
Age, Sex, and Length.....	8
Laboratory Analysis	9
Assaying Genotypes.....	9
Laboratory Failure Rates and Quality Control.....	9
Statistical Analysis	9
Data Retrieval and Quality Control.....	9
Subsampling for Genetic Mixed-Stock Analysis	9
Genetic Mixed-Stock Analysis.....	10
DISCUSSION	11
Representation of Harvests.....	11
Making Inferences Outside Study Years	11
Relative Size of Stocks and Unknown Hatchery Contribution.....	12
ACKNOWLEDGMENTS.....	13
REFERENCES CITED.....	14
TABLES	17

TABLE OF CONTENTS (Continued)

FIGURES	35
APPENDIX A: STOCK COMPOSITION AND STOCK-SPECIFIC HARVEST ESTIMATES OF FINE-SCALE TEMPORAL STRATA, 2016 COMMERCIAL FISHERIES.....	43
APPENDIX B: STOCK COMPOSITION AND STOCK-SPECIFIC HARVEST ESTIMATES OF FINE-SCALE TEMPORAL STRATA, 2016 TYONEK SUBSISTENCE FISHERY	47

LIST OF TABLES

Table	Page
1 Populations of Chinook salmon in the Upper Cook Inlet genetic baseline, including the sampling location, collection years, number of individuals included in baseline from each population (<i>n</i>), and reporting groups for mixed stock analysis of the Tyonek subsistence fishery and Northern District commercial fishery harvests.	18
2 Chinook salmon collection details by time and area used in genetic mixed-stock analysis (MSA) of northern Cook Inlet marine fisheries, 2016–2017.....	20
3 Age, sex, and length composition of Chinook salmon harvested in the entire Northern District commercial set gillnet fishery, 2016.	21
4 Age, sex, and length composition of Chinook salmon harvested in the entire Northern District commercial set gillnet fishery, 2017.	22
5 Age, sex, and length composition of Chinook salmon harvested in the Eastern Subdistrict of Northern District commercial set gillnet fishery, 2016.	23
6 Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (north) of Northern District commercial set gillnet fishery, 2016.	24
7 Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (south) of Northern District commercial set gillnet fishery, 2016.	25
8 Age, sex, and length composition of Chinook salmon harvested in the Eastern Subdistrict of Northern District commercial set gillnet fishery, 2017.	26
9 Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (north) of Northern District commercial set gillnet fishery, 2017.	27
10 Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (south) of Northern District commercial set gillnet fishery, 2017.	28
11 Age, sex, and length composition of Chinook salmon harvested in the Tyonek Subsistence set gillnet fishery, 2016.	29
12 Age, sex, and length composition of Chinook salmon harvested in the Tyonek subsistence set gillnet fishery, 2017.	30
13 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016.....	31
14 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2017.....	32
15 Annual stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017.....	33
16 Annual stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017....	34

LIST OF FIGURES

Figure	Page
1 Map of statistical areas for set gillnet commercial fishing in the Northern District of Upper Cook Inlet.	36
2 Sampling locations for Chinook salmon populations from Cook Inlet included in the Cook Inlet genetic baseline.	37
3 Graphical depictions of temporal and area strata (boxes) in the a) Northern District set gill net and b) Tyonek Subsistence set gillnet fisheries, in which stock composition and stock-specific harvest was estimated and c) keys for the colors indicating percentage of the harvest analyzed in each analysis, acronyms of each area stratum, and subscripts used to indicate temporal strata	38
4 Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017	39
5 Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017	40
6 Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017	41

LIST OF APPENDICES

Appendix	Page
A1 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Trading Bay and Tyonek statistical areas of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016.	44
A2 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (north) area of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016.	45
A3 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Eastern Subdistrict area of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016	46
B1 Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016	48

ABSTRACT

This is a draft report for the 2020 Upper Cook Inlet Board of Fisheries meeting that will be superseded by a future Fisheries Data Series report undergoing peer review. Genetic tissue samples were collected from Chinook salmon harvested in the Northern District commercial set gillnet fishery and the Tyonek subsistence fishery during 2016–2017 to determine stock composition of marine harvests in Northern (Upper) Cook Inlet. Samples represented 100% of commercial harvests and 98% of subsistence harvests in both 2016 and 2017. Genetic mixed-stock analysis produced stock composition and stock-specific harvest estimates for each fishery by 6 reporting groups: 1) *West*, 2) *Susitna*, 3) *Deshka*, 4) *Yentna*, 5) *Knik-Turnagain*, and 6) *Kenai Peninsula*. For the General Subdistrict (south), harvest contributions were 34–36% *Susitna* in both 2016 and 2017, 24% *West* in 2016, followed by 19% *Yentna* and 14% *Deshka*. In 2017, harvest contributions were similar among *West* (19%), *Yentna* (17%), and *Deshka* (18%). For the General Subdistrict (north), harvest contributions were most (66–72%) from *Knik-Turnagain* in both years. For the Eastern Subdistrict, harvest contributions in both years were greatest from *Knik-Turnagain* (48–60%) followed by *Susitna* (21–23%). Overall Northern District commercial harvest contributions in 2016 and 2017 were greatest from *Knik-Turnagain* (32–35%), followed by *Susitna* (27–28%). Remaining harvest contributions were similar among *West* (13–14%), *Deshka* (11–14%), and *Yentna* (12–13%). In the 2016 Tyonek subsistence fishery, the largest contributions to harvest were from *Susitna* (32%) and *West* (29%) followed by *Yentna* (17%), *Deshka* (12%), and *Knik-Turnagain* (10%). In 2017, Tyonek subsistence harvest was dominated by *Susitna* (38%) followed by *West* (17%), *Yentna* (18%), and *Knik-Turnagain* (16%); *Deshka* contributed 10%. In 2016 and 2017, harvest contributions of the *Kenai Peninsula* reporting group ranged from 0–2% across all 3 commercial areas and across commercial and subsistence fisheries.

Key words: Chinook salmon, *Oncorhynchus tshawytscha*, genetic tissue, mixed-stock analysis, Northern District of Upper Cook Inlet, set gillnet, subsistence, Tyonek

INTRODUCTION

Chinook salmon (*Oncorhynchus tshawytscha*) returning to Cook Inlet, Alaska streams are harvested in the mixed-stock marine fisheries of Northern Cook Inlet (NCI), primarily the Tyonek subsistence set gillnet and Northern District commercial set gillnet fisheries (Figure 1). There are 7 Chinook salmon stocks of concern that are susceptible to harvest in NCI marine fisheries (Chuitna, Theodore, and Lewis rivers, and Alexander, Willow, Goose, and Sheep creeks). The *Northern District King Salmon Management Plan* (5 AAC 21.366) was created by the Alaska Board of Fisheries in 1986 and was most recently modified in 2014. The plan provides direction to the Alaska Department of Fish and Game (ADF&G) regarding management of the Northern District for the directed commercial harvest of Chinook salmon. The directed Chinook salmon commercial fishing season opens on the first Monday on or after May 25 and remains open for all Mondays through June 24. Fishing periods are 12 hours per day from 7:00 AM to 7:00 PM unless altered by emergency order. The commercial fishery is limited to an annual harvest not to exceed 12,500 Chinook salmon. Each permit holder is allowed to fish one 35-fathom, maximum 6-inch mesh set gillnet, with a minimum separation of 1,200 feet between nets.

Under the *Northern District King Salmon Management Plan*, if sport fishing restrictions are imposed on Theodore, Ivan, Lewis, Chuitna, or Deshka rivers, time and area restrictions will be placed on the commercial fishery. The Northern District set gillnet fishery has been closed from the wood chip dock, adjacent to Tyonek, to the Susitna River due to a sport fishing closure on the Chuitna River since 2011 (Shields and Frothingham 2018). Following the directed Chinook salmon fishery, the set gillnet fishery in the Northern District opens by regulation on or after June 25 for regular Monday and Thursday 12-hour periods to target other salmon species. This fishery is managed primarily by the *Northern District Salmon Management Plan* (5 AAC 21.358).

Under the statewide subsistence fishery regulations for Upper Cook Inlet (5 AAC 01.560), the Tyonek subsistence fishery is open for 2 seasons per year. The early season, May 15 through June

15, is open 3 days per week (Tuesday, Thursday, and Friday) for 16 hours per day (4:00 AM to 8:00 PM). The late season, June 16 through October 15, is open for 1 day per week (Saturday) for 12 hours (7:00 AM to 7:00 PM). Subsistence fishing targeting Chinook salmon occurs from May 15 until approximately June 30. The fishery extends from a point 1 mile south of the southern edge of the Chuitna River to the easternmost tip of Granite Point (Figure 1).

Genetic mixed-stock analysis (MSA) has been used to estimate the stock composition of sockeye salmon in the Cook Inlet commercial fishery since the 1990s (Seeb et al. 2000; Habicht et al. 2007; Barclay et al. 2010a, 2010b, 2013, 2014). With the development of comprehensive genetic baselines for Upper Cook Inlet Chinook salmon (Barclay et al. 2012; Barclay and Habicht 2015), MSA has more recently been used to estimate the stock composition of Chinook salmon harvested in the Upper Subdistrict set gillnet fishery (Eskelin 2013; Eskelin and Barclay 2015, 2016) and the Cook Inlet marine sport fishery (Barclay et al. 2016).

For the first time, stock composition estimates of the Tyonek subsistence and Northern District commercial set gillnet Chinook salmon fisheries were produced from harvest samples collected during 2014 and 2015 using the Barclay and Habicht (2015) baseline (St. Saviour et al. 2019). Four groups of populations (reporting groups) were chosen for MSA:

- 1) *UCI Northwest* (West Cook Inlet and Yentna River populations)
- 2) *Susitna-Matanuska* (Susitna River mainstem and Matanuska River populations)
- 3) *Knik-Turnagain* (Knik Arm and Turnagain Arm populations)
- 4) *Kenai Peninsula* (Kenai Peninsula populations from the Kenai River south to the Anchor River)

Although the estimates for the 4 reporting groups represented the finest-scale possible and the best available information at the time, estimates were needed for Yentna River, West Cook Inlet, Susitna River, and Deshka River to more precisely manage the sport and commercial fisheries of NCI. The inability of the Barclay and Habicht (2015) baseline to distinguish fine-scale NCI Chinook salmon reporting groups prompted an effort to improve the baseline with additional populations and genetic markers.

In 2016, the Pacific States Marine Fisheries Commission (PSMFC) appropriated Cook Inlet disaster relief funding for a collaborative study between the University of Washington and ADF&G with the primary objective of increasing the genetic differentiation of Chinook salmon reporting groups in the Cook Inlet baseline. This was done by developing a higher resolution baseline with hundreds of genetic markers and a subset of key populations. In addition to the Cook Inlet disaster relief funding, the previous MSA project (St. Saviour et al. 2019; Alaska Sustainable Salmon Fund project 44908) was leveraged by using remaining project funds to genotype additional baseline populations and the 2016 Tyonek subsistence and Northern District commercial samples for the same set of single nucleotide polymorphisms (SNPs). The genotypic data from both projects was combined and analyzed to produce a new baseline of 67 populations genotyped for 413 SNPs with the capability of distinguishing finer-scale reporting groups in NCI (Barclay et al. 2019). This new baseline made it possible to produce the fine-scale reporting group estimates for this PSMFC project (16-107G). All methods used to produce estimates are detailed below.

Here we present a draft report of the MSA results of Chinook salmon harvested in the Tyonek subsistence and Northern District commercial marine fisheries in 2016–2017 for the 2020 Upper

Cook Inlet Board of Fisheries meeting. This report will be superseded by a future Fisheries Data Series report that will undergo regional and interregional peer review.

OBJECTIVES

Primary Objectives

- 1) Estimate the proportion of Chinook salmon harvested in the Tyonek subsistence fishery and Northern District set gillnet commercial fishery by reporting group (*West, Susitna, Deshka, Yentna, Knik-Turnagain*, and *Kenai Peninsula*) for each geographic stratum such that the estimated proportions are within 10 percentage points of the true values 90% of the time.
- 2) Estimate the age composition of Chinook salmon harvested by the Tyonek subsistence and Northern District set gillnet commercial fisheries such that the estimates are within 10 percentage points of the true values 95% of the time.

Secondary Objectives

- 1) Sample a minimum of 100 Chinook salmon per stratum harvested in the Northern District commercial set gillnet fishery and the Tyonek subsistence fishery for tissue, age, sex, and length in proportion to reported harvest.
- 2) Estimate the number of Chinook salmon harvested by the Tyonek subsistence and Northern District set gillnet commercial fisheries by reporting group (*West, Susitna, Deshka, Yentna, Knik-Turnagain*, and *Kenai Peninsula*) for each geographic stratum.
- 3) Estimate the sex and length compositions of Chinook salmon harvested in the Tyonek subsistence and Northern District set gillnet commercial fisheries.

METHODS

SAMPLE COLLECTION

Genetic tissue samples and age, sex, and length data were collected from Chinook salmon harvested in the Northern District set gillnet commercial fishery and the Tyonek subsistence fishery in 2016 and 2017 to estimate stock composition of marine harvest in NCI. Samples were collected at ports, processors, buying stations, and set gillnet sites in Anchorage, Tyonek, and the Soldotna area during and after fishery openings. Crews were directed to maximize the number of samples collected to meet sample size requirements. Target sample sizes were a minimum of 100 fish per spatiotemporal stratum within each fishery in proportion to reported harvest.

Similar to previous years, 4 targeted Chinook salmon commercial fishing periods occurred in 2017 during May 29–June 19; each was intensively sampled. The first 2 “regular” commercial fishing periods (June 26 and 29) that target primarily sockeye salmon but incidentally harvest a small number of Chinook salmon were opportunistically sampled.

Tissue Sampling

Tissue samples were preserved for DNA analysis by stapling them onto numbered Whatman¹ (GE Healthcare Life Sciences) paper cards in numbered grid locations; cards were then placed in an

¹ Product names used in this publication are included for completeness but do not constitute product endorsement.

airtight case with desiccant beads to preserve samples. The Whatman paper card and grid numbers were recorded on data sheets. Genetic tissues were sent to the ADF&G Gene Conservation Laboratory (GCL) for long-term storage and genetic analysis.

Age, Sex, and Length Sampling

Sampled fish were measured from mid eye to tail fork (METF) to the nearest 5 mm. Sex was determined by external physical characteristics, such as kype development (males) or a protruding ovipositor (females). When sex could not be determined by external characteristics, samplers would request permission to make a small incision from the vent forward and sex was determined by inspection of gonads.

During both commercial and subsistence sampling, 4 scales were removed from the preferred area of each fish and placed on an adhesive-coated card (Clutter and Whitesel 1956; Welander 1940). Impressions from scales mounted on the adhesive-coated cards were made in cellulose acetate as described in Clutter and Whitesel (1956) and Scarnecchia (1979). The impressions were magnified 40×, viewed on a microfiche reader, and the ages were determined from growth patterns of the circuli. Ages were recorded in European notation (Jearld Jr. 1983). All age data were recorded on the same data sheets described in the previous “Tissue Sampling” section and entered into electronic spreadsheets.

LABORATORY ANALYSIS

Assaying Genotypes

Genomic DNA was extracted from tissue samples using a NucleoSpin 96 Tissue Kit by Macherey-Nagel (Düren, Germany).

Samples were genotyped for 588 amplicons as reported in Barclay et al. (2019). These amplicons were partitioned into 2 panels hereafter referred to as the UW (289 amplicons) and CRITFC (299 amplicons) panels. Of the 289 UW amplicons, 230 contained 1 SNP locus and 59 contained 2–4 SNP loci (366 SNPs total) (Dann et al. 2018; McKinney et al. 2019). All 299 CRITFC amplicons contained 1 SNP locus (Janowitz-Koch et al. 2019). For each panel, sequencing followed the GT-seq methods described in Campbell et al. (2014) other than deviations as follows. During PCR2, the volume was increased to 2 µL of 10 µM well-specific i5 tag primers per well, bringing the final reaction volume to 11 µL. During the purification step with magnetic beads the final elution volume was increased to 17 µL and no additional TE (pH 8.0 with 1% TWEEN 20) was added. The quantification by qPCR was completed using triplicate dilutions of 1:1000, 1:5000, and 1:10000. Four microliters of each dilution was used as template in a 10 µL reaction using the 6 µL Kapa Library Quantification Kit - Illumina/ROX Low (Kapa Biosystems, Wilmington, MA). The qPCRs were performed in 384-well plates on a QuantStudio 12K Flex Real-Time PCR System (Life Technologies). Final dilutions of each plate library were normalized to 4 nM. The final pooled library went through an additional purification step via magnetic beads. This involved adding 46.4 µL of Agencourt AMPure XP magnetic beads to 58 µL of pooled library in a 1.5 mL tube. The tube was incubated at room temperature for 7 minutes, placed in a magnetic stand for 5 minutes, and the supernatant was discarded. A double wash of 80% ETOH was performed, each for 30 seconds. The tube incubated at room temperature for 5 minutes to evaporate residual ETOH. The elution was performed with 30 µL of 1X Low-EDTA TE, pH 8.0, and incubated for 5 minutes before final transfer to a new 1.5 mL tube. The elution product was quantified for DNA yield via the manufacturer’s direction for the Qubit 3.0 (Thermo Fisher Scientific). The final pooled library

was sequenced at a final concentration of 3.5 pM on an Illumina NextSeq 500 with single-end read flow cells using 150 cycles.

Locus genotypes (single SNPs or microhaplotypes) for each sample were called using the GTscore software (https://github.com/csjaalbert/GTscore_ADFG) with 1 modification (likelihood threshold $pval < 0.001$; McKinney et al. 2019). Single SNPs not conforming to expected allelic ratios (e.g., polyploid or off-target amplification) were removed before allele calls were assigned. Alleles from multiple SNPs known to be linked (i.e., on the same amplicon), were combined to form microhaplotype loci. Genotypes were imported and archived in the GCL Oracle database, LOKI. From this point forward, each single SNP or microhaplotype is referred to as a locus.

Laboratory Failure Rates and Quality Control

QC analyses were conducted to identify laboratory errors and to measure the background discrepancy rate of the genotyping process. Separate QC methods were used for 2016 samples that had previously been genotyped using TaqMan SNP assays and the 2017 samples that had never been genotyped previously.

The QC protocol for the 2016 samples consisted of comparing old TaqMan SNP genotypes (old genotypes) in the database with the new GT-seq genotypes (new genotypes) for the same loci and individuals. Inconsistencies between the old and new genotypes were checked for laboratory errors, laboratory errors were corrected, and the old genotypes were replaced with the new genotypes.

The QC protocol for the 2017 samples consisted of re-extracting 8% of project fish and genotyping them following the GT-seq genotyping protocol above. Laboratory errors found during the QC process were corrected, and genotypes were corrected in the database. Inconsistencies not attributable to laboratory error were recorded, but original genotype scores were retained in the database.

For all genotyped samples, the overall failure rate was calculated by dividing the number of failed single-locus genotypes by the number of assayed single-locus genotypes. Discrepancy rates were calculated for the 2017 samples as the number of conflicting genotypes divided by the total number of genotypes compared. Assuming that discrepancies were due equally to errors during both genotyping events (original and QC), and that these analyses are unbiased, the error rate in the genotyping was estimated as half the overall rate of discrepancies. This QC method is the best representation of the error rate of the GCL's current genotype production.

STATISTICAL ANALYSIS

Data Retrieval and Quality Control

Genotypes were retrieved from LOKI and imported into R^2 with the *RJDBC* package³. Only the final set of 413 loci used in the Barclay et al. (2019) baseline were imported into R . All subsequent analyses were performed in R , unless otherwise noted. Prior to statistical analysis, 2 analyses were performed to confirm the quality of the data. First, the 80% rule (missing data at 20% or more of loci; Dann et al. 2009) was used to identify individuals missing substantial genotypic data. These

² R Development Core Team. 2016. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

³ Urbanek, S. 2014. *RJDBC*: Provides access to databases through the JDBC interface. *R* package version 0.2-5. <http://CRAN.R-project.org/package=RJDBC>.

individuals were removed from further analyses. The inclusion of individuals with poor quality DNA might introduce genotyping errors and reduce the accuracy of the MSA.

The final QC analysis identified individuals with duplicate genotypes and removed them from further analyses. Duplicate genotypes, which can occur by sampling or extracting the same individual twice, were defined as pairs of individuals sharing the same alleles in 95% of screened loci. The sample with the most missing genotypic data from each duplicate pair was removed from further analyses. If both samples had the same amount of genotypic data, the first sample was removed from further analyses.

Stratification and Subsampling for Genetic Mixed-Stock Analysis

The commercial fishery was stratified into 3 geographic areas: 1) General Subdistrict (south)⁴, including stat areas 247-10 (Trading Bay) and 247-20 (Tyonek); 2) General Subdistrict (north), including stat areas 247-41 (Susitna Flats), 247-42 (Pt. McKenzie), and 247-43 (Fire Island); and 3) Eastern Subdistrict, including stat areas 247-70 (Pt. Possession), 247-80 (Birch Hill), and 247-90 (#3 Bay) (Figure 1).

In 2016, finer-scale temporal and geographic strata were used to produce MSA mixtures. This resolution was unnecessary for fisheries management and in 2017 mixtures were at the subdistrict and annual scale.

The subsistence fishery for Chinook salmon only occurs in 1 geographic area so only 1 geographic stratum from the subsistence fishery was analyzed each year. In 2016, subsistence samples were collected by ADF&G staff and 2 local research assistants from May 17 to June 25. In 2017, subsistence samples were collected from May 16 to June 24. The vast majority of samples were collected during peak harvests May 24–June 14, 2016 and May 26–June 15, 2017.

Genetic Baseline

The genetic baseline used in this analysis was derived from nearly 7,800 samples collected from Chinook salmon spawning locations throughout Cook Inlet. The baseline consisted of 413 genetic markers and 67 Cook Inlet populations, with a minimum sample size of 40 fish per population (Table 1 and Figure 2; Barclay et al. 2019). A Cook Inlet–only baseline was chosen because marine harvests in NCI are believed to contain only fish of Cook Inlet origin.

Mixed-Stock Analysis Reporting Groups

Six reporting groups of interest to management were chosen for the MSA in this study (Table 1 and Figure 2). These groups are as follows:

- 1) *West* (Western Cook Inlet and Alexander Creek populations)
- 2) *Yentna* (Yentna River populations)
- 3) *Susitna* (Susitna River populations)
- 4) *Deshka* (the Deshka River population)
- 5) *Knik-Turnagain* (Cook Inlet populations from Turnagain Arm and Knik Arm)
- 6) *Kenai Peninsula* (Kenai Peninsula populations from the Kenai River south to the Anchor River)

⁴ Stat area 247-30 (Beluga) is closed to commercial fishing for Chinook salmon.

The first 5 reporting groups were all tested in the Barclay et al. (2019) baseline study and were found to be sufficiently identifiable for MSA. Although the 6th reporting group was not tested as a single unit in the baseline study, the stocks from the Kenai Peninsula that make up this group (*Kenai tributary*, *Kenai mainstem*, *Kasilof tributary*, *Kasilof mainstem*, *South Kenai Peninsula*) were tested, and all stocks were found to be sufficiently identifiable for MSA.

Genetic Mixed-Stock Analysis

The stock composition of the geographically and temporally stratified commercial and subsistence fishery samples selected for MSA (mixtures) were estimated using the *R* package *rubias* (Moran and Anderson 2019). The *rubias* package is a Bayesian approach to the conditional genetic stock identification model based upon computationally efficient C code implemented in *R*. It uses cross-validation and simulation to quantify and correct for biases in reporting group estimates. Each mixture was analyzed for 1 Markov chain Monte Carlo (MCMC) chain with 25,000 iterations and the first 5,000 iterations were discarded to remove the influence of starting values. The prior parameters for each reporting group were defined to be equal (i.e., a flat prior). Within each reporting group, the population prior parameters were divided equally among the populations within that reporting group. Stock proportion estimates and the 90% credibility intervals for mixture were calculated by taking the mean and 5% and 95% quantiles of the posterior distribution from the single chain output. After the MCMC analysis, 100 parametric bootstrap simulations were performed to correct for biases in the stock proportion estimates.

Stock Specific Harvest Estimates

Estimates of stock specific harvest were derived by applying the stock composition proportions p_g to the fishery harvest H following methods of Habicht et al. (2012):

$$H_g = Hp_g \quad (1)$$

The estimate \hat{H}_g and distribution of stock specific harvest for each reporting group (g) were obtained by Monte Carlo simulation. Independent realizations of the reporting group-specific harvest $H_g^{(k)}$ were drawn randomly from the joint distribution of the harvest $H^{(k)}$ and stock composition $p_g^{(k)}$ for each fishery mixture (K observations):

$$H_g^{(k)} = H^{(k)}p_g^{(k)} \quad (2)$$

Descriptive statistics were estimated directly from the K observations of $H_g^{(k)}$ with the mean used as the estimate of stock specific harvest \hat{H}_g and the 5th and 95th quantiles determining the bounds of the 90% CI.

Generation of stock-specific catch distributions required an estimate of the distribution of each component. The distributions of the stock compositions $p_g^{(k)}$ were the Bayesian posterior distributions of stock proportions from the mixed stock analysis. The harvest $H^{(k)}$ from the Tyonek subsistence fishery was assumed to follow a lognormal distribution with the mean and SD supplied by the Division of Subsistence staff (D. S. Koster, Division of Subsistence, Research Analyst, ADF&G, Anchorage, personal communication).

Combining MSA Mixtures

Estimates from spatiotemporal mixtures in 2016 were combined into annual stock-specific harvest estimates for each commercial fishing subdistrict (General Subdistrict [south], General Subdistrict [north], Eastern Subdistrict) and the Tyonek subsistence fishery by weighting them by their respective harvests (Table 2) following the methods of Dann et al. (2009). These harvest estimates, including their upper and lower bounds, were divided by the total harvest from each area to derive the overall proportion and credibility interval of each reporting group in the harvest. This same method was used to combine mixture estimates from each area into annual stock-specific harvest estimates for the entire Northern District in 2016 and 2017.

Age, Sex, and Length Composition of Harvest

Age, sex, and length composition estimates were calculated using standard statistical techniques as detailed in Eskelin and Barclay (2016).

RESULTS

SAMPLE COLLECTION

Tissue Sampling

Sampling rates were high because in past years crews had learned the best times and locations to intercept fish and build relationships with processors and fisherman (St. Saviour et al. 2019). In 2016, a total of 1,458 samples were collected from the commercial fishery and 289 were collected from the subsistence fishery. In 2016, 68.2% of reported commercial harvest was sampled and 28.5% of estimated subsistence harvest was sampled (Table 2). The minimum sample size of 100 fish was met for all spatiotemporal strata and ranged from 128 to 267 samples. After laboratory QC procedures and subsampling for MSA, the lowest number of genetic samples used for an individual stratum was 125 for the General Subdistrict (North) (Table 2).

In 2017, a total of 1,786 samples were collected from the commercial fishery and 252 were collected from the subsistence fishery (Table 2). In 2017, 83.0% of reported commercial harvest was sampled and 19.3% of estimated subsistence harvest was sampled (calculated from Table 2). The minimum sample size of 100 fish was met for all spatiotemporal strata and ranged from 252 to 994 samples. After laboratory QC procedures and subsampling for MSA, the lowest number of genetic samples used for an individual stratum was 247 for the Tyonek subsistence fishery (Table 2).

Age, Sex, and Length

Commercial

In 2016, the estimated age composition of Chinook salmon sampled from the entire Northern District commercial set gillnet fishery was 1.4% age 1.1 (SE 0.2%), 34.1% age 1.2 (SE 1.1%), 56.9% age 1.3 (SE 1.2%), and 7.5% age 1.4 (SE 0.7%). The average length-at-age for each age group was as follows: 506 mm (SE 16 mm) for age 1.1, 623 mm (SE 2 mm) for age 1.2, 782 mm (SE 2 mm) for age 1.3, and 878 mm (SE 8 mm) for age 1.4. The estimated sex ratio of sampled Chinook salmon was 41.5% (SE 1.2%) female (Table 3).

In 2017, the estimated age composition of Chinook salmon sampled from the entire Northern District commercial set gillnet fishery was 0.2% age 1.0 (SE 0.1%), 4.7% age 1.1 (SE 0.4%),

25.8% age 1.2 (SE 0.8%), 51.3% age 1.3 (SE 0.9%), 17.8% age 1.4 (SE 0.7%), and 0.2% age 1.5 (SE 0.1%). The average length-at-age for each age group was as follows: 332 mm (SE 0 mm) for age 1.0, 519 mm (SE 8 mm) for age 1.1, 617 mm (SE 2 mm) for age 1.2, 797 mm (SE 2 mm) for age 1.3, 892 mm (SE 4 mm) for age 1.4, and 915 mm (SE 0 mm) for age 1.5. The estimated sex ratio of sampled Chinook salmon was 42.9% (SE 1.0%) female (Table 4).

Age, sex, and length estimates by area can be found in Tables 5–10.

Subsistence

In 2016, the estimated age composition of Chinook salmon sampled from the Tyonek subsistence set gillnet fishery was 5.2% age 1.1 (SE 1.4%), 18.0% age 1.2 (SE 2.4%), 63.0% age 1.3 (SE 3.0%), 13.3% age 1.4 (SE 2.1%), and 0.5% age 1.5 (SE 0.4%). The average length-at-age for each age group was as follows: 595 mm (SE 15 mm) for age 1.1, 645 mm (SE 10 mm) for age 1.2, 796 mm (SE 5 mm) for age 1.3, 862 mm (SE 19 mm) for age 1.4, and 900 mm (SE 0 mm) for age 1.5. The estimated sex ratio of sampled Chinook salmon was 53.1% (SE 3.1%) female (Table 11).

In 2017, the estimated age composition of Chinook salmon sampled from the Tyonek subsistence set gillnet fishery was 1.6% age 1.1 (SE 0.9%), 18.1% age 1.2 (SE 2.7%), 50.0% age 1.3 (SE 3.4%), 30.2% age 1.4 (SE 3.2%). The average length-at-age for each age group was as follows: 552 mm (SE 52 mm) for age 1.1, 618 mm (SE 11 mm) for age 1.2, 816 mm (SE 6 mm) for age 1.3, and 912 mm (SE 9 mm) for age 1.4. The estimated sex ratio of sampled Chinook salmon was 53.3% (SE 3.4%) female (Table 12).

LABORATORY ANALYSIS

Assaying Genotypes

Of the 3,785 fish sampled from the 2016 and 2017 Northern District commercial and Tyonek subsistence fisheries harvests, 2,940 fish (2,401 commercial and 539 subsistence fish) were genotyped for 665 SNP markers.

Laboratory Failure Rates and Quality Control

Genotyping failure rates among the commercial and subsistence collections ranged from 2.74% to 3.74%. For the 2016 samples, no inconsistencies were found that were attributable to laboratory errors. For the 2017 samples, discrepancy rates between original and QC analyses were 2.74% for commercial and 2.91% for subsistence. Assuming equal error rates in the original and the QC analyses, estimated error rate in the samples is half of the discrepancy rate (1.37% and 1.46%).

STATISTICAL ANALYSIS

Data Retrieval and Quality Control

Genotypic data from the 2016 and 2017 Northern District commercial and Tyonek subsistence fisheries harvest samples for the final 413 baseline markers was read into R. Thirty-four of the genotyped samples (1.16%) were removed from further analyses based upon the 80% rule. Seven samples were identified as duplicates and were removed from further analyses. After removing fish based on the 80% rule and duplicates, 2,899 fish were available to subsample for MSA.

Subsampling for Genetic Mixed-Stock Analysis

Of the 2,899 genotyped fish available (from 2016 and 2017), 2,702 were subsampled in proportion to harvest to create 13 mixtures for MSA: 10 commercial and 3 subsistence mixtures comprising

all annual and geographic strata (Table 2 and Figure 3). Enough samples ($n \geq 100$) were available to represent annual harvest by commercial fishing subdistrict (General Subdistrict [south], General Subdistrict [north], and Eastern Subdistrict) and fishery (Northern District commercial and Tyonek subsistence).

Genetic Mixed-Stock Analysis

MSA was performed on the 13 mixtures created during the subsampling process to produce stock composition and stock-specific harvest estimates (Tables 13–16; Appendices A1–A3 and B1). Stock composition and stock-specific harvests estimates were produced for finer-scale temporal strata (i.e., mixtures) for the 2016 Northern District commercial and Tyonek subsistence harvests (Table 2); however, for consistency with the 2017 estimates, only the fishery subdistrict and district scale estimates are reported in the results section of this report. The estimates for fine-scale 2016 mixtures can be found in Appendices A1–A3 and B1.

Figure 3 illustrates the levels of strata used in the MSA and how the fine-scale strata were combined into fishery subdistrict and district scale estimates using the stratified estimator. This figure also guides the reader to the tables containing results for each level of analysis. The 2016 MSA estimates from the fine-scale MSA mixtures were combined to produce annual stock composition and stock-specific harvest estimates for the General Subdistrict (south), General Subdistrict (north), Eastern Subdistrict, the entire Northern District commercial fishery, and the entire Tyonek subsistence fishery (Figure 3). The MSA estimates from the 3 commercial subdistrict MSA mixtures were combined to produce stock composition and stock-specific harvest estimates for the entire Northern District commercial fishery (Figure 3). The estimates reported in the following sections include these fishery subdistrict and district scale results.

Commercial Estimates by Subdistrict

In 2016 and 2017, annual Chinook salmon harvests in the General Subdistrict (south) area (Trading Bay and Tyonek commercial) were 1,150 and 1,188, respectively (Tables 2, 13, and 14). Stock composition estimates were greatest for the *Susitna* (point estimate range: 33.7–35.8%) reporting group. The *West* reporting group had the second highest harvest in 2016 at 24.4% with similar contributions by *Yentna* (18.6%) and *Deshka* (14.4%). In 2017, harvest contributions were similar among *West* (18.9%), *Yentna* (16.5%), and *Deshka* (18.4%). In 2016 and 2017, harvest contributions were 8.9–10.2% for the *Knik-Turnagain* reporting group (Tables 13 and 14; Figure 4).

In 2016 and 2017, annual Chinook salmon harvests in the General Subdistrict (north) were 464, and 545 respectively (Tables 2, 13, and 14). Harvests were dominated by the *Knik-Turnagain* reporting group (point estimate range: 65.9–71.5%; Tables 4 and 5, Figure 4). The remaining harvests were represented by smaller contributions of *Susitna* (range: 16.0–18.1%), *Deshka* (5.5–7.5%), *Yentna* (2.7–6.3%), and *West* (range: 2.3–4.3%; Tables 13 and 14; Figure 4).

In 2016 and 2017, annual Chinook salmon harvests in the Eastern Subdistrict were 523 and 418 respectively (Tables 2, 13, and 14). Harvest contributions were greatest from the *Knik-Turnagain* (point estimate range: 47.7–60.3%) reporting group, followed by *Susitna* (20.9–23.2%; Tables 4 and 5; Figure 4). Remaining harvest contributions were similar among *West* (2.3–9.7%), *Deshka* (7.9–9.9%), and *Yentna* groups (7.7–8.2%; Tables 13 and 14; Figure 4).

The *Kenai Peninsula* reporting group contributed less than 2% to the harvest in all 3 commercial areas in 2016 and 2017 (Tables 13 and 14; Figure 4).

Annual Northern District Commercial Estimates

In 2016 and 2017, overall annual Chinook salmon harvests in the Northern District commercial fishery were 2,137 and 2,151 respectively (Tables 2 and 15). In both years, *Knik-Turnagain* was the largest contributor to the harvest (point estimate range: 31.6–35.1%), followed by *Susitna* (range: 27.2–28.3%) reporting group. The remaining harvest was made up of similar contributions from *West* (13.4–14.3%), *Deshka* (10.9–14.0%), and *Yentna* (12.2–12.6%) reporting groups. The *Kenai Peninsula* reporting group contributed less than 1% to overall annual harvest in both years (Table 15; Figure 5).

Annual Tyonek Subsistence Estimates

In 2016 and 2017, estimated annual Chinook salmon harvests by Tyonek subsistence users during sampling periods were 1,013 and 1,304 fish respectively (Tables 2 and 16). In 2016, the *Susitna* (point estimate: 32.3%) and *West* (28.8%) reporting groups were the largest contributors to harvest (Table 7 and Figure 6). *Yentna* contributed 17.1%, followed by *Deshka* (11.6%) and *Knik-Turnagain* (9.9%). In 2017, *Susitna* was the dominant stock present in Tyonek subsistence harvest at 38.3%. *West* (16.5%), *Yentna* (17.9%), and *Knik-Turnagain* (16.0%) all contributed similarly, with *Deshka* (10.4%) contributing less. The *Kenai Peninsula* reporting group contributed less than 1% to annual harvest in both years (Table 16; Figure 6). Credibility intervals were greater for the Tyonek subsistence estimates than the commercial estimates because, in addition to genetic error, error from the subsistence harvest estimates was also incorporated into the estimates.

DISCUSSION

This report includes the MSA of harvest samples collected from the Northern District commercial and Tyonek subsistence fisheries in 2016 and 2017. These results represent the second MSA of Chinook salmon captured in NCI fisheries (St. Saviour et al. 2019).

REPRESENTATION OF HARVESTS

Only 17 out of 1,030 fish harvested in the 2016 Tyonek subsistence fishery could not be represented because they fell outside of the sampled date range or dates of harvest were unknown (D. Koster, ADF&G Subsistence Division, personal communication, September 20, 2019). All other harvests were fully represented in all strata analyzed in 2016 and 2017. Due to the small harvests of the Northern District commercial and Tyonek subsistence Chinook salmon fisheries relative to other commercial fisheries sampled by ADF&G, acquiring samples in proportion to daily harvests was sometimes difficult to achieve. To reach harvest-representative samples size goals, we allowed the use of surplus samples from 1 collection day to represent harvest on other days provided it was within 7 days and within the same stratum. The general stability in stock composition estimates over entire season gave us confidence that this procedure accurately represented harvest (St. Saviour et al. 2019).

MAKING INFERENCES OUTSIDE STUDY YEARS

These analyses are derived from samples collected during a specific period of time during particular environmental and fishery conditions. Nonetheless, this study can be used to inform future scientific and regulatory activities. Currently, these results are derived from the most

comprehensive data set available to examine stock composition of Chinook salmon captured in the Northern District commercial and Tyonek subsistence fisheries. Along with the previous study (St. Saviour et al. 2019), this 2-year data set provides some measure of interannual variability in stock compositions; however, some caution must be exercised when extrapolating these results to unanalyzed years because changes in relative abundance among reporting groups, prosecution of fisheries, or migratory behavior due to ocean conditions might affect the distribution of stock-specific harvests by time and area.

RELATIVE SIZE OF STOCKS AND UNKNOWN HATCHERY CONTRIBUTION

In this and the previous study (St. Saviour et al. 2019), there were some unexpected results in harvest by reporting group. The relative magnitude of stocks originating from Susitna River Basin including Susitna, Yentna, and Deshka rivers is thought to be considerably greater than those originating from Knik and Turnagain arms (Decovich et al. 2020, *In prep*; Oslund et al. 2017; Bosch 2010; Baumer and Blain-Roth 2020). Yet harvests of the *Knik-Turnagain* reporting group were similar to or greater than the *Susitna*, *Deshka*, or *Yentna* in 2016 and 2017 (Tables 5 and 6 and Figure 5).

Chinook salmon escapements to the Susitna, Deshka, and Yentna rivers have been monitored by relatively precise mark-recapture or weir projects in recent years (Decovich et al. *in prep*^{1,2}; Oslund et al. 2017). The Chinook salmon producing systems of the *Knik-Turnagain* reporting group are more numerous and much smaller in size. Escapements to most of these systems are monitored by relatively less-precise foot surveys or not at all (Baumer and Blain-Roth 2020). Additionally, unknown numbers of adult Chinook salmon of hatchery origin return to Ship Creek and the Eklutna Tailrace. These fish would have allocated to the *Knik-Turnagain* reporting group and likely contributed a substantial but unknown proportion of harvest.

Future Chinook salmon stock identification work in NCI should include increased escapement monitoring and precision of stocks in the *Knik-Turnagain* reporting group and identification and quantification of hatchery-produced Chinook salmon. Vegetative cover and glacial turbidity would likely interfere in some cases, but comprehensive aerial surveys may increase our understanding of the relative size of the *Knik-Turnagain* reporting group. The proportion of hatchery-produced Chinook salmon in NCI harvests could be estimated by collection of thermally marked otoliths or coded wire tags. Collecting all otoliths would likely be challenging for samplers due to time-constraints imposed by fisherman and market forces that give preference to fully intact Chinook salmon. Adipose fin clips administered to hatchery smolt would aid in identifying which fish to sample for otoliths or coded wire tags.

The updated baseline reported in Barclay et al. (*In prep*) now allows for the identification of hatchery stocks through parentage-based tagging techniques (Anderson and Garza 2006) as an alternative to the collection of coded-wire tags. This technique involves collecting samples from fishery harvests and hatchery broodstock from potential brood years contributing to the fishery and genotyping them. Genotypes from the broodstock samples (parents) could then be used to identify hatchery fish in a harvest sample by assigning hatchery fish to their parents through genetic parentage analysis. Genetic tissue samples are collected each year from hatchery brood stocks in Cook Inlet, and GCL has archived tissue samples from Cook Inlet hatchery brood stocks dating back to 2012.

ACKNOWLEDGMENTS

Funding for this project was provided by the Alaska Sustainable Salmon Fund and the Pacific States Marine Fishery Commission. The authors would like to thank the following Alaska Department of Fish and Game field staff and Tyonek local hires for their hard work in obtaining samples from the fisheries: Donnie Arthur, Phil Stacey, Paul Kuriscak, Jay Baumer, Bruce Whelan, Ben Cohen, Brittany Blain, Erica Chenowith, Katelyn Zoneville, Annalise Theisen, Madeline Fox, Amanda Alaniz, Capra Smith, Cody Watkins, Bronwyn Jones, Philip Martinez, Leonard Allowan, Gwen Chickalusion, and John Standifer. Thanks to commercial fisherman, subsistence fishers, and seafood processors for making catch samples available. The authors would also like to acknowledge the work of the people in the ADF&G's Gene Conservation Laboratory: Dan Prince, Chase Jalbert, Heather Hoyt, Nick Ellickson, Marie Filteau, Paul Kuriscak, Zach Pechacek, Mariel Terry, Erica Chenoweth, Zac Grauvogel, Judy Berger, Eric Lardizabal. Chase Jalbert provided logistical support, training for field crews, and a review of this manuscript. Tim McKinley, Tania Vincent, and an anonymous reviewer are also thanked for help editing and publishing this report.

REFERENCES CITED

- Anderson, E. C., and J. C. Garza. 2006. The power of single-nucleotide polymorphisms for large-scale parentage inference. *Genetics* 172(4):2567–2582.
- Barclay, A. W., D. F. Evenson, and C. Habicht. 2019. New Genetic Baseline for Upper Cook Inlet Chinook Salmon Allows for the Identification of More Stocks in Mixed-Stock Fisheries: 413 Loci and 67 Populations. Alaska Department of Fish and Game, Fishery Manuscript Series No. 19-06, Anchorage.
- Barclay, A. W., B. J. Failor, and C. Habicht. 2016. Report to the Alaska Board of Fisheries: Progress report on genetic and coded wire tag mixed stock analysis of Chinook salmon harvested in Cook Inlet marine sport fishery, 2014–2016. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J16-09, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/RIR.5J.2016.09.pdf>
- Barclay, A. W., and C. Habicht. 2015. Genetic baseline for Upper Cook Inlet Chinook salmon: 42 SNPs and 7,917 fish. Alaska Department of Fish and Game, Fishery Manuscript Series No. 15-01, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMS15-01.pdf>
- Barclay, A. W., C. Habicht, R. A. Merizon, and R. J. Yanusz. 2012. Genetic baseline for Upper Cook Inlet Chinook salmon: 46 SNPs and 5,279 fish. Alaska Department of Fish and Game, Fishery Manuscript Series No. 12-02, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FMS12-02>
- Barclay, A. W., C. Habicht, W. D. Templin, H. A. Hoyt, T. Tobias, and T. M. Willette. 2010a. Genetic stock identification of Upper Cook Inlet sockeye salmon harvest, 2005–2008. Alaska Department of Fish and Game, Fishery Manuscript No. 10–01, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FMS10-01.pdf>
- Barclay, A. W., C. Habicht, T. Tobias, E. L. Chenoweth, and T. M. Willette. 2014. Genetic stock identification of Upper Cook Inlet sockeye salmon harvest, 2011. Alaska Department of Fish and Game, Fishery Data Series No. 14-43, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS14-43.pdf>
- Barclay, A. W., C. Habicht, T. Tobias, and T. M. Willette. 2010b. Genetic stock identification of Upper Cook Inlet sockeye salmon harvest, 2009. Alaska Department of Fish and Game, Fishery Data Series No. 10–93, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FDS10-93.pdf>
- Barclay, A. W., C. Habicht, T. Tobias, and T. M. Willette. 2013. Genetic stock identification of Upper Cook Inlet sockeye salmon harvest, 2010. Alaska Department of Fish and Game, Fishery Data Series No. 13-56, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FDS13-56>
- Baumer, J., and B. J. Blain-Roth. 2020. Area management report for the sport fisheries of Anchorage, 2016–2018. Alaska Department of Fish and Game, Fishery Management Report No. 20-03, Anchorage.
- Bosch, D. 2010. Area management report for the recreational fisheries of Anchorage, 2009 and 2010. Alaska Department of Fish and Game, Fishery Management Report No. 10-53, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/Fmr10-53.pdf>
- Campbell, N. R., S. A. Harmon, and S. R. Narum. 2014. Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources* 15(4):855–867.
- Clutter, R., and L. Whitesel. 1956. Collection and interpretation of sockeye salmon scales. International Pacific Salmon Commission, Bulletin 9. Westminster, British Columbia, Canada.
- Dann, T. H., C. Habicht, J. R. Jasper, H. A. Hoyt, A. W. Barclay, W. D. Templin, T. T. Baker, F. W. West, and L. F. Fair. 2009. Genetic stock composition of the commercial harvest of sockeye salmon in Bristol Bay, Alaska, 2006–2008. Alaska Department of Fish and Game, Fishery Manuscript Series No. 09-06, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMS09-06.pdf>
- Dann, T. H., C. Habicht, W. D. Templin, L. W. Seeb, G. McKinney, and J. E. Seeb. 2018. Identification of genetic markers useful for mixed stock analysis of Chinook salmon in Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J18-04, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/RIR.5J.2018.04.pdf>

REFERENCES CITED (Continued)

- DeCovich, N.A., J. Campbell, S. Dotomain, and D. Evans. 2020. Susitna River Chinook salmon abundance and spawning distribution, 2017. Alaska Department of Fish and Game, Fishery Data Series 20-04, Anchorage.
- DeCovich, N.A., J. Campbell, S. Dotomain, and D. Evans. *In prep.* Susitna River Chinook salmon abundance and spawning distribution, 2016. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Eskelin, A., and A. W. Barclay. 2016. Mixed stock analysis and age, sex, and length composition of Chinook salmon in Upper Cook Inlet, Alaska, 2015. Alaska Department of Fish and Game, Fishery Data Series No. 16-16, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS16-16.pdf>
- Eskelin, T. 2013. Upper Cook Inlet Commercial Eastside set gillnet Chinook salmon sampling study. Alaska Department of Fish and Game, Division of Sport Fish, Regional Operational Plan ROP.SF.2A.2013.17, Soldotna. <http://www.adfg.alaska.gov/FedAidPDFs/ROP.SF.2A.2013.17.pdf>
- Eskelin, T., and A. W. Barclay. 2015. Mixed stock analysis and age, sex, and length composition of Chinook salmon in Upper Cook Inlet, Alaska, 2014. Alaska Department of Fish and Game, Fishery Data Series No. 15-19, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS15-19.pdf>
- Fall, James A., A. Godduhn, G. Halas, L. Hutchinson-Scarborough, B. Jones, B. McDavid, E. Mikow, L.A. Sill, A. Wiita, and T. Lemons. 2020. Alaska Subsistence and Personal Use Salmon Fisheries 2017 Annual Report. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 0-05, Anchorage.
- Habicht, C., J. R. Jasper, T. H. Dann, N. DeCovich, and W. D. Templin. 2012. Western Alaska Salmon Stock Identification Program Technical Document 11: Defining reporting groups. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J12-16, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/RIR.5J.2012.16>
- Habicht, C., W. D. Templin, T. M. Willette, L. F. Fair, S. W. Raborn, and L. W. Seeb. 2007. Post-season stock composition analysis of Upper Cook Inlet sockeye salmon harvest, 2005-2007. Alaska Department of Fish and Game, Fishery Manuscript No. 07-07, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/fms07-07.pdf>
- Janowitz-Koch, I., C. Rabe, R. Kinzer, D. Nelson, M. A. Hess, and S. R. Narum. 2019. Long-term evaluation of fitness and demographic effects of a Chinook Salmon supplementation program. *Evolutionary Applications* 12(3):456–469.
- Jearld Jr., A. 1983. Age determination. Pages 301-324 in L. A. Nielsen, editors. *Fisheries techniques*. The American Fisheries Society, Bethesda, Maryland
- Jones, B.E. and D. Koster. 2018. Subsistence harvests and uses of salmon in Tyonek, 2015 and 2016. ADF&G Division of Subsistence, Technical Paper No. 439.
- McKinney, G. J., C. E. Pascal, W. D. Templin, S. E. Gilk-Baumer, T. H. Dann, L. W. Seeb, and J. E. Seeb. 2019. Dense SNP panels resolve closely related Chinook salmon populations. *Canadian Journal of Fisheries and Aquatic Sciences*, Published on the web 9 August 2019, <https://doi.org/10.1139/cjfas-2019-0067>.
- Moran, B. M., and E. C. Anderson. 2019. Bayesian inference from the conditional genetic stock identification model. *Canadian Journal of Fisheries and Aquatic Sciences* 76(4):551–560.
- Oslund, S., S. Ivey, and D. Lescanec. 2017. Area management report for the recreational fisheries of northern Cook Inlet, 2014–2015. Alaska Department of Fish and Game, Fishery Management Report No. 17-07, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR17-07.pdf>
- Scamecchia, D. L. 1979. Variation of scale characteristics of coho salmon with sampling location on the body. *Progressive Fish Culturist* 41(3):132-135.
- Seeb, L. W., C. Habicht, W. D. Templin, K. E. Tarbox, R. Z. Davis, L. K. Brannian, and J. E. Seeb. 2000. Genetic diversity of sockeye salmon of Cook Inlet, Alaska, and its application to management of populations affected by the Exxon Valdez oil spill. *Transactions of the American Fisheries Society* 129(6):1223-1249.

REFERENCES CITED (Continued)

- Shields, P., and A. Frothingham. 2018. Upper Cook Inlet commercial fisheries annual management report, 2017. Alaska Department of Fish and Game, Fishery Management Report No. 18-10, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR18-10.pdf>
- St. Saviour, A., A. W. Barclay, and N. Logelin. 2019. Northern Cook Inlet Chinook salmon marine harvest stock composition, 2014–2015. Alaska Department of Fish and Game, Fishery Data Series No. 19-03, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS19-03.pdf>
- Welanders, A. D. 1940. A study of the development of the scale of Chinook salmon *Oncorhynchus tshawytscha*. Master's thesis. University of Washington, Seattle.

TABLES

Table 1.—Populations of Chinook salmon in the Upper Cook Inlet genetic baseline, including the sampling location, collection years, number of individuals included in baseline from each population (*n*), and reporting groups for mixed stock analysis of the Tyonek subsistence fishery and Northern District commercial fishery harvests.

Pop. no.	Reporting group	Geographic region	Location	Collection year(s)	<i>n</i>	
1	<i>West</i>	West Side Cook Inlet	Straight Creek	2010	93	
2			Nikolai Creek	2012,2013	81	
3			Chuitna River	2009	92	
4			Coal Creek	2009,2010,2011	119	
5			Theodore River	2010,2011,2012	119	
6			Lewis River	2011,2012,2014	96	
7			Sucker Creek	2011,2012	144	
8	<i>Susitna</i>	Susitna River	Cheechako Creek	2014	57	
9			Portage Creek	2009,2010,2011,2013	164	
10			Indian River	2013,2014	98	
11			Chulitna River east fork	2013,2014	97	
12			Chulitna River middle fork	2009,2010,2013	229	
13			Honolulu Creek	2013,2014	106	
14			Pass Creek	2013,2014	104	
15			Byers Creek	2013,2014	109	
16			Spink Creek	2013,2014	74	
17			Bunco Creek	2013	103	
18			Troublesome Creek	2013,2014	119	
19			Talkeetna River - no name #1	2013,2014	84	
20			Talkeetna River - no name #2	2013,2014	53	
21			Prairie Creek	2008,2013	142	
22			Iron Creek	2013,2014	102	
23			Disappointment Creek	2013,2014	133	
24			Chunilna Creek	2009,2012,2013	104	
25			Montana Creek	2009,2010	120	
26			Sheep Creek	2013,2014	60	
27			Kashwitna River	2013,2014	62	
28			Little Willow Creek	2013,2014	104	
29			Willow Creek	2005	70	
30			Deception Creek	2009	100	
31	<i>Deshka</i>	Yentna River	Deshka River	1995,2005,2012,2015	302	
32	<i>Yentna</i>		Red Creek	2012,2013	111	
33			Happy River	2012,2014	45	
34			Hayes River	2012,2013,2014	74	
35			Canyon Creek	2012,2013	91	
36			Talachulitna River	2008,2010	120	
37			Sunflower Creek	2009,2011	123	
39	<i>Knik-Turnagain</i>		Knik Arm	Little Susitna River	2009,2010	125
40				Granite Creek	2013,2014,2015	83
41				Moose Creek	1995,2009,2012	120

-continued-

Table 1.–Page 2 of 2.

Pop. no.	Reporting group	Geographic region	Location	Collection year(s)	<i>n</i>
42	<i>Knik-Turnagain</i>	Knik Arm	Eagle River	2009, 2011, 2012	78
43			Ship Creek	2009	172
44		Turnagain Arm	Campbell Creek	2011, 2012	96
45			Bird Creek	2009, 2011, 2015	83
46			Carmen River	2011, 2012, 2013	74
47			Resurrection Creek	2010, 2011, 2012	98
48			Chickaloon River	2008, 2010, 2011	74
49	<i>Kenai Peninsula</i>	Kenai River	Grant Creek	2011, 2012, 2013	87
50			Quartz Creek	2006, 2008, 2010, 2011	82
51			Crescent Creek	2006	165
52			Russian River	2005, 2006, 2007, 2008	211
53			Juneau Creek	2005, 2006, 2007	116
54			Kenai River - upper mainstem	2009	92
55			Benjamin Creek	2005	54
56			Killey River	2005, 2006	167
57			Funny River	2005, 2006	128
58			Kenai River - middle mainstem	2003, 2004, 2006	299
59			Kenai River - lower mainstem	2010, 2011	125
60			Slikok Creek	2004, 2008	81
61		Kasilof River	Kasilof River - middle mainstem	2005	190
62			Kasilof River - lower mainstem	2005	132
63		Coastal Kenai Peninsula	Crooked Creek	1992, 2005, 2011	305
64			Ninilchik River	2010, 2015	144
65			Deep Creek	2010	41
66			Stariski Creek	2011, 2012	106
67			Anchor River	2006, 2010	145

Note: Populations and reporting groups match those in Figure 1.

Table 2.—Chinook salmon collection details by time and area used in genetic mixed-stock analysis (MSA) of northern Cook Inlet marine fisheries, 2016–2017.

Year	Geographic area ^a	Date range	Harvest		Sampled		Analyzed for MSA	
			Reported	Estimated	Number	Proportion	Number	Proportion
2016	Trading Bay	May 29–June 30	581	-	202	0.347676	143	24.6%
	Tyonek commercial	May 29–June 12	346	-	198	0.572254	193	55.8%
	Tyonek commercial	June 13–30	223	-	212	0.950673	156	70.0%
	General Subdistrict (south)	May 29–June 30	1,150	-	612	53.2%	492	42.8%
	General Subdistrict (north)	May 29–June 12	166	-	128	77.1%	125	75.3%
	General Subdistrict (north)	June 13–27	298	-	267	89.6%	216	72.5%
	General Subdistrict (north)	May 29–June 27	464	-	395	85.1%	341	73.5%
	Eastern Subdistrict	May 29–June 12	257	-	196	76.3%	186	72.4%
	Eastern Subdistrict	June 13–30	266	-	255	95.9%	218	82.0%
	Eastern Subdistrict	May 29–June 30	523	-	451	86.2%	404	77.2%
	Commercial subtotals, averages	May 29–June 30	2,137		1,458	68.2%	1,237	57.9%
	Tyonek subsistence	May 16–31	362	412	140	34.0%	137	33.3%
	Tyonek subsistence	June 1–25	525	601	149	24.8%	142	23.6%
	Tyonek subsistence ^b	May 16–June 25	887	1,014	289	28.5%	279	27.5%
	Totals, averages		3,024		1,747	57.8%	1,516	86.8%
2017	General Subdistrict (south)	May 29–June 29	1,188	-	994	83.7%	423	35.6%
	General Subdistrict (north)	May 29–June 19	545	-	423	77.6%	257	47.2%
	Eastern Subdistrict	May 29–June 29	418	-	369	88.3%	259	62.0%
	Tyonek subsistence	May 16–June 24	825	1,304	252	19.3%	247	18.9%
	Totals, averages		3,455		2,038	59.0%	1,186	58.2%

Note: Gray highlighted rows indicate mixture samples analyzed in the *rubias* R package. Nonhighlighted rows are the overall totals for General Subdistrict (south), General Subdistrict (north), Eastern Subdistrict, and Tyonek subsistence in 2016.

^a Refers to the Northern District commercial fishery unless subsistence is explicitly stated.

^b Fifteen reported fish and 16 estimated fish were excluded from season totals reported in Jones and Koster (2018) and Fall et al. (2020) because they fell outside of the sampled date range or dates of harvest were unknown and therefore could not be accurately represented in MSA.

Table 3.—Age, sex, and length composition of Chinook salmon harvested in the entire Northern District commercial set gillnet fishery, 2016.

Sex	Parameter	Age class					All ages
		1.1	1.2	1.3	1.4	1.5	
Females							
	Sample size by age	0	39	332	37	0	408
	Age composition	0.0%	4.1%	33.2%	4.1%	0.0%	41.5%
	SE (%)	0.0%	0.5%	1.1%	0.5%	0.0%	1.2%
	Harvest by age	0	89	710	88	0	887
	SE (harvest by age)	0	11	24	11	0	25
	Mean length (mm METF)	0	646	789	878	0	783
	SE (mm)	0	6	3	10	0	4
Males							
	Sample size by age	17	334	243	30	0	624
	Age composition	1.4%	30.0%	23.7%	3.4%	0.0%	58.5%
	SE (%)	0.2%	1.0%	1.0%	0.5%	0.0%	1.2%
	Harvest by age	30	641	506	73	0	1250
	SE (harvest by age)	5	22	22	10	0	25
	Mean length (mm METF)	506	620	773	877	0	694
	SE (mm)	16	3	4	13	0	4
Both							
	Sample size by age	17	373	575	67	0	1,032
	Age composition	1.4%	34.1%	56.9%	7.5%	0.0%	100.0%
	SE (%)	0.2%	1.1%	1.2%	0.7%	0.0%	0.0%
	Harvest by age	30	730	1,217	161	0	2,137
	SE (harvest by age)	5	23	25	15	0	0
	Mean length (mm METF)	506	623	782	878	0	731
	SE (mm)	16	2	2	8	0	3

Table 4.—Age, sex, and length composition of Chinook salmon harvested in the entire Northern District commercial set gillnet fishery, 2017.

Sex	Parameter	Age class						All ages
		1.0	1.1	1.2	1.3	1.4	1.5	
Females								
	Sample size by age	0	1	48	382	119	1	551
	Age composition	0.0%	0.1%	4.0%	29.4%	9.4%	0.1%	42.9%
	SE (%)	0.0%	0.0%	0.4%	0.9%	0.6%	0.0%	1.0%
	Harvest by age	0	1	86	632	203	1	924
	SE (harvest by age)	0	1	9	19	13	1	21
	Mean length (mm METF)	0	590	640	793	886	1005	799
	SE (mm)	0	0	7	3	6	0	4
Males								
	Sample size by age	3	60	281	277	109	2	732
	Age composition	0.2%	4.7%	22.0%	21.6%	8.5%	0.1%	57.2%
	SE (%)	0.1%	0.4%	0.7%	0.7%	0.5%	0.1%	0.9%
	Harvest by age	5	100	473	466	184	3	1,231
	SE (harvest by age)	2	8	16	16	11	1	19
	Mean length (mm METF)	332	518	613	804	899	866	720
	SE (mm)	0	8	3	4	6	0	5
Both								
	Sample size by age	3	61	329	659	228	3	1283
	Age composition	0.2%	4.7%	25.8%	51.3%	17.8%	0.2%	100.0%
	SE (%)	0.1%	0.4%	0.8%	0.9%	0.7%	0.1%	0.0%
	Harvest by age	5	102	554	1,103	382	5	2,151
	SE (harvest by age)	2	8	17	19	15	2	0
	Mean length (mm METF)	332	519	617	797	892	915	754
	SE (mm)	0	8	2	2	4	0	3

Table 5.—Age, sex, and length composition of Chinook salmon harvested in the Eastern Subdistrict of Northern District commercial set gillnet fishery, 2016.

Sex	Parameter	Age class					All ages
		1.1	1.2	1.3	1.4	1.5	
Females							
	Sample size by age	0	9	105	7	0	121
	Age composition	0.0%	2.4%	28.4%	1.9%	0.0%	32.7%
	SE (%)	0.0%	0.4%	1.3%	0.4%	0.0%	1.3%
	Harvest by age	0	13	148	10	0	171
	SE (harvest by age)	0	2	7	2	0	7
	Mean length (mm METF)	0	673	787	911	0	785
	SE (mm)	0	7	4	8	0	5
Males							
	Sample size by age	10	154	81	4	0	249
	Age composition	2.7%	41.6%	21.9%	1.1%	0.0%	67.3%
	SE (%)	0.5%	1.4%	1.2%	0.3%	0.0%	1.3%
	Harvest by age	14	218	114	6	0	352
	SE (harvest by age)	2	7	6	2	0	7
	Mean length (mm METF)	489	616	773	896	0	667
	SE (mm)	23	3	5	48	0	6
Both sexes							
	Sample size by age	10	163	186	11	0	370
	Age composition	2.7%	44.1%	50.3%	3.0%	0.0%	100.0%
	SE (%)	0.5%	1.4%	1.4%	0.5%	0.0%	0.0%
	Harvest by age	14	230	263	16	0	523
	SE (harvest by age)	2	7	7	3	0	0
	Mean length (mm METF)	489	619	781	906	0	706
	SE (mm)	23	3	3	17	0	5

Table 6.—Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (north) of Northern District commercial set gillnet fishery, 2016.

Sex	Parameter	Age class					All ages
		1.1	1.2	1.3	1.4	1.5	
Females							
	Sample size by age	0	6	65	3	0	74
	Age composition	0.0%	2.6%	28.4%	1.3%	0.0%	32.3%
	SE (%)	0.0%	0.8%	2.1%	0.5%	0.0%	2.2%
	Harvest by age	0	12	132	6	0	150
	SE (harvest by age)	0	3	10	2	0	10
	Mean length (mm METF)	0	639	781	865	0	773
	SE (mm)	0	29	5	30	0	7
Males							
	Sample size by age	5	87	61	2	0	155
	Age composition	2.2%	38.0%	26.6%	0.9%	0.0%	67.7%
	SE (%)	0.7%	2.3%	2.1%	0.4%	0.0%	2.2%
	Harvest by age	10	176	124	4	0	314
	SE (harvest by age)	3	11	10	2	0	10
	Mean length (mm METF)	507	614	771	840	0	675
	SE (mm)	33	5	8	15	0	8
Both sexes							
	Sample size by age	5	93	126	5	0	229
	Age composition	2.2%	40.6%	55.0%	2.2%	0.0%	100.0%
	SE (%)	0.7%	2.3%	2.3%	0.7%	0.0%	0.0%
	Harvest by age	10	188	255	10	0	464
	SE (harvest by age)	3	11	11	3	0	0
	Mean length (mm METF)	507	616	776	855	0	707
	SE (mm)	33	5	5	18	0	7

Table 7.—Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (south) of Northern District commercial set gillnet fishery, 2016.

Sex	Parameter	Age class					All ages
		1.1	1.2	1.3	1.4	1.5	
Females							
	Sample size by age	0	24	162	27	0	213
	Age composition	0.0%	5.5%	37.4%	6.2%	0.0%	49.2%
	SE (%)	0.0%	0.9%	1.8%	0.9%	0.0%	1.9%
	Harvest by age	0	64	430	72	0	566
	SE (harvest by age)	0	10	21	11	0	22
	Mean length (mm METF)	0	642	792	875	0	785
	SE (mm)	0	6	4	12	0	5
Males							
	Sample size by age	2	93	101	24	0	220
	Age composition	0.5%	21.5%	23.3%	5.5%	0.0%	50.8%
	SE (%)	0.3%	1.6%	1.6%	0.9%	0.0%	1.9%
	Harvest by age	5	247	268	64	0	584
	SE (harvest by age)	3	18	18	10	0	22
	Mean length (mm METF)	550	627	774	878	0	721
	SE (mm)	10	5	7	14	0	7
Both sexes							
	Sample size by age	2	117	263	51	0	433
	Age composition	0.5%	27.0%	60.7%	11.8%	0.0%	100.0%
	SE (%)	0.3%	1.7%	1.9%	1.2%	0.0%	0.0%
	Harvest by age	5	311	698	135	0	1,150
	SE (harvest by age)	3	19	21	14	0	0
	Mean length (mm METF)	550	630	785	876	0	753
	SE (mm)	10	4	3	9	0	5

Table 8.—Age, sex, and length composition of Chinook salmon harvested in the Eastern Subdistrict of Northern District commercial set gillnet fishery, 2017.

Sex	Parameter	Age class						All ages
		1.0	1.1	1.2	1.3	1.4	1.5	
Females								
	Sample size by age	0	1	7	99	31	0	138
	Age composition	0.0%	0.4%	2.5%	34.9%	10.9%	0.0%	48.6%
	SE (%)	0.0%	0.2%	0.5%	1.6%	1.0%	0.0%	1.7%
	Harvest by age	0	1	10	146	46	0	203
	SE (harvest by age)	0	1	2	7	4	0	7
	Mean length (mm METF)	0	590	649	793	880	0	804
	SE (mm)	0	NA	5	5	10	0	6
Males								
	Sample size by age	1	14	53	57	20	1	146
	Age composition	0.4%	4.9%	18.7%	20.1%	7.0%	0.4%	51.4%
	SE (%)	0.2%	0.7%	1.3%	1.3%	0.9%	0.2%	1.7%
	Harvest by age	1	21	78	84	29	1	215
	SE (harvest by age)	1	3	5	6	4	1	7
	Mean length (mm METF)	300	502	610	802	880	880	711
	SE (mm)	0	15	7	9	14	0	12
Both sexes								
	Sample size by age	1	15	60	156	51	1	284
	Age composition	0.4%	5.3%	21.1%	54.9%	18.0%	0.4%	100.0%
	SE (%)	0.2%	0.8%	1.4%	1.7%	1.3%	0.2%	0.0%
	Harvest by age	1	22	88	230	75	1	418
	SE (harvest by age)	1	3	6	7	5	1	0
	Mean length (mm METF)	300	508	614	796	880	880	756
	SE (mm)	0	15	6	4	8	0	7

Table 9.—Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (north) of Northern District commercial set gillnet fishery, 2017.

Sex	Parameter	Age class						All ages
		1.0	1.1	1.2	1.3	1.4	1.5	
Females								
	Sample size by age	0	0	13	79	24	0	116
	Age composition	0.0%	0.0%	4.1%	24.8%	7.5%	0.0%	36.4%
	SE (%)	0.0%	0.0%	0.7%	1.6%	1.0%	0.0%	1.7%
	Harvest by age	0	0	22	135	41	0	198
	SE (harvest by age)	0	0	4	8	5	0	9
	Mean length (mm METF)	0	0	627	783	891	0	788
	SE (mm)	0	0	11	5	8	0	8
Males								
	Sample size by age	1	20	85	68	29	0	203
	Age composition	0.3%	6.3%	26.6%	21.3%	9.1%	0.0%	63.6%
	SE (%)	0.2%	0.9%	1.6%	1.5%	1.0%	0.0%	1.7%
	Harvest by age	2	34	145	116	50	0	347
	SE (harvest by age)	1	5	9	8	6	0	9
	Mean length (mm METF)	340	514	612	811	926	0	712
	SE (mm)	0	17	5	7	12	0	10
Both sexes								
	Sample size by age	1	20	98	147	53	0	319
	Age composition	0.3%	6.3%	30.7%	46.1%	16.6%	0.0%	100.0%
	SE (%)	0.2%	0.9%	1.7%	1.8%	1.3%	0.0%	0.0%
	Harvest by age	2	34	167	251	91	0	545
	SE (harvest by age)	1	5	9	10	7	0	0
	Mean length (mm METF)	340	514	614	796	910	0	740
	SE (mm)	0	17	4	5	8	0	7

Table 10.—Age, sex, and length composition of Chinook salmon harvested in the General Subdistrict (south) of Northern District commercial set gillnet fishery, 2017.

Sex	Parameter	Age class						All ages
		1.0	1.1	1.2	1.3	1.4	1.5	
Females								
	Sample size by age	0	0	28	204	64	1	297
	Age composition	0.0%	0.0%	4.1%	30.0%	9.4%	0.1%	43.7%
	SE (%)	0.0%	0.0%	0.5%	1.2%	0.7%	0.1%	1.2%
	Harvest by age	0	0	49	356	112	2	519
	SE (harvest by age)	0	0	6	14	9	1	15
	Mean length (mm METF)	0	0	645	796	885	1005	801
	SE (mm)	0	0	8	3	7	0	5
Males								
	Sample size by age	1	26	143	152	60	1	383
	Age composition	0.1%	3.8%	21.0%	22.4%	8.8%	0.1%	56.3%
	SE (%)	0.1%	0.5%	1.0%	1.0%	0.7%	0.1%	1.2%
	Harvest by age	2	45	250	266	105	2	669
	SE (harvest by age)	1	6	12	12	8	1	15
	Mean length (mm METF)	350	529	615	802	891	855	727
	SE (mm)	0	9	4	5	8	0	7
Both sexes								
	Sample size by age	1	26	171	356	124	2	680
	Age composition	0.1%	3.8%	25.1%	52.4%	18.2%	0.3%	100.0%
	SE (%)	0.1%	0.5%	1.1%	1.3%	1.0%	0.1%	0.0%
	Harvest by age	2	45	299	622	217	3	1,188
	SE (harvest by age)	1	6	13	15	12	2	0
	Mean length (mm METF)	350	529	620	799	888	930	759
	SE (mm)	0	9	3	3	5	0	4

Table 11.—Age, sex, and length composition of Chinook salmon harvested in the Tyonek Subsistence set gillnet fishery, 2016.

Sex	Parameter	Age class					All ages
		1.1	1.2	1.3	1.4	1.5	
Females							
	Sample size by age	2	12	78	19	1	112
	Age composition	0.9%	5.7%	37.0%	9.0%	0.5%	53.1%
	SE (%)	0.6%	1.4%	3.0%	1.8%	0.4%	3.1%
	Harvest by age	10	58	375	91	5	538
	SE (harvest by age)	7	16	35	20	5	38
	Mean length (mm METF)	670	667	797	842	900	790
	SE (mm)	10	20	6	26	0	8
Males							
	Sample size by age	9	26	55	9	0	99
	Age composition	4.3%	12.3%	26.1%	4.3%	0.0%	46.9%
	SE (%)	1.2%	2.0%	2.7%	1.2%	0.0%	3.1%
	Harvest by age	43	125	264	43	0	476
	SE (harvest by age)	14	23	31	14	0	37
	Mean length (mm METF)	579	635	794	906	0	743
	SE (mm)	13	11	10	16	0	12
Both							
	Sample size by age	11	38	133	28	1	211
	Age composition	5.2%	18.0%	63.0%	13.3%	0.5%	100.0%
	SE (%)	1.4%	2.4%	3.0%	2.1%	0.4%	0.0%
	Harvest by age	53	183	639	135	5	1,014
	SE (harvest by age)	15	27	38	24	5	30
	Mean length (mm METF)	595	645	796	862	900	768
	SE (mm)	15	10	5	19	0	7

Table 12.—Age, sex, and length composition of Chinook salmon harvested in the Tyonek subsistence set gillnet fishery, 2017.

Sex	Parameter	Age class						All ages
		1.0	1.1	1.2	1.3	1.4	1.5	
Females								
	Sample size by age	0	1	15	49	32	0	97
	Age composition	0.0%	0.5%	8.2%	26.9%	17.6%	0.0%	53.3%
	SE (%)	0.0%	0.5%	1.9%	3.1%	2.6%	0.0%	3.4%
	Harvest by age	0	7	107	351	229	0	695
	SE (harvest by age)	0	7	26	43	36	0	49
	Mean length (mm METF)	0	655	642	824	911	0	823
	SE (mm)	0	0	18	8	14	0	11
Males								
	Sample size by age	0	2	18	42	23	0	85
	Age composition	0.0%	1.1%	9.9%	23.1%	12.6%	0.0%	46.7%
	SE (%)	0.0%	0.7%	2.1%	2.9%	2.3%	0.0%	3.4%
	Harvest by age	0	14	129	301	165	0	609
	SE (harvest by age)	0	10	28	40	32	0	49
	Mean length (mm METF)	0	500	599	806	913	0	784
	SE (mm)	0	15	13	10	12	0	14
Both								
	Sample size by age	0	3	33	91	55	0	182
	Age composition	0.0%	1.6%	18.1%	50.0%	30.2%	0.0%	100.0%
	SE (%)	0.0%	0.9%	2.7%	3.4%	3.2%	0.0%	0.0%
	Harvest by age	0	21	236	652	394	0	1,304
	SE (harvest by age)	0	12	37	49	44	0	29
	Mean length (mm METF)	0	552	618	816	912	0	804
	SE (mm)	0	52	11	6	9	0	9

Table 13.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016. Sample sizes (*n*), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
General Subdistrict (south) <i>n</i> = 492								
	<i>West</i>	24.4	20.1	28.9	2.7	281	231	332
	<i>Susitna</i>	33.7	29.3	38.1	2.6	388	337	438
	<i>Deshka</i>	14.4	11.4	17.5	1.8	166	131	202
	<i>Yentna</i>	18.6	15.0	22.2	2.2	214	173	256
	<i>Knik-Turnagain</i>	8.9	6.4	11.5	1.6	102	73	132
	<i>Kenai Peninsula</i>	0.0	0.0	0.3	0.2	0	0	3
Harvest represented						1,150		
Harvest not represented						0		
Total harvest						1,150		
General Subdistrict (north) <i>n</i> = 341								
	<i>West</i>	2.3	0.3	4.8	1.4	11	1	22
	<i>Susitna</i>	18.1	14.6	21.8	2.2	84	68	101
	<i>Deshka</i>	5.5	3.5	8.0	1.4	25	16	37
	<i>Yentna</i>	2.7	1.2	4.6	1.0	12	6	21
	<i>Knik-Turnagain</i>	71.5	67.2	75.6	2.5	332	312	351
	<i>Kenai Peninsula</i>	0.0	0.0	0.6	0.3	0	0	3
Harvest represented						464		
Harvest not represented						0		
Total harvest						464		
Eastern Subdistrict <i>n</i> = 403								
	<i>West</i>	2.7	0.7	5.1	1.4	14	4	27
	<i>Susitna</i>	20.9	17.5	24.2	2.0	109	92	127
	<i>Deshka</i>	7.9	5.5	10.3	1.5	41	29	54
	<i>Yentna</i>	8.2	5.9	10.7	1.5	43	31	56
	<i>Knik-Turnagain</i>	60.3	56.1	64.3	2.5	315	294	336
	<i>Kenai Peninsula</i>	0.1	0.0	0.7	0.3	0	0	4
Harvest represented						523		
Harvest not represented						0		
Total harvest						523		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

Table 14.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2017. Sample sizes (*n*), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
General Subdistrict (south) <i>n</i> = 423								
	<i>West</i>	18.9	15.2	22.7	2.2	224	181	270
	<i>Susitna</i>	35.8	31.7	39.9	2.5	425	377	474
	<i>Deshka</i>	18.4	15.2	21.6	1.9	219	181	257
	<i>Yentna</i>	16.5	13.2	20.1	2.1	196	156	239
	<i>Knik-Turnagain</i>	10.2	7.8	12.8	1.5	122	92	152
	<i>Kenai Peninsula</i>	0.2	0.0	0.7	0.2	3	0	8
Harvest represented						1,188		
Harvest not represented						0		
Total harvest						1,188		
General Subdistrict (north) <i>n</i> = 257								
	<i>West</i>	4.3	1.6	7.7	1.8	24	9	42
	<i>Susitna</i>	16.0	11.9	20.2	2.5	87	65	110
	<i>Deshka</i>	7.5	4.8	10.4	1.7	41	26	57
	<i>Yentna</i>	6.3	3.8	9.3	1.7	34	21	51
	<i>Knik-Turnagain</i>	65.9	60.5	71.0	3.2	359	330	387
	<i>Kenai Peninsula</i>	0.0	0.0	0.3	0.2	0	0	2
Harvest represented						545		
Harvest not represented						0		
Total harvest						545		
Eastern Subdistrict <i>n</i> = 259								
	<i>West</i>	9.7	5.9	13.9	2.5	41	25	58
	<i>Susitna</i>	23.2	18.7	28.3	2.9	97	78	118
	<i>Deshka</i>	9.9	6.7	13.5	2.0	42	28	57
	<i>Yentna</i>	7.7	4.8	11.2	2.0	32	20	47
	<i>Knik-Turnagain</i>	47.7	42.3	53.2	3.3	199	177	223
	<i>Kenai Peninsula</i>	1.7	0.5	3.3	0.9	7	2	14
Harvest represented						418		
Harvest not represented						0		
Total harvest						418		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

Table 15.—Annual stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017. Estimates were calculated using a stratified estimator for combined area strata. Within each year, sample sizes (*n*), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
Year 2016, <i>n</i> =1,236								
	<i>West</i>	14.3	11.8	16.9	1.5	306	252	360
	<i>Susitna</i>	27.2	24.6	29.8	1.6	581	525	637
	<i>Deshka</i>	10.9	9.1	12.8	1.1	232	194	273
	<i>Yentna</i>	12.6	10.5	14.6	1.3	269	224	312
	<i>Knik-Turnagain</i>	35.1	33.1	37.0	1.2	749	707	791
	<i>Kenai Peninsula</i>	0.0	0.0	0.3	0.1	0	0	6
Harvest represented						2,137		
Harvest not represented						0		
Total harvest						2,137		
Year 2017, <i>n</i> = 939								
	<i>West</i>	13.4	11.2	15.8	1.4	289	241	340
	<i>Susitna</i>	28.3	25.7	31.0	1.6	609	554	667
	<i>Deshka</i>	14.0	12.0	16.0	1.2	301	258	345
	<i>Yentna</i>	12.2	10.2	14.3	1.3	262	219	308
	<i>Knik-Turnagain</i>	31.6	29.4	33.9	1.3	680	633	728
	<i>Kenai Peninsula</i>	0.5	0.1	0.9	0.2	10	3	18
Harvest represented						2,151		
Harvest not represented						0		
Total harvest						2,151		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

Table 16.—Annual stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017. Within each year, sample sizes (*n*), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
Year 2016, <i>n</i> = 279 ^a								
	<i>West</i>	28.8	23.4	35.0	3.5	292	237	355
	<i>Susitna</i>	32.3	27.1	37.8	3.4	327	274	384
	<i>Deshka</i>	11.6	8.1	15.4	2.2	118	82	156
	<i>Yentna</i>	17.1	12.8	21.8	2.8	174	130	221
	<i>Knik-Turnagain</i>	9.9	7.0	13.2	1.9	100	71	134
	<i>Kenai Peninsula</i>	0.2	0.0	0.9	0.4	2	0	9
Harvest represented						1,013		
Harvest not represented ^a						17		
Total harvest						1,030		
Year 2017, <i>n</i> = 247								
	<i>West</i>	16.5	11.6	21.9	3.2	216	152	285
	<i>Susitna</i>	38.3	32.4	44.3	3.7	499	422	577
	<i>Deshka</i>	10.4	7.0	14.2	2.2	136	92	185
	<i>Yentna</i>	17.9	13.3	22.9	2.9	234	174	298
	<i>Knik-Turnagain</i>	16.0	11.8	20.3	2.6	208	154	265
	<i>Kenai Peninsula</i>	0.8	0.0	2.1	0.7	11	1	27
Harvest represented						1,304		
Harvest not represented						0		
Total harvest						1,304		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

^a Seventeen fish were not represented in 2016 because they fell outside of the sampled date range or dates of harvest were unknown and therefore could not be accurately represented in MSA.

FIGURES




Figure 1.—Map of statistical areas for set gillnet commercial fishing in the Northern District of Upper Cook Inlet.




Figure 2.—Sampling locations for Chinook salmon populations from Cook Inlet included in the Cook Inlet genetic baseline.

Note: Numbers correspond to map numbers in Table 1. Location dot color matches reporting group assignment.




Figure 3.—Graphical depictions of temporal and area strata (boxes) in the a) Northern District set gill net and b) Tyonek Subsistence set gillnet fisheries, in which stock composition and stock-specific harvest was estimated and c) keys for the colors indicating percentage of the harvest analyzed in each analysis, acronyms of each area stratum, and subscripts used to indicate temporal strata. Lines connecting smaller strata to larger strata indicate which estimates were stratified to calculate larger strata.

Note: Superscript numbers next the name or acronym of each temporal and area stratum indicate the table number where their estimates can be found.




Figure 4.—Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the General Subdistrict (south), General Subdistrict (north), and Eastern Subdistrict of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017.




Figure 5.—Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017.

Note: Estimates were calculated using a stratified estimator for combined area strata




Figure 6.—Annual stock-specific harvest estimates and 90% credibility intervals based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016 and 2017.

**APPENDIX A: STOCK COMPOSITION AND STOCK-
SPECIFIC HARVEST ESTIMATES OF FINE-SCALE
TEMPORAL STRATA, 2016 COMMERCIAL FISHERIES**

Appendix A1.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Trading Bay and Tyonek statistical areas of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016. Sample sizes (n), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
Trading Bay, May 29–June 30, $n = 143$								
	<i>West</i>	25.1	17.8	32.9	4.5	146	103	191
	<i>Susitna</i>	39.0	31.5	46.4	4.6	226	183	269
	<i>Deshka</i>	11.2	6.8	16.0	2.9	65	39	93
	<i>Yentna</i>	15.0	9.4	21.0	3.5	87	55	122
	<i>Knik-Turnagain</i>	9.7	5.7	14.5	2.7	56	33	84
	<i>Kenai Peninsula</i>	0.0	0.0	0.5	0.3	0	0	3
Harvest represented						581		
Harvest not represented						0		
Total harvest						581		
Tyonek, May 29–June 12, $n = 193$								
	<i>West</i>	20.5	14.7	26.3	3.6	71	51	91
	<i>Susitna</i>	30.5	24.4	36.5	3.7	105	84	126
	<i>Deshka</i>	16.6	12.1	21.7	2.9	57	42	75
	<i>Yentna</i>	22.2	16.8	28.0	3.5	77	58	97
	<i>Knik-Turnagain</i>	10.3	6.4	14.8	2.5	35	22	51
	<i>Kenai Peninsula</i>	0.0	0.0	0.3	0.2	0	0	1
Harvest represented						346		
Harvest not represented						0		
Total harvest						346		
Tyonek, June 13–30, $n = 156$								
	<i>West</i>	28.5	21.5	36.2	4.5	64	48	81
	<i>Susitna</i>	25.1	19.0	31.6	3.8	56	42	70
	<i>Deshka</i>	19.4	13.6	25.6	3.6	43	30	57
	<i>Yentna</i>	22.3	15.7	29.2	4.0	50	35	65
	<i>Knik-Turnagain</i>	4.7	2.0	8.1	1.9	10	4	18
	<i>Kenai Peninsula</i>	0.0	0.0	0.5	0.3	0	0	1
Harvest represented						223		
Harvest not represented						0		
Total harvest						223		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

Appendix A2.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the General Subdistrict (north) area of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016. Sample sizes (n), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
General Subdistrict (north), May 29–June 12, $n = 125$								
	<i>West</i>	6.1	1.6	11.4	2.9	10	3	19
	<i>Susitna</i>	26.2	19.9	33.1	4.0	43	33	55
	<i>Deshka</i>	8.2	4.4	12.8	2.6	14	7	21
	<i>Yentna</i>	2.4	0.0	6.4	2.0	4	0	11
	<i>Knik-Turnagain</i>	57.1	49.5	64.2	4.5	95	82	107
	<i>Kenai Peninsula</i>	0.0	0.0	1.1	0.5	0	0	2
Harvest represented						166		
Harvest not represented						0		
Total harvest						166		
General Subdistrict (north), June 13–27, $n = 216$								
	<i>West</i>	0.2	0.0	2.7	1.3	1	0	8
	<i>Susitna</i>	13.5	9.4	18.0	2.6	40	28	54
	<i>Deshka</i>	4.0	1.8	6.8	1.6	12	5	20
	<i>Yentna</i>	2.8	1.2	4.8	1.1	8	4	14
	<i>Knik-Turnagain</i>	79.5	74.4	84.3	3.1	237	222	251
	<i>Kenai Peninsula</i>	0.0	0.0	0.8	0.3	0	0	2
Harvest represented						298		
Harvest not represented						0		
Total harvest						298		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

Appendix A3.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Eastern Subdistrict area of the Northern District set gillnet fishery of Cook Inlet, Alaska, in 2016. Sample sizes (n), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting Group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
Eastern Subdistrict, May 29–June 12, $n = 186$								
	<i>West</i>	0.3	0.0	4.1	2.0	1	0	11
	<i>Susitna</i>	25.9	20.7	31.5	3.3	67	53	81
	<i>Deshka</i>	9.0	5.5	12.7	2.2	23	14	33
	<i>Yentna</i>	10.1	6.4	14.2	2.4	26	16	37
	<i>Knik-Turnagain</i>	54.7	48.7	60.6	3.7	141	125	156
	<i>Kenai Peninsula</i>	0.0	0.0	0.4	0.2	0	0	1
Harvest represented						257		
Harvest not represented						0		
Total harvest						257		
Eastern Subdistrict, June 13–30, $n = 218$								
	<i>West</i>	5.0	2.1	8.5	1.9	13	6	23
	<i>Susitna</i>	16.0	12.1	20.3	2.5	43	32	54
	<i>Deshka</i>	6.8	3.9	10.1	1.9	18	10	27
	<i>Yentna</i>	6.3	3.7	9.4	1.8	17	10	25
	<i>Knik-Turnagain</i>	65.7	60.2	71.3	3.4	175	160	190
	<i>Kenai Peninsula</i>	0.2	0.0	1.3	0.6	0	0	4
Harvest represented						266		
Harvest not represented						0		
Total harvest						266		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

**APPENDIX B: STOCK COMPOSITION AND STOCK-
SPECIFIC HARVEST ESTIMATES OF FINE-SCALE
TEMPORAL STRATA, 2016 TYONEK SUBSISTENCE
FISHERY**

Appendix B1.—Stock composition (%) and stock-specific harvest estimates based on genetic data for Chinook salmon harvested in the Tyonek subsistence set gillnet fishery of Cook Inlet, Alaska, in 2016. Sample sizes (*n*), means, 90% credibility intervals (CI), and standard deviation of the proportions (SD) are provided.

Harvest	Reporting group	Stock composition (%)				Stock-specific harvest		
		Mean	90% CI		SD	Mean	90% CI	
			5%	95%			5%	95%
Tyonek subsistence, May 16–31, <i>n</i> = 137								
	<i>West</i>	29.2	21.4	37.5	4.9	120	88	155
	<i>Susitna</i>	32.9	24.9	41.2	5.0	136	103	170
	<i>Deshka</i>	7.3	3.3	12.2	2.7	30	14	50
	<i>Yentna</i>	21.3	14.8	28.5	4.2	88	61	117
	<i>Knik-Turnagain</i>	8.6	4.7	13.3	2.6	36	20	55
	<i>Kenai Peninsula</i>	0.5	0.0	2.0	0.8	2	0	8
Harvest represented						412		
Harvest not represented ^a						0		
Total harvest						412		
Tyonek subsistence, June 1–25, <i>n</i> = 142								
	<i>West</i>	28.6	21.0	37.2	4.8	172	126	224
	<i>Susitna</i>	31.9	24.7	39.4	4.5	191	149	237
	<i>Deshka</i>	14.5	9.5	20.1	3.2	87	57	121
	<i>Yentna</i>	14.3	8.5	21.1	3.8	86	51	127
	<i>Knik-Turnagain</i>	10.8	6.8	15.4	2.6	65	41	93
	<i>Kenai Peninsula</i>	0.0	0.0	0.5	0.3	0	0	3
Harvest represented						601		
Harvest not represented ^a						0		
Total harvest						601		

Note: Stock-specific harvest estimates may not sum to the total harvest represented due to rounding error and their 90% credibility intervals may not include the point estimate for the very low stock-specific harvest numbers because fewer than 5% of iterations had values above zero.

^a Seventeen fish were not represented in 2016 because they fell outside of the sampled date range or dates of harvest were unknown and therefore could not be accurately represented in MSA.