

Gulkana River Chinook Salmon Distribution

by

Corey J. Schwanke

March 2015

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H _A
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	<i>e</i>
hectare	ha	at	@	catch per unit effort	CPUE
kilogram	kg	compass directions:		coefficient of variation	CV
kilometer	km	east	E	common test statistics	(F, t, χ^2 , etc.)
liter	L	north	N	confidence interval	CI
meter	m	south	S	correlation coefficient (multiple)	R
milliliter	mL	west	W	correlation coefficient (simple)	r
millimeter	mm	copyright	©	covariance	cov
		corporate suffixes:		degree (angular)	°
Weights and measures (English)		Company	Co.	degrees of freedom	df
cubic feet per second	ft ³ /s	Corporation	Corp.	expected value	<i>E</i>
foot	ft	Incorporated	Inc.	greater than	>
gallon	gal	Limited	Ltd.	greater than or equal to	≥
inch	in	District of Columbia	D.C.	harvest per unit effort	HPUE
mile	mi	et alii (and others)	et al.	less than	<
nautical mile	nmi	et cetera (and so forth)	etc.	less than or equal to	≤
ounce	oz	exempli gratia		logarithm (natural)	ln
pound	lb	(for example)	e.g.	logarithm (base 10)	log
quart	qt	Federal Information Code	FIC	logarithm (specify base)	log ₂ , etc.
yard	yd	id est (that is)	i.e.	minute (angular)	'
		latitude or longitude	lat. or long.	not significant	NS
Time and temperature		monetary symbols (U.S.)	\$, ¢	null hypothesis	H ₀
day	d	months (tables and figures): first three letters	Jan, ..., Dec	percent	%
degrees Celsius	°C	registered trademark	®	probability	P
degrees Fahrenheit	°F	trademark	™	probability of a type I error (rejection of the null hypothesis when true)	α
degrees kelvin	K	United States (adjective)	U.S.	probability of a type II error (acceptance of the null hypothesis when false)	β
hour	h	United States of America (noun)	USA	second (angular)	"
minute	min	U.S.C.	United States Code	standard deviation	SD
second	s	U.S. state	use two-letter abbreviations (e.g., AK, WA)	standard error	SE
				variance	
Physics and chemistry				population	Var
all atomic symbols				sample	var
alternating current	AC				
ampere	A				
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

REGIONAL OPERATIONAL PLAN SF.3F.2014.11

GULKANA RIVER CHINOOK SALMON DISTRIBUTION

by

Corey J. Schwanke

Alaska Department of Fish and Game, Division of Sport Fish, Fairbanks

Alaska Department of Fish and Game
Division of Sport Fish

March 2015

The Regional Operational Plan Series was established in 2012 to archive and provide public access to operational plans for fisheries projects of the Divisions of Commercial Fisheries and Sport Fish, as per joint-divisional Operational Planning Policy. Documents in this series are planning documents that may contain raw data, preliminary data analyses and results, and describe operational aspects of fisheries projects that may not actually be implemented. All documents in this series are subject to a technical review process and receive varying degrees of regional, divisional, and biometric approval, but do not generally receive editorial review. Results from the implementation of the operational plan described in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author if you have any questions regarding the information provided in this plan. Regional Operational Plans are available on the Internet at: <http://www.adfg.alaska.gov/sf/publications/>

*Corey J. Schwanke,
Alaska Department of Fish and Game, Division of Sport Fish,
1300 College Road, Fairbanks, AK 99701-1599*

This document should be cited as:

Schwanke, C. J. 2014. Gulkana River Chinook salmon distribution. Alaska Department of Fish and Game, Regional Operational Plan ROP.SF.3F.2014.11, Fairbanks.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3464 or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

SIGNATURE PAGE

Project Title: Gulkana River Chinook Salmon Distribution

Project leader(s): Corey Schwanke, *Fishery Biologist II*

Division, Region, and Area: Division of Sport Fish, Region III, Fairbanks

Project Nomenclature: AKSSF Grant # 44724

Period Covered: March 2013–August 2015

Field Dates: 15 May-1 September 2013–2015

Plan Type: Category III

Approval







Title	Name	Signature	Date
Project leader	Corey Schwanke		
Biometrician	Jieqi Huang		
Research Coordinator	Matthew Evenson		
Regional Supervisor Chief Fisheries	Don Roach		3/5/15
Scientist Sport Fish Chief Fisheries	James Hasbrouck		3/6/2015
Scientist Commercial Fish	Eric Volk		3/6/2015
CSRI Coordinator	Ed Jones		3/6/15

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	ii
LIST OF FIGURES	iii
LIST OF APPENDICES	iii
PURPOSE.....	1
OBJECTIVES.....	2
BACKGROUND.....	4
Introduction	4
METHODS.....	6
Study Area	6
Study Design	6
Overview	6
Fish capture.....	6
Telemetric Procedures	9
Data Collection and Reduction.....	12
Data Analysis.....	12
SCHEDULE AND DELIVERABLES	14
RESPONSIBILITIES	16
REFERENCE CITED.....	17
APPENDIX A	18

LIST OF TABLES

Table	Page
1. Tagging schedule for Chinook salmon, Gulkana River, 2013.....	11
2. List of all possible fates of radiotagged Chinook salmon in the Gulkana River drainage.....	13

LIST OF FIGURES

Figure	Page
1. Map of the Gulkana River drainage within the Upper Copper Upper Susitna Management Area.....	3
2. Number of Chinook salmon harvested and released (the sum is catch), and fishing effort for all species, Gulkana River, 1992-2011	4
3. Historic expanded counts of Chinook salmon from the counting tower, Gulkana River, 2002-2012.....	5
4. Map of the Gulkana River with radiotagging and tracking station locations	8

LIST OF APPENDICES

Appendix	Page
A1. Example of the field form for Chinook salmon sampling, Gulkana River, 2013	18

ABSTRACT

The primary objective of this study is to estimate the proportions of Chinook salmon *Oncorhynchus tshawytscha* that spawn above and below the ADF&G counting tower which is located about 71 river kilometers from its confluence with the Copper River. The counting tower is placed upstream of the West Fork Gulkana River because it runs turbid intermittently throughout the summer and only enumerates an unknown proportion of the total escapement for the Gulkana River drainage. Past estimates of drainage-wide Chinook salmon distribution study for the Copper River has shown that between 50 and 80% of fish bound for the Gulkana River spawned above the counting tower. This is the first study ever to specifically address what proportion of the total spawning escapement is passing the counting tower, and the results will help determine if a spawning escapement goal is warranted. This will be achieved by deploying radio tags at the mouth of the Gulkana River and monitoring their movements using aerial surveys and ground-based tracking stations. Additionally, this study will describe inriver run timing as well as age, sex and length composition of the Gulkana River Chinook salmon run.

Key words: Chinook salmon, *Oncorhynchus tshawytscha*, Gulkana River, spawning distribution, spawning escapement, radiotelemetry, counting tower, run timing.

PURPOSE

The Copper River is 1 of 12 indicator stocks chosen by the ADF&G in the *Chinook Salmon Stock Assessment and Research Plan* (ADFG Chinook Research Team 2013) as a stock for which additional information on stock productivity is desired, and the lack of escapement estimates from the six major spawning tributaries has been identified as an information gap. To establish a sustainable escapement goal, long-term information about the number of recruits per spawner is required. To better understand this process, Region III Sport Fish Division will conduct a radiotelemetry study to estimate spawning distribution and run timing patterns of Chinook salmon above and below an established counting tower on the Gulkana River. Studies have shown that under certain river conditions (i.e., low water, high temperatures) a substantial proportion (>40%) of the escapement may not be assessed (Savereide 2005). The Gulkana River Chinook salmon distribution study, in conjunction with the counting tower, will provide a method to estimate the total escapement of Chinook salmon on a highly utilized stock. Gulkana River Chinook salmon are genetically distinct from other Copper River stocks and exhibit an early run-timing pattern. They comprise an average of 21% of the Copper River drainage Chinook salmon spawning population (Savereide 2005) and over 29% of the Chinook salmon available to Upper Copper River (Glennallen Subdistrict) subsistence fisheries.

Chinook salmon will be captured and implanted with radio transmitters over the course of the run in the lower end of the Gulkana River (Figure 1). Migrations of radiotagged fish will be monitored with a combination of tracking stations and aerial tracking surveys. Run timing patterns will be estimated from final spawning locations and the date and time of initial capture. The study will also collect age, sex, and length (ASL) information from Chinook salmon to characterize the composition of the Chinook salmon escapement and use that information in an age-structured model to evaluate the escapement goal and forecast returns.

OBJECTIVES

The objectives of this study during 2012–2014 are to:

- 1) estimate the proportions of spawning Chinook salmon (\geq than 20") above and below the counting tower on the Gulkana River such that all estimates are within 6% of the true values 95% of the time;
- 2) describe inriver timing patterns of Chinook salmon spawning (\geq than 20") in the Gulkana River; and,
- 3) estimate ASL composition of the escapement of Chinook salmon (\geq than 20") in the Gulkana River such that estimated proportions are within 0.10 of the true proportions 95% of the time.




Figure 1.—Map of the Gulkana River drainage within the Upper Copper Upper Susitna Management Area.

BACKGROUND

INTRODUCTION

The Gulkana River Chinook salmon stock is subject to a substantial commercial fishery at the mouth of the Copper River, significant subsistence and personal-use (PU) fisheries in the mainstem Copper River, and the largest Chinook salmon sport fishery in the drainage. There are no stock specific estimates of harvest available for the subsistence or personal use fisheries, but there is for the sport fishery. Annual sport fishing catch and harvest has been substantial in the past, peaking in 1997, but has steadily declined over the last 15 years (Figure 2; Somerville 2011). Recently, catch and harvest rates have decreased due to poor returns and associated sport fishing restrictions. Similar trends in overall harvest in the PU fishery exist; however, harvest has remained relatively stable in the subsistence fisheries (i.e., state and federal) as no restrictions have taken place on these fisheries (Botz et al. 2011 and Somerville 2011).




Figure 2.—Number of Chinook salmon harvested and released (the sum is catch), and fishing effort for all species, Gulkana River, 1992–2011.

Since 2002, a salmon counting tower has been cooperatively operated on the Gulkana River by ADF&G and the Bureau of Land Management (BLM). Operational costs for ADF&G have been funded by the Alaska Sustainable Salmon Fund (AKSSF) since 2011. The counting tower is located in the mainstem about 2.5 km above the confluence of the West Fork Gulkana River. Since 2002, the number of Chinook salmon spawners past the tower has declined each year for a total decline > 50% from 2002–2012 (Figure 3; Maclean 2013). In 2012, the smallest escapement of Chinook salmon was recorded at 1,730 fish. This telemetry study, funded operationally by AKSSF through 2015, will estimate the spawning distribution above and below the counting

tower and investigate the relationship between inriver habitat conditions (e.g., river height, water temperature) and spawning distribution, both of which are needed to establish an escapement goal for Gulkana River Chinook salmon. The Chinook Salmon Research Initiative is funding the research biologists that conduct these vital projects.




Figure 3.—Historic expanded counts of Chinook salmon from the counting tower, Gulkana River, 2002–2012.

Also located at the counting tower, since 2011, is a passive integrated transponder (PIT) tag array that the Native Village of Eyak (NVE) is using to estimate run-timing patterns from Chinook salmon that are PIT tagged during their inriver abundance project in the Lower Copper River (Baird and Wood canyons). Since 2003, estimates of inriver harvest (subsistence and personal-use) have been subtracted from the inriver abundance estimates derived by NVE to estimate the drainage-wide escapement. This array in combination with the counting tower and telemetry study can be used to supplement the second event of NVE’s inriver abundance study by determining the marked to unmarked ratio of the Gulkana River escapement. Currently, ADF&G personnel download the data from the PIT tag array for NVE and discussions are ongoing with NVE to continue the inriver abundance and PIT tag studies cooperatively in the future.

Past drainage-wide telemetry studies have shown that not all Chinook salmon spawn above the tower with factors such as water temperature, water height, and likely just general run timing affecting not only swim speeds, but the proportions of the overall escapement above the tower (Savereide 2005; Maclean 2013). Telemetry studies have demonstrated that 14%–50% of the run may spawn below the tower site annually. In 2004, nearly 50% of all radiotagged Gulkana River Chinook salmon appeared to have spawned below the counting tower. That year the Gulkana River was low and clear, and abnormally high water temperatures were recorded at the counting tower site (Maclean 2013). Larger sample sizes of radiotagged fish within the Gulkana River will be attained with this study. It is also hoped that a variety of water conditions will be experienced during this study to further investigate those effects on the proportions of fish spawning above and below the counting tower.

The Gulkana River Chinook salmon distribution study, in conjunction with the counting tower, will provide a method to estimate the total escapement of Chinook salmon on a highly utilized stock. Gulkana River Chinook salmon are genetically distinct from other Copper River stocks and exhibit an early run-timing pattern. They comprise an average of 21% of the Copper River drainage Chinook salmon spawning population (Savereide 2005) and over 29% of the Chinook salmon available to Upper Copper River (Glennallen Subdistrict) subsistence fisheries. On average, over 70% of the average (2002–2011) annual subsistence harvest of 2,710 Chinook salmon is harvested before 4 July. The Gulkana River Chinook salmon counting tower coupled with distribution estimates of spawning will provide the only method to assess the inseason progression of escapement for any individual Chinook salmon stock in the Copper River drainage. This inseason monitoring combined with anecdotal harvest data from the various Upper Copper River fisheries is vital to managers when making decisions about the Copper River’s important subsistence, personal-use, commercial, and sport fisheries. The estimates of total escapement will be used to establish a SEG for Gulkana River Chinook salmon.

METHODS

STUDY AREA

The Gulkana River is one of six major spawning tributaries for Chinook salmon in the Copper River drainage (Savereide 2005). The mainstem Gulkana River is fed by the East Fork, Middle Fork, and West Fork Gulkana rivers. The study area will include most of the Gulkana River drainage, from its confluence with the Copper River upriver to spawning locations of radiotagged Chinook salmon (Figure 4). Radiotagging will take place in the lower river from the Richardson Highway downstream to the confluence with the Copper River, a distance of approximately 7 rkm. The majority of the sampling will take place at the confluence with the Copper River.

STUDY DESIGN

Overview

This study will estimate the spawning distribution and run timing of Chinook salmon in the Gulkana River. Migrations of radiotagged fish will be monitored with 5 tracking stations positioned at strategic points throughout the Gulkana River drainage, as well as a series of aerial tracking surveys from a fixed-wing aircraft. Spawning distribution will be estimated as the ratio of numbers of radiotagged fish migrating above and below the counting tower to the total number of radiotagged fish surviving and migrating to all spawning areas. Run timing will be identified from final locations and the date and time of initial capture.

Fish capture

Some technical information (Savereide 2005) is available as to when Chinook salmon reach the mouth of the Gulkana River, but we feel run timing has changed substantially the last few years, and that the handling effects (i.e., fish wheel capture and tagging) might have influenced the arrival times during these studies. For this study, the initial tagging schedule will be based partially on previous telemetry work, run timing curves from the counting tower, personal experiences of the principal investigator and anecdotal information on angling at the mouth of the Gulkana River. First, the date of 3 June was chosen as when fish are expected to arrive in catchable numbers at the mouth of the Gulkana River. Second, a run curve was constructed from 11 years of data from the counting tower. This was calculated by averaging the mean daily proportion of fish passing the counting tower each day. Finally, this run curve was shifted to a start date of June 3 to project an expected passage rate at the tagging area (the mouth of the Gulkana River). The proportion of expected passage at the mouth was then summed by week and




Figure 4.–Map of the Gulkana River with radiotagging and tracking station locations.

multiplied by the total number of tags to be deployed (150) to determine weekly tagging goals (Table 1). The tagging schedule will be evaluated at specified times during the run and increased or decreased based on the perceived run strength and timing. Sampling will only take place during the workweek (Mon-Fri) and possibly at night to avoid potential conflicts with sport fisherman using the area.

Chinook salmon will be captured with rod-and-reel, dip nets, and possibly beach seine. Tackle used to capture Chinook salmon will be flies, wiggle warts, and spinning glow/salmon roe combinations. A total of 150 Chinook salmon will be radiotagged from 3 June through 28 July in the lower portion of the Gulkana River to ensure radio transmitters are distributed across the entire run. Every healthy fish will receive a radio tag. When the quota for a tagging period (i.e., a work-week) has been reached, sampling will cease. A sampling day or two may take place prior to 3 June to ensure that no fish are missed.

All captured fish will be measured to the nearest 5-mm mid-eye to fork length (MEF) and sex will be determined from external characteristics. After capture, Chinook salmon will be placed in a tagging cradle filled with water. Two types of radio tags will be used the first year of the study: esophageal and external mount models. Seventy-five of each will be used with the intent of using the model that works the best (ease of application, tag retention, fish response, etc.) in future years. Esophageal tags will be inserted through the esophagus and into the upper stomach as described in Saveriede (2005). External model tags will be attached using standard procedures outlined in Schwanke and Reed (2011). All fish captured will also receive a uniquely numbered, brightly colored Floy™ tag with an encased distinctly identifiable pit tag¹. All tagging will be performed without the use of anesthesia.

Telemetric Procedures

Radio tags will be Model Five pulse encoded transmitters made by Advanced Telemetry Systems (ATS). Each transmitter will be distinguishable by its frequency and encoded pulse pattern. Five frequencies chosen for this project are 149.420, 149.440, 149.480, 149.510 and 149.550 kHz. Each frequency will have 30 radio tags each with distinguishable encoded pulse patterns. The pulse rate for these tags will be about once every 1.3 seconds. Radiotagged Chinook salmon will be tracked along the course of the Gulkana River using a network of 4 ground-based tracking stations (Figure 4). Each station includes 2 deep-cycle batteries, a solar array, an antenna switch box, a steel housing box, 2 Yagi antennas, and an ATS Model R4500. The receiver will be programmed to scan through the frequencies at 2 s intervals, and receive from

¹ Pit tags are being applied to all captured Chinook salmon to aid a study which will evaluate the efficacy of a pit tag detection array placed near the counting tower. This is a separate project being conducted by the Native Village of Eyak.

both antennas simultaneously. When a signal of sufficient strength is encountered, the receiver will pause for 12 seconds on each antenna, and then transmitter frequency, transmitter code, signal strength, date, time, and antenna number will be recorded on the data logger. The relatively short cycle period will minimize the chance that a radiotagged fish will swim past the receiver site without being detected. Cycling through all frequencies will require 2–3 minutes depending on the number of active tags in the reception range and level of background noise. Recorded data will be downloaded to a laptop computer every 10–12 days.

Table 1.–Tagging schedule for Chinook salmon, Gulkana River.

Calendar Week (Mon-Sun)	Sampling Dates (Mon-Fri)	Number of Tags Deployed	Cumulative Tags Deployed
3–9 June	3–7 June	12	12
10–16 June	10–14 June	24	36
17–23 June	17–21 June	32	68
24–30 June	24–28 June	26	94
1–7 July	1–5 July	20	114
8–14 July	8–12 July	15	129
15–21 July	15–19 July	14	143
22–28 July	22–26 July	7	150

The first tracking station will be placed downstream of the Richardson Hwy Bridge (Figure 4), approximately 6 km above the capture site. This station will be used to determine the number of radiotagged fish that survived handling and tagging and continued to migrate up the Gulkana River. The next station will be placed on the mainstem Gulkana River downstream from Sourdough Landing, near the perceived lower spawning boundary. The next station will be on the West Fork Gulkana approximately 1 km upstream from the confluence with the mainstem river. This station will enumerate radiotagged Chinook salmon spawning in the tributaries and mainstem of the West Fork. The final station in the Gulkana River will be placed at the counting tower and will enumerate all radiotagged Chinook salmon spawning upstream of the counting tower. A fifth station will be placed on the mainstem Copper River approximately 2 km downstream from the mouth of the Gakona River (Figure 4). This station will be used to enumerate all radiotagged fish that were bound for Upper Copper River drainage tributaries, but were captured during their upstream migration near the mouth of the Gulkana River.

The distribution of radiotagged Chinook salmon throughout the Gulkana River drainage will be further determined by aerial tracking from small aircraft. A series (3–5) of aerial-tracking surveys of the entire drainage will be conducted in late June, July, and August. Tracking flights will be conducted with one aircraft and one person (in addition to the pilot) utilizing one R4500 receiver. All frequencies will be loaded into the receiver prior to each flight. Dwell time on each frequency will be 2 s. Flight altitude will range from 100 to 300 m above ground. Two antennas, one on each wing strut, will be mounted such that the antennas receive signals perpendicular to the direction of travel. Once a transmitter is identified, its frequency, code, and GPS location will be recorded. The purpose of the aerial tracking will be to locate fish that tracking stations failed to record, and to validate that fish recorded at the tracking stations did migrate into that particular area.

DATA COLLECTION AND REDUCTION

For each Chinook salmon captured, data collected will include: 1) measurement of fish length to the nearest 5 mm MEF length; 2) type of capture gear used; 3) location (river kilometer and GPS coordinate); 4) number printed on the Floy™ tag; and, 5) date. For all fish receiving a radio tag, additional data collected will include the radio transmitter frequency, code, and type of tag (e.g., esophageal or external; Appendix A).

Ages will be determined from scale patterns as described by Mosher (1969). Three scales will be removed from the left side of the fish approximately 2 rows above the lateral line along a diagonal line downward from the posterior insertion of the dorsal fin to the anterior insertion of the anal fin (Welander 1940).

Each tracking station will record date, time, frequency, code, signal strength, and antenna number for each time a signal of sufficient strength is encountered. Tracking stations will be periodically downloaded using a laptop computer (weekly). During each aerial tracking survey, data collected for each fish will include frequency, code, latitude, longitude, and a general description of its location (e.g., approximately 1 mile upstream of Sourdough Creek).

Coordinates of located fish will be plotted on USGS maps (represented in Alaska NAD27 Datum) using Arcview™. Final copies of the Excel and Arcview files will be provided with the completed report when it is submitted for review to be archived in the Sport Fish Division Docushare website. The file name and directory location will be presented in the final report and can be made available upon request.

DATA ANALYSIS

To facilitate data analysis, all radiotagged Chinook salmon will be assigned a “fate” (Table 2). The known fate of all radiotagged fish is required to attain unbiased estimates of parameters. Information attained from radio tag/Floy™ tag returns, tracking stations, and aerial surveys will be used to define fates. In previous studies (1999–2004) fates were assigned to all radiotagged Chinook salmon. All radiotagged fish located and designated as “spawners” will be used to estimate distribution and timing.

Table 2.–List of all possible fates of radiotagged Chinook salmon in the Gulkana River drainage.

Fate	Description
Fishery Mortality	A fish harvested in the sport or subsistence fisheries.
Spawner	A fish that migrates into a spawning area of the Gulkana River. Within this category, spawning locations will be determined, not only for tributaries, but also river segments (e.g., above and below the counting tower).
Upstream Migrant	A fish that migrates upriver of the Gulkana River.
Failure	A fish that is never recorded upstream of the Lower Gulkana River or Copper River tracking stations.

Among fish that migrate past a given tracking station, the proportion of fish that have fate j will be estimated as:

$$\hat{P}_j = \frac{\sum_i^{\text{days}} R_{ij}}{n} \quad (1)$$

with variance (Cochran 1977):

$$\hat{V}[\hat{p}_j] = \frac{\hat{p}_j(1 - \hat{p}_j)}{n - 1} \quad (2)$$

where R_{ij} is the number of fish tagged on day i having fate j and n equals the total number of radiotagged fish deemed Gulkana River spawners. Fate j can encompass a variety of fates, such as the proportion of fish that spawned above and below the counting tower.

Run timing patterns will be described as time-density functions, where the relative abundance of stock j that enters into the Gulkana River during time interval t will be described by (Mundy 1979):

$$f_j(t) = \frac{R_{tj}}{\sum_i^{\text{days}} R_{ij}} \quad (3)$$

where:

$f_j(t)$ = the empirical temporal probability distribution over the total span of the run for fish spawning in a tributary or mainstem (or portion thereof) j ; and,

R_{ij} = the subset of radiotagged Chinook salmon bound for tributary/area j that will be caught and tagged during day t .

The mean date of passage (\bar{t}_j) from the lower river tracking station for fish spawning in tributary/area j will be estimated as:

$$\bar{t}_j = \sum_t t f_j(t), \quad (4)$$

and the variance of the run timing distribution will be estimated as:

$$Var(t_j) = \sum_t (t - \bar{t}_j)^2 f_j(t). \quad (5)$$

The proportion of fish at age, sex, or length category k by sex s for samples collected solely for age, sex, and length will be calculated as:

$$\hat{p}_{sk} = \frac{y_{sk}}{n_s} \quad (6)$$

where: \hat{p}_{sk} = the estimated proportion of Chinook salmon that are age or length category k and sex s ; y_{sk} = the number of Chinook salmon sampled that are age or length category k ; and, n_s = the total number of Chinook salmon sampled by sex.

The variance of this proportion will be estimated as:

$$\hat{V}[\hat{p}_{sk}] = \frac{\hat{p}_{sk}(1 - \hat{p}_{sk})}{n_s - 1} \quad (7)$$

Escapement at age or length category k for each sex is then estimated by:

$$\hat{N}_{sk} = \hat{p}_{sk} \hat{N}_s \quad (8)$$

The variance for \hat{N}_{sk} in this case is (Goodman 1960):

$$\hat{V}(\hat{N}_{sk}) = \hat{V}(\hat{p}_{sk}) \hat{N}_s^2 + \hat{V}(\hat{N}_s) \hat{p}_{sk}^2 - \hat{V}(\hat{p}_{sk}) \hat{V}(\hat{N}_s). \quad (9)$$

SCHEDULE AND DELIVERABLES

Dates of sampling events, milestones, and other activities for 2013–2015 are summarized in the following table.

Date(s)	Sampling Activity/Milestone
26 April 2013	Operational Plan completed
Late May 2013–2015	Set up tracking stations
3 June-28 July 2013–2015	Deploy radio tags
Periodic Summer Dates	Download tracking stations
Late June 2013–2015	Aerial tracking survey
Early July 2013–2015	Aerial tracking survey
Late July 2013–2015	Aerial tracking survey
Early Aug 2013–2015	Aerial tracking survey
Mid-August 2013–2015	Aerial tracking survey
September 2013–2015	Disassemble tracking stations
December 2015	FDS Final Report Complete

RESPONSIBILITIES

List of Personnel and Duties:

ADF&G

Corey Schwanke:	Fishery Biologist II;	Overall supervision of project. Coordinate sampling schedules with project personnel. Analyze data and prepare reports with technical assistance.
James Savereide:	Fishery Biologist III;	Supervise project leader, assist with fixed-station set up, assist with capture, sampling, data collection and review all reports.
Jiaqi Huang:	Biometrician II;	Assist in preparation of statistical design of field investigation for operational plan, and review data analysis and final report.
Klaus Wuttig:	Fishery Biologist III;	Supervise project leader, assist with capture, sampling, data collection, and review all reports.
Matt Albert:	Fishery Biologist I;	Assist with capture, sampling, and data collection.
Chad Bear:	F&W Tech III;	Assist with fixed-station set up, capture, sampling, and data collection.
Unknown:	F&W Tech III;	Assist with fixed-station set up, capture, sampling, and data collection. Serve as crew leader and provide support with operating and downloading tracking stations as well as data entry and editing.

REFERENCE CITED

- ADFG Chinook Research Team. 2013. Chinook salmon stock assessment and research plan, 2013. Alaska Department of Fish and Game, Special Publication No. 13-01, Anchorage.
- Botz, J., G. Hollowell, T. Sheridan, R. Brenner, and S. Moffitt. In prep. Prince William Sound Management Area, 2011. Annual Finfish Management Report. Alaska Department of Fish and Game, Fisheries Management Report No. 13-XX, Anchorage.
- Cochran, W. G. 1977. Sampling Techniques. 3rd edition, John Wiley, New York.
- Goodman, L. A. 1960. On the exact variance of products. Journal of the American Statistical Association. 66:708-713.
- Maclean, S. H. 2013. Chinook salmon escapement and run timing in the Gulkana River, 2011-2012. Alaska Department of Fish and Game, Fishery Data Series No. 13-07, Anchorage.
- Mosher, K. H. 1969. Identification of Pacific salmon and steelhead trout by scale characteristics. United States Department of the Interior, U.S. Fish and Wildlife Service, Bureau of Commercial Fisheries, Washington, D.C., Circular 317.
- Mundy, P. R. 1979. A quantitative measure of migratory timing illustrated by application to the management of commercial fisheries. Phd. Dissertation. University of Washington.
- Savereide, J. W. 2005. Inriver Abundance, Spawning Distribution and Run Timing of Copper River Chinook Salmon, 2002-2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-50, Anchorage.
- Welander, A. D. 1940. A study of the development of the scale of Chinook salmon *Oncorhynchus tshawytscha*. Master's Thesis. University of Washington, Seattle.
- Schwanke, C. J. and D. J. Reed. 2011. Stock assessment of Chinook salmon in the Naknek River, 2003-2004. Alaska Department of Fish and Game, Fishery Data Series No. 11-64, Anchorage.
- Somerville, M. A. 2011. Fishery management report for the recreational fisheries of the Upper Copper/Upper Susitna River management area, 2009. Alaska Department of Fish and Game, Fishery Management Report No. 11-38, Anchorage.

APPENDIX A

