

**Fishery Data Series No. 10-71**

---

---

**Escapements of Chinook Salmon in Southeast Alaska  
and Transboundary Rivers in 2008**

by

**Keith A. Pahlke**

November 2010

---

---

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries





***FISHERY DATA SERIES NO. 10-71***

**ESCAPEMENTS OF CHINOOK SALMON IN SOUTHEAST ALASKA  
AND TRANSBOUNDARY RIVERS IN 2008**

by  
Keith A. Pahlke  
Division of Sport Fish, Douglas

Alaska Department of Fish and Game  
Division of Sport Fish, Research and Technical Services  
333 Raspberry Road, Anchorage, Alaska, 99518-1599

November 2010

Development and publication of this manuscript were partially financed by the Federal Aid in Sport Fish Restoration Act (16 U.S.C.777-777K) under Project F-10-24.

ADF&G Fishery Data Series was established in 1987 for the publication of Division of Sport Fish technically oriented results for a single project or group of closely related projects, and in 2004 became a joint divisional series with the Division of Commercial Fisheries. Fishery Data Series reports are intended for fishery and other technical professionals and are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm> This publication has undergone editorial and peer review.

*Keith A. Pahlke<sup>a</sup>*

*Alaska Department of Fish and Game, Division of Sport Fish  
802 3<sup>rd</sup> St., Douglas, AK 99824, P.O. Box 110024, Juneau, AK 99811, USA*

<sup>a</sup> Author to whom all correspondence should be addressed: [keith.pahlke@alaska.gov](mailto:keith.pahlke@alaska.gov)

*This document should be cited as:*

*Pahlke, K. A. 2010. Escapements of Chinook salmon in Southeast Alaska and transboundary rivers in 2008. Alaska Department of Fish and Game, Fishery Data Series No. 10-71, Anchorage.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375..

# TABLE OF CONTENTS

	<b>Page</b>
LIST OF TABLES.....	ii
LIST OF FIGURES.....	ii
LIST OF APPENDICES.....	iii
ABSTRACT.....	1
INTRODUCTION.....	1
Description of Study Sites.....	2
METHODS.....	5
Escapement Goals.....	5
Indices of Escapement.....	6
Age, Sex, and Length Composition of Escapements.....	9
RESULTS.....	9
Taku River.....	10
Stikine River.....	10
Andrew Creek.....	13
Alsek River.....	13
Unuk River.....	16
Chickamin River.....	16
Blossom River.....	16
Keta River.....	20
King Salmon River.....	23
Situk River.....	23
Chilkat River.....	24
Other Systems.....	24
Coded Wire Tag Recovery.....	24
Observer Training.....	24
DISCUSSION.....	24
ACKNOWLEDGMENTS.....	32
REFERENCES CITED.....	32
APPENDIX A.....	39
APPENDIX B.....	77
APPENDIX C.....	91

## LIST OF TABLES

Table	Page
1. Peak survey counts, survey expansion factors, estimated total escapement from expanded survey counts, mark–recapture projects or weir, for large Chinook salmon returning to Southeast Alaska and transboundary rivers in 2008.....	8
2. Distribution of spawning Chinook salmon among index areas of the Taku River during years when all index areas were surveyed.....	11
3. Counts of spawning Chinook salmon in index areas of the Taku River, 1973–2008.....	12
4. Counts of large spawning Chinook salmon in the Little Tahltan River, Stikine River, 1975–2008.....	13
5. Counts of spawning Chinook salmon in selected rivers in central Southeast Alaska, 1960–2008.....	15
6. Count of Chinook salmon through the Kluksu River weir, harvest above and below the weir, estimated proportion of large fish through the weir, and counts of spawning adults in other tributaries of the Alsek River, 1976–2008.....	18
7. Peak escapement counts of Chinook salmon to index areas of the Unuk River, 1972–2008.....	19
8. Distribution of spawning Chinook salmon among index areas of the Unuk River for years when all index areas were surveyed.....	20
9. Counts of Chinook salmon in index areas of the Chickamin River, 1972–2008.....	22
10. Distribution of spawning Chinook salmon among index areas of the Chickamin River for years when all index areas were surveyed.....	23
11. Counts of Chinook salmon for selected rivers in Behm Canal, 1961–2008.....	25
12. Peak escapement counts and weir counts of spawning Chinook salmon in the King Salmon River, 1971–2008.....	26
13. Estimated harvests and escapement, by size class, of Situk River Chinook salmon, 1976–2008.....	28
14. Mark–recapture estimates of large Chinook salmon escapement in Chilkat River, 1991–2008.....	29
15. Comparison between primary and two alternate observer counts in survey training flights conducted in 2008.....	31

## LIST OF FIGURES

Figure	Page
1. Location of selected Chinook salmon systems in Southeast Alaska, Yakutat, and transboundary rivers.....	2
2. Counts of Chinook salmon in index areas of the Taku River, 1975–2008 and mark–recapture estimates divided by expansion factor of 5.2.....	11
3. Counts of Chinook salmon at the Little Tahltan River weir, Stikine River, 1975–2008, and mark–recapture estimates divided by expansion factor of 5.36.....	14
4. Counts of Chinook salmon at the Andrew Creek weir, 1976–1984, 1997, and in aerial/foot surveys, 1975, 1985–2008.....	14
5. Weir count of Chinook salmon to the Kluksu River tributary of the Alsek River, 1976–2008, and mark–recapture estimates divided by expansion factor of 4.03.....	17
6. Counts of large Chinook salmon in index areas of the Unuk River, 1975–2008, and mark–recapture estimates divided by expansion factor (4.87).....	17
7. Counts of Chinook salmon in index areas of the Chickamin River, 1975–2008 and mark–recapture estimates divided by expansion factor (4.75).....	21
8. Counts of Chinook salmon into the Blossom River, 1975–2008 and mark–recapture estimates.....	21
9. Counts of Chinook salmon to the Keta River, 1975–2008 and mark–recapture estimates for 1998–2000.....	27
10. Counts of Chinook salmon at a weir and in survey counts in the index area of the King Salmon River, 1975–2008.....	27
11. Counts of large Chinook salmon at the Situk River weir, 1975–2008.....	29
12. Mark–recapture estimates of large Chinook salmon escapement to the Chilkat River, 1991–2008.....	30

## LIST OF APPENDICES

Appendix	Page
A1. Survey escapement goals and system goals for large Chinook salmon, Southeast Alaska and transboundary rivers, as accepted by the Alaska Department of Fish and Game, Canadian Department of Fisheries and Oceans, Chinook Technical Committee, and Transboundary Technical Committee, 2009.....	40
A2. Coordinates of Chinook salmon survey areas in Southeast Alaska and transboundary rivers. ....	41
A3. Descriptions of Chinook salmon escapement index areas in Southeast Alaska and northern British Columbia and peak spawning survey dates.....	43
A4. Estimated total escapements of large Chinook salmon to escapement indicator systems and to Southeast Alaska and transboundary rivers, 1975–2008. ....	54
A5. Variance of estimated total escapements of large Chinook salmon to escapement indicator systems and to Southeast Alaska and transboundary rivers, 1975–2008.....	56
A6. Detailed 2008 Southeast Alaska Chinook salmon escapement surveys as entered into Commercial Fisheries Division Integrated Fisheries Database (IFDB/ALEX). ....	57
A7. Estimated abundance and composition by age and sex of the escapement of Chinook salmon to select systems in Southeast Alaska and transboundary rivers, 2007. ....	60
A8. Average length (MEF), by age, of Chinook salmon in selected systems in Southeast Alaska and transboundary rivers, 2008. ....	67
A9. Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.2 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008.....	72
A10. Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.3 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008.....	73
A11. Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.4 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008.....	74
A12. Age composition and average length by age and sex of Chinook salmon sampled in the Farragut River, 2008.....	75
A13. Numbers of Chinook salmon examined for coded wire tags (CWT) and numbers of tags recovered in rivers in Southeast Alaska and transboundary rivers, 2007–2008.....	76
B1. Predicting escapement from index counts using an expansion factor. ....	78
B2. Peak aerial survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Keta River 1975–2008. ....	81
B3. Peak aerial survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Blossom River 1975–2008. ....	82
B4. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Chickamin River 1975–2008.....	83
B5. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Unuk River 1977–2008. ....	84
B6. Peak survey counts, weir counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Stikine River 1975–2008. ....	85
B7. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in Andrew Creek 1975–2008.....	86
B8. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in King Salmon River, 1971–2008. ....	87
B9. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Taku River, 1973–2008. ....	88
B10. Peak survey counts, estimated total spawning abundance $\hat{N}_L$ with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Alek River, 1976–2008.....	89
C1. Computer files used to complete this report.....	92



## ABSTRACT

As part of a continuing stock assessment program in Southeast Alaska, the Division of Sport Fish obtained indices of escapement for Chinook salmon *Oncorhynchus tshawytscha* in designated streams and transboundary rivers. The estimated total escapement in 2008 was 69,109 large (age .3 and older) Chinook salmon, a 21% increase from the escapement of 57,244 fish estimated in 2007. Ten of 11 escapement indices were within or above escapement goal ranges; the Alsek River was below. Estimated age and sex composition and mean length at age of all stocks sampled in 2008 are presented.

Key words: Chinook, *Oncorhynchus tshawytscha*, escapement, escapement goals, Taku River, Stikine River, Alsek River, Chilkat River, Unuk River, Chickamin River, Blossom River, Keta River, King Salmon River, Situk River, Andrew Creek, U.S./Canada Treaty, transboundary rivers

## INTRODUCTION

Chinook salmon *Oncorhynchus tshawytscha* are known to occur in 34 rivers in, or draining into, the Southeast Region of Alaska from British Columbia or Yukon Territory, Canada (Kissner 1978). In the mid-1970s it became apparent that many of the Chinook salmon stocks in this region were depressed relative to historical levels of production (Kissner 1974), and a fisheries management program was implemented to rebuild stocks in Southeast Alaska streams and in transboundary rivers (rivers that originate in Canada and flow into Southeast Alaska coastal waters; ADF&G 1981). Initially, this management program closed commercial and recreational fisheries in terminal and near-terminal areas in U.S. waters.

In 1981, this program was formalized and expanded to a 15-year (roughly 3 life-cycles) rebuilding program for the transboundary Taku, Stikine, Alsek, Unuk, Chickamin, and Chilkat rivers and the non-transboundary Blossom, Keta, Situk, and King Salmon rivers (ADF&G 1981) (Figure 1). The program used regionwide, all-gear catch ceilings for Chinook salmon, designed to rebuild spawning escapements by 1995 (ADF&G 1981). In 1985, the Alaskan program was incorporated into a comprehensive coastwide rebuilding program for all wild stocks of Chinook salmon, under the auspices of the U.S./Canada Pacific Salmon Treaty (PST).

To track the spawning escapement, the Alaska Department of Fish and Game (ADF&G), the Canadian Department of Fisheries and Oceans (DFO), the Taku River Tlingit First Nation (TRTFN), and the Tahltan First Nation (TFN)

count spawning Chinook salmon in a designated set of 11 watersheds (Appendix A1). These streams were selected on the basis of their historical importance to fisheries, size of the population, geographic distribution, extent of the historical database, and ease of data collection. Counts from each of these streams are considered to be indicators of relative abundance, based on the assumption that counts are a relatively constant proportion of the annual escapement in an index area or watershed.

Programs to estimate total escapement and survey count expansion factors have been implemented for all 11 index stocks. Long-term annual programs are in place on the Situk, Chilkat, Taku, Stikine and Unuk rivers. Short-term (2–3 year) projects were used to estimate expansion factors for the other six systems. Estimates of escapement from these mark–recapture and weir studies are generally superior to expanded survey count estimates, and are preferentially employed whenever they are available.

This project obtained indices of spawner abundance for major Chinook salmon stocks in Southeast Alaska. Objectives for 2008 were to count large ( $\geq 660$  mm MEF, or ocean-age 3 and older) spawning Chinook salmon during the time of peak abundance in tributaries and mainstem areas of the Stikine, Taku, Alsek, Situk, Unuk, Chickamin, Keta, Blossom and King Salmon rivers and in Andrew Creek, and to compile and compare the indices to those from past years.

Escapement data are provided annually to the Joint Chinook Technical Committee (CTC) of the Pacific Salmon Commission (PSC), who use them to evaluate status of indicator stocks (PSC 1997).




Figure 1.—Location of selected Chinook salmon systems in Southeast Alaska, Yakutat, and transboundary rivers.

Estimates of the total escapement of large spawners for 6 stocks (Situk, Chilkat, Taku, Stikine, and King Salmon rivers and Andrew Creek) and index counts for the remaining 5 stocks are provided to the CTC to determine trends in escapement.

In addition to these applications, biological escapement goals (BEGs) (5 AAC 39.222) have been established for all 11 systems and fisheries are managed to achieve those escapement goal ranges.

### DESCRIPTION OF STUDY SITES

Many individual spawning areas are surveyed annually in a designated set of watersheds. Detailed descriptions and maps of these areas are

found in Mecum and Kissner (1989); locations and descriptions of the index areas are found in Appendices A2 and A3, and general descriptions of the watersheds are below.

*The Taku River* originates in northern British Columbia and flows into the ocean 48 km east of Juneau, Alaska. The Taku River drainage covers over 17,000 km<sup>2</sup>; average monthly flows range from 60 m<sup>3</sup>/sec in February to 1,097 m<sup>3</sup>/sec in June (Bigelow et al. 1995). Principal tributaries are the Sloko, Nakina, Sheslay, Inklin, and Nahlin rivers. The clearwater Nakina and Nahlin rivers contribute less than 25% of the total drainage discharge; most is from glacier-fed streams on the eastern slope of the Coast Range

of British Columbia. Upstream of the abandoned mining community of Tulsequah, British Columbia, the drainage remains in pristine condition, with very few mining, logging, or other development activities. The upper Taku River area is extremely remote, with no road access and few year-round residents. All of the important Chinook salmon spawning areas are in tributaries in the upper drainage in British Columbia.

Stock assessment of Chinook salmon has been conducted intermittently on the Taku River since the 1950s, and standardized helicopter surveys of the index areas have been conducted annually since 1973. Survey index areas include portions of the Nakina, Nahlin, Dudidontu, Tatsamenie, and Kowatua rivers. In addition, since 1973 the DFO, TRTFN, and ADF&G have operated a carcass collection weir below the major spawning area on the Nakina River, which provides an estimate of the age and size composition of the escapement. Mark-recapture experiments have provided annual independent estimates of total escapement since 1995 (McPherson et al. 1998a; Jones III et al. 2010).

*The Stikine River* originates in British Columbia and flows to the sea approximately 32 km south of Petersburg, Alaska. Its drainage covers about 52,000 km<sup>2</sup>, much of which is inaccessible to anadromous fish because of natural barriers and velocity blocks. The Stikine River's principal tributaries include the Tahltan, Chutine, Scud, Iskut, and Tuya rivers. The lower river and most tributaries are glacially occluded (e.g., Chutine, Scud, and Iskut rivers).

Only 2% of the Stikine River drainage is in Alaska (Beak Consultants Limited 1981), and the majority of the Chinook salmon spawning areas in the Stikine River are located in British Columbia, Canada, in the mainstem Tahltan and Little Tahltan rivers (including Beatty Creek). However, Andrew Creek, in the U.S. portion of the lower Stikine River, supports a significant run of Chinook salmon. The upper drainage of the Stikine is accessible via the Telegraph Creek Road.

Helicopter surveys of the Little Tahltan River index area have been conducted annually since 1975, and the DFO and TFN have operated a fish counting weir at the mouth of the Little Tahltan

River since 1985. Counts from the weir represent the total escapement to that tributary. Since 1996, mark-recapture experiments have provided independent estimates of total escapement to the Stikine River (Pahlke and Etherton 1997, 1999, 2000; Pahlke et al. 2000; Der Hovanisian et al. 2001; 2003-5; Richards et al. 2008, *in prep a-b*).

*Andrew Creek* flows into the lower Stikine River in Alaska, not far from the limit of tidal influence. The drainage covers about 200 km<sup>2</sup> with 2 main tributaries. Only a small portion of the North fork is accessible to salmon and most spawning occurs in the South fork. From 1976 to 1984, a weir was operated on Andrew Creek to provide brood stock for hatcheries. Foot, aerial fixed-wing and helicopter surveys to count Chinook salmon have been conducted annually since 1985. A weir was operated on Andrew Creek in 1997 and 1998.

*The Alsek River* originates in Yukon Territory, Canada, and flows in a southerly direction into the Gulf of Alaska approximately 75 km southeast of Yakutat, Alaska. Its largest tributaries are the Dezadeash and Tatshenshini rivers. The Alsek River drainage covers about 28,000 km<sup>2</sup> (Bigelow et al. 1995), but much of it, including the mainstem of the Alsek River itself, is inaccessible to anadromous salmonids because of velocity barriers. The significant spawning areas for Chinook salmon are found mostly in tributaries of the Tatshenshini River, including the Klukshu, Blanchard, and Takhanne rivers and in Village and Goat creeks. The Klukshu and upper Tatshenshini rivers are accessible by road near Dalton Post, Yukon Territory.

Counts of Chinook salmon have been collected on the Alsek River since 1962. Beginning in 1976, the DFO has operated a weir at the mouth of the Klukshu to count Chinook, sockeye *O. nerka*, and coho *O. kisutch* salmon. The count of Chinook salmon through the Klukshu River weir is used as the index for the Alsek River. Some aboriginal harvest takes place above the weir. Aerial surveys to count spawning Chinook salmon were conducted by ADF&G with a helicopter from 1981 to 1999. The escapement to the Klukshu River is difficult to count by aerial, boat or foot surveys because of deep pools and overhanging vegetation. However, surveys of the Klukshu River were conducted

periodically to provide some continuity in estimates in the event that funding for the weir is discontinued. The Blanchard and Takhanne rivers and Goat Creek, three smaller tributaries of the Tatshenshini River, are also surveyed annually, but are not used to index escapements. Mark–recapture studies were conducted during 1988–2004 to estimate the escapement of spawning Chinook salmon in the Alsek River and radiotelemetry studies were conducted in 1998 and 2002 to estimate the distribution of spawning Chinook salmon (Pahlke et al. 1999; Pahlke and Etherton 2001a-b, 2002; Pahlke and Waugh 2003, 2004, 2006).

The Unuk, Chickamin, Blossom, and Keta river drainages all feed into Behm Canal, a narrow passage of water east of Ketchikan, Alaska. Misty Fiords National Monument/Wilderness Area surrounds the eastern or “back” Behm Canal and includes the Boca de Quadra fjords. Many of the mainland rivers in the area support Chinook salmon; the Unuk, Chickamin, Blossom and Keta rivers are designated Chinook salmon escapement index systems.

*The Unuk River* originates in a glaciated area of British Columbia and flows 129 km to Burroughs Bay, 85 km northeast of Ketchikan, Alaska; only the lower 39 km of the river are in Alaska. The Unuk is a large braided, glacially occluded river with a drainage of approximately 3,885 km<sup>2</sup>. Most (approximately 85%) spawning occurs in tributaries of the Alaska portion of the river (Pahlke et al. 1996). The escapement index areas are all small clearwater tributaries: Eulachon River and Cripple, Genes Lake, Clear, Lake, and Kerr creeks. Cripple Creek and Genes Lake Creek cannot be surveyed by air because of heavy vegetation, so fish are counted by foot survey. Chinook salmon have been counted annually by foot or helicopter surveys in these areas since 1977. Chinook salmon have been periodically counted in Boundary Creek, but survey conditions there are often poor and the counts are not included in the index. Total escapement was estimated by a mark–recapture project in 1994 (Pahlke et al. 1996) and annually since 1997 (Jones III et al. 1998a; Jones III and McPherson 1999, 2000, 2002; Weller and McPherson 2003a-b, 2006a-b; Weller and Evans 2009; Weller et al. *In prep.*

*The Chickamin River* is a large, glacial river with a drainage of approximately 2,000 km<sup>2</sup>. It originates in British Columbia and flows into Behm Canal approximately 32 km southeast of Burroughs Bay and 65 km northeast of Ketchikan. Although it is technically a transboundary river, there are no Chinook spawning areas on the Chickamin River upstream from the Canadian border (Pahlke 1997a). Important spawning tributaries are the South Fork of the Chickamin and Barrier, Butler, Indian, Leduc, Humpy, King, and Clear Falls creeks. Chinook salmon have been counted by foot or helicopter surveys in index areas of the Chickamin River each year since 1975. Total escapement was estimated by mark–recapture projects in 1995, 1996 and 2001–2005, and spawning distribution was estimated by radiotelemetry in 1996 (Pahlke 1996, 1997a; Freeman and McPherson 2003–2005; Freeman et al. 2007; Weller et al. 2007b).

*The Blossom, Keta, Wilson, and Marten rivers* are clearwater rivers that flow into Behm Canal approximately 45 km east of Ketchikan. These rivers lie inside the boundaries of the Misty Fiords National Monument in southern Behm Canal but are within an area that has been specifically excluded from wilderness designation because of the potential development of a large-scale molybdenum mine (Quartz Hill) near the divide of the Blossom and Keta rivers. The mine is presently undeveloped, but an access road has been completed that terminates at salt water near the mouth of the Blossom River.

The Keta River drainage covers about 192 km<sup>2</sup> and the Blossom about 176 km<sup>2</sup> (Bigelow et al. 1995) and have been surveyed by helicopter annually since 1975. Chinook salmon escapements to the Wilson and Marten rivers have been monitored on an intermittent basis in recent years. Mark–recapture experiments were conducted in 1998 to estimate the escapement of Chinook salmon in the Blossom and Keta rivers (Brownlee et al. 1999) and were repeated on the Keta River in 1999 and 2000 (Freeman et al. 2000, 2001) and on the Blossom from 2004 to 2006 (Pahlke and Magnus 2005, 2006; Weller et al. 2007a).

*The King Salmon River* drains an area of approximately 100 km<sup>2</sup> on Admiralty Island, flowing into King Salmon Bay on the eastern side

of Stephens Passage about 48 km south of Juneau. The King Salmon River is the only island river system in Southeast Alaska to support more than 100 spawning Chinook salmon. ADF&G operated a weir on the King Salmon River from 1983 through 1992 to count Chinook salmon and collect broodstock for Snettisham Hatchery. Helicopter surveys have been conducted annually since 1975 and foot surveys since 1992.

*The Chilkat River* is a large glacial river which originates in Yukon Territory, Canada, and flows into Chilkat Inlet at the head of northern Lynn Canal near Haines, Alaska. The basin encompasses an area approximately 2,600 km<sup>2</sup> (Bugliosi 1988), and 1,667 km<sup>2</sup> are considered accessible to anadromous fish (Ericksen and McPherson 2004). Helicopter and foot surveys are an ineffective index of abundance for this system (Johnson et al. 1992) and were suspended in 1993 in favor of annual estimates of escapement using mark–recapture methods. Total escapement has been estimated annually since 1991 (Ericksen 2005; Ericksen and Chapell 2006; Chapell *In prep a-b*).

*The Situk River* is a small drainage (176 km<sup>2</sup>) located about 16 km east of Yakutat, Alaska. The Situk supports a large run of sockeye salmon that are harvested in commercial and subsistence set gillnet fisheries concentrated at the mouth of the Situk River. Situk River Chinook salmon are harvested both incidentally and targeted in the set gillnet fisheries, depending on run strength, and in a recreational fishery in the river. A weir was operated on the Situk River at the upper limit of the intertidal area from 1928 to 1955 to count all 5 species of Pacific salmon spawning in the river. Since 1976, a weir has been operated primarily to count Chinook and sockeye salmon. The proportion of the recreational harvest above the weir varies from year-to-year (Howe et al. 2001).

## METHODS

There are 34 river systems in the region (Figure 1) with populations of wild Chinook salmon. Three transboundary rivers, the Taku, Stikine, and Alek, are classed as major producers, each with potential production (harvest plus escapement) greater than 10,000 fish (Kissner 1974). Nine rivers are classed as medium producers, each with production of 1,500 to 10,000 fish. The remaining

22 rivers are minor producers, with production less than 1,500 fish. Small numbers of Chinook salmon occur in other streams of the region but they are not included in the above list because successful spawning has not been documented. Chinook salmon are counted via aerial surveys or at weirs each year in all three major producing systems, in six of the medium producers, and in one minor producer (Appendices A4 and A5). Abundance in the Chilkat River is estimated only by a mark–recapture program. These index systems, along with the Chilkat River, are believed to account for about 90% of the total Chinook salmon escapement in Southeast Alaska and transboundary rivers (Pahlke 1998).

## ESCAPEMENT GOALS

The initial rebuilding program established interim escapement goals in 1981 for 9 systems: the Alek, Taku, Stikine, Situk, King Salmon, Unuk, Chickamin, Keta and Blossom/Wilson rivers. Although the aim was to have escapement goals that provided the optimal level of harvest, little data were available to produce such goals. As a result, escapement goals were originally set based on the highest observed escapement count prior to 1981 (Pahlke 1997b). Goals for the Chilkat River and Andrew Creek were added in 1985, bringing the total number of regularly monitored river systems to 11. Pahlke (1997b) provides detailed descriptions of the escapement goals and their origins. Escapement goals have been revised when sufficient new information warrants. Most of the revised escapement goals have been developed with spawner-recruit analysis as ranges of optimum escapement rather than a single point estimate (Appendix A1). Spawner-recruit analysis requires not only a long series of escapement estimates, but also annual age and sex-specific estimates of escapement (McPherson and Carlile 1997). The United States Section of the CTC developed data standards in 1997 for stock specific assessments of escapement, terminal runs, and forecasts of abundance that are used to evaluate existing stock assessment programs (PSC 1997). One of those standards is the collection of annual age and sex-specific estimates of total escapement. These data have been collected routinely at weirs and during mark–recapture studies and recently specific programs have been

implemented to collect age, sex and length data from Chinook salmon in the Blossom, Keta, and King Salmon rivers and Andrew Creek.

## INDICES OF ESCAPEMENT

Spawning Chinook salmon are counted at 26 designated index areas in 9 of the systems (Appendix A3); total escapement in the other two systems are estimated by complete counts of Chinook salmon at the Situk River weir and by annual mark-recapture estimates on the Chilkat River. Counts are made during aerial or foot surveys during periods of peak spawning, or at weirs. Peak spawning times, defined as the period when the largest number of adult Chinook salmon actively spawn in a particular stream or river, have been well-documented from surveys of these index areas conducted since 1976 (Kissner 1982; Pahlke 1997b). The proportion of fish in pre-spawning, spawning and post-spawning condition is used to judge whether the survey timing is correct to encompass peak spawning. Index areas are surveyed at least twice unless turbid water or unsafe conditions preclude the second survey. Survey conditions during each index survey are rated as poor, normal or excellent for that particular index area, and coded as to whether that survey is potentially useful for indexing or estimating escapement. Factors that affect the rating include water level, clarity, light conditions, and weather.

Only large Chinook salmon  $\geq 660$  mm MEF are counted during aerial or foot surveys. No attempt is made to accurately count Chinook salmon  $< 660$  mm MEF (typically age-.1 and -.2; Mecum 1990). These Chinook salmon, also called jacks, are early maturing, precocious males. They are distinct from their older age counterparts under most conditions because of their short, compact bodies and lighter color. They are, however, difficult to distinguish from other smaller species such as pink *O. gorbuscha* and sockeye salmon. In some systems age-1.2 fish may be larger than 660 mm MEF and be difficult to avoid counting.

Aerial surveys are conducted from a Bell 206 or Hughes 500D helicopter. Pilots are directed to fly the helicopter from 6 to 15 m above the river bed at a speed of 6–16 km/h. The helicopter door on the side of the observer is removed, and the

helicopter is flown sideways while observations of spawning Chinook salmon are made. Foot surveys are conducted by at least 2 people walking in the creek bed or on the riverbank.

Weather, distances involved, run timing, etc., can make it difficult for a single surveyor to complete all the index surveys annually under normal or excellent conditions. Thus, alternate surveyors are selected to conduct the counts when the primary surveyor is unavailable. Also, new surveyors take on primary responsibilities at infrequent intervals. Because between-observer variability and bias can be significant (Jones III et al. 1998b), new surveyors must be trained and calibrated against the primary surveyor to provide consistency and continuity in the data. Alternate observers accompany the primary observer on regularly scheduled surveys to learn survey methods and counting techniques (back seat, training flights). Each alternate observer also accompanies the primary observer on additional regularly scheduled surveys to independently count Chinook salmon (replicate, calibration flights). Each calibration flight consists of 2 passes over the index area so the two observers in turn sit in the preferred location in the helicopter during 1 pass along the river. Counts are not shared during the calibration surveys, but are shared and discussed following the completion of the second pass of each flight. Calibration data will be collected annually for several years. The relationship between observer escapement counts will be determined from accumulated data and applied to counts.

Several index areas are routinely surveyed by more than one method; e.g. Andrew Creek is surveyed from airplanes, helicopters and by foot. The various surveys are conducted as close as possible to each other to promote comparison and calibration of the different methods.

Counts and other observations from the 2008 surveys (Appendix A6) are entered into the ADF&G Division of Commercial Fisheries Integrated Fisheries Database (IFDB) in Juneau for archiving and general distribution.

Estimates of total escapement are needed to model total production, exploitation rates and other population parameters. Counts from index areas

are increased by an expansion factor (Table 1). An expansion factor is an estimate of the proportion of the total escapement counted in a river system during the peak spawning period. Expansion factors are based on comparisons with weir counts, mark–recapture estimates, and spawning distribution studies. They vary among rivers according to how complete the coverage of spawning areas is and difficulties encountered in observing spawners, such as overhanging vegetation, turbid water conditions, presence of other salmon species (i.e., pink and chum *O. keta* salmon), or protraction of run timing. Expansion factors range from 1.5 for the King Salmon River to 5.2 for the Taku River (Table 1).

Escapement counts are obtained from a fish-counting weir on the Situk River and a mark–recapture program on the Chilkat River. Survey expansions are not necessary for those streams where weirs or other estimation programs are used to count all migrating Chinook salmon.

Finally, to estimate total regional escapement, escapement estimates from the 11 index systems are expanded to account for the unsurveyed systems (Appendices A4 and A5). The total estimated escapement in the index areas represents approximately 90% of the region total (Pahlke 1998). Escapement estimates for the Chilkat River are not available prior to 1991. From 1991 to 1997, the estimated escapement to the Chilkat River averaged 6% of the estimated regionwide total. Therefore, prior to 1991 the expanded index counts represent approximately 84% of the estimated Southeast Alaska total escapement.

Expansion factors for individual rivers in this report have been revised, based on results from experiments to date that estimate total escapement and spawning distribution. For example, estimated total escapement and radiotracking distribution data were used to revise tributary expansion factors for the Taku and Unuk rivers (Pahlke and Bernard 1996; Pahlke et al. 1996; McPherson et al. 1998a). Mark–recapture studies to estimate spawning abundance on the Unuk River in 1994 (Pahlke et al. 1996) and on the Chickamin River

in 1995 and 1996 (Pahlke 1996, 1997a) were used to revise expansion factors for those two rivers in 1996; results were also applied to the nearby Blossom and Keta rivers. More mark–recapture studies were conducted on all four rivers and the expansion factors for the Behm Canal systems were revised again (Pahlke 2007). On Andrew Creek, a weir was operated over 4 years (1979, 1981, 1982, and 1984), during which index counts were also made, establishing a new expansion factor for that system in 1995. Also in 1997, 10 years (1983–1992) of matched weir and index counts were used to revise the expansion factor for the King Salmon River (McPherson and Clark 2001). The expansion factors for the Taku River were revised in 1996 and again in 1999 based on the results of mark–recapture studies (Pahlke and Bernard 1996; McPherson et al. 2000).

These studies have helped to estimate total escapement in the region and have shown that, in most cases, the surveyed index area counts are reasonably accurate in assessing trends in escapements. However, Johnson et al. (1992) demonstrated that expansion factors used before 1991 on the Chilkat River system were highly inaccurate because the index areas received less than 5% of the escapement. Consequently, since 1991, escapement to the Chilkat River has been estimated annually by mark–recapture experiments (Ericksen 2005). Studies on the Taku, Stikine, Alsek, Unuk, Chickamin, Blossom, Keta and King Salmon rivers, as well as on Andrew Creek, have shown that the index expansion factors used on those systems were much more accurate than those used on the Chilkat (PSC 1991; Pahlke 1996, 1997a). Expansion factors will continue to be revised as additional data become available (Appendix B1). Ongoing research projects should provide more information on the expansion factors for the Taku, Stikine, and Unuk rivers. Estimates of escapement from expanded counts are included in this document to provide relative estimates of total spawner abundance over time, with the caveat that expansion factors may produce incorrect estimates or be revised in the future.

Table 1.–Peak survey counts, survey expansion factors, estimated total escapement from expanded survey counts, mark–recapture projects or weir, for large Chinook salmon returning to Southeast Alaska and transboundary rivers in 2008.

Survey area	Survey count	Survey expansion factor (SE)	Survey expansion estimated escapement <sup>a</sup>	Estimated total escapement <b>M–R or weir<sup>b</sup></b> (SE)	Reference <sup>c</sup>
Major producers					
Alsek River	Klukshu weir	465	4.03 (1.71)	1,339 <sup>d</sup>	1,339 <sup>d</sup> (548)
Taku River	5 tributaries	4,817	5.20 (1.78)	25,048	<b>27,383</b> , (2,454) Jones III et al. (2010)
Stikine River	Little Tahltan weir	2,657	5.36 (1.35)	14,242	<b>18,843</b> (3,153) Richards et al. <i>In prep b</i>
Medium producers					
Situk River	NA	NA	NA	NA	<b>453<sup>e</sup></b>
Chilkat River	NA	NA	NA	NA	<b>2,833</b> (530) Chapell <i>In prep b</i>
Andrew Cr.	All	503	1.95 (0.45)		981 (225)
Unuk River	6 tributaries		4.87 (0.60)	incomplete counts	<b>3,104</b> (390) Weller et al. <i>In prep b</i>
Chickamin River	8 tributaries	1,111	4.75 (0.70)	5,277	5,277 (777) Johnson et al. (2009)
Blossom River	All	257	3.01 (1.03)	773	773 (265)
Keta River	All	363	3.01 (0.56)	1,093	1,093 (203)
Minor producers					
King Salmon River	All	79	1.52 (0.27)	120	120 (21)
Index system total					M–R plus survey expansions
Region total <sup>f</sup>			1/0.9	62,198	
				69,109	

<sup>a</sup> Estimated by multiplying survey count by expansion factor, Appendix B.

<sup>b</sup> Estimated from mark–recapture program or weir count in bold, others expanded survey counts. Final numbers used for ADF&G management.

<sup>c</sup> Reference document for mark–recapture estimate.

<sup>d</sup> Alsek River escapement estimated as per Pahlke and Waugh (2006).

<sup>e</sup> Situk River weir count, minus estimated sport harvest above weir (zero in 2008).

<sup>f</sup> Index count total estimated to represent 90% of regionwide total.

## AGE, SEX, AND LENGTH COMPOSITION OF ESCAPEMENTS

Estimates of escapement by age and sex for all 11 systems having Chinook salmon stock assessment projects in Southeast Alaska in 2008 were compiled to provide a basic statistical summary for managers and researchers. Estimates for the Unuk, Stikine, Taku, and Chilkat rivers were the results of mark–recapture experiments (Weller et al. *In prep*; Richards et al. *In prep b*; Chapell *In prep b*; Jones III et al. 2010). Results compiled from each of these projects are the reported unbiased estimates of escapement of medium- and large-sized Chinook salmon, except for the Stikine River, where the unbiased estimates include small fish. Size classification of small and medium fish varies slightly between projects. Estimates for medium and large fish from the Situk River are based on age sampling and a total census of the escapement at a weir. Age composition estimates for the Keta and King Salmon rivers and Andrew Creek were calculated by dividing the peak survey count by the escapement expansion factor (Table 1), and multiplying the result by the age composition of the escapement sampled on the spawning grounds of each drainage in 2008. Standard errors include variance of the estimated escapements and proportions by age from sampling (Pahlke and Waugh 2003). Note that the survey index counts for the Blossom and Keta rivers include many age-1.2 Chinook salmon because of their large size at age (65% to 75% of age-1.2 fish in these systems are  $\geq 660$  mm MEF), which makes them part of the large-fish population counted in surveys. All fish in the medium and large-size categories sampled on the spawning grounds and aged (most are age-1.2 and older) are used in the calculations. Also note that there may be slight biases for some systems without mark–recapture estimates in 2008; however, we have employed sampling gear to minimize size- or sex-selective sampling in these spawning ground samples. The estimates for systems with mark–recapture or weir (Situk) projects are the result of batteries of tests and stratification to produce unbiased estimates of age and sex structure.

Estimates of mean length by sex and age and their estimated variances were also calculated for each system. These estimates are either the unbiased estimates reported in the publications cited above, or made using the spawning ground samples as noted above.

All Chinook salmon sampled for age, sex and length data were also examined for missing adipose fins, which may indicate the presence of a coded wire tag (CWT). In most cases fish with missing adipose fins were sacrificed to recover the tag. On the Taku, Chilkat, Stikine, Chickamin and Unuk rivers, most of the CWT fish were wild fish tagged earlier in those rivers during ongoing projects. Other tags were recovered from both non-natal wild and hatchery stocks.

## RESULTS

In 2008, 34 locations, 26 of which were designated index areas, were surveyed specifically for Chinook salmon escapement (Appendix A3). Surveys generally progressed as planned with the exception of the Unuk River, where bad weather prevented surveys on several important index areas.

The estimated escapement of large Chinook salmon for all Southeast Alaska and transboundary rivers in 2008 was 69,109 (Table 1), a 21% increase from the estimated 57,244 fish in 2007. Escapement indices for 10 of 11 index areas were within or above escapement goal ranges.

From 1984 to 1993, the estimated escapement of Chinook salmon in Southeast Alaska increased, peaking in 1993 (Appendix A4). This was primarily attributed to strong returns to the Taku, Stikine, and Chilkat rivers, which together make up over 75% of the summed escapement goals in the region. Escapements declined in 1994 and 1995 and then peaked again in 1996 and 1997 as a result of record high escapements in the Taku River. In 1998 and 1999, escapements to the Taku River declined dramatically and since then have remained below the 1990–1999 average, but still well above the escapement goal. Escapement to the Stikine River increased greatly since 1999, including the highest on record in 2001.

## TAKU RIVER

Survey conditions in 2008 were normal on most of the index areas of the Taku River. The counts of large Chinook salmon in 3 of 5 index areas of the Taku River were below recent 10-year averages (Tables 2 and 3). Counts increased from 1983 to 1993, and exceeded the upper limit of the survey goal range 5 times in the 1990s (Figure 2). Expansion of the 2008 survey counts of 4,817 large Chinook salmon by the survey expansion factor of 5.20 (SE = 1.78; Appendix B9) produced a total Taku River escapement estimate of 25,048 (SE = 8,574, Table 1). A mark-recapture experiment conducted in 2008 resulted in a escapement estimate 27,383 large; (SE = 2,454; Jones III et al. 2010). The expansion factor was revised in 1999 based on 5 years of mark-recapture experiments on the Taku River (Appendix B9; McPherson et al. 2000). McPherson et al. (2000) recommended an escapement goal range of 30,000 to 55,000 large spawners. These changes were adopted by the Transboundary River Technical Committee (TTC) and the CTC of the PSC. The revised PSC goal uses counts in 5 index areas expanded by 5.2 (SE 1.78) which corresponds to an index goal range of 5,800 to 10,600 fish.

Commercial fisheries targeting Taku River Chinook salmon were opened in 2005 and 2006 for the first time in 27 years. The combined U.S. and Canadian fleets harvested about 28,000 fish in 2005, and about 21,000 Chinook salmon in 2006. The preseason forecast for 2008 was below the threshold agreed to by the U.S. and Canada in 2005 to allow targeted Chinook fisheries.

Age, sex and length data were collected from carcasses at the Nahlin and Tatsamenie rivers, and live fish were sampled with angling gear at Nahlin and Tatsamenie rivers (Appendices A7, panel H and A8, panel H).

## STIKINE RIVER

In 2008, 2,657 large Chinook salmon were counted at the Little Tahltan River weir. The weir count was about 5 times the record low count of 562 in 2007, but still below the 10-year average (Table 4).

Surveys of the Little Tahltan River have continued in order to maintain the time series of data and to train surveyors. The peak aerial survey above the Little Tahltan River weir was 837 large fish in 2008. Survey conditions were normal, and the count was also below average. From 1985 to 2005, the proportion of the total escapement of Chinook salmon counted during peak aerial surveys has ranged from 27.9% to 56.6% and averaged 34.8% during 1997–2007 (Table 4). The proportion of the total escapement observed in a single survey often declined after the peak of spawning as fish died or were removed by predators. In 1998, 1999, 2003, and 2005, survey conditions were not unusual and there is no explanation for the lower than average proportion of escapement observed. Age, sex and length data were collected from 327 fish sampled at the Little Tahltan River weir and Verrett Creek (Appendices A7, panel E and A8, panel E).

Based on a stock-recruit model, the BEG was revised in 1999 to a range of 14,000 to 28,000 large Chinook total in the Stikine River drainage, or 2,700 to 5,300 at the Little Tahltan weir (Bernard et al. 2000). The 2008 weir count was below the escapement goal range for the second time since the weir was installed in 1985 (Figure 3). The expansion factor includes the annual estimates through 2005. Expansion of the 2008 weir count of 2,657 large Chinook salmon by the survey expansion factor of 5.36 (SE = 1.35; Appendix B6) produced a total Stikine River escapement estimate of 14,242 (SE = 3,587, Table 1) large Chinook salmon. The estimate of total escapement to the Stikine River from a mark-recapture experiment conducted in 2008 is 18,843 large Chinook (SE = 3,153; Richards et al. *In prep b*), which is within the escapement goal range for the drainage.

Commercial fisheries targeting Stikine River Chinook salmon were opened in 2005 for the first time in 27 years (Richards et al. 2008) and again in 2006 and 2007. The combined U.S. and Canadian fleets harvested about 50,000 fish in 2005, 44,000 in 2006, and 25,000 in both 2007 and 2008.




Figure 2.—Counts of Chinook salmon in index areas of the Taku River, 1975–2008 and mark–recapture estimates divided by expansion factor of 5.2. Lines show upper and lower limits of index escapement goal range.

Table 2.—Distribution of spawning Chinook salmon among index areas of the Taku River during years when all index areas were surveyed.

Year	Nakina River	%	Nahlin River	%	Kowatua River	%	Tatsamenie River	%	Dudidontu River	%	Tseta Creek	%	Total
1981	5,110	52	2,945	30	560	6	839	9	74	1	258	3	9,786
1982	2,533	53	1,246	26	289	6	387	8	130	3	228	5	4,813
1983	968	47	391	19	171	8	236	11	117	6	179	9	2,062
1985	2,647	37	2,236	31	699	10	848	12	475	7	303	4	7,208
1986	3,868	51	1,612	21	548	7	886	12	413	5	193	3	7,520
1987	2,906	51	1,122	20	570	10	678	12	287	5	180	3	5,743
1988	4,500	52	1,535	18	1,010	12	1,272	15	243	3	66	1	8,626
1989	5,141	54	1,812	19	601	6	1,228	13	204	2	494	5	9,480
1990	7,917	65	1,658	14	614	5	1,068	9	820	7	172	1	12,249
1991	5,610	55	1,781	18	570	6	1,164	11	804	8	224	2	10,153
1992	5,750	52	1,821	16	782	7	1,624	15	768	7	313	3	11,058
1993	6,490	49	2,128	16	1,584	12	1,491	11	1,020	8	497	4	13,210
1994	4,792	48	2,418	24	410	4	1,106	11	573	6	614	6	9,913
1995	3,943	45	2,069	24	550	6	678	8	731	8	786	9	8,757
1996	7,720	39	5,415	27	1,620	8	2,011	10	1,810	9	1,201	6	19,777
1997	6,095	44	3,655	26	1,360	10	1,148	8	943	7	648	5	13,849
1998	2,720	43	1,294	20	473	7	675	11	807	13	360	6	6,329
1999	1,900	46	532	13	561	13	431	10	527	13	221	5	4,172
2000	2,907	49	728	12	702	12	953	16	482	8	160	3	5,932
2001	1,552	30	935	18	1,050	20	1,024	20	479	9	202	4	5,242
2002	4,066	49	1,099	13	945	11	1,145	14	834	10	192	2	8,281
2003	2,126	36	861	15	850	14	1,000	17	644	11	436	7	5,917
2004	4,091	41	1,787	18	828	8	1,396	14	1,036	10	906	9	10,044
2005	1,213	29	471	11	833	20	1,146	27	318	8	215	5	4,196
2006	1,900	34	955	17	1,180	21	908	16	395	7	199	4	5,537
2007	-	-	277	-	262	-	390	-	4	-	-	-	-
Average	3,790	45	1,646	20	755	11	990	14	575	7	370	5	8,110
2008	1,437	27	1,185	22	632	12	1,083	20	480	9	497	9	5,314

Table 3.—Counts of spawning Chinook salmon in index areas of the Taku River, 1973–2008.

Year <sup>a</sup>	Nakina River		Nahlin River		Kowatua River		Tatsamenie River		Dudidontu River		5 trib. total	Tseta Creek <sup>b</sup>
1973	2,000	N(H)	300	E(H)	100	N(H)	200	E(H)	200	E(H)	2,800	4 (A)
1974	1,800	E(H)	900	E(H)	235	(A)	120	(A)	24	(A)	3,079	4 (A)
1975	1,800	E(H)	274	E(H)	–	–	–	–	15	N(H)	2,089	–
1976	3,000	E(H)	725	E(H)	341	P(A)	620	E(H)	40	(H)	4,726	–
1977	3,850	E(H)	650	E(H)	580	E(A)	573	E(H)	18	(H)	5,671	–
1978	1,620	E(H)	624	E(H)	490	N(H)	550	E(H)	–	–	3,284	21 E(H)
1979	2,110	E(H)	857	E(H)	430	N(H)	750	E(H)	9	E(H)	4,156	–
1980	4,500	E(H)	1,531	E(H)	450	N(H)	905	E(H)	158	E(H)	7,544	–
1981	5,110	E(H)	2,945	E(H)	560	N(H)	839	E(H)	74	N(H)	9,528	258 N(H)
1982	2,533	E(H)	1,246	E(H)	289	N(H)	387	N(H)	130	N(H)	4,585	228 N(H)
1983	968	E(H)	391	N(H)	171	E(H)	236	E(H)	117	E(H)	1,883	179 N(H)
1984 <sup>c</sup>	1,887	(H)	951	(H)	279	E(H)	616	E(H)	–	–	3,733	176 (H)
1985	2,647	N(H)	2,236	E(H)	699	E(H)	848	E(H)	475	(H)	6,905	303 E(H)
1986	3,868	(H)	1,612	E(H)	548	E(H)	886	E(H)	413	E(H)	7,327	193 E(H)
1987	2,906	E(H)	1,122	E(H)	570	E(H)	678	E(H)	287	E(H)	5,563	180 E(H)
1988	4,500	E(H)	1,535	E(H)	1,010	E(H)	1,272	E(H)	243	E(H)	8,560	66 E(H)
1989	5,141	E(H)	1,812	E(H)	601 <sup>d</sup>	(W)	1,228	E(H)	204	E(H)	8,986	494 E(H)
1990	7,917	E(H)	1,658	E(H)	614 <sup>d</sup>	(W)	1,068	N(H)	820	E(H)	12,077	172 N(H)
1991	5,610	E(H)	1,781	E(H)	570	N(H)	1,164	E(H)	804	E(H)	9,929	224 N(H)
1992	5,750	E(H)	1,821	E(H)	782	E(H)	1,624	N(H)	768	N(H)	10,745	313 N(H)
1993	6,490	E(H)	2,128	N(H)	1,584	E(H)	1,491	E(H)	1,020	E(H)	12,713	491 N(H)
1994	4,792	N(H)	2,418	E(H)	410	P(H)	1,106	N(H)	573	N(H)	9,299	614 E(H)
1995	3,943	E(H)	2,069	E(H)	550	N(H)	678	N(H)	731	E(H)	7,971	786 E(H)
1996	7,720	E(H)	5,415	E(H)	1,620	N(H)	2,011	N(H)	1,810	N(H)	18,576	1,201 N(H)
1997	6,095	E(H)	3,655	E(H)	1,360	N(H)	1,148	N(H)	943	N(H)	13,201	648 N(H)
1998	2,720	E(H)	1,294	N(H)	473	N(H)	675	E(H)	807	E(H)	5,969	360 E(H)
1999	1,900	N(H)	532	N(H)	561	E(H)	431	N(H)	527	E(H)	3,951	221 N(H)
2000	2,907	N(H)	728	P(H)	702	N(H)	953	N(H)	482	N(H)	5,772	160 N(H)
2001	1,552	P(H)	935	N(H)	1,050	N(H)	1,024	N(H)	479	N(H)	5,040	202 N(H)
2002	4,066	E(H)	1,099	N(H)	945	N(H)	1,145	N(H)	834	N(H)	8,089	192 N(H)
2003	2,126	N(H)	861	E(H)	850	E(H)	1,000	N(H)	644	E(H)	5,481	436 N(H)
2004	4,091	N(H)	1,787	N(H)	828	N(H)	1,396	N(H)	1,036	N(H)	9,138	906 N(H)
2005	1,213	N(H)	471	P(H)	833	E(H)	1,146	N(H)	318	N(H)	3,981	215 N(H)
2006	1,900	N(H)	955	N(H)	1,180	N(H)	908	N(H)	395	N(H)	5,338	199 P(H)
2007	-	none	277	P(H)	262	E(H)	390	N(H)	4	P(H)	-	- none
2008	1,437	N(H)	1,185	N(H)	632	E(H)	1,083	N(H)	480	N(H)	4,817	497 N(H)
98–07	2,497		984		768		907		553		5,377	321
Avg.												

Note: (F) = foot survey; – = no survey conducted; (A) = fixed-wing aircraft; (H) = helicopter; (W) = weir; P = poor survey conditions hampered by glacial or turbid waters; N = normal survey conditions; E = conditions excellent.

<sup>a</sup> Counts before 1975 may not be comparable due to changes in methods; foot surveys may include jacks.

<sup>b</sup> Tseta Creek removed from index areas in 1999.

<sup>c</sup> Surveys in 1984 conducted by DFO; partial survey of Tseta Creek and Nahlin River.

<sup>d</sup> Carcass weir at Kowatua River used to partially count escapement due to poor survey conditions, 1989, 1990.

Table 4.—Counts of large spawning Chinook salmon in the Little Tahltan River, Stikine River, 1975–2008.

Year	Weir count	Above weir catch	Spawning escapement	Aerial survey	
				Peak count <sup>a</sup>	Percent counted
1975	-			700	E(H)
1976	-			400	N(H)
1977	-			800	P(H)
1978	-			632	E(H)
1979	-			1,166	E(H)
1980	-			2,137	N(H)
1981	-			3,334	E(H)
1982	-			2,830	N(H)
1983	-			594	E(H)
1984	-			1,294	E(H)
1985	3,114	0	3,114	1,598	E(H) 51.3
1986	2,891	0	2,891	1,201	E(H) 41.5
1987	4,783	0	4,783	2,706	E(H) 56.6
1988	7,292	0	7,292	3,796	E(H) 52.1
1989	4,715	0	4,715	2,527	E(H) 53.6
1990	4,392	0	4,392	1,755	E(H) 40.0
1991	4,506	0	4,506	1,768	E(H) 39.2
1992	6,627	12	6,615	3,607	E(H) 54.4
1993	11,449	12	11,437	4,010	P(H) 35.1
1994	6,387	14	6,373	2,422	N(H) 38.0
1995	3,072	0	3,072	1,117	N(H) 36.4
1996	4,821	0	4,821	1,920	N(H) 39.8
1997	5,557	10	5,547	1,907	N(H) 34.4
1998	4,879	6	4,873	1,385	N(H) 28.4
1999	4,738	5	4,733	1,379	N(H) 27.9
2000	6,640	9	6,631	2,720	N(H) 41.0
2001	9,738	0	9,730	4,158	N(H) 42.7
2002	7,490	14	7,476	No survey	
2003	6,492	0	6,492	1,903	N(H) 29.3
2004	16,381	0	16,381	6,014	E(H) 36.7
2005	7,253	0	7,253	2,157	N(H) 29.7
2006	3,845	0	3,845	1,372	N(H) 35.7
2007	562	0	562	213	P(H) 34.9
98–07	6,803	2	6,803	2,367	34.8
Avg.					
2008	2,657	0	2,657	837	N(H) 31.5

Note: N = normal survey conditions; P = poor survey conditions; E = excellent survey conditions, H = helicopter

<sup>a</sup> Peak count equals peak survey above weir plus count below weir on that date.

## ANDREW CREEK

The 2008 survey count of Chinook salmon in Andrew Creek was 503 fish, compared to 890 in 2007 (Table 5). In 1998, a spawner-recruit analysis was completed and a biological escapement goal range of 650 to 1,500 total (325–750 index count) large spawners was adopted (Clark et al. 1998). Since 1985, Andrew Creek escapements have exceeded the lower limit of the goal in all but 2 years (Figure 4).

From 1976 to 1984 a weir was operated on Andrew Creek to provide broodstock for hatcheries (Pahlke 1995). Surveys were also conducted on the system during 4 of those years and, on the basis of those paired counts, the survey expansion factor was revised in 1995 from 1.6 to 2.0 (SE = 0.409). A weir was operated and surveys were also conducted in 1997 and the expansion factor was revised again to 1.95 (SE = 0.45; Appendix B7). No survey expansion was necessary for the years when the weir provided total escapement counts (Appendix A4).

Five surveys were conducted between 31 July and 12 August 2008. The peak count was by helicopter on 7 August. The count of 503 was used as the peak count based on experience of the surveyors and what was most representative of normal survey conditions. Expansion of the aerial count of 503 large Chinook salmon by the survey expansion factor (1.95) produced a total Andrew Creek escapement estimate of 981 (SE = 225) large Chinook salmon (Table 1; Appendix B7).

Age, sex, and length data was collected from 50 pre-spawning fish in Andrew Creek, using angling gear and dip nets, 39 of which were successfully aged (Appendices A7, panel F and A8, panel F).

## ALSEK RIVER

The count of large Chinook salmon through the Klukshu River weir in 2008 was 465 fish, down from the count of 675 in 2007 (Table 6; Figure 5). The escapement to the Klukshu River, estimated by subtracting the aboriginal fishery harvest (0) and sport harvest (0) above the weir from the weir count, was 465 fish. This was below the escapement goal range of 1,100 to 2,300, for the third year in a row (McPherson et al. 1998b). All the sport and aboriginal harvest in 2008 was below the weir.

No aerial survey of the Klukshu River was conducted in 2008. However, in helicopter surveys 41 large Chinook salmon were counted in the Takhanne River, 11 in Goat Creek, and 65 in the Blanchard River. There is no agreement in the PSC on use of expansion factors for the Alsek River; expansion factors used in the past have ranged from 1.56 to 2.5, based on assumptions that the Klukshu River represented 40 to 64 percent of the escapement to the entire drainage (Pahlke 1997b).




Figure 3.—Counts of Chinook salmon at the Little Tahltan River weir, Stikine River, 1975–2008, and mark–recapture estimates divided by expansion factor of 5.36. Data for 1985–2000 from weir counts, 1975–1984 estimated by doubling index count. Lines show upper and lower limits of escapement goal range.




Figure 4.—Counts of Chinook salmon at the Andrew Creek weir, 1976–1984, 1997, and in aerial/foot surveys, 1975, 1985–2008. Lines show upper and lower bounds of index escapement goal range.

Table 5.—Counts of spawning Chinook salmon in selected rivers in central Southeast Alaska, 1960–2008.

Year	Andrew Creek <sup>a</sup>		North Arm		Clear Creek		Harding River		Aaron Creek		Bradfield River	
											North Fork	East Fork
1960	287	(F)	200	(F)N	—	—	—	—	—	—	—	—
1961	103	(F)	138	(F)	—	—	—	—	—	—	—	—
1962	300	(A)	80	(A)N	—	—	—	—	—	—	—	—
1963	500	(A/H)	187	(F)	—	—	—	—	—	—	—	—
1964	400	(H)	—	—	—	—	—	—	—	—	—	—
1965	100	(A)	—	—	—	—	25	—	—	—	—	—
1966	75	(A)	—	—	—	—	—	—	—	—	—	—
1967	30	(A)	—	—	—	—	—	—	—	—	—	—
1968	15	—	—	—	—	—	—	—	—	—	—	—
1969	12	(A)	—	—	—	—	—	—	—	—	—	—
1970	—	—	—	—	—	—	—	—	—	—	—	—
1971	305	(A)	—	—	—	—	—	—	—	—	—	—
1972	—	—	—	—	—	—	—	—	—	—	—	—
1973	40	(A)	—	—	—	—	10	—	—	—	—	—
1974	129	(A)	—	—	—	—	35	—	—	—	—	—
1975	260	(F)	—	—	—	—	—	—	—	—	—	—
1976	404	(W/F)	—	—	—	—	12	N(A)	24	—	—	13 P(A)
1977	456	(W/F)	—	—	—	—	410	E(A)	—	—	—	—
1978	388	(W/F)	24	E(F)	—	—	12	N(H)	—	—	—	63 P(A)
1979	327	(W/F)	16	E(F)	—	—	—	—	—	—	—	10 P(A)
1980	282	(W/F)	68	F(N)	—	—	—	—	—	30	P(H)	—
1981	536	(W/F)	84	E(F)	4	P(F)	28	P(H)	12	84	P(H)	—
1982	672	(W/F)	138	N(F)	188	N(F)	8	E(A)	—	—	—	—
1983	366	(W/F)	15	N(F)	—	—	15	P(A)	—	55	N(H)	—
1984	389	(W/F)	31	N(F)	—	—	35	N(B)	—	—	—	—
1985	320	E(F)	44	E(F)	—	—	243	N(F)	179	58	N(A)	85 N(A)
1986	708	N(F)	73	N(F)	45	E(A)	240	N(B)	178	104	E(A)	215 E(A)
1987	788	E(H)	71	E(F)	122	N(F)	40	E(A)	51	186	P(A)	175 P(A)
1988	564	N(F)	125	N(F)	167	N(F)	70	P(A)	325	680	N(A)	410 N(A)
1989	530	E(F)	150	N(A)	49	N(H)	80	P(A)	135	193	P(A)	132 P(A)
1990	664	E(F)	83	N(F)	33	P(H)	24	P(A)	—	—	—	—
1991	400	N(A)	38	N(A)	46	N(A)	42	N(F)	—	81	P(A)	320 P(A)
1992	778	E(H)	40	E(F)	31	N(A)	48	P(A)	30	P(A)	—	—
1993	1,060	E(F)	53	E(F)	—	—	40	N(A)	—	33	P(A)	118 P(A)
1994	572	E(H)	58	E(F)	10	N(A)	87	N(H)	27	P(H)	15	P(H)
1995	343	P(A)	28	P(A)	1	E(A)	38	N(H)	65	N(H)	16	P(A)
1996	335	N(F)	35	N(F)	21	N(A)	75	N(A)	15	N(H)	78	N(A)
1997	293	N(F)	—	—	—	—	—	—	55	N(H)	—	30 A(P)
1998	487	E(F)	35	N(A)	28	N(A)	75	N(A)	69	P(A)	—	66 P(A)
1999	605	E(A)	22	N(A)	—	—	—	—	550	N(A)	—	5 P(A)
2000	690	N(A)	35	N(A)	—	—	—	—	16	P(A)	—	33 N(A)
2001	1,054	N(F)	28	N(F)	—	—	150	N(H)	130	N(A)	248	E(A)
2002	876	N(F)	34	N(F)	8	N(A)	33	A	15	A	—	—
2003	595	N(H)	39	N(F)	19	N(F)	5	P(A)	24	P(A)	—	95 N(A)
2004	1,534	N(H)	27	N(F)	65	P(F)	69	N(H)	115	N(A)	26	N(A)
2005	1,015	N(H)	78	N(F)	102	N(F)	15	P(A)	79	N(A)	—	122 N(A)
2006	1,089	N(H)	51	N(A)	83	N(F)	18	N(A)	74	N(A)	67	N(H)
2007	890	N(A)	50	A(N)	60	F(N)	70	A(N)	95	A(N)	—	175 A(E)
98–07	884	—	43	—	52	—	60	—	117	—	114	96
2008	503	E(H)	14	A(N)	—	—	12	A(N)	11	A(P)	—	58 A(P)

Note: (A) = fixed-wing aircraft; — = no survey conducted; (B) = boat; (F/A) = combined foot and fixed-wing; (F) = foot; (H) = helicopter; (W/F) = weir and foot; N = normal conditions; E = excellent conditions; P = poor conditions.

<sup>a</sup> Andrew Creek total escapement equals sum of weir count, counts below weir, and on North Fork, minus egg take, 1976–1984.

Results from tagging studies in 1998–2000 indicated that the Klukshu River accounts for about 16–25% of the Chinook salmon escapement to the Alsek River drainage (Pahlke et al. 1999; Pahlke and Etherton 2001b, 2002). After the conclusion of the mark–recapture program in 2004 the expansion factor was revised with 7 years of data (Pahlke and Waugh 2006). The revised expansion factor, based on the estimate of large fish at the weir and the harvest immediately below the weir, is 4.17 (SE = 1.71; Appendix B10). This expansion factor has not been through the approval process with the PSC. The sum of the total weir count of 465 plus the harvest immediately below the weir of 14 in the aboriginal and sport fisheries was multiplied by the proportion of large fish in the sample collected at the weir (0.670) to get an estimate of large Chinook salmon returning to the Klukshu River (321), which was then multiplied by 4.17 to produce an estimate of escapement to the Alsek drainage of 1,339 (SE = 548) large Chinook salmon (Table 1; Appendix B10).

Age, sex and length data were collected from 230 live fish sampled at the Klukshu River weir, 161 of which were successfully aged (Appendices A7, panel J, and A8, panel J).

### **UNUK RIVER**

Poor weather and bad survey conditions prevented us from completing all the index surveys on the Unuk River in 2008 (Table 7). Cripple Creek and Genes Lake Creek typically make up over 60% of the total index counts and were either incomplete or poor in 2008 (Table 8).

Based on results of mark–recapture and radiotracking studies, the expansion factor for the Unuk and Chickamin rivers was revised in 1996 from 1.6 to 4.0 times the summed tributary counts (Pahlke et al. 1996, 1997a-b). After additional mark–recapture estimates were obtained, the expansion factor was revised in 2002 to 5.0 (McPherson et al. 2003) and again in 2007 to 4.87 (SE 0.60; Pahlke 2007; Appendix B5).

The ongoing mark–recapture program estimated an escapement of 3,104 (SE = 390) large Chinook salmon (Table 1; Weller et al. *in prep*). The mark–recapture estimate divided by expansion factor was equivalent to an index count of about 637

large Chinook salmon to the Unuk River in 2007, just below the index goal range of 650 to 1,400 (McPherson and Carlile 1997). Index counts have been below the lower end of the escapement goal range only 3 times since 1981 (Figure 6).

As part of the mark-recapture project, sport gear was used to sample live fish and spears were used to collect carcasses for age, sex and size data; 389 fish were sampled (Appendices A7, panel D and A8, panel D).

### **CHICKAMIN RIVER**

In index areas on 8 tributaries of the Chickamin River, 1,111 large Chinook salmon were counted in 2008, compared to 893 in 2007 (Tables 9 and 10). Counts in 2008 were above the 10-year average in 4 out of 8 Chickamin River tributaries, with very high counts in Humpy Creek and King Creek (Table 9). Those two tributaries have the latest run timing of any of the Unuk or Chickamin tributaries, with Chinook commonly spawning into September. The 2008 count was above the index survey escapement goal range of 450 to 900 fish (Figure 7; McPherson and Carlile 1997). The summed counts for 2008 were multiplied by a survey expansion factor of 4.75 to produce a total escapement estimate of 5,277 (SE = 777) fish to the system (Johnson et al. 2009) (Table 1; Appendix B4).

Sport gear and spears were used to collect age, sex and length data from 454 fish in 2008 (Johnson *In prep*; Appendices A7, panel C, and A8, panel C).

### **BLOSSOM RIVER**

In index areas of the Blossom River, 257 large Chinook salmon were counted in 2008, up from 135 fish counted in 2007 (Table 11). The 2008 count was within the index survey goal range of 250 to 500 (McPherson and Carlile 1997). Counts had exceeded the point goal of 300 in 1982–1989, but since 1991 they have frequently been below the escapement goal range (Figure 8). Based on results of mark–recapture studies, the expansion factor for the Blossom River was revised in 1996 from 1.6 to 2.5 (Pahlke 1997b), in 2002 to 4.0 (McPherson et al. 2003), and again in 2006 to 3.01 (Appendix B3; Weller et al. 2007a).




Figure 5.—Weir count of Chinook salmon to the Klukshu River tributary of the Alsek River, 1976–2008, and mark–recapture estimates divided by expansion factor of 4.03. Lines show upper and lower limits of escapement goal range.




Figure 6.—Counts of large Chinook salmon in index areas of the Unuk River, 1975–2008, and mark–recapture estimates divided by expansion factor (4.87). Lines show upper and lower limits of index escapement goal range.

Table 6.—Count of Chinook salmon through the Klukshu River weir, harvest above and below the weir, estimated proportion of large fish through the weir, and counts of spawning adults in other tributaries of the Alsek River, 1976–2008.

Year	Klukshu River							Blanchard River	Takhanne River		Goat Creek			
	Aerial count <sup>a</sup>	Weir count	Below weir harvest	Total	% Large	$\hat{C}_L$ <sup>b</sup>	Above weir harvest							
1976	—	1,278	130	1,408	0.98	1,382	214	—	—	—	—			
1977	—	3,144	195	3,339	0.75	2,517	446	—	—	—	—			
1978	—	2,976	195	3,171	0.89	2,819	446	—	—	—	—			
1979	—	4,404	422	4,826	0.93	4,477	1,300	—	—	—	—			
1980	—	2,673	130	2,767	0.70	1,937	150	—	—	—	—			
1981	—	2,113	150	2,263	0.88	1,997	150	35	(H)	11	(H)	—		
1982	633	N(H)	2,369	183	2,552	0.86	2,200	400	59	(H)	241	(H)	13	(H)
1983	917	N(H)	2,537	202	2,739	0.97	2,645	300	108	(H)	185	(H)	—	—
1984	—	1,672	275	1,947	0.92	1,797	100	304	(H)	158	(H)	28	(H)	
1985	—	1,458	170	1,628	—	1,381	175	232	(H)	184	(H)	—	—	
1986	738	P(H)	2,709	125	2,834	0.84	2,394	102	556	(H)	358	(H)	142	(H)
1987	933	E(H)	2,616	326	2,942	0.93	2,733	125	624	(H)	395	(H)	85	(H)
1988	—	2,037	249	2,286	0.86	1,973	43	437	E(H)	169	E(H)	54	E(H)	
1989	893	E(H)	2,456	215	2,671	0.82	2,183	254	—	158	E(H)	34	E(H)	
1990	1,381	E(H)	1,915	468	2,383	0.88	2,109	217	—	325	E(H)	32	E(H)	
1991	—	2,489	652	3,141	0.97	3,051	266	121	N(H)	86	E(H)	63	E(H)	
1992	261	P(H)	1,367	139	1,506	0.88	1,323	124	86	P(H)	77	N(H)	16	N(H)
1993	1,058	N(H)	3,303	258	3,561	0.85	3,043	82	326	N(H)	351	E(H)	50	N(H)
1994	1,558	N(H)	3,727	387	4,114	0.72	2,952	107	349	N(H)	342	E(H)	67	N(H)
1995	1,053	E(H)	5,678	921	6,599	0.92	6,072	281	338	P(H)	260	P(H)	—	—
1996	788	N(H)	3,599	656	4,255	0.81	3,464	217	132	N(H)	230	N(H)	12	N(H)
1997	718	P(H)	2,989	267	3,256	0.94	3,045	160	109	P(H)	190	P(H)	—	—
1998	—	1,364	266	1,630	0.69	1,131	17	71	P(H)	136	N(H)	39	N(H)	
1999	500	P(H)	2,193	337	2,530	0.76	1,918	27	371	N(H)	194	N(H)	51	N(H)
2000	—	1,365	53	1,416	0.89	1,263	44	168	N(H)	152	N(H)	33	N(H)	
2001	—	1,825	152	1,977	0.85	1,679	87	543	N(H)	287	N(H)	21	N(H)	
2002	—	2,241	185	2,426	0.92	2,237	100	351	N(H)	220	N(H)	86	E(H)	
2003	—	1,737	136	1,873	0.76	1,416	76	127	N(H)	105	N(H)	10	N(H)	
2004	—	2,523	113	2,636	0.94	2,481	68	84	P(H)	46	P(H)	—	—	
2005	—	1,070	78	1,148	0.93	1,070	36	112	E(H)	47	N(H)	7	N(H)	
2006	—	568	17	578	0.77	451	0	98	N(H)	28	P(H)	9	N(H)	
2007	—	676	41	717	0.88	628	0	39	P(H)	32	P(H)	45	N(H)	
98–07	—	1,556	138	1,694	0.84	1,427	62	203	—	141	—	32	—	
Avg.	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2008	—	465	14	479	0.67	321	0	65	E(H)	41	N(H)	11	N(H)	

Note: (A) = fixed-wing aircraft; — = no survey; (H) = helicopter; N = normal conditions; E = excellent conditions; P = poor conditions.

<sup>a</sup> Counts prior to 1975 may not be comparable due to differences in survey dates and counting methods.

<sup>b</sup>  $\hat{C}_L$  = weir count plus catch immediately below weir multiplied by estimated proportion of large fish at weir (Pahlke and Waugh 2006).

Table 7.—Peak escapement counts of Chinook salmon to index areas of the Unuk River, 1972–2008.

Year <sup>a</sup>	Cripple Creek	Genes Lake Creek	Eulachon Creek	Clear Creek	Lake Creek	Kerr Creek	Total
1972	95 (A)	35 (A)	450 (A)	90 (A)	55 (A)	—	725
1973	—	—	64 (H)	—	—	—	64
1974	—	—	68 (H)	—	—	—	68
1975	—	—	17 (H)	—	—	—	17
1976	— <sup>b</sup>	—	3 (A)	—	—	—	3
1977	529 <sup>b</sup> (F)	339 (F)	57 (H)	34 (H)	—	15 (H)	974
1978	394 <sup>b</sup> (F)	374 (F)	218 (H)	85 (H)	20 (H)	15 (H)	1,106
1979	363 (F)	101 (F)	48 (H)	14 (H)	30 (H)	20 (H)	576
1980	748 (F)	122 (F)	95 (H)	28 (H)	5 (H)	18 (H)	1,016
1981	324 (F)	112 (F)	196 (H)	54 (H)	20 (H)	25 (H)	731
1982	538 (F)	329 (F)	384 (H)	24 (H)	48 (H)	28 (H)	1,351
1983	459 (F)	338 (F)	288 (H)	24 (H)	12 (H)	4 (H)	1,125
1984	644 (F)	647 (F)	350 (H)	113 (H)	32 (H)	51 (H)	1,837
1985	284 (F)	553 (F)	275 (H)	37 (H)	22 (H)	13 (H)	1,184
1986	532 (F)	838 (F)	486 (H)	183 (F)	25 (H)	62 (H)	2,126
1987	860 (F)	398 (F)	520 (H)	107 (H)	37 (H)	51 (H)	1,973
1988	1,068 (F)	154 (F)	146 (F)	292 (H)	60 (H)	26 (H)	1,746
1989	351 (F)	302 (F)	298 (H)	128 (H)	27 (F)	43 (H)	1,149
1990	86 (F)	284 (F)	81 (H)	103 (F)	26 (F)	11 (H)	591
1991	358 (W/F)	123 (F)	43 (H)	96 (F)	23 (F)	12 (H)	655 <sup>c</sup>
1992	327 (W/F)	360 (F)	57 (F)	69 (F)	31 (H)	30 (H)	874 <sup>c</sup>
1993	448 N(F)	330 N(F)	132 E(F)	137 N(F)	8 N(F)	13 P(H)	1,068
1994	161 P(F)	300 N(F)	52 N(H)	128 E(F)	18 N(F)	52 N(F)	711
1995	211 N(F)	347 N(F)	74 N(H)	66 E(H)	35 E(H)	39 N(H)	772
1996	417 N(F)	400 N(F)	79 N(F)	148 E(F)	25 E(H)	98 E(F)	1,167
1997	244 P(F)	154 N(F/H)	53 N(F)	113 N(F)	13 N(H)	59 E(F)	636
1998	311 N(F)	283 N(F)	39 N(H)	81 N(F)	22 N(F)	104 N(F)	840
1999	202 N(F)	307 N(F)	54 N(H)	67 N(F)	9 N(F)	41 N(F)	680
2000	450 N(F)	565 N(F)	116 N(H)	86 N(H)	56 E(H)	68 N(H)	1,341
2001	701 N(F)	806 N(F/H)	217 E(H)	167 N(H)	84 N(H)	44 P(H)	2,019
2002	156 P(F)	455 N(F/H)	78 N(H)	87 N(H)	61 N(H)	60 E(F)	897
2003	232 P(F)	448 N(F)	95 N(H)	198 E(F)	68 E(F)	80 N(F)	1,121
2004	237 N(F)	388 E(F)	78 N(F)	191 E(F)	47 N(H)	67 N(F)	1,008
2005	314 N(F)	338 N(F)	99 N(H)	132 E(F)	33 N(H)	13 P(F)	929
2006	210 N(F)	551 N(F)	30 P(H)	88 N(F)	55 N(H)	6 P(H)	940
2007	204 N(F)	232 N(F)	81 N(H)	167 E(F)	28 E(F)	8 P(H)	720
98–07							
Avg.	302	437	89	126	46	49	1,050
2008	- No survey	100 P(H)	7 P(F)	42 N(F)	64 N(H)	29 N(H)	242

Note: (A) = fixed-wing aircraft; — = no survey conducted; (F) = foot; (H) = helicopter; (W/F) = weir and foot; (F/H) = foot and helicopter; N = normal conditions; E = excellent conditions; P = poor conditions.

<sup>a</sup> Counts prior to 1975 may not be comparable due to differences in survey dates and counting methods.

<sup>b</sup> Not including 35 fish for egg take in 1976; 132 in 1977; 85 in 1978.

<sup>c</sup> Cripple Creek weir count reduced by /0.625 to be comparable with foot surveys.

Table 8.—Distribution of spawning Chinook salmon among index areas of the Unuk River for years when all index areas were surveyed.

Year	Cripple Creek		Genes Lake Creek		Eulachon Creek		Clear Creek		Lake Creek		Kerr Creek		Total
	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	
1978	394	36	374	34	218	20	85	8	20	2	15	1	1,106
1979	363	63	101	18	48	8	14	2	30	5	20	3	576
1980	748	74	122	12	95	9	28	3	5	0	18	2	1,016
1981	324	44	112	15	196	27	54	7	20	3	25	3	731
1982	538	40	329	24	384	28	24	2	48	4	28	2	1,351
1983	459	41	338	30	288	26	24	2	12	1	4	0	1,125
1984	644	35	647	35	350	19	113	6	32	2	51	3	1,837
1985	284	24	553	47	275	23	37	3	22	2	13	1	1,184
1986	532	25	838	39	486	23	183	9	25	1	62	3	2,126
1987	860	44	398	20	520	26	107	5	37	2	51	3	1,973
1988	1,068	61	154	9	146	8	292	17	60	3	26	1	1,746
1989	351	31	302	26	298	26	128	11	27	2	43	4	1,149
1990	86	15	284	48	81	14	103	17	26	4	11	2	591
1991	358	55	123	19	43	7	96	15	23	4	12	2	655
1992	327	37	360	41	57	7	69	8	31	4	30	3	874
1993	448	42	330	31	132	12	137	13	8	0	13	1	1,068
1994	161	23	300	42	52	7	128	18	18	3	52	7	711
1995	211	27	347	45	74	10	66	9	35	5	39	5	772
1996	417	36	400	34	79	7	148	13	25	2	98	8	1,167
1997	244	38	154	24	53	8	113	18	13	2	59	9	636
1998	311	37	283	34	39	5	81	10	22	3	104	12	840
1999	202	30	307	45	54	8	67	10	9	1	41	6	680
2000	450	34	565	42	116	9	86	6	56	4	68	5	1,341
2001	701	35	806	40	217	11	167	8	84	4	44	2	2,019
2002	156	17	455	51	78	9	87	10	61	7	60	7	897
2003	232	21	448	40	95	8	198	18	68	6	80	7	1,121
2004	237	24	388	38	78	8	191	19	47	5	67	7	1,008
2005	314	34	338	36	99	11	132	14	33	4	13	1	929
2006	210	22	551	59	30	3	88	9	55	6	6	1	940
2007	204	28	232	32	81	11	167	23	28	4	8	1	720
Avg.	394	36	365	34	159	13	107	10	33	3	39	4	1,096

The count for 2008 was multiplied by the expansion factor of 3.01 to produce a total escapement estimate of 773 (SE = 265) large fish. Sport gear was used to sample age, sex and length data and 68 samples were collected in 2008 (Appendices A7, panel B and A8, panel B).

## KETA RIVER

In 2008, 363 Chinook salmon were counted in the Keta River, up from 311 in 2007 (Table 11) and within the 1996 revised index goal range of 250 to 500 large fish (McPherson and Carlile

1997). Prior to 1990, counts of Chinook salmon in the Keta River increased steadily after implementation of the 1980 rebuilding program (Figure 9). Based on results of mark-recapture studies in 1998–2000, the expansion factor for the Keta River was revised in 2001 from 2.5 to 3.01 (SE 0.56; Appendix B2; Freeman et al. 2001). The peak count for 2008 was multiplied by 3.01 to produce a total escapement estimate of 1,093 (SE = 203) large fish (Table 1; Appendix B2). Sport gear was used to collect 101 age, sex and length samples from live fish (Appendices A7, panel A and A8, panel A).




Figure 7.—Counts of Chinook salmon in index areas of the Chickamin River, 1975–2008 and mark–recapture estimates divided by expansion factor (4.75). Lines show upper and lower limits of index escapement goal range.




Figure 8.—Counts of Chinook salmon into the Blossom River, 1975–2008 and mark–recapture estimates. Lines show upper and lower limits of index escapement goal range.

Table 9.—Counts of Chinook salmon in index areas of the Chickamin River, 1972–2008.

Year <sup>a</sup>	South Fork Creek		Barrier Creek		Butler Creek		Leduc Creek		Indian Creek		Humpy Creek		King Creek		Clear Falls Creek		Total <sup>b</sup>
1972	350	(A)	25	(A)	—	—	85	(A)	—	—	65	(A)	510	(A)	—	—	1,035
1973	—	—	—	—	—	—	—	—	—	—	14	(A)	65	(A)	—	—	79
1974	144	(H)	—	—	—	—	—	—	—	—	—	—	11	(H)	—	—	155
1975	141	(H)	9	(H)	66	(H)	6	(H)	90	(H)	7	(H)	30	(H)	—	—	370
1976	46	(H)	10	(H)	15	(H)	12	(H)	9	(H)	—	—	—	—	—	—	157
1977	52	(H)	66	(H)	30	(H)	26	(H)	53	(H)	0	(H)	—	—	—	—	363
1978	21	(H)	94	(H)	4	(H)	42	(H)	20	(H)	—	—	—	—	—	—	308
1979	63	(H)	17	(H)	29	(H)	0	(H)	31	(H)	—	—	—	—	—	—	239
1980	56	(H)	62	(H)	104	(H)	17	(H)	22	(H)	—	—	—	—	—	—	445
1981	51	(H)	105	(H)	51	(H)	25	(H)	12	(H)	4	(F)	105	(F)	31	(H)	384
1982	84	(H)	149	(H)	37	(H)	36	(H)	30	(F)	37	(F)	165	(F)	33	(H)	571
1983	28	(H)	138	(H)	91	(H)	30	(H)	47	(H)	—	—	212	(F)	30	(H)	599
1984	185	(H)	171	(H)	124	(H)	15	(H)	103	(H)	88	(F)	388	(F)	28	(H)	1,102
1985	163	(H)	129	(H)	92	(H)	8	(H)	125	(H)	50	(H)	377	(H)	12	(H)	956
1986	562	(H)	168	(H)	203	(H)	20	(H)	120	(H)	—	—	564	(H)	40	(H)	1,745
1987	261	(H)	76	(H)	120	(H)	19	(H)	115	(H)	26	(H)	310	(H)	48	(H)	975
1988	280	(F/H)	82	(F/H)	159	(H)	25	(F/H)	32	(H)	19	(F/H)	164	(H)	25	(H/F)	786
1989	226	(F/H)	90	(H)	137	(H)	57	(H)	84	(H)	22	(F/H)	224	(H)	94	(H)	934
1990	135	(F)	107	(H)	27	(H)	20	(H)	24	(H)	35	(H)	163	(H)	53	(H)	564
1991	125	(H)	18	(H)	49	(H)	14	(H)	38	(H)	13	(H)	185	(H)	45	(H)	487
1992	87	(H)	4	(H)	68	(H)	4	(H)	20	(H)	8	(H)	131	(H)	24	(H)	346
1993	67	N(H)	46	E(H)	68	N(H)	11	N(H)	29	N(H)	13	N(H)	80	N(H)	75	N(H)	389
1994	31	N(H)	29	E(H)	64	E(H)	18	E(H)	16	N(H)	44	N(H)	129	E(H)	57	E(H)	388
1995	87	E(H)	12	E(F)	59	E(F)	60	E(H)	36	N(F)	13	N(F)	62	N(H)	27	E(H)	356
1996	72	N(H)	13	N(F)	74	E(H)	23	E(H)	48	N(F)	30	N(F)	106	E(F)	56	E(H)	422
1997	28	P(H)	10	N(H)	43	N(H)	7	N(H)	24	N(H)	15	N(H)	95	N(H)	50	N(H)	272
1998	46	N(H)	0	N(H)	124	E(H)	16	P(H)	46	N(H)	28	N(H)	123	N(H)	8	P(H)	391
1999	54	N(H)	18	N(H)	106	N(H)	33	N(H)	52	N(F)	16	N(F)	200	N(H)	22	N(H)	501
2000	109	N(H)	27	N(H)	230	E(H)	61	N(H)	63	N(H)	20	N(H)	251	N(H)	40	P(H)	801
2001	264	E(H)	27	N(H)	270	E(H)	59	N(H)	61	N(H)	78	N(F)	221	N(H)	30	N(H)	1,010
2002	329	N(H)	20	N(H)	102	N(H)	23	N(H)	146	E(H)	9	P(H)	361	E(H)	23	N(H)	1,013
2003	183	E(H)	13	N(H)	172	N(H)	37	E(H)	21	N(H)	119	E(H)	363	N(H)	56	N(H)	964
2004	109	N(H)	17	N(H)	143	N(H)	35	E(F)	56	E(F)	162	E(F)	272	N(H)	4	P(H)	798
2005	106	P(H)	46	E(H)	115	N(H)	69	N(H)	49	N(H)	38	N(H)	450	E(H)	53	N(H)	926
2006	179	E(H)	10	N(H)	325	N(H)	52	N(H)	55	N(H)	37	E(H)	620	N(H)	52	N(H)	1,330
2007	197	N(H)	19	N(H)	133	N(H)	15	N(F)	66	N(F)	96	F(N)	315	N(H)	52	N(H)	893
98–07																	
Avg.	157		20		172		40		62		60		318		34		863
2008	87	N(H)	3	N(H)	68	N(H)	5	P(H)	76	N(F)	190	E(H)	622	E(H)	60	N(H)	1,111

Note: (A) = fixed-wing aircraft; — = no survey conducted; (F) = foot; (H) = helicopter; (F/H) = foot and helicopter; N = normal conditions; E = excellent conditions; P = poor conditions.

<sup>a</sup> Escapement counts conducted prior to 1975 may not be comparable due to differences in survey dates and counting methods.

<sup>b</sup> Totals for 1975–1980, 1983 and 1986 expanded for unsurveyed index areas by 1981–1992 average %.

Table 10.—Distribution of spawning Chinook salmon among index areas of the Chickamin River for years when all index areas were surveyed.

Year	South Fork		Barrier		Butler		Leduc		Indian		Humpy		King		Clear Falls		Total
	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	Creek	%	
1981	51	13	105	27	51	13	25	7	12	3	4	1	105	27	31	8	384
1982	84	15	149	26	37	6	36	6	30	5	37	6	165	29	33	6	571
1984	185	17	171	16	124	11	15	1	103	9	88	8	388	35	28	3	1,102
1985	136	14	156	16	93	10	8	0	125	13	50	5	377	39	12	1	957
1987	261	27	76	8	120	12	19	2	115	12	26	3	310	32	48	5	975
1988	280	36	82	10	159	20	25	3	32	4	19	2	164	21	25	3	786
1989	226	24	90	10	137	15	57	6	84	9	22	2	224	24	94	10	934
1990	135	24	107	19	27	5	20	4	24	4	35	6	163	29	53	9	564
1991	125	26	18	4	49	10	14	3	38	8	13	3	185	38	45	9	487
1992	87	25	4	1	68	20	4	1	20	6	8	2	131	38	24	7	346
1993	67	17	46	12	68	17	11	3	29	7	13	3	80	21	75	19	389
1994	31	8	29	7	64	16	18	5	16	4	44	11	129	33	57	15	388
1995	87	24	12	3	59	17	60	17	36	10	13	4	62	17	27	8	356
1996	72	17	13	3	74	18	23	5	48	11	30	7	106	25	56	13	422
1997	28	10	10	4	43	16	7	3	24	9	15	6	95	35	50	18	272
1998	46	12	0	0	124	32	16	4	46	12	28	7	123	31	8	2	391
1999	54	11	18	4	106	21	33	7	52	10	16	3	200	40	22	4	501
2000	109	14	27	3	230	29	61	8	63	8	20	2	251	31	40	5	801
2001	264	26	27	3	270	27	59	6	61	6	78	8	221	22	30	3	1,010
2002	329	32	20	2	102	10	23	2	146	14	9	1	361	36	23	2	1,013
2003	183	19	13	1	172	18	37	4	21	2	119	12	363	38	56	6	964
2004	109	14	17	2	143	18	35	4	56	7	162	20	272	34	4	1	798
2005	106	11	46	5	115	12	69	7	49	5	38	4	450	49	53	6	926
2006	179	13	10	1	325	24	52	4	55	4	37	3	620	47	52	4	1,330
2007	197	22	19	2	133	15	15	2	66	7	96	11	315	35	52	6	893
Avg.	149	20	58	8	118	16	29	4	56	8	41	6	246	33	40	5	737
2008	87	8	3	0	68	6	5	0	76	7	190	17	622	56	60	5	1,111

## KING SALMON RIVER

Two helicopter and 2 foot surveys were completed on King Salmon River in 2008. The peak count during the helicopter survey was 15 large Chinook salmon, and 79 were counted during the foot survey, both under normal conditions. The peak count was below the 119 fish counted in 2007 and below the 10-year average of 106 (Table 12). The escapement goal was revised in 1997 to a range of 120 to 240 total large fish (McPherson and Clark 2001). The resulting index goal range is 80–160 large fish observed. Counts exceeded the lower bound of the index goal range since 1993 and the 2007 count was just below the range (Figure 10). The peak count of 79 was multiplied by 1.52 to produce a total escapement estimate of 120 (SE = 21) large fish (Table 1; Appendix B8).

Angling gear was used to collect age, sex and length data from 26 Chinook salmon in 2008, (Appendices A7, panel G, and A8, panel G).

## SITUK RIVER

The count of all Chinook salmon through the Situk River weir in 2008 was 798 Chinook salmon, of which 453 were large (Tables 1 and 13). There was no harvest above the weir. The preseason forecast of large Chinook salmon returning to the Situk in 2008 was below the threshold to allow directed harvest of Chinook (5AAC30.365). Sport harvest of fish  $\geq 20$  in was prohibited and the commercial fishery opening was delayed.

Escapements have met or exceeded the escapement goal range of 450–1,050 large spawners each year since 1983 (Figure 11). The proportion of the recreational harvest that is caught above the weir varies from year to year and is estimated, from the Statewide Harvest Survey (Howe et al. 2001) and a creel survey. The escapement counts from the base period of 1976–1981 all exceeded the revised escapement goal, indicating the Situk Chinook salmon stock may not have been depressed (McPherson et al. 2005).

Age, sex and length data was collected from 79 live fish sampled at the weir, 63 of which were successfully aged (Appendices A7, panel K and A8, panel K).

## **CHILKAT RIVER**

The 2008 escapement to the Chilkat River was estimated by a mark–recapture experiment to be 2,833 large Chinook salmon (SE = 530), over twice the escapement estimated in 2007 and below the 10-year average of 3,341 (Chapell *In prep b*; Table 14). The escapement goal was reviewed in 2003 and revised slightly to a range of 1,750 to 3,500 large fish (Ericksen and McPherson 2004). Except in 2007, estimated escapements have been within, or exceeded the escapement goal ranges since the start of the program in 1991 (Figure 12). The mark–recapture experiment also provided age, sex, and size data (Appendices A7, panel I, and A8, panel I).

## **OTHER SYSTEMS**

Counts of Chinook salmon in the Marten and Wilson rivers are not included in the regional index program, and no official escapement goals have been set for these systems. However, periodic counts have been made in the two rivers since 1982 because of their proximity to other surveyed systems (Table 11). Grant and Klahini rivers are small Chinook systems near the Unuk River in Behm Canal that have been surveyed sporadically (Table 11). In 2008, no surveys were conducted on any of these systems. Occasional surveys have been flown on the Harding River and Aaron Creek to determine the feasibility of adding these medium and small systems to the program (Table 5). The remaining systems are too remote and funds are not currently available for these surveys. However, several are routinely surveyed by the local management biologists and in 2008, 58 Chinook were counted in the East Fork of the Bradfield River, 12 in Harding River, 11 in Aaron Creek (Table 5), and 26 in Farragut River.

Trips to collect genetic samples from Chinook salmon in the Farragut River were conducted in 2008. Fifteen fish were sampled, 14 of which were aged (Appendix A12).

## **CODED WIRE TAG RECOVERY**

One hatchery fish marked with a CWT was recovered in the Farragut River in 2008 and 1 fish

tagged as a smolt in the Taku River was recovered in the Stikine River (Appendix A13).

## **OBSERVER TRAINING**

Eight calibration surveys with 2 alternate observers were conducted in 2008 (Table 15).

Four surveys were flown with the alternate observer on index areas in Behm Canal where pink and chum salmon are mixed with the Chinook. The four counts varied widely ranging from 42% to over 200% of the primary observer's. Four surveys conducted in systems without pink or chum salmon were more consistent, ranging from 66% to 111% of the primary observer's.

## **DISCUSSION**

The utility of the index method as a measure of escapement is based on the assumption that the number of fish counted in an index area is a constant proportion of the escapement in the index area or watershed. Therefore, a change in the escapement is assumed to cause a proportional change in the index count. Consequently, if this assumption holds, even though index counts are not estimates of total escapement, multi-year trends in escapement are correct. Two types of error affect the accuracy of the survey counts.

First, features intrinsic to each area interfere with the ability to count fish. Examples include heavily shaded areas or topography that prevent close approach with a helicopter, presence of other species that could be confused with Chinook salmon, overhanging brush, and deep or occluded water. Also, not all spawning areas in a tributary or drainage are surveyed. These features are accounted for by survey expansion factors.

Second, factors that affect counting efficiency may vary greatly from year to year and survey to survey. These include annual changes in migratory timing, changes in the distribution of spawners among the tributaries of a watershed between years, inclement weather, turbidity events, or changes in pilot and/or observer experience. Also, the proportion of fish counted in an index area may vary with the number of fish in the index area, e.g., a lower proportion of fish may be counted when abundance is extremely high.

Table 11.—Counts of Chinook salmon for selected rivers in Behm Canal, 1961–2008.

Year <sup>a</sup>	Keta River		Blossom River		Wilson River		Marten River		Grant River		Klahini River		Total
1961	44	(F)	68	(F)	—	—	22	(F)	40	(A)	—	—	174
1962	—	—	—	—	—	—	—	—	6	(A)	100	(A)	106
1963	—	—	450	(A)	375	(A)	—	—	15	(A)	—	—	840
1964	—	—	—	—	—	—	—	—	—	—	—	—	—
1965	—	—	—	—	50	(A)	43	(H)	—	—	—	—	93
1966	75	(A)	200	(A)	60	(A)	10	(A)	100	(A)	3	(A)	448
1967	86	(H)	—	—	8	(H)	7	(H)	15	(H)	—	—	116
1968	—	—	—	—	—	—	—	—	4	(H)	—	—	4
1969	200	(A)	—	—	10	(A)	10	(A)	69	(H)	3	(H)	292
1970	—	—	100	(H)	—	—	—	—	—	—	—	—	100
1971	—	—	—	—	—	—	—	—	—	—	—	—	—
1972	255	(A)	225	(A)	275	(A)	—	—	25	(A)	150	(A)	930
1973	—	—	—	—	30	(A)	—	—	38	(A)	7	(H)	75
1974	25	(H)	166	(H)	—	—	—	—	—	—	—	—	191
1975	203	(H)	146	(H)	7	(H)	15	(H)	—	—	—	—	371
1976	84	(H)	68	(H)	—	—	—	—	—	—	—	—	152
1977	230	(H)	112	(H)	—	—	—	—	—	—	—	—	342
1978	392	(H)	143	(H)	—	—	2	(A)	—	—	—	—	537
1979	426	(H)	54	(H)	36	(H)	—	—	—	—	—	—	516
1980	192	(H)	89	(H)	—	—	—	—	—	—	—	—	281
1981	329	(H)	159	(H)	76	(F)	—	—	25	(H)	42	(F)	631
1982	754	(H)	345	(H)	300	(B)	75	(F)	33	(F)	79	(F)	1,586
1983	822	(H)	589	(H)	178	(B)	138	(B)	8	(A)	10	(H)	1,745
1984	610	(H)	508	(H)	133	(F)	12	(B)	124	(F)	54	(F)	1,441
1985	624	(H)	709	(H)	420	(H)	69	(F)	55	(F)	20	(F)	1,897
1986	690	(H)	1,278	(H)	—	—	—	—	—	—	—	—	1,968
1987	768	(H)	1,349	(H)	—	—	270	(H)	33	(A)	—	—	2,420
1988	575	(H)	384	(H)	—	—	543	(H)	—	—	40	(H)	1,542
1989	1,155	(H)	344	(H)	—	—	133	(H)	—	—	—	—	1,632
1990	606	(H)	257	(H)	—	—	283	(H)	—	—	—	—	1,146
1991	272	N(H)	239	N(H)	—	—	135	N(H)	—	—	—	—	646
1992	217	N(H)	150	N(H)	109	E(H)	76	(H)	25	N(H)	19	(H)	596
1993	362	E(H)	303	N(H)	63	P(H)	229	E(H)	—	—	—	—	957
1994	306	E(H)	161	N(H)	—	—	178	E(H)	—	—	—	—	645
1995	175	E(H)	217	N(H)	58	N(H)	171	E(H)	—	—	—	—	621
1996	297	N(H)	220	E(H)	23	P(H)	62	N(H)	—	—	—	—	602
1997	246	N(H)	132	N(H)	16	N(H)	56	N(H)	9	N(H)	—	—	459
1998	180	N(H)	91	N(H)	—	—	—	—	—	—	—	—	271
1999	276	E(H)	212	N(H)	—	—	—	—	—	—	—	—	488
2000	300	N(H)	231	N(H)	—	—	—	—	—	—	—	—	531
2001	343	E(H)	204	N(H)	79	E(H)	—	—	—	—	83	E(H)	626
2002	411	E(H)	224	E(H)	—	—	—	—	—	—	—	—	635
2003	322	N(H)	203	E(H)	—	—	—	—	—	—	—	—	525
2004	376	E(H)	333	E(H)	—	—	—	—	—	—	—	—	709
2005	497	E(H)	445	E(H)	—	—	—	—	—	—	—	—	942
2006	747	E(H)	339	N(H)	—	—	—	—	—	—	—	—	—
2007	311	N(H)	135	N(H)	—	—	—	—	—	—	—	—	—
98–07													
Avg.	376		242		79						83		513
2008	363	N(H)	257	E(H)	—	—	—	—	—	—	—	—	—

Note: (A) = fixed-wing aircraft; — = no survey conducted; (F) = foot; (H) = helicopter; (B) = boat; N = normal conditions; E = excellent conditions; P = poor conditions.

<sup>a</sup> Escapement counts prior to 1975 may not be comparable due to differences in survey dates or methods.

Table 12.—Peak escapement counts and weir counts of spawning Chinook salmon in the King Salmon River, 1971–2008.

Year	Survey count		Survey as percent of weir count	Total egg take (adults)	Total weir count (adults)	Total weir count (jacks) <sup>a</sup>	Adults below weir (foot count)	Total inriver (adults)	Total natural spawning
	Below weir	Above weir							
	A	B	B/(D-C)	C	D	E	F	D+F	D+F-C
1971	–	94 (F)	–	–	–	–	–	–	–
1972	–	90 (F)	–	–	–	–	–	–	–
1973	–	211 (F)	–	–	–	–	–	–	–
1974	–	104 (F)	–	–	–	–	–	–	–
1975	–	42 (H)	–	–	–	–	–	–	–
1976	–	65 (H)	–	–	–	–	–	–	–
1977	–	134 (H)	–	–	–	–	–	–	–
1978	–	57 (H)	–	–	–	–	–	–	–
1979	–	88 (H)	–	17	–	–	–	–	–
1980	–	70 (H)	–	–	–	–	–	–	–
1981	–	101 (H)	–	11	–	–	–	101	90
1982	–	259 (H)	–	30	–	–	–	259	229
1983	25	183 (H)	85%	37	252	20	30	282	245 <sup>b</sup>
1984	14	184 (H)	71%	46	299	82	12	311	265 <sup>b</sup>
1985	12	105 (H)	64%	29	194	45	10	204	175 <sup>b</sup>
1986	9	190 (H)	80%	26	264	72	17	281	255 <sup>b</sup>
1987	19	128 (H)	73%	31	207	62	20	227	196 <sup>b</sup>
1988	5	94 (H)	50% <sup>c</sup>	35	231	54	12	243	208 <sup>b</sup>
1989	34	133 (H)	63%	38 <sup>e</sup>	249	71	29	278	240 <sup>b</sup>
1990	34	98 (H)	57%	29	190	32	8	198	179 <sup>b</sup>
1991	6	91 (H)	72%	20	146	89	8	154	134 <sup>b</sup>
1992	–	58 (H)	59% <sup>e</sup>	18	47	16	70	117	99 <sup>b</sup>
1993	–	175 E(H)			no weir or egg take				
1994	–	140 N(F)			no weir or egg take				
1995	–	97 P(H)			no weir or egg take				
1996	–	192 E(F)			no weir or egg take				
1997		238 N(F)			no weir or egg take				
1998		88 E(F)			no weir or egg take				
1999		200 E(F)			no weir or egg take				
2000		91 N(F)			no weir or egg take				
2001		98 N(F)			no weir or egg take				
2002		102 N(F)			no weir or egg take				
2003		78 N(F)			no weir or egg take				
2004		89 E(F)			no weir or egg take				
2005		94 P(F)			no weir or egg take				
2006		99 N(F)			no weir or egg take				
2007		119 N(F)			no weir or egg take				
98-07		106							
Avg.									
2008		79 N(F)							

Notes: – = no survey conducted or data not comparable; (F) = foot; (H) = helicopter; N = survey conditions normal; E = excellent; P = poor.

<sup>a</sup> Minimum count as jacks could pass through weir.

<sup>b</sup> Natural spawning (adults) = (total inriver egg take; 1983–1992).

<sup>c</sup> Four females and two males were held but not spawned for egg take; % = 94/(231-37-6) = 50%.

<sup>d</sup> Includes holding mortality of 4 males and 6 females for egg take.

<sup>e</sup> Peak survey was after weir was removed 58/99 = 59%.




Figure 9.—Counts of Chinook salmon to the Keta River, 1975–2008 and mark–recapture estimates for 1998–2000. Lines show upper and lower limits of index escapement goal range.




Figure 10.—Counts of Chinook salmon at a weir and in survey counts in the index area of the King Salmon River, 1975–2008. Lines show upper and lower limits of index escapement goal range.

Table 13.—Estimated harvests and escapement, by size class, of Situk River Chinook salmon, 1976–2008.

Year	Harvests below weir				Weir count				Harvest above weir				Estimated escapement <sup>a</sup>			
	182–70 Gillnet	Subsistence	Sport	Total	Small	Medium	Large	Total	Small	Medium	Large	Total	Small <sup>b</sup>	Medium	Large	Total
1976	1,002	41	200	1,243		520	1,421	1,941						520	1,421	1,941
1977	833	24	244	1,101		148	1,732	1,880						148	1,732	1,880
1978	382	50	210	642		295	808	1,103						295	808	1,103
1979	1,028	25	282	1,335		470	1,284	1,754						470	1,284	1,754
1980	969	57	233	1,259		220	905	1,125						220	905	1,125
1981	858	62	130	1,050		105	702	807						105	702	807
1982	248	27	63	338		177	434	611						177	434	611
1983	349	50	52	451		257	592	849						257	592	849
1984	512	89	151	752		475	1,726	2,201						475	1,726	2,201
1985	484	156	511	1,151		461	1,521	1,982						461	1,521	1,982
1986	202	99	37	338		505	2,067	2,572						505	2,067	2,572
1987	891	24	395	1,310		505	1,379	1,884						505	1,379	1,884
1988	299	90	132	521		193	885	1,078		39	17	56		154	868	1,022
1989	1	496 <sup>c</sup>	0	497	972	243	637	1,852		0	0	0	991	243	637	1,871
1990	0	516	0	516	147	499	628	1,274		0	0	0	236	499	628	1,363
1991	786	220	67	1,073	584	132	897	1,613	2	19	8	29	582	114	889	1,585
1992	1,504	341	127	1,972	131	236	1,618	1,985	3	28	23	54	129	207	1,595	1,931
1993	790	202	50	1,042	2,730	490	980	4,200	92	13	28	133	2,638	477	952	4,067
1994	2,656	367	397	3,420	1,634	1,471	1,311	4,416	50	80	40	170	1,584	1,391	1,271	4,246
1995	8,106	528	1,180	9,814	2,914	617	4,700	8,231	84	52	370	506	2,830	565	4,330	7,725
1996	3,717	478	1,270	5,465	1,374	602	2,175	4,151	568	107	375	1,050	1,061	495	1,800	3,356
1997	2,339	352	802	3,493	1,729	582	2,690	5,001	467	148	812	1,427	1,521	434	1,878	3,833
1998	2,101	594	494	3,189	3,125	851	1,353	5,329	405	206	429	1,040	2,902	645	924	4,471
1999	3,810	588	605	5,003	473	301	1,947	2,721	150	112	486	748	396	189	1,461	2,046
2000	1,318	594	352	2,264	413	161	2,518	3,092	211	60	733	1,004	381	101	1,785	2,267
2001	1,087	402	45	1,534	463	102	696	1,261	300	5	40	345	163	97	656	916
2002	1,078	416	63	1,557	300	448	1,024	1,772	18	24	24	66	282	424	1,000	1,706
2003	2,342	600	414	3,356	334	329	2,615	3,278	108	30	498	636	226	299	2,117	2,642
2004	1,222	396	294	1,912	348	419	796	1,563	3	7	41	51	345	412	755	1,512
2005	1	140	101	242	178	263	613	1,054	0	0	0	0	178	263	613	1,054
2006	19	192	0	211	307	348	749	1,404	0	0	0	0	307	348	749	1,404
2007	83	158	0	241	1,268	228	677	2,173	0	0	0	0	1,268	228	677	2,173
97–06	1,532	427	317	2,276	767	380	1,500	2,648	166	59	306	532	670	321	1,194	2,185
2008	91	325	0	416	124	221	453	798	0	0	0	0	124	221	453	798

<sup>a</sup> Escapement from McPherson et al. (2005), based on age composition.

<sup>b</sup> Small Chinook escapement includes 1- and 2-ocean jacks from 1990 to 1996; 1-ocean fish not counted before 1990. <sup>c</sup> Non-retention regulation in effect in 1989 and 1990; 2006–2008, estimated personal use harvest of 400 large Chinook in 1989, 415 in 1990, and 109 in 1991.




Figure 11.—Counts of large Chinook salmon at the Situk River weir, 1975–2008. Lines show upper and lower limits of escapement goal range.

Table 14.—Mark–recapture estimates of large Chinook salmon escapement in Chilkat River, 1991–2008.

Year	Escapement estimate	SE
1991	5,897	1,005
1992	5,284	949
1993	4,472	851
1994	6,795	1,057
1995	3,790	805
1996	4,920	751
1997	8,100	1,193
1998	3,675	565
1999	2,271	408
2000	2,035	334
2001	4,517	722
2002	4,051	429
2003	5,657	690
2004	3,422	456
2005	3,366	780
2006	3,039	454
2007	1,452	228
98–07 Avg.	3,341	507
2008	2,833	530

Source: From Chapell *In prep b*.

Weather, logistics, run timing, etc., can make it difficult for a single surveyor to complete all the index surveys annually under good or excellent conditions. Thus, alternate surveyors are selected to conduct the counts when the primary surveyor

cannot. Also, new surveyors take on primary responsibilities at infrequent intervals. Because between-observer variability and bias can be significant (Jones III et al. 1998b), new surveyors must be trained and calibrated against the primary surveyor to provide consistency and continuity in the data.

Estimates of total escapement (direct estimates or expanded counts) are needed when comparing escapements among watersheds or for estimating exploitation rates and spawner/recruit relationships. Though survey and tributary expansion factors have been endorsed by the PSC since 1981, the original expansion factors were developed on the basis of judgment rather than on empirical data (Pahlke 1997b), and error associated with these expansions can be large. Johnson et al. (1992) showed that expansion factors for the Chilkat River, for example, greatly underestimated escapement to that watershed. ADF&G recognized the need to develop better expansions throughout the region, and has independently estimated distribution and escapement for Chinook salmon in the Unuk (Pahlke et al. 1996; Jones III and McPherson 1999, 2000), Chickamin (Pahlke 1996, 1997a), Stikine (Pahlke and Etherton 1999; Bernard et al. 2000), Taku (Pahlke and Bernard 1996; McPherson et al. 1998a; Jones III et al. 2010), Keta (Brownlee et al. 1999), Blossom

(Pahlke and Magnus 2005-2006), and Alsek rivers (Pahlke et al. 1999; Pahlke and Waugh 2006). Total escapement projects are continuing on many of those rivers.

On the basis of information collected on the Unuk and Chickamin rivers, expansion factors for the four Behm Canal systems were revised in 1996 and again in 2002. After 3 mark-recapture experiments, the expansion factor for the Keta River was revised again in 2001, and the Blossom River in 2007. The expansion factor for the King Salmon River was based on 10 years of weir counts compared with aerial surveys, and the expansion factor for Andrew Creek was based on 4 years of paired weir and survey counts. The expansion factor for the Taku River was revised in 1999 after 5 years of mark-recapture data. The expansion factor for the Alsek River was revised in 2002 based on 4 years of mark-recapture studies

and again in 2004. The most current estimates for the expansion factors and variances around them are presented in Appendices B2–B10. Variances around the estimates of total escapement in Appendix A4 are shown in Appendix A5. Some of these expansions are different from those reported in previous years, as they are revised each time another year of data is collected. Mark-recapture information was collected on the Stikine, Taku and Unuk rivers in 2008, and will continue annually.

Changing the escapement goals, however, requires a formal review by ADF&G and the CTC of the PSC, as was done for the Situk River in 1991 and 2005, the Behm Canal systems in 1994, and King Salmon River in 1997. The Andrew Creek escapement goal was also revised in 1998 to a range of 650 to 1,500 total large spawners (Clark et al. 1998). The DFO and the TTC are included in any review of Taku, Stikine or Alsek river goals.




Figure 12.—Mark-recapture estimates of large Chinook salmon escapement to the Chilkat River, 1991–2008. Lines show upper and lower limits of escapement goal range.

Table 15.—Comparison between primary and two alternate observer counts in survey training flights conducted in 2008.

Index area	Cond	Primary observer	Alternate observer 1	Percent	Comments
Keta River	N	363	450	41.6%	backseat
Blossom R.	E	257	242	94.2%	backseat
South Fork	N	43	87	202.3%	backseat
Andrew Creek	N	402	240	59.7%	replicate
Systems with Chinook, chum and pink salmon				Avg. 99.4%	
				Med. 76.9%	
		Primary observer	Alternate observer 2		
Tatsamenie	N	1083	880	81.3%	replicate
Tatsamenie	N	710	630	88.7%	backseat
Kowatua	E	632	417	66.0%	backseat
Kowatua	N	620	690	111.3%	backseat
Systems with Chinook and sockeye only				Avg. 86.8%	
				Med. 85.0%	

Notes: Conditions (cond.) - E = excellent, N = normal, P = poor.

In 1998, a revised stock-recruitment analysis by ADF&G and DFO staff estimated that the escapement goal for the Klukshu River should range between 1,100 and 2,300 spawners (McPherson et al. 1998b). Escapement goals for the Taku and Stikine rivers were approved in 1999 (McPherson et al. 2000; Bernard et al. 2000) and for the Chilkat River in 2003 (Ericksen and McPherson 2004).

Expansion factors and escapement goals will continue to be revised as more studies are completed that include both index counts and estimates of total escapement. Any change in survey methods or observers must take into account the comparability of historical data with new data. Year-to-year consistency and repeatability of index counts may be more important than their absolute accuracy to agencies that compare escapement estimates between years.

Currently, only one of the 22 minor producers in the region and 7 of 9 medium producing watersheds are included in the index survey program. Prior to 1997, counts from these streams were expanded to represent the escapement of all streams in minor and medium producing categories. The King Salmon River is unique among Southeast Alaska Chinook populations as the only island system, and using it to represent the other 21 small systems most likely produces inaccurate estimates of total escapement. However, because escapements to

small and medium systems are a small proportion of the total regional escapement, errors in those estimates have little effect on estimates of regional escapement. In 1997, the method used to expand the index counts to a total regional escapement estimate was revised based on over 20 years of systematic escapement surveys in Southeast Alaska and the transboundary rivers (Pahlke 1998). The revised method assumes the sum of the expanded indices accounts for approximately 90% of the total escapement and that number is expanded to account for the remaining 10%. This method is believed to more accurately reflect the contribution to regionwide escapement of the unsurveyed systems.

Escapement goal revisions based on spawner-recruit analysis require a long-time series of age and sex composition data along with total escapement estimates. Age, sex, and length composition estimates for all sampled Chinook stocks in Southeast Alaska and transboundary rivers are presented in Appendices A7 and A8. An interesting pattern became apparent in 1999, when the largest fish were observed in the southern systems and average size decreased toward the north. In 2000 and 2001, the largest fish were again seen in the southern systems, but fish in 2 of the northern systems, the Chilkat and Alsek rivers, were larger than Chinook salmon in the central systems. The trend has continued since 2002, with the smallest fish in the region returning to the Taku River and Andrew

Creek. Many (up to 75%) of the 2-ocean fish sampled on the Blossom, Keta and Chickamin rivers were of legal size (28 in TL, or approximately 625 mm MEF), which is uncommon in other systems in Southeast Alaska. Another interesting pattern is that the variance in mean length-at-age is consistently less for females than males. Mean lengths at age were tested for differences between systems (Appendices A9-A11). Estimated age compositions varied greatly between systems.

Sampling strategies were designed to make the estimated age and sex distributions relatively unbiased for age-.2 to age-.5 fish. A weir was used to sample the Situk and Alsek rivers; stratified mark-recapture studies were used on the Chilkat, Taku, Stikine, and Unuk rivers; and non-selective rod and reel and/or carcass sampling was used on the Chickamin, Blossom, Keta, Andrew Creek and King Salmon systems. Therefore, comparisons of length or age compositions between stocks within the age-.2. to age-.5 should be relatively unbiased. The Situk River is the only Chinook system in Southeast Alaska where the escapement of age-.1 jacks is estimated annually. The mean length-at-age data are unbiased for all stocks.

## ACKNOWLEDGMENTS

Many people provided valuable assistance in this study. David Magnus, Phil Richards, Todd Johnson, Tom Rockne, Alex Blaine, Roger Hayward and Dale Brandenburger conducted foot surveys; Sarah Power provided biometric support and reviewed the draft manuscript; John Der Hovanisian reviewed and edited the draft manuscript; Scott McPherson provided the Situk River data and the age and length summaries; Joanne MacClellan edited the manuscript for publication; Amy Holm provided logistical help. Bill Waugh of DFO provided weir counts from transboundary systems.

## REFERENCES CITED

- ADF&G (Alaska Department of Fish and Game). 1981. Proposed management plan for Southeast Alaska Chinook salmon runs in 1981. Southeast Region, Alaska Department of Fish and Game Division of Commercial Fisheries. Regional Report 1J81-3, Juneau.
- Beak Consultants Limited. 1981. Preliminary analysis of the potential impact of hydroelectric development of the Stikine River system on biological resources of the Stikine River estuary. Report for the British Columbia Hydro and Power Authority. Richmond, B. C.
- Bernard, D. R., S. A. McPherson, K. A. Pahlke, and P. Etherton. 2000. Optimal production of Chinook salmon from the Stikine River. Alaska Department of Fish and Game, Fishery Manuscript No. 00-1, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fms00-01.pdf>
- Bigelow, B. B., B. J. Bailey, M. M. Hiner, M. F. Schellekens, and K. R. Linn. 1995. Water resources data Alaska water year 1994. U. S. Geological Survey Water Data Report AK-94-1, Anchorage.
- Brownlee, K. M., S. A. McPherson, and D. L. Magnus. 1999. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Blossom and Keta rivers, 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-45, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds99-45.pdf>
- Bugliosi, E. F. 1988. Hydrologic reconnaissance of the Chilkat River Basin, Southeast Alaska. U. S. Geological Survey Water Resources Investigation Report 88-4021, Anchorage.
- Chapell, R. S. *In prep a.* Production and spawning distribution of Chilkat River Chinook salmon in 2006. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Chapell, R. S. *In prep b.* Production, escapement, and juvenile tagging of Chilkat River Chinook salmon in 2007. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Clark, J. H., S. A. McPherson, and D. M. Gaudet. 1998. Biological escapement goal for Andrew Creek Chinook salmon. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J98-08, Juneau. <http://www.sf.adfg.state.ak.us/FedAidPDFs/RIR.5J.1998.08.pdf>
- Der Hovanisian, J. A., P. Etherton, and K. A. Pahlke. 2005. Abundance of the Chinook salmon escapement on the Stikine River, 2003. Alaska Department of Fish and Game, Fishery Data Series No. 05-25, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/Fds05-25.pdf>

## REFERENCES CITED (Continued)

- Der Hovanisian, J. A., K. A. Pahlke, and P. Etherton. 2001. Abundance of the Chinook salmon escapement on the Stikine River, 2000. Alaska Department of Fish and Game, Fishery Data Series No. 01-18, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds01-18.pdf>
- Der Hovanisian, J. A., K. A. Pahlke, and P. Etherton. 2003. Abundance of the Chinook salmon escapement on the Stikine River, 2001. Alaska Department of Fish and Game, Fishery Data Series No. 03-09, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds03-09.pdf>
- Der Hovanisian, J. A., K. A. Pahlke, and P. Etherton. 2004. Abundance of the Chinook salmon escapement on the Stikine River, 2002. Alaska Department of Fish and Game, Fishery Data Series No. 04-08, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds04-08.pdf>
- Efron, B., and R. J. Tibshirani. 1993. 1st edition. An introduction to the bootstrap. Chapman and Hall, New York, NY
- Ericksen, R. P. 2005. Escapement, terminal harvest, and juvenile tagging of Chilkat River Chinook salmon in 2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-68, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds05-68.pdf>
- Ericksen, R. P., and R. S. Chapell. 2006. Production and spawning distribution of Chilkat River Chinook salmon in 2005. Alaska Department of Fish and Game, Fishery Data Series No. 06-76, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds06-76.pdf>
- Ericksen, R. P., and S. A. McPherson. 2004. Optimal production of Chinook salmon from the Chilkat River. Alaska Department of Fish and Game, Fishery Manuscript No. 04-01, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fms04-01.pdf>
- Freeman, G. M., and S. A. McPherson. 2003. Spawning abundance of Chinook salmon in the Chickamin River in 2001. Alaska Department of Fish and Game, Fishery Data Series 03-14, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds03-14.pdf>
- Freeman, G. M., and S. A. McPherson. 2004. Spawning abundance of Chinook salmon in the Chickamin River in 2002. Alaska Department of Fish and Game, Fishery Data Series No. 04-09, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds04-09.pdf>
- Freeman, G. M., and S. A. McPherson. 2005. Spawning abundance of Chinook salmon in the Chickamin River in 2003. Alaska Department of Fish and Game, Fishery Data Series No. 05-63, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds05-63.pdf>
- Freeman, G. M., S. A. McPherson, and D. L. Magnus. 2000. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Keta River, 1999. Alaska Department of Fish and Game, Fishery Data Series No. 00-36, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds00-36.pdf>
- Freeman, G. M., S. A. McPherson, and D. L. Magnus. 2001. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Keta River, 2000. Alaska Department of Fish and Game, Fishery Data Series No. 01-19, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds01-19.pdf>
- Freeman, G. M., S. A. McPherson, and D. J. Reed. 2007. Spawning abundance of Chinook salmon in the Chickamin River in 2004. Alaska Department of Fish and Game, Fishery Data Series No. 07-13, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds07-13.pdf>
- Hendrich, C. F., J. L. Weller, S. A. McPherson, and D. R. Bernard. 2008. Optimal production of Chinook salmon from the Unuk River. Alaska Department of Fish and Game, Fishery Manuscript No. 08-03, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fms08-03.pdf>
- Howe, A. L., R. J. Walker, C. Olnes, K. Sundet, and A. E. Bingham. 2001. Revised Edition. Harvest, catch, and participation in Alaska sport fisheries during 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-25 (revised), Anchorage. [http://www.sf.adfg.state.ak.us/FedAidPDFs/fds98-25\(revised\).pdf](http://www.sf.adfg.state.ak.us/FedAidPDFs/fds98-25(revised).pdf)

## REFERENCES CITED (Continued)

- Johnson, R. E., R. P. Marshall, and S. T. Elliott. 1992. Chilkat River Chinook salmon studies, 1991. Alaska Department of Fish and Game, Fishery Data Series No. 92-49, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds92-49.pdf>
- Johnson, T. *In prep.* Age, sex, and length composition of Chinook salmon in the Chickamin River, 2006–2007. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Jones III, E. L., and S. A. McPherson. 1999. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Unuk River, 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-14, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds99-14.pdf>
- Jones III, E. L., and S. A. McPherson. 2000. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Unuk River, 1999. Alaska Department of Fish and Game, Fishery Data Series No. 00-22, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds00-22.pdf>
- Jones III, E. L., and S. A. McPherson. 2002. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Unuk River, 2000. Alaska Department of Fish and Game, Fishery Data Series No. 02-17, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds02-17.pdf>
- Jones III, E. L., S. A. McPherson, D. R. Bernard, and I. Boyce. 2010. Spawning abundance of Chinook salmon in the Taku River 1999-2007. Alaska Department of Fish and Game, Fishery Data Series 10-70, Anchorage. <http://www/sf.adfg.state.ak.us/FedAidPDFs/fds10-70.pdf>
- Jones III, E. L., S. A. McPherson, and D. L. Magnus. 1998a. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Unuk River, 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-23, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds98-23.pdf>
- Jones III, E. L., T. J. Quinn, and B. W. Van Alen. 1998b. Observer accuracy and precision in aerial and foot survey counts of pink salmon in a Southeast Alaska stream. *North American Journal of Fisheries Management* 18:832-846.
- Kissner, P. D. 1974. A study of Chinook salmon in Southeast Alaska. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1973-1974, Project AFS-41-2, Juneau.
- Kissner, P. D. 1978. Status of important native Chinook salmon stocks in Southeastern Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Performance Report, 1976-1977, Project F-9-9(18) AFS 41-5, Juneau. [http://www.sf.adfg.state.ak.us/FedAidPDFs/FRED F-9-9\(18\)AFS41-5.pdf](http://www.sf.adfg.state.ak.us/FedAidPDFs/FRED F-9-9(18)AFS41-5.pdf)
- Kissner, P. D. 1982. Status of important native Chinook salmon stocks in Southeastern Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Performance Report, 1981-1982, Project F-9-14, 23 (AFS 41-10), Juneau.
- McPherson, S., D. Bernard, J. H. Clark, K. Pahlke, E. Jones, J. Der Hovanisian, J. Weller, and R. Ericksen. 2003. Stock status and escapement goals for Chinook salmon stocks in Southeast Alaska. Alaska Department of Fish and Game, Special Publication No. 03-01, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/sp03-01.pdf>
- McPherson, S., and J. H. Clark. 2001. Biological escapement goal for King Salmon River Chinook salmon. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 1J01-40, Juneau.
- McPherson, S. A., D. R. Bernard, and J. H. Clark. 2000. Optimal production of Chinook salmon from the Taku River. Alaska Department of Fish and Game, Fishery Manuscript No. 00-2, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fms00-02.pdf>
- McPherson, S. A., D. R. Bernard, M. S. Kelley, P. A. Milligan, and P. Timpany. 1998a. Spawning abundance of Chinook salmon in the Taku River in 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-41, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds98-41.pdf>
- McPherson, S. A., and J. Carlile. 1997. Spawner-recruit analysis of Behm Canal Chinook salmon stocks. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 1J97-06, Juneau.

## REFERENCES CITED (Continued)

- McPherson, S. A., P. Etherton, and J. H. Clark. 1998b. Biological escapement goal for Klukshu River Chinook salmon. Alaska Department of Fish and Game, Fishery Manuscript No. 98-2, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fms98-02.pdf>
- McPherson, S. A., R. E. Johnson, and G. F. Woods. 2005. Optimal production of Chinook salmon from the Situk River. Alaska Department of Fish and Game, Fishery Manuscript No. 05-04, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/Fms05-04.pdf>
- Mecum, R. D. 1990. Escapements of Chinook salmon in southeast Alaska and transboundary rivers in 1989. Alaska Department of Fish and Game, Fishery Data Series No. 90-52, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds90-52.pdf>
- Mecum, R. D., and P. D. Kissner. 1989. A study of Chinook salmon in southeast Alaska. Alaska Department of Fish and Game, Fishery Data Series No. 117, Juneau. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds-117.pdf>
- Mood, A. M., F. A. Graybill, and D. C. Boes. 1974. Introduction to the theory of statistics, 3rd edition. McGraw-Hill Book Co., New York.
- Neter, J., W. Wasserman, and M. H. Kutner. 1990. Applied linear statistical models, 3rd edition. Irwin Publishing Company. Homewood, IL.
- Pahlke, K. A. 1995. Escapement of Chinook salmon in Southeast Alaska and Transboundary Rivers in 1994. Alaska Department of Fish and Game, Fishery Data Series No. 95-35, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds95-35.pdf>
- Pahlke, K. A. 1996. Abundance of the Chinook salmon escapement on the Chickamin River, 1995. Alaska Department of Fish and Game, Fishery Data Series No. 96-37, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds96-37.pdf>
- Pahlke, K. A. 1997a. Abundance and Distribution of the Chinook salmon escapement on the Chickamin River, 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-28, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds97-28.pdf>
- Pahlke, K. A. 1997b. Escapements of Chinook salmon in Southeast Alaska and transboundary rivers in 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-33, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds97-33.pdf>
- Pahlke, K. A. 1998. Escapements of Chinook salmon in Southeast Alaska and transboundary rivers in 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-33, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds98-33.pdf>
- Pahlke, K. A. 2007. Escapements of Chinook salmon in Southeast Alaska and Transboundary rivers in 2005. Alaska Department of Fish and Game, Fishery Data Series No. 07-62, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidpdfs/fds07-62.pdf>
- Pahlke, K. A., and D. R. Bernard. 1996. Abundance of the Chinook salmon escapement in the Taku River, 1989 to 1990. Alaska Fishery Research Bulletin 3(1):8-19, Juneau.
- Pahlke, K. A., and P. Etherton. 1997. Chinook salmon research on the Stikine River, 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-37, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds97-37.pdf>
- Pahlke, K. A., and P. Etherton. 1999. Abundance and distribution of the Chinook salmon escapement on the Stikine River, 1997. Alaska Department of Fish and Game, Fishery Data Series No. 99-6, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds99-06.pdf>
- Pahlke, K. A., and P. Etherton. 2000. Abundance of the Chinook salmon escapement on the Stikine River, 1998. Alaska Department of Fish and Game, Fishery Data Series No. 00-24, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds00-24.pdf>
- Pahlke, K. A., and P. Etherton. 2001a. Abundance of the Chinook salmon escapement on the Alsek River, 1999. Alaska Department of Fish and Game, Fishery Data Series No. 01-11, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds01-11.pdf>

## REFERENCES CITED (Continued)

- Pahlke, K. A., and P. Etherton. 2001b. Abundance of the Chinook salmon escapement on the Alsek River, 2000. Alaska Department of Fish and Game, Fishery Data Series No. 01-30, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds01-30.pdf>
- Pahlke, K. A., and P. Etherton. 2002. Abundance of the Chinook salmon escapement on the Alsek River, 2001. Alaska Department of Fish and Game, Fishery Data Series 02-20, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds02-20.pdf>
- Pahlke, K. A., P. Etherton, and J. A. Der Hovanisian. 2000. Abundance of the Chinook salmon escapement on the Stikine River, 1999. Alaska Department of Fish and Game, Fishery Data Series No. 00-25, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds00-25.pdf>
- Pahlke, K. A., P. Etherton, R. E. Johnson, and J. E. Andel. 1999. Abundance and distribution of the Chinook salmon escapement on the Alsek River, 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-44, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds99-44.pdf>
- Pahlke, K. A., and D. L. Magnus. 2005. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Blossom River, 2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-70, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds05-70.pdf>
- Pahlke, K. A., and D. L. Magnus. 2006. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Blossom River, 2005. Alaska Department of Fish and Game, Fishery Data Series No. 06-75, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds06-75.pdf>
- Pahlke, K. A., S. A. McPherson, and R. P. Marshall. 1996. Chinook salmon research on the Unuk River, 1994. Alaska Department of Fish and Game, Fishery Data Series No. 96-14, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds96-14.pdf>
- Pahlke, K. A., and B. Waugh. 2003. Abundance and distribution of the Chinook salmon escapement on the Alsek River, 2002. Alaska Department of Fish and Game, Fishery Data Series No. 03-20, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds03-20.pdf>
- Pahlke, K. A., and B. Waugh. 2004. Abundance of the Chinook salmon escapement on the Alsek River, 2003. Alaska Department of Fish and Game, Fishery Data Series No. 04-27, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds04-27.pdf>
- Pahlke, K. A., and B. Waugh. 2006. Abundance of the Chinook salmon escapement on the Alsek River in 2004. Alaska Department of Fish and Game, Fishery Data Series No. 06-12, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidpdfs/fds06-12.pdf>
- PSC (Pacific Salmon Commission). 1991. Escapement goals for Chinook salmon in the Alsek, Taku, and Stikine Rivers. Transboundary River Technical Report, TCTR (91)-4, Vancouver, B. C.
- PSC (Pacific Salmon Commission, U. S. Chinook Technical Committee (USCTC)). 1997. A review of stock assessment data and procedures for U. S. Chinook salmon stocks. Pacific Salmon Commission Report USTCHINOOK (97)-1, Vancouver, B. C.
- Richards, P., K. A. Pahlke, and P. Etherton. *In prep b*. Abundance of the Chinook salmon escapement on the Stikine River, 2007, and production and harvest of fish from brood year 2000. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Richards, P. J., K. A. Pahlke, J. A. Der Hovanisian, J. L. Weller, and P. Etherton. 2008. Abundance and distribution of the Chinook salmon escapement on the Stikine River in 2005, and production of fish from brood year 1998. Alaska Department of Fish and Game, Fishery Data Series No. 08-33, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds08-33.pdf>
- Weller, J. L., D. Evans, M. Brunette, and S. A. McPherson. *In prep*. Production of Chinook salmon in the Unuk River in 2007. Alaska Department of Fish and Game, Fisheries Data Series, Anchorage.
- Weller, J. L., and D. G. Evans. 2009. Estimation of the Escapement of Chinook Salmon in the Unuk River in 2006. Alaska Department of Fish and Game, Fishery Data Series No. 09-02, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds09-02.pdf>

## REFERENCES CITED (Continued)

- Weller, J. L., D. L. Magnus, D. J. Reed, and K. A. Pahlke. 2007a. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Blossom River, 2006. Alaska Department of Fish and Game, Fishery Data Series No. 07-66, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds07-66.pdf>
- Weller, J. L., and S. A. McPherson. 2003a. Estimation of the escapement of Chinook salmon in the Unuk River in 2001. Alaska Department of Fish and Game, Fishery Data Series 03-13, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds03-13.pdf>
- Weller, J. L., and S. A. McPherson. 2003b. Estimation of the escapement of Chinook salmon in the Unuk River in 2002. Alaska Department of Fish and Game, Fishery Data Series 03-15, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds03-15.pdf>
- Weller, J. L., and S. A. McPherson. 2006a. Estimation of the escapement of Chinook salmon in the Unuk River in 2004. Alaska Department of Fish and Game, Fishery Data Series No. 06-07, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds06-07.pdf>
- Weller, J. L., and S. A. McPherson. 2006b. Estimation of the escapement of Chinook salmon in the Unuk River in 2005. Alaska Department of Fish and Game, Fishery Data Series No. 06-59, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds06-59.pdf>
- Weller, J. L., D. J. Reed, and G. M. Freeman. 2007b. Spawning abundance of Chinook salmon in the Chickamin River in 2005. Alaska Department of Fish and Game, Fishery Data Series No. 07-63, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds07-63.pdf>



## **APPENDIX A**

Appendix A1.—Survey escapement goals and system goals for large Chinook salmon, Southeast Alaska and transboundary rivers, as accepted by the Alaska Department of Fish and Game, Canadian Department of Fisheries and Oceans, Chinook Technical Committee, and Transboundary Technical Committee, 2009.

River	Index areas	Index survey goal <sup>a</sup>			System goal <sup>b</sup>		
		Point est.	Range		Point est.	Range	
			Lower	Upper		Lower	Upper
Alsek <sup>c</sup>	Klukshu		1,100	2,300			
Taku <sup>d</sup>	5 tributaries				25,500	19,000	36,000
Stikine <sup>e</sup>	Little Tahltan	3,300	2,700	5,300	17,500	14,000	28,000
Situk <sup>f</sup>	All				730	450	1,050
Chilkat <sup>g</sup>	All				2,200	1,750	3,500
Andrew Cr. <sup>h</sup>	All	400	325	750	800	650	1,500
Unuk <sup>i</sup>	6 tributaries	800	375	800	2,764	1,800	3,800
Chickamin <sup>j</sup>	8 tributaries	525	450	900			
Blossom <sup>j</sup>	All	300	250	500			
Keta <sup>j</sup>	All	300	250	500			
King Salmon R. <sup>k</sup>	All	100	80	160	150	120	240

<sup>a</sup> Index survey goal corresponds to the peak or highest single day count of large spawners in annual survey counts. However, the Alsek River survey goal is germane to fish of all sizes, counted at the Klukshu River weir.

<sup>b</sup> System goal corresponds to the estimated total escapement of large spawners in the river system, estimated from mark–recapture studies, weir counts or expanded survey counts.

<sup>c</sup> McPherson et al. (1998b). This goal awaits approval by the Centre for Scientific Advice – Pacific.

<sup>d</sup> McPherson et al. (2000).

<sup>e</sup> Bernard et al. (2000).

<sup>f</sup> McPherson et al. (2005).

<sup>g</sup> Ericksen and McPherson (2004).

<sup>h</sup> Clark et al. (1998).

<sup>i</sup> Hendrich et al. (2008).

<sup>j</sup> McPherson and Carlile (1997).

<sup>k</sup> McPherson and Clark (2001).

Appendix A2.–Coordinates of Chinook salmon survey areas in Southeast Alaska and transboundary rivers.

Waypoint	Description	Latitude	Longitude
King Salmon River			
1	King Salmon River top of index area	N58 04.662	W134 24.073
Taku River drainage			
2	Windy Lake fuel cache, near Nakina	N59 05.262	W132 55.529
3	Nakina, Grizzly Bar, bottom of IA1	N59 03.494	W133 01.789
4	Nakina, Top of IA1, Taku	N59 04.581	W133 01.264
5	Top of IA2, Nakina River, weir site	N59 05.866	W133 00.646
6	Top of IA3, Nakina River	N59 07.560	W132 55.143
7	Top of IA4, Nakina Canyon, telegraph trail	N59 11.048	W132 50.210
8	Top of Tseta Creek, Taku River	N59 02.011	W132 13.255
9	Long Lake fuel cache, near Nahlin River	N58 44.557	W131 30.607
10	Top of IA3, Nahlin River	N58 39.557	W131 10.259
250	Top of IA2, Nahlin River	N 58 43 432	W131 17 501
11	Top of IA1, Nahlin River	N58 48.541	W131 28.027
12	Bottom of IA1, Nahlin River	N58 53.126	W131 45.054
73	Nahlin Cabin riffles	N58 45.866	W131 21.299
249	Old smolt camp, Nahlin River	N58 44 494	W131 18 796
13	Bottom of Dudidontu Index Area	N58 38.816	W131 48.707
14	Fork with Matsatu Creek, Dudidontu	N58 35.358	W131 47.002
15	Top of Dudidontu IA, maybe need to be revised	N58 31.005	W131 50.585
32	Bottom of Kowatua River IA, Taku	N58 30.324	W132 32.512
33	Bottom of Tatsamenie IA, Taku	N58 28.647	W132 23.273
227	Big Trapper fuel	N58 27.869	W132 38.379
252	Hackett River weir site	N 58 15 544	W 131 48 411
251	Hackett River Chinook spawning observed	N 58 13 570	W 131 45 430
Stikine River drainage			
18	Top end of Little Tahltan River IA, Stikine	N58 11.896	W131 28.876
19	Saloon Lake, near Tahltan	N58 07.473	W131 22.752
20	Little Tahltan River weir	N58 07.328	W131 19.239
91	Chutine Chinook spawning	N57 41.496	W132 18.082
160	Verrett Cr	N56 41.956	W130 59.565
50	Andrew Creek, top IA	N56 36.008	W132 09.408
51	Andrew Creek, mouth	N56 38.398	W132 12.002
	Christina Creek	N57 14.432	W131 52.179
	Johnny Tashoots Cr, outlet to Tahltan Lk.	N58 00.720	W131 34.763
Alsek River drainage			
254	Klukshu Weir	N60 06.979	W137 01.978
	Blanchard R. Mouth	N60 00.843	W136 52.318
253	Blanchard R. top survey	N60 00.843	W136 52.318
255	Tatsamenie/Goat Cr.	N59 50.618	W136 39.248
	Tat/Low Fog	N59 36.015	W137 14.637
21	Bottom Takhanne River IA, Alsek	N60 05.687	W136 59.386
22	Top Takhanne River IA, Alsek	N60 06.493	W136 56.838

-continued-

Appendix A2.–Page 2 of 2.

Waypoint	Description	Latitude	Longitude
Unuk River drainage			
23	Bottom of Eulachon River IA, Unuk	N56 06.597	W131 07.293
24	Top of Eulachon River IA, 2 <sup>nd</sup> avalanche chute	N56 09.216	W131 07.884
258	Clear Creek, Unuk	N 56 08 104	W 130 58 347
165	Unuk fuel	N56 05.151	W131 05.363
260	Unuk Camp	N 56 07 683	W 130 48 824
166	Genes Lake	N56 12.654	W130 51.733
167	Kerr Creek	N56 11.003	W130 55.792
Chickamin River drainage			
25	Chickamin River camp	N55 49.493	W130 52.826
26	Bottom King Creek IA, Chickamin River	N55 50.507	W130 51.162
27, 28	Top of King Creek IA, Chickamin	N55 49.149	W130 48.006
37	Top of King Creek king distribution, Chickamin	N55 48.523	W130 46.940
38	Mouth of King Creek	N55 50.441	W130 50.848
39	Bottom Humpy Creek IA, Chickamin	N55 50.812	W130 52.309
40	Top Humpy Creek IA, Chickamin	N55 52.076	W130 53.638
53	Indian Creek, Chickamin, mouth	N55 57.355	W130 41.532
54	Indian Creek, Chickamin, top	N55 59.534	W130 40.017
55	Lucky Jake Creek, Chickamin	N55 59.207	W130 38.001
56	Ranger Paige Creek, Chickamin	N55 59.701	W130 36.985
57	Butler Creek mouth	N56 02.357	W130 43.354
58	Butler Creek, top	N56 02.870	W130 43.359
59	Clear Falls, Chickamin	N55 58.812	W130 45.560
60	Top of King Creek foot survey	N55 49.262	W130 48.449
168	Chickamin fuel	N55 49.610	W130 54.445
Blossom and Keta River drainages			
41	Apparent barrier on Blossom River, top IA	N55 30.285	W130 28.708
43	Bottom of Keta River	N55 19.880	W130 29.099
47	Top of Index area Keta River	N55 27.430	W130 20.946
226	Blossom Camp	N55 25.802	W130 33.260
B	Blossom Fuel	N55 21.995	W130 37.499

## **TAKU RIVER DRAINAGE**

### **NAKINA RIVER**

Stream Code: 111-32-220                      Anadromous Stream Number: 111-32-10320-2999

Peak Spawning: August 4

Survey Dates: August 1–7

Survey duration: 1.5hr

In years of good escapement several hundred Chinook salmon can be observed from the junction of the Sloko and Nakina Rivers upstream to Grizzly Bar, a distance of about 5.5 miles. This area is not surveyed because of the few Chinook present. Sockeye and pink salmon in survey area.

Fuel cache at Windy Lake.

#### Survey Index Area I

50 meters below Grizzly Bar (sport cabins and tent frames on gravel bar) to the heavy rapids and small gorge 2.4 km upstream.

The area from 50 m below Grizzly Bar upstream for about 650 m is always well seeded, while the area above to the small gorge is only well utilized during years of good escapement. Count by 10s.

#### Survey Index Area II

From the heavy rapids and small gorge upriver to the weir site.

The area has never been well utilized (except from old cabins to weir), however use increases in years of good escapements.

#### Survey Index Area III

Weir to major gorge 3.2 km upstream. This is an excellent spawning area with largest spawning concentration just below the gorge. Count by 10s.

#### Survey Index Area IV

Gorge to barrier approximately 2 km below Telegraph Trail crossing (old cabin). In years of large escapements or high water significant numbers of Chinook salmon spawn in this area. Survey light-windy, narrow canyon.

### **NAHLIN RIVER**

Stream Code: 111-32-270                      Anadromous Stream Number: 111-32-10320-2998

Peak Spawning: July 24

Survey Dates: July 22–28

Survey Duration: 2.2 hrs

Spawning occurs earliest in headwaters above Beaver Dam Valley. Chinook do spawn above and below the survey areas but not in large numbers. Fuel cache is at Long Lake.

---

-continued-

Survey Index Area I

Nahlin Crossing (Outlet of Tedideech Ck, cabin, cable crossing of Telegraph Trail) upriver to Beaver Dam Valley (start of slow moving water, three large rocks in river, old weir site). From Nahlin Crossing to the junction of Kawdy Creek Chinook Salmon spawning is sparse, usually less than several hundred fish. From the junction of Kawdy Creek upriver to the three large rocks, dense spawning occurs in years of large escapements.

Survey Index Area II

Three large rocks at beginning of slow water (Beaver Dam Valley) upriver for about 13 km to faster moving water. This area is very difficult to survey, except on bright sunny days, because of deep, dark water and many meanders. Only one regular spawning area near old trapper cabin riffles (Waypoint 73).

Survey Index Area III

Beginning of faster moving and shallower water upriver for about 8 km to the area where the river forks, up each fork about 2 km. Highest percentage of spawning occurs in this area.

In some years as many sockeye as Chinook are present in this area, and they often have not colored up yet.

**TATSAMENIE RIVER**

Stream Code: 111-32-240 Anadromous Stream Number: 111-32-10320-2997

Peak Spawning: August 23

Survey Dates: August 20–26

Survey Duration: 45 minutes

Latest spawning in Taku River drainage. Sometimes semi-glacial. Survey early to avoid glacial melt. Chinook spawn above Survey Area II but not in large numbers (at outlet to Big Tatsamenie Lake). Sockeye in area. Old sockeye weir site at cabins below little lake. New sockeye weir at outlet to big lake. Fuel at Big Trapper Lake.

Survey Index Area I

Tatsatua River confluence to Little Tatsamenie Lake. Largest concentration of spawning Chinook opposite meadow about 200 m above Tatsatua confluence. Carcass weir goes in right below meadow. Count by 10s.

Survey Index Area II

Inlet stream to Little Tatsamenie Lake upstream to confluence of the two forks. Then fly fast to top of rapids and count outlet to Big Lake, below sockeye weir.

---

-continued-

**DUDIDONTU RIVER**

Stream Code: 111-32-280 Anadromous Stream Number: 111-32-10320-2996

Peak Spawning: August 2

Survey Dates: July 30–August 4

Survey Duration: 45 minutes

Spawning well distributed over large area. Many trout in upper waters near swamp. One of the easiest surveys, no other species of salmon to worry about, no big trees or cliffs.

Fuel at Long Lake.

Survey Index Area

Upper end of large canyon upstream to approximately 18 km past confluence with Matsatu Creek, near Alkali Pond. Survey lower 2 k of Matsatu Cr, both forks. Survey upper end of index area at 30–40 mph, slowing when concentrations of fish observed, usually on riffles from old beaver dams. Large beaver dam swamp in the middle of the survey area. Chinook continue on upriver for long way beyond index area.

**KOWATUA RIVER**

Stream Code: 111-32-240 Anadromous Stream Number: 111-32-10320-2994

Peak Spawning: August 20

Survey Dates: August 18–24

Survey Duration: 30 minutes

Late spawning Chinook run, just slightly earlier than Tatsamenie. Spawning occurs below Index Area, but not in large numbers. Many sockeye salmon in area. River is semi-glacial at best.

Fuel at Big Trapper Lake.

Survey Index Area

Little Trapper Lake outlet to confluence with small glacial stream that flows into Kowatua River from the South (River Right) about 8 km below Little Trapper Lake. Sockeye salmon weir at outlet to Little Trapper.

**TSETA CREEK**

Stream Code: 111-32-275 Anadromous Stream Number: 111-32-10320-2993

Peak Spawning: July 29

Survey Dates: July 28–August 2

Survey Duration: 1hr

---

-continued-

Densest spawning occurs for 3 km below barrier falls at upper end. Spawning scattered in the rest of the index area. Most of the river is surveyed at 20–30 mph, regular speed at upper end.

Trapper cabin at small lake near upper end. Tseta was removed from Index Surveys in 1999. Fuel at Windy Lake or Long Lake, survey on the way from Nakina to Nahlin.

Survey Area

From barrier falls downriver to start of canyon just above confluence with Nahlin River.

**STIKINE RIVER DRAINAGE**

**LITTLE TAHLTAN RIVER**

Stream Code: 108-80-120 Anadromous Stream Number: 108-40-10150-2999

Peak Spawning: August 3

Survey Dates: July 28–August 6

Survey Duration: 1hr

Spawning is most intense from Clay Corner (high muddy bank that usually causes fairly poor visibility downriver) upriver to confluence with outlet to Saloon Lake. In years of high escapement spawning continues in high density above this area. Some spawning occurs above index area. Weir has been operated by DFO at confluence with Tahltan River since 1985. Fuel cache was at Saloon Lake, but is now provisioned by truck from Dease Lake, in cooperation with DFO.

Survey Index Area

From confluence with mainstem Tahltan River upriver for about 18 km to steep walled canyon.

Count by 10s.

**MAINSTEM TAHLTAN RIVER**

Stream Code: 108-80-100 Anadromous Stream Number: *unassigned (Canadian stream)*

Peak Spawning: August 8

Survey Dates: August 5–10

Survey Duration: 1hr

Most concentrated spawning occurs below confluence with Little Tahltan River and for 2 km above confluence with Beatty Creek. Chinook salmon spawn above index area and in Johnny Tashoots Creek. Very glacial, try to survey early in morning after cold nights.

Survey area

From canyon 1.5 km above Little Tahltan downriver to junction with Stikine. Removed from annual surveys after telemetry study in 1997.

---

-continued-

**BEATTY CREEK**

Stream Code: 108-80-115 Anadromous Stream Number: *unassigned (Canadian stream)*  
Peak Spawning: August 2  
Survey Dates: July 28–August 6  
Survey Duration: 15 min

Uniform spawning in survey area. Chinook spawn for 15 km above survey area in small numbers. Narrow windy canyon, survey light. Removed from annual surveys after telemetry study in 1997.

Survey Area

From confluence with Tahltan upstream through first canyon (approximately 4 km).

**ANDREW CREEK**

Stream Code: 108-40-020 Anadromous Stream Number: 108-40-10150-2008  
Peak Spawning: August 15  
Survey Dates: August 10–August 17  
Survey Duration: 20 min

Spawning throughout survey area, concentrated in lower river. Pinks, chums and sockeye present. Refuel in Wrangell.

Survey Area

Slough to barrier. Count both forks, keep North Fork separate.

**ALSEK RIVER DRAINAGE**

**KLUKSHU RIVER**

Stream Code: 182-30-020 Anadromous Stream Number: *unassigned (Canadian stream)*  
Peak Spawning: August 1  
Survey Dates: July 30–August 3  
Survey Duration: 1hr 15 min

Little spawning in lower 5 km and meander area further upriver–survey these areas at faster speed. Difficult survey stream because of overhanging trees and sockeye salmon. Very windy in afternoon, so survey as rapidly as possible. Fuel transported by DFO and stored at weir site at Dalton Post. Do not fly on weekends if possible, because parking lot where fuel is stored will be full of fishermen. Proportion observed was always very low, so surveys have been discontinued since the weir looks like it will be a long term program.

Survey Area

Weir upriver to Klukshu Lake

---

-continued-

**TAKHANNE RIVER**

Stream Code: 182-30-043 Anadromous Stream Number: *unassigned (Canadian stream)*

Peak Spawning: August 1

Survey Dates: July 30–August 3

Survey Duration: 15 min

Most fish concentrated at lower end. Survey after Blanchard about 10:30 am. Can be very windy in afternoon, tight canyon, survey light. Some sockeye in area.

Survey Area

Confluence with Tatshenshini River upriver to waterfall.

**BLANCHARD RIVER**

Stream Code: 182-30-050 Anadromous Stream Number: 182-30-10100-2999

Peak Spawning: August 1

Survey Dates: July 30–August 3

Survey Duration: 1 hr

Most concentrated spawning occurs below bridge to confluence with Tatshenshini. Survey in early morning because of glacial melt. Some sockeye in area.

Survey Area I

Bridge downriver to confluence with Tatshenshini.

Survey Area II

Bridge upriver to Blanchard Lake. Spawning scattered and mostly just below lake in rock piles. Survey fast, slowing down when concentrations of fish occur and at outlet to Lake. Sockeye spawning in upper area.

**GOAT CREEK**

Stream Code: 182-30-045 Anadromous Stream Number: *unassigned (Canadian stream)*

Peak Spawning: August 1

Survey Dates: July 30–August 3

Survey Duration: 15 min

Survey Area

From just above the bridge at beginning of canyon, downriver to glacial Tatshenshini.

---

-continued-

## UNUK RIVER DRAINAGE

### CRIPPLE CREEK

Stream Code: 101-75-30Q      Anadromous Stream Number: 101-75-10300-2030  
Peak Spawning: August 6  
Survey Dates: August 3–9  
Survey Duration: Foot survey, all day

Most intensive spawning occurs in long straight stretch about .8 km upstream from confluence with glacial water. Many brown bears in area. Overhanging trees make aerial survey difficult, stream should be surveyed by foot. Many chum salmon and some pinks in area. Fuel at private property near mouth of Unuk River. Helicopter landings in the Wilderness Area restricted to only those allowed under permit.

#### Survey Index Area

From confluence with glacial Unuk upriver to top of area of very extensive braiding.

### GENES LAKE CREEK

Stream Code: 101-75-30G      Anadromous Stream Number: 101-75-10300-2022  
Peak Spawning: August 27  
Survey Dates: August 15–27  
Survey Duration: Foot survey, all day

Because of overhanging trees this creek should be surveyed by foot. Spawning is very protracted. Because fish hold in clear pools it should be surveyed before peak spawning for best count. Lake should be surveyed at the same time, can be done by boat or helicopter. Many sockeye in system.

Many brown bears.

#### Survey Index Area

Lake inlet to small lake outlet upstream about 9 km

.

### KERR CREEK

Stream Code: 101-75-30K      Anadromous Stream Number: 101-75-10300-2019  
Peak Spawning: August 10  
Survey Dates: August 7–14  
Survey Duration: Foot survey, 4 hrs, helicopter 15 min.

#### Survey Index Area

Falls downstream to glacial water. In recent years visibility has got much worse due to influx of muddy river water.

**EULACHON RIVER**

Stream Code: 101-75-015 Anadromous Stream Number:101-75-10150

Peak Spawning: August 18

Survey Dates: August 14–21

Survey Duration: Foot survey, all day, helicopter 45 min.

Chinook hold in large numbers in the first two large pools below the fork. Heaviest spawning occurs just below and in the west fork. East Fork gets fair numbers in high water years.

Jet boat can get almost to the holding pools. Pinks, chums, and cohos may be present. Many bears.

Survey Index Area

From upper end of boat access to barrier falls.

**CLEAR CREEK**

Stream Code: 101-75-30C Anadromous Stream Number:101-75-10300-2014-3004

Peak Spawning: August 10

Survey Dates: August 7–14

Survey Duration: Foot survey, 2 hrs, helicopter 15 min.

Uniform spawning above confluence with Lake Creek. Chinook hold at mouth and in small narrow canyon (grotto) .5 km upstream. Very difficult to see into grotto from the air. Pinks, chums, sockeye present. Bears. Also called Kingsbury Creek.

Survey Index Area

Mouth of Creek to barrier falls.

**LAKE CREEK**

Stream Code: 101-75-30L Anadromous Stream Number: 101-75-10300-2014

Peak Spawning: August 10

Survey Dates: August 7–14

Survey Duration: helicopter 15 min.

Survey Index Area

Confluence with Clear Creek to falls. Spawning on riffles in lower river, near the big bend and in the falls pool. Pinks and chums present

---

-continued-

---

**CHICKAMIN RIVER DRAINAGE**

**SOUTH FORK**

Stream Code: 101-71-04S Anadromous Stream Number: 101-71-10040-2018  
Peak Spawning: August 18  
Survey Dates: August 14–21  
Survey Duration: helicopter 25 min.

Mainstem spawning. Survey early in day (first stream of day) as river is semi-glacial at best. Can vary in survey conditions dramatically in short period of time. Many pinks and chums. Fuel cache at private land at tidewater. Helicopter landings limited in Wilderness Area.

Survey Index Area

Confluence of middle fork of Chickamin and South Fork upriver to mouth of Barrier Creek.

**BARRIER CREEK**

Stream Code: 101-71-04A Anadromous Stream Number: 101-71-10040-2018-3010  
Peak Spawning: August 12  
Survey Dates: August 7–14  
Survey Duration: helicopter 10 min.

Survey Index Area

From confluence with South Fork to barrier falls 1.6 km upstream. Survey both forks.  
Pinks and chums in area.

**INDIAN CREEK**

Stream Code: 101-71-04I Anadromous Stream Number: 101-71-10040-2025  
Peak Spawning: August 10  
Survey Dates: August 7–14  
Survey Duration: helicopter 20 min.

Survey Index Area

From confluence with middle fork of Chickamin upstream to barrier falls. Spawning evenly distributed; many overhanging trees, pinks and chums.

---

-continued-

**BUTLER CREEK**

Stream Code: 101-71-04B Anadromous Stream Number: 101-71-10040-2015-3013

Peak Spawning: August 10

Survey Dates: August 7–14

Survey Duration: helicopter 15 min.

Small clear water tributary of upper Leduc River. Pinks and chums in system.

Survey Index Area: From mouth to falls.

**CLEAR FALLS**

Stream Code: 101-71-04C Anadromous Stream Number: 101-71-10040-2015-3009

Peak Spawning: August 10

Survey Dates: August 7–14

Survey Duration: helicopter 5 min.

Survey Index Area : Mouth to falls.

**LEDUC CREEK**

Stream Code: 101-71-04L Anadromous Stream Number: 101-71-10040-2015-3003

Peak Spawning: August 10

Survey Dates: August 7–14

Survey Duration: helicopter 10 min.

Survey Index Area.

Mouth to falls. Look carefully at mixing zone between Clearwater and muddy river.

**KING CREEK**

Stream Code: 101-71-04K Anadromous Stream Number: 101-71-10040-2006

Peak Spawning: September 1

Survey Dates: August 21–28

Survey Duration: helicopter 30 min.

Spawning occurs far upriver; latest system in Southeast. Chinook school in holes in lower river and are easiest to count there before spawning. Count by 10s. Pinks and chums in system.

Survey Index Area.

Mouth upriver about 7 km. Creek gets shallow and swifter, valley on left goes through to South Fork. Coho salmon go further up.

---

-continued-

**HUMPY CREEK**

Stream Code: 101-71-04H Anadromous Stream Number: 101-71-10040-20005  
Peak Spawning: September 1  
Survey Dates: August 28–Sept 3  
Survey Duration: helicopter 20 min.  
Survey Index Area

Mouth upriver to forks, up each fork 100m. Lots of pinks in creek, so best survey is as late as possible.

**BLOSSOM RIVER**

Stream Code: 101-55-040 Anadromous Stream Number:101-55-10400  
Peak Spawning: August 28  
Survey Dates: August 21–28  
Survey Duration: helicopter 1 hr.

Spawning very protracted, many schooling fish will be observed. Spawning occurs from lower river to very far upriver. Many pinks, chums and coho. Fuel cache at gear shed on road to mine.

Survey Index Area.: Mouth to barrier.

**KETA RIVER**

Stream Code: 101-30-030 Anadromous Stream Number:101-30-10300  
Peak Spawning: August 21  
Survey Dates: August 18–23  
Survey Duration: helicopter 1 hr.

Spawning very protracted, many schooling fish will be observed. Spawning occurs from lower river to very far upriver. Several possible barriers that Chinook make it past. Many pinks, chums and coho. Fuel cache at gear shed on road to mine.

Survey Index Area.: Mouth to barrier.

**KING SALMON RIVER**

Stream Code: 111-17-010 Anadromous Stream Number: 111-17-10100  
Peak Spawning: July 28  
Survey Dates: July 23–August 1  
Survey Duration: 1hr

Early system to survey, many chums in river at the same time. Most Chinook below large tributary on river right.

Survey Index Area: Mouth to barrier falls

---

Appendix A4.—Estimated total escapements of large Chinook salmon to escapement indicator systems and to Southeast Alaska and transboundary rivers, 1975–2008. Numbers may be revised annually as data are collected. Index escapements are expanded by average expansion factors, numbers in bold type are weir counts or mark–recapture estimates and are not expanded (region total expanded for 84% without Chilkat River, 90% with Chilkat escapement included).

Year	MAJOR SYSTEMS				MEDIUM SYSTEMS							King Salmon	Total all systems	Expanded region total	
	Alsek	Taku	Stikine	Major subtotal	Situk	Chilkat	Andrew	Unuk	Chickamin	Blossom	Keta				Medium subtotal
1975		12,917	7,571				508		1,758	439	611		64		
1976	5,765	24,575	5,723	36,063	<b>1,421</b>		<b>404</b>		745	205	253		99		
1977	10,496	29,489	11,445	51,430	<b>1,732</b>		<b>456</b>	4,739	1,722	337	692	9,679	204	61,313	72,992
1978	11,754	17,118	6,835	35,707	<b>808</b>		<b>388</b>	5,382	1,465	430	1,180	9,653	87	45,447	54,103
1979	18,670	21,611	12,610	52,891	<b>1,284</b>		<b>327</b>	2,803	1,133	162	1,283	6,992	134	60,016	71,448
77–79	13,640	22,740	10,297	46,676	1,275		390	4,308	1,440	310	1,052	8,775	141	55,592	66,181
Avg.															
1980	8,077	39,229	30,573	77,879	<b>905</b>		<b>282</b>	4,944	2,112	268	578	9,089	106	87,074	103,659
1981	8,327	49,546	36,057	93,929	<b>702</b>		<b>536</b>	3,557	1,824	478	990	8,088	153	102,170	121,631
1982	9,174	23,842	40,488	73,504	<b>434</b>		<b>672</b>	6,574	2,712	1,038	2,270	13,700	393	87,597	104,282
1983	11,028	9,792	6,424	27,243	<b>592</b>		<b>366</b>	5,474	2,847	1,772	2,475	13,526	<b>245</b>	41,014	48,826
1984	7,494	20,774	13,995	42,263	<b>1,726</b>		<b>389</b>	8,939	5,235	1,528	1,836	19,653	<b>265</b>	62,181	74,025
1985	5,758	35,906	16,691	58,336	<b>1,521</b>		625	5,761	4,541	2,133	1,879	16,460	<b>175</b>	74,970	89,251
1986	9,981	38,100	15,496	63,559	<b>2,067</b>		1,383	10,345	8,289	3,844	2,077	28,006	<b>255</b>	91,820	109,310
1987	11,395	28,928	25,637	65,929	<b>1,379</b>		1,540	9,601	4,631	4,058	2,312	23,520	<b>196</b>	89,645	106,721
1988	8,227	44,512	39,085	91,778	<b>868</b>		1,102	8,496	3,734	1,155	1,731	17,086	<b>208</b>	109,072	129,848
1989	9,105	<b>40,329</b>	25,272	74,676	<b>637</b>		1,036	5,591	4,437	1,035	3,477	16,212	<b>240</b>	91,129	108,486
Avg.	8,856	33,096	24,958	66,910	1,083		793	6,928	4,036	1,731	1,963	16,534	224	83,667	99,604
1990	8,794	<b>52,142</b>	23,541	84,449	<b>628</b>		1,298	2,876	2,679	773	1,824	10,078	<b>179</b>	94,706	112,745
1991	12,722	51,645	24,152	88,491	<b>889</b>	<b>5,897</b>	782	3,187	2,313	719	819	14,606	<b>134</b>	103,231	114,701
1992	5,519	55,889	35,521	96,887	<b>1,595</b>	<b>5,284</b>	1,520	4,253	1,644	451	653	15,400	<b>99</b>	112,386	124,874
1993	12,688	66,125	61,367	140,108	<b>952</b>	<b>4,472</b>	2,071	5,197	1,848	911	1,090	16,541	266	156,915	174,350
1994	12,312	48,368	34,234	95,083	<b>1,271</b>	<b>6,795</b>	1,118	<b>4,623</b>	1,843	484	921	17,055	213	112,351	124,834
1995	25,322	<b>33,805</b>	16,466	76,575	<b>4,330</b>	<b>3,790</b>	670	3,757	<b>2,309</b>	653	527	16,035	147	92,758	103,064
1996	14,443	<b>79,019</b>	<b>23,886</b>	122,411	<b>1,800</b>	<b>4,920</b>	655	5,679	<b>1,587</b>	662	894	16,196	292	138,899	154,332
1997	12,697	<b>114,938</b>	<b>28,185</b>	154,631	<b>1,878</b>	<b>8,100</b>	<b>478</b>	<b>2,970</b>	1,292	397	741	15,856	361	170,848	189,831
1998	<b>4,969</b>	31,039	<b>25,968</b>	61,976	<b>924</b>	<b>3,675</b>	952	<b>4,132</b>	1,857	<b>364</b>	<b>446</b>	12,350	134	74,460	82,733
1999	<b>13,617</b>	<b>19,734</b>	<b>19,947</b>	53,298	<b>1,461</b>	<b>2,271</b>	1,182	<b>3,914</b>	2,380	638	<b>968</b>	12,771	304	66,373	73,747
Avg.	12,308	55,270	29,812	97,391	1,573	5,023	1,073	4,059	1,971	605	888	14,689	213	112,293	125,521

-continued-

Year	MAJOR SYSTEMS				MEDIUM SYSTEMS								King Salmon	Total all systems	Expanded region total
	Alsek	Taku	Stikine	Major subtotal	Situk	Chilkat	Andrew	Unuk	Chickamin	Blossom	Keta	Medium subtotal			
2000	<b>6,835</b>	<b>30,529</b>	<b>27,531</b>	64,895	<b>1,785</b>	<b>2,035</b>	1,348	<b>5,872</b>	3,805	695	<b>914</b>	16,454	138	81,487	90,541
2001	<b>6,111</b>	<b>42,980</b>	<b>63,523</b>	112,614	<b>656</b>	<b>4,517</b>	2,060	<b>10,541</b>	<b>5,177</b>	614	1,033	24,597	149	137,360	152,622
2002	<b>5,396</b>	<b>52,409</b>	<b>50,875</b>	108,680	<b>1,000</b>	<b>4,050</b>	1,712	<b>6,988</b>	<b>5,007</b>	674	1,237	20,668	155	129,503	143,892
2003	<b>4,782</b>	<b>36,435</b>	<b>46,824</b>	88,041	<b>2,117</b>	<b>5,657</b>	1,163	<b>5,546</b>	<b>4,579</b>	611	969	20,642	118	108,801	120,890
2004	<b>6,995</b>	<b>68,199</b>	<b>48,900</b>	124,094	<b>755</b>	<b>3,422</b>	2,998	<b>3,963</b>	<b>4,268</b>	<b>734</b>	1,132	17,272	135	141,501	157,223
2005	4,462	<b>38,806</b>	<b>39,833</b>	83,101	<b>613</b>	<b>3,366</b>	1,979	<b>4,742</b>	<b>4,257</b>	<b>926</b>	1,496	17,379	143	101,291	112,546
2006	1,881	<b>41,831</b>	<b>24,405</b>	68,119	<b>749</b>	<b>3,039</b>	2,124	<b>5,645</b>	6,318	<b>1,270</b>	2,248	21,393	150	89,657	99,618
2007	2,619	<b>17,516</b>	<b>15,953</b>	36,087	<b>677</b>	<b>1,452</b>	1,736	<b>5,718</b>	4,242	406	936	15,167	181	51,435	57,150
2008	1,339	<b>27,383</b>	<b>18,843</b>	47,563	<b>453</b>	<b>2,833</b>	981	<b>3,104</b>	5,277	774	1,093	14,515	120	59,498	66,109
00–08	4,491	39,565	37,410	81,466	978	3,375	1,789	5,791	4,707	745	1,229	18,676	143	100,285	111,428
Avg.															
CHANGE FROM 2007 to 2008															
Number	(1,280)	9,867	2,890	11,476	(224)	1,381	(755)	(2,614)	1,035	368	157	(652)	(61)	10,763	11,959
Percent	-49%	56%	18%	32%	-33%	95%	-43%	-46%	24%	91%	17%	-4%	-34%	21%	21%
Escapement goals:															
Lower	5,500	30,000	14,000	49,500	450	1,750	650	3,250	2,325	750	750	9,926	120	59,546	66,162
Point	8,500	36,000	17,500	62,000	730	2,200	800	4,000	2,700	1,125	1,125	13,420	150	75,570	83,967
Upper	11,500	55,000	28,000	94,500	1,050	3,500	1,500	7,000	4,650	1,500	1,500	20,703	240	115,443	128,270
Average percent of goal:															
77–79	160%	63%	59%	75%	175%		52%	108%	41%	27%	93%	65%	94%	73%	
80–89	104%	92%	143%	108%	148%		106%	173%	116%	153%	174%	125%	149%	111%	
90–99	145%	154%	170%	157%	215%	228%	143%	101%	56%	53%	79%	108%	142%	148%	
00–06	61%	123%	247%	150%	150%	169%	255%	155%	137%	70%	115%	144%	94%	149%	

Appendix A5.—Variance of estimated total escapements of large Chinook salmon to escapement indicator systems and to Southeast Alaska and transboundary rivers, 1975–2008. Region total expanded for 84% without Chilkat River, 90% with Chilkat escapement included.

Year	Alsek	Taku	Stikine	Chilkat	Andrew	Unuk	Chickamin	Blossom	Keta	KSR	Region expansion	SE
1975	16,799,847	19,549,839	2,634,129		13,555	0	66,965	22,664	12,921	129	31,604,595	5,622
1976	8,757,302	70,766,455	870,489		0	0	12,009	4,917	2,212	308	109,444,233	10,462
1977	10,725,774	101,896,508	3,478,225		0	341,523	64,264	13,338	16,587	1,309	176,119,466	13,271
1978	10,005,510	34,336,787	2,146,225		0	440,365	46,484	21,743	48,181	237	85,297,739	9,236
1979	20,941,987	54,725,669	7,311,616		0	119,439	27,810	3,100	56,901	565	170,977,856	13,076
1980	3,940,947	180,319,778	24,820,324		0	371,612	96,655	8,422	11,559	357	306,913,695	17,519
1981	20,829,547	287,636,173	59,768,361		0	192,370	72,075	26,880	33,939	744	509,277,480	22,567
1982	19,613,392	66,606,818	43,533,604		0	657,072	159,365	126,555	178,256	4,890	177,674,429	13,329
1983	82,967,656	11,234,161	1,896,129		0	455,625	175,597	368,868	211,858	0	49,202,184	7,014
1984	26,991,714	50,567,743	9,000,000		0	1,214,845	593,586	274,390	116,670	0	100,875,003	10,044
1985	20,859,316	151,066,223	17,672,775		20,533	504,668	446,721	534,481	122,087	0	249,323,105	15,790
1986	2,875,936	170,095,329	15,232,238		100,512	1,627,155	1,488,468	1,736,608	149,279	0	293,537,339	17,133
1987	14,199,257	98,052,377	41,693,495		124,510	1,401,382	464,654	1,934,924	184,937	0	234,706,556	15,320
1988	3,591,482	232,160,074	96,908,274		63,784	1,097,466	301,971	156,784	103,666	0	484,879,665	22,020
1989	6,341,021	29,069,351	40,516,408		56,325	475,272	426,397	125,822	418,278	0	120,430,456	10,974
1990	11,260,887	52,507,414	35,155,413		88,408	125,741	155,481	70,227	115,145	0	143,385,911	11,974
1991	4,512,840	312,356,844	37,004,106	1,010,025	32,083	154,449	115,925	60,735	23,197	0	535,534,764	23,142
1992	13,844,531	365,807,701	80,038,968	900,601	121,370	274,995	58,516	23,923	14,765	0	558,448,926	23,632
1993	2,576,025	512,077,977	238,892,573	724,201	225,302	410,625	73,964	97,617	41,088	2,232	962,365,378	31,022
1994	457,652	273,975,987	75,257,360	1,117,249	65,606	1,602,756	73,584	27,561	29,359	1,429	466,106,882	21,590
1995	887,364	22,873,263	19,356,920	648,025	23,591	214,554	522,729	50,068	9,602	686	186,576,631	13,659
1996	231,842	124,224,399	7,689,529	564,001	22,503	490,280	39,601	51,462	27,658	2,687	207,483,151	14,404
1997	16,799,847	319,980,544	8,862,529	1,423,249	0	73,441	36,162	18,526	18,975	4,129	441,267,256	21,006
1998	8,757,302	112,886,800	15,452,761	319,225	47,557	155,236	74,726	5,929	2,500	565	163,789,012	12,798
1999	10,725,774	15,657,849	10,497,600	166,464	73,395	230,400	118,318	47,787	13,456	2,916	46,330,150	6,807
2000	10,005,510	29,343,889	10,036,224	111,556	95,467	414,736	313,607	56,737	14,884	604	55,602,472	7,457
2001	20,941,987	41,951,529	34,257,609	521,284	222,758	1,394,761	944,784	44,249	36,888	700	108,129,632	10,399
2002	3,940,947	120,077,764	34,951,744	184,041	153,873	583,696	544,644	53,350	52,965	758	211,337,756	14,537
2003	20,829,547	41,075,281	36,942,084	476,100	70,988	187,489	350,464	43,816	32,510	444	104,966,001	10,245
2004	19,613,392	84,437,721	15,178,816	207,936	471,850	105,625	797,449	5,073	44,328	577	147,130,196	12,130
2005	82,967,656	20,502,784	6,441,444	608,400	206,579	156,816	349,281	9,801	77,449	644	39,121,814	6,255
2006	26,991,714	30,713,764	48,135,844	206,116	237,798	318,096	864,616	29,584	174,962	714	100,338,357	10,017
2007	20,859,316	19,210,689	7,717,284	51,984	158,830	186,624	389,783	102,840	5,714	1,032	26,953,201	5,192
2008	2,875,936	73,517,935	9,941,409	280,900	50,733	0	603,321	70,227	41,316	455	104,614,984	10,228

Appendix A6.—Detailed 2008 Southeast Alaska Chinook salmon escapement surveys as entered into Commercial Fisheries Division Integrated Fisheries Database (IFDB/ALEX).

Stream no.	Stream	Date	Mouth	Live	Dead	Total	Survey	Obs. <sup>a</sup>	Use <sup>b</sup>	Comment <sup>c</sup>
101-30-030	Keta River	8/30/08	0	300	0	300	H	TAJ	1	
101-30-030	Keta River	9/3/08	0	363	0	363	H	KAP	3	lots cohos
101-30-030	Keta River	9/3/08	0	450	0	450	H	TAJ	3	backseat
101-55-040	Blossom River	8/30/08	0	102	0	102	H	TAJ	1	
101-55-040	Blossom River	9/3/08	0	242	0	242	H	TAJ	3	backseat
101-55-040	Blossom River	9/3/08	0	257	0	257	H	KAP	3	lots cohos
101-71-04A	Barrier Creek	8/5/08	0	3	0	3	H	KAP	2	
101-71-04A	Barrier Creek	8/6/08	0	1	0	1	H	KAP	2	
101-71-04B	Butler Creek	8/5/08	0	68	0	68	H	KAP	2	
101-71-04C	Clear Creek	8/5/08	0	60	0	60	H	KAP	3	
101-71-04H	Humpy Creek	8/30/08	0	20	0	20	H	TAJ	1	
101-71-04H	Humpy Creek	9/5/08	0	190	0	190	H	TAJ	2	
101-71-04I	Indian Creek	8/5/08	0	14	0	14	H	KAP	1	vis poor
101-71-04I	Indian Creek	8/9/08	0	76	0	76	F	TAJ	2	
101-71-04K	King Creek	8/30/08	0	187	0	187	H	TAJ	1	
101-71-04K	King Creek	9/5/08	0	622	0	622	H	TAJ	2	
101-71-04L	Leduc River	8/5/08	0	5	0	5	H	KAP	2	
101-71-04S	South Fork Chickamin	8/5/08	0	65	0	65	H	KAP	2	
101-71-04S	South Fork Chickamin	8/6/08	0	43	0	43	H	KAP	2	
101-71-04S	South Fork Chickamin	8/6/08	0	87	0	87	H	TAJ	2	backseat
101-75-015	Eulachon River	8/12/08	0	7	0	7	H	KAP	1	poor vis
101-75-015	Eulachon River	8/29/08	0	7	0	7	F	DWD	2	
101-75-30C	Clear Creek-Unuk R	8/5/08	0	10	0	10	H	KAP	2	
101-75-30C	Clear Creek-Unuk R	8/8/08	0	42	0	42	F	DWD	2	poor vis
101-75-30C	Clear Creek-Unuk R	8/12/08	0	30	0	30	H	KAP	2	poor vis
101-75-30G	Genes Lake Cr. Unuk	8/12/08	100	0	0	100	H	KAP	1	in lake
101-75-30K	Kerr Creek-Unuk R	8/5/08	0	27	0	27	H	KAP	2	
101-75-30K	Kerr Creek-Unuk R	8/12/08	0	0	0	0	H	KAP	1	poor vis
101-75-30K	Kerr Creek-Unuk R	8/30/08	0	29	0	29	F	DWD	2	
101-75-30L	Lake Creek-Unuk R	8/5/08	0	64	0	64	H	KAP	2	43 at riffles
101-75-30L	Lake Creek-Unuk R	8/12/08	0	20	0	20	H	KAP	1	only 7 at riffles
101-75-30L	Lake Creek-Unuk R	8/22/08	0	44	0	44	F	ADB	2	
101-80-070	Hatchery Ck-Yes Bay	9/11/08	0	0	1	1	F	SCH	2	
101-80-070	Hatchery Ck-Yes Bay	9/17/08	0	0	1	1	F	SCH	2	

-continued-

Stream no.	Stream	Date	Mouth	Live	Dead	Total	Survey	Obs. <sup>a</sup>	Use <sup>b</sup>	Comment <sup>c</sup>
106-44-031	Crystal Creek	6/17/08	0	0	0	100	A	WRB	2	50 abv rapids & 50 blw
106-44-031	Crystal Creek	7/2/08	0	0	1	91	A	WRB	2	all in pool abv rapids
106-44-031	Crystal Creek	7/6/08	0	20	0	220	A	TST	2	no fish blw rapids
106-44-031	Crystal Creek	7/9/08	20	0	0	640	A	WRB	2	240 abv rapids, 120@ cr, 250 float rks
106-44-031	Crystal Creek	7/12/08	30	0	0	640	A	WRB	2	40 blw rapid, 330 abv, 60@ cr, 180 fltng rks
106-44-031	Crystal Creek	7/18/08	0	0	0	380	A	TST	2	poor light, clear water
106-44-031	Crystal Creek	8/1/08	0	60	0	130	A	WRB	2	none @ rapids or floating rocks
106-44-031	Crystal Creek	8/16/08	0	100	0	250	A	WRB	2	50 abv rpds 20 fltng rks 80 blw str 300 pens
107-40-024	Aaron Creek	8/16/08	0	11	0	11	A	WRB	2	mostly glacial not good peak
107-40-049	Harding River	8/16/08	0	12	0	12	A	WRB	2	
107-40-053	Bradfield River E Fk	7/24/08	0	0	0	0	A	TST	2	
107-40-053	Bradfield River E Fk	8/16/08	0	58	0	58	A	WRB	2	mostly glacial
108-40-017	Goat Ck Stikine R	8/13/08	0	16	0	16	F	SNF	1	partial survey
108-40-020	Andrews Creek	7/31/08	30	320	0	350	A	WRB	2	inc 20 n fk
108-40-020	Andrews Creek	8/7/08	0	501	2	503	H	KAP	3	36 in n. fork
108-40-020	Andrews Creek	8/12/08	0	240	0	240	H	PJR	2	replicate survey
108-40-020	Andrews Creek	8/12/08	0	402	0	402	H	KAP	2	60 in n. fork
108-40-020	Andrews Creek	8/12/08	0	486	8	494	F	SNF	2	heavy rain high water: + 10 jacks
108-40-020	Andrews Creek	9/6/08	0	2	0	2	A	WRB	2	
108-41-010	North Arm Creek	7/31/08	0	14	0	14	A	WRB	2	
108-80-120	Little Talhtan River	7/29/08	0	600	0	600	H	KAP	2	lots below weir
108-80-120	Little Talhtan River	8/4/08	0	837	0	837	H	KAP	3	lots in mainstem
110-14-007	Farragut River	8/7/08	0	26	0	26	H	KAP	1	top only
110-32-009	Chuck R	7/16/08	0	2	0	2	A	TST	2	
110-32-009	Chuck R	7/30/08	0	8	0	8	A	WRB	2	
110-32-009	Chuck R	8/7/08	0	0	0	7	A	WRB	2	partly foggy
110-32-009	Chuck R	8/8/08	0	8	0	8	A	TST	2	
111-17-010	King Salmon River	7/22/08	0	12	0	12	H	KAP	1	early?
111-17-010	King Salmon River	7/25/08	0	15	0	15	H	KAP	1	
111-17-010	King Salmon River	7/25/08	0	31	0	31	F	KAP	1	vis poor
111-17-010	King Salmon River	8/5/08	0	79	0	79	F	PJR	3	
111-32-220	Nakina River	7/28/08	0	150	0	150	H	KAP	2	IA3
111-32-220	Nakina River	7/28/08	0	530	0	530	H	KAP	2	IA1
111-32-220	Nakina River	7/28/08	0	80	0	80	H	KAP	2	IA2
111-32-220	Nakina River	8/4/08	0	1437	0	1437	H	KAP	3	peak total

-continued-

Appendix A6.–Page 3 of 3.

Stream no.	Stream	Date	Mouth	Live	Dead	Total	Survey	Obs. <sup>a</sup>	Use <sup>b</sup>	Comment <sup>c</sup>
111-32-220	Nakina River	8/4/08	0	26	0	26	H	KAP	2	IA4
111-32-220	Nakina River	8/4/08	0	418	0	418	H	KAP	2	IA3
111-32-220	Nakina River	8/4/08	0	200	0	200	H	KAP	2	IA2
111-32-220	Nakina River	8/4/08	0	793	0	793	H	KAP	2	IA1
111-32-240	Kowatua Creek	8/21/08	0	417	0	417	H	PJR	2	back seat
111-32-240	Kowatua Creek	8/21/08	0	632	0	632	H	KAP	3	
111-32-240	Kowatua Creek	8/29/08	0	690	0	690	H	DKH	2	back seat
111-32-240	Kowatua Creek	8/29/08	0	620	0	620	H	PJR	3	
111-32-255	Tatsamenie River	8/21/08	0	573	0	573	H	KAP	3	IA1
111-32-255	Tatsamenie River	8/21/08	0	880	0	880	H	PJR	2	total
111-32-255	Tatsamenie River	8/21/08	0	1083	0	1083	H	KAP	3	peak total
111-32-255	Tatsamenie River	8/21/08	0	450	0	450	H	PJR	2	IA2
111-32-255	Tatsamenie River	8/21/08	0	430	0	430	H	PJR	2	IA1
111-32-255	Tatsamenie River	8/21/08	0	510	0	510	H	KAP	3	IA2
111-32-255	Tatsamenie River	8/29/08	0	710	0	710	H	PJR	2	
111-32-255	Tatsamenie River	8/29/08	0	630	0	630	H	DKH	2	
111-32-270	Nahlin River	7/21/08	0	86	0	86	H	KAP	2	IA2
111-32-270	Nahlin River	7/21/08	0	1066	0	1066	H	KAP	2	IA3
111-32-270	Nahlin River	7/28/08	0	1121	0	1121	H	KAP	2	peak total
111-32-270	Nahlin River	7/28/08	0	841	0	841	H	KAP	2	IA3
111-32-270	Nahlin River	7/28/08	0	51	0	51	H	KAP	2	IA2
111-32-270	Nahlin River	7/28/08	0	229	0	229	H	KAP	2	IA1
111-32-275	Tseta Creek	8/4/08	0	497	0	497	H	KAP	2	
111-32-280	Dudidontu River	7/28/08	0	384	0	384	H	KAP	2	201 below matatsu
111-32-280	Dudidontu River	8/7/08	0	480	0	480	H	KAP	3	202 below matatsu
111-50-052	Montana Creek	8/28/08	0	7	0	7	F	SBR	2	21,33
111-50-069	Fish Creek-Douglas I	8/7/08	0	55	2	60	F	SPH	2	21,32,42
182-30-043	Takhanni River (CAN)	7/31/08	0	41	0	41	H	KAP	2	
182-30-045	Goat Creek	7/31/08	0	11	0	11	H	KAP	2	
182-30-050	Blanchard Ck (CAN)	7/31/08	0	65	0	65	H	KAP	2	15 above bridge

<sup>a</sup> Observer initials on file in Commercial Fisheries IFDB/ALEX database.

<sup>b</sup> IFDB Standard Usage Codes: 1= not useful for indexing or estimating escapement; 2= potentially useful for indexing or estimating escapement; 3= potentially useful as the “peak” survey count for this species.

<sup>c</sup> Includes all surveys where Chinook salmon were observed; many are not used to estimate escapement.

Appendix A7.—Estimated abundance and composition by age and sex of the escapement of Chinook salmon to select systems in Southeast Alaska and transboundary rivers, 2007. Note: includes medium and in some cases, small fish, so total will vary from escapement estimates of large fish.

PANEL A. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE KETA RIVER IN 2008															
		BROOD YEAR AND AGE CLASS													Total
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n					13		3	25		2	10			53
	%					12.9		3.0	24.8		2.0	9.9			52.5
	SE of %					3.3		1.7	4.3		1.4	3.0			5.0
	Escapement					141		32	270		22	108			573
	SE of esc.					44		19	68		15	38			119
Females	n							1	31		1	14		1	48
	%							1.0	30.7		1.0	13.9		1.0	47.5
	SE of %							1.0	4.6		1.0	3.4		1.0	5.0
	Escapement							11	335		11	151		11	519
	SE of esc.							11	80		11	46		11	110
Combined	n					13		4	56		3	24		1	101
	%					12.9		4.0	55.4		3.0	23.8		1.0	100.0
	SE of %					3.3		1.9	4.9		1.7	4.2		1.0	0.0
	Escapement					141		43	606		32	260		11	1,093
	SE of esc.					44		22	125		19	66		11	203
PANEL B. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE BLOSSOM RIVER IN 2008															
Males	n					5		3	17		1	7			33
	%					7.4		4.4	25.0		1.5	10.3			48.5
	SE of %					3.2		2.5	5.3		1.5	3.7			6.1
	Escapement					73		44	249		15	102			483
	SE of esc.					33		25	65		15	40			98
Females	n					1		8	19		1	5		1	35
	%					1.5		11.8	27.9		1.5	7.4		1.5	51.5
	SE of %					1.5		3.9	5.4		1.5	3.2		1.5	6.1
	Escapement					15		117	278		15	73		15	512
	SE of esc.					15		43	70		15	33		15	101
Combined	n					6		11	36		2	12		1	68
	%					8.8		16.2	52.9		2.9	17.6		1.5	100.0
	SE of %					3.4		4.5	6.1		2.0	4.6		1.5	0.0
	Escapement					88		161	527		29	176		15	995
	SE of esc.					37		51	103		21	53		15	159

-continued-

PANEL C. AGE COMPOSITION OF SMALL, MEDIUM AND LARGE CHINOOK SALMON IN THE CHICKAMIN RIVER IN 2008																
BROOD YEAR AND AGE CLASS																
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	Total
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
Males	n	1	27			95			133			31			1	288
	%	0.2	6.0			21.0			29.2			6.8			0.2	63.5
	SE of %	0.2	1.5			2.8			2.4			1.2			0.2	2.7
	Escapement	15	403			1,412			1,961			457			15	4,262
	SE of esc.	15	99			215			313			103			15	514
Females	n					2			111			53				166
	%					0.4			24.0			11.6				36.0
	SE of %					0.3			2.3			1.6				2.7
	Escapement					29			1,607			781				2,418
	SE of esc.					21			270			151				382
Combined	n	1	27			97			244			84			1	454
	%	0.2	6.0			21.5			53.6			18.5			0.2	100.0
	SE of %	0.2	1.5			2.7			3.2			2.0			0.2	
	Escapement	15	403			1,441			3,598			1,238				6,710
	SE of esc.	15	99			217			536			217			15	822
PANEL D. AGE COMPOSITION OF SMALL, MEDIUM AND LARGE CHINOOK SALMON IN THE UNUK RIVER IN 2008 <sup>a</sup>																
Males	n		16			98			72			54				240
	%		4.1			25.1			18.5			13.9				61.8
	SE of %		1.2			3.1			2.1			1.8				2.9
	Escapement		163			998			735			551				2,458
	SE of esc.		46			143			118			97				266
Females	n								49			99			1	149
	%								12.6			25.4			0.3	38.2
	SE of %								1.7			2.4			0.3	2.9
	Escapement								500			1,011			10	1,521
	SE of esc.								90			152			10	211
Combined	n		16			98			121			153			1	389
	%		4.1			25.1			31.0			39.3			0.5	100.0
	SE of %		1.2			3.1			2.6			2.9			0.4	
	Escapement		163			998			1,235			1,562			20	3,979
	SE of esc.		46			143			177			215			15	419

<sup>a</sup> Weller et al. *In prep.*

-continued-

PANEL E. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE STIKINE RIVER IN 2008 <sup>b</sup>																
BROOD YEAR AND AGE CLASS																
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	Total
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
Males	n		3			11			48	1		75				138
	%		9.0%			3.4%			14.7%	0.3%		22.9%				42.2%
	SE of %		1.0%			2.9%			2.2%	0.3%		2.6%				4.2%
	Escapement		175			640			2,794	58		4,366				8,034
	SE of esc.		169			560			631	58		845				1,638
Females	n					1			61			126			1	189
	%					0.3%			18.7%			38.5%			0.3%	57.8%
	SE of %					0.3%			2.4%			3.4%			0.3%	4.2%
	Escapement					58			3,551			7,335			58	11,003
	SE of esc.					58			714			1,317			58	1,894
Combined	n		3			12			109	1		201			1	327
	%		0.9%			3.7%			33.3%	0.3%		61.5%			0.3%	100.0%
	SE of %		1.0%			2.9%			2.6%	0.3%		4.3%			0.3%	0.0%
	Escapement		175			699			6,345	58		11,701			58	19,036
	SE of esc.		169			563			1,155	58		2,004			58	3,163

<sup>b</sup> Richards et al. *In prep b.*

PANEL F. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN ANDREW CREEK IN 2008																
Males	n								11			6				17
	%								25.9%			15.9%				41.8%
	SE of %								7.3%			6.0%				8.2%
	Escapement								282			173				455
	SE of esc.								93			75				126
Females	n								3			19				22
	%								7.9%			50.3%				58.2%
	SE of %								4.5%			8.3%				8.2%
	Escapement								87			548				635
	SE of esc.								51			151				166
Combined	n								14			25				39
	%								33.8							
	SE of %								7.9%			66.2%				100.0%
	Escapement								369			721				1,090
	SE of esc.								110			182				233

-continued-

PANEL G. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE KING SALMON RIVER IN 2008

		BROOD YEAR AND AGE CLASS													Total	
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002		2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5		1.5
Males	n								7			2				9
	%								41.2			11.8				52.9
	SE of %								11.9			7.8				12.1
	Escapement								49			14				64
	SE of esc.								17			10				18
Females	n								3			5				8
	%								17.6			29.4				47.1
	SE of %								9.2			11.1				12.1
	Escapement								21			35				56
	SE of esc.								12			14				17
Combined	n								10			7				17
	%								58.8			41.2				100.0
	SE of %								11.9			11.9				
	Escapement								71			49				120
	SE of esc.								19			17				21

PANEL H. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE TAKU RIVER IN 2008<sup>c</sup>

Males	n	20	1	334	2	355	2	45								759
	%	1.8%	0.1%	0.1%	0.2%	25.8%	0.1%	3.2%								60.2
	SE of %	0.4%	0.1%	0.1%	0.1%	1.4%	0.1%	0.5%								1.9
	Escapement	726	36	11,619	73	10,348	57	1,289								24,147
	SE of esc.	179	36	1,234	52	954	40	218								2,879
Females	n	1		22	2	441	5	82						1		554
	%	0.1%		1.8%	0.1%	31.5%	0.4%	5.8%						0.1%		39.4
	SE of %	0.1%		0.4%	0.1%	1.7%	0.2%	0.7%						0.1%		1.9
	Escapement	28		720	57	12,655	142	2,342						28		15,972
	SE of esc.	28		162	40	1,184	65	322						28		1,937
Combined	n	21	1	356	4	796	7	127						1		1,313
	%	1.9%	0.1%	0.1%	0.3%	57.3%	0.5%	9.0%						0.1%		100
	SE of %	0.4%	0.1%	0.1%	0.2%	2.4%	0.2%	0.9%						0.1%		
	Escapement	754	36	12,339	129	23,002	199	3,631						28		40,119
	SE of esc.	181	36	1,286	66	1,974	77	438						28		2,886

<sup>c</sup> Jones III et al. (2010).

-continued-

PANEL I. AGE COMPOSITION OF LARGE CHINOOK SALMON IN THE CHILKAT RIVER IN 2008<sup>d</sup>

		BROOD YEAR AND AGE CLASS														
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	Total
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
Males	n							45				8			1	54
	%							32%				6%			1%	38.8%
	SE of %							4.0%				2.0%			0.7%	
	Escapement							917				163			20	1,100
	SE of esc.							204				63			20	215
Females	n							58				27				85
	%							42%				19%				61.2%
	SE of %							4.2%				3.4%				
	Escapement							1,182				550				1,732
	SE of esc.							250				139				286
Combined	n							103				35			1	139
	%							74%				25%			1%	100.0%
	SE of %							3.7%				3.7%			0.7%	
	Escapement							2,099				713			20	2,833
	SE of esc.							406				168			20	530

<sup>d</sup> Chapell *In prep b.*

PANEL J. AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE ALSEK RIVER IN 2008

Males	n			39		1		1	29		2		30			102
	%			24.1%		0.6%		0.6%	18.3%		1.3%		19.0%			63.9%
	SE of %			8.7%		0.7%		0.7%	3.2%		0.9%		4.3%			
	Escapement			471		12		12	357		25		371			1,248
	SE of esc.			149		12		12	127		19		161			
Females	n			4					20		1		30	1	1	57
	%			2.5%					12.7%		0.6%		19.0%	0.6%	0.6%	36.1%
	SE of %			1.3%					2.8%		0.6%		4.3%	0.6%	0.6%	
	Escapement			49					248		12		371	12	12	705
	SE of esc.			26					111		12		161	12	12	
Combined	n			43		1		1	49		3		60	1	1	159
	%			26.6%		0.6%		0.6%	31.0%		1.9%		38.0%	0.6%	0.6%	100.0%
	SE of %			9.2%		0.7%		0.7%	4.5%		1.1%		7.0%	0.6%	0.6%	
	Escapement			520		12		12	604		37		743	12	12	1,953
	SE of esc.			159		12		12	224		25		309	12	12	102

-continued-

PANEL K. AGE COMPOSITION OF SMALL, MEDIUM AND LARGE CHINOOK SALMON IN THE SITUK RIVER IN 2008															
BROOD YEAR AND AGE CLASS															
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	Total
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
Males	n	1		18	3	2	9	1							34
	%	1.4%		26.0%	4.1	3.4%	15.2%	1.7%							51.8%
	SE of %	1.4%		4.7%	2.3	2.4%	4.6%	1.7%							
	Escapement	9		175	28	23	102	12							349
	SE of esc.														
Females	n			9		3	17								29
	%			13.7%		5.2%	29.3%								48.2%
	SE of %			4.3%		2.9%	5.4%								
	Escapement			93		35	197								325
	SE of esc.														
Combined	n	1		27	3	5	26	1							63
	%	1.4%		39.7%	4.1	8.6%	44.4%	1.7%							100.0%
	SE of %	1.4%		5.8%	2.3	3.6%	5.4%	1.7%							
	Escapement	9		268	28	58	300	12							674
	SE of esc.														34

65

SUMMARY. PERCENTAGE AGE COMPOSITION ESTIMATED FROM CHINOOK SALMON SAMPLED IN SOUTHEAST ALASKA RIVERS IN 2008 <sup>c</sup>															
BROOD YEAR AND AGE CLASS															
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
1. Keta	NE	NE			13		4	55		3	24				1
2. Blossom	NE	NE			9		16	53		3	18				1
3. Chickamin	NE	6			22			54			19				0
4. Unuk	NE	4			25			31			39				1
5. Stikine	NE	NE			4			33			61				0
6. Andrew Cr	NE	NE						34			66				0
7. King Salmon	NE	NE						59			41				0
8. Taku	NE	2			31			57			9				0
9. Chilkat	NE	29			12			44			15				0
10. Alsek	NE	NE			27	1	1	31	2		38	1			1
11. Situk	1	0		40	4	9	44	2							

<sup>c</sup> Small fish not included (NE) in experimental design, 2008.

-continued-

SUMMARY. ESTIMATED NUMBERS OF CHINOOK SALMON BY AGE CLASS IN ESCAPEMENTS TO SOUTHEAST ALASKA RIVERS IN 2008.

	BROOD YEAR AND AGE CLASS														Total
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
1. Keta	0	0	0	0	141	0	43	606	0	32	260	0	0	11	1,093
2. Blossom	0	0	0	0	88	0	161	527	0	29	176	0	0	15	995
3. Chickamin	15	403	0	0	1,441	0	0	3,598	0	0	1,238	0	0	15	6,710
4. Unuk	0	163	0	0	998	0	0	1,235	0	0	1,562	0	0	20	3,979
5. Stikine	0	175	0	0	699	0	0	6,345	58	0	11,701	0	0	58	19,036
6. Andrew Cr	0	0	0	0	0	0	0	369	0	0	721	0	0	0	1,090
7. King Salmon	0	0	0	0	0	0	0	71	0	0	49	0	0	0	120
8. Taku	0	754	36	0	12,339	129	0	23,002	199	0	3,631	0	0	28	40,119
9. Chilkat	0	0	0	0	0	0	0	2,099	0	0	713	0	0	20	2,832
10. Alsek	0	0	0	0	520	12	12	604	37	0	743	12	0	12	1,953
11. Situk	9	0	0	268	28	58	300	12	0	0	0	0	0	0	674

SUMMARY. PERCENTAGE SEX COMPOSITION OF MALES BY AGE CLASS ESTIMATED FROM CHINOOK SALMON SAMPLED IN SOUTHEAST ALASKA RIVERS IN 2008.

	BROOD YEAR AND AGE CLASS													
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
1. Keta					100		75	45		67	42			
2. Blossom					83		27	47		50	58			
3. Chickamin		100			98			55			37			
4. Unuk		100			100			60			35			
5. Stikine					92			44			37			
6. Andrew Cr								77			24			
7. King Salmon								70			29			
8. Taku		96			94	56		45	29		35			
9. Chilkat								44			23			
10. Alsek					91	100		59	67		50			
11. Situk	100			65			34							
Average		99			92	78	45	54	48	58	37			

Appendix A8.—Average length (MEF), by age, of Chinook salmon in selected systems in Southeast Alaska and transboundary rivers, 2008.

PANEL A. AVERAGE LENGTH OF CHINOOK SALMON IN THE KETA RIVER IN 2008															
BROOD YEAR AND AGE CLASS															
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n		4		2	28		3	25		2	10			
Average length			428		585	648		760	809		885	907			
	SD		37		42	52		17	67		21	79			
	SE		18		30	10		10	13		15	25			
Females	n							1	32		1	14			1
Average length								800	822		870	894			955
	SD								62			47			
	SE								11			13			
Combined	n		4		2	28		4	57		3	24			1
Average length			428		585	648		770	816		880	899			955
	SD		37		42	52		24	64		17	61			
	SE		18		30	10		12	9		10	13			
Panel B. AVERAGE LENGTH OF CHINOOK SALMON IN THE BLOSSOM RIVER IN 2008															
Males	n		1		2	16		3	17		1	7			
Average length			495		635	635		762	836		915	937			
	SD				7	43		62	84			94			
	SE				5	11		36	20			35			
Females	n					2		8	19		1	5			1
Average length						660		820	832		829	904			940
	SD					42		54	52			32			
	SE					30		19	12			14			
Combined	n		1		2	18		11	36		2	12			1
Average length			495		635	638		804	834		872	923			940
	SD				7	43		60	68		61	74			
	SE				5	10		18	11		43	21			
PANEL C. AVERAGE LENGTH OF CHINOOK SALMON IN THE CHICKAMIN RIVER IN 2008 <sup>a</sup>															
Males	n	1	27			96			133			31			1
Average length		275	445			635			788			899			1,000
	SD		40			57			70			62			
	SE		8			6			6			11			
Females	n					2			111			53			
Average length						678			807			870			
	SD					11			55			38			
	SE					8			5			5			
Combined	n	1	27			97			244			84			1
Average length		275	445			635			797			880			1,000
	SD		40			55			64			50			
	SE		8			6			4			5			

<sup>a</sup> Johnson et al. (2009).

-continued-

PANEL D. AVERAGE LENGTH OF CHINOOK SALMON IN THE UNUK RIVER IN 2008 <sup>b</sup>															
		BROOD YEAR AND AGE CLASS													
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n		16			98			72			54			1
Average length			393			623			780			896			955
	SD		34			56			56			62			
	SE		8			6			7			8			
Females	n								49			99			1
Average length									802			876			855
	SD								47			39			
	SE								7			4			0
Combined	n		16			98			121			153			2
Average length			393			623			789			883			905
	SD		34			56			54			49			71
	SE		8			6			5			4			50

<sup>b</sup> Weller et al. *In prep.*

PANEL E. AVERAGE LENGTH OF CHINOOK SALMON IN THE STIKINE RIVER IN 2008 <sup>c</sup>															
Males	n		3			11			48	1		75			
Average length			473			568			801	932		886			
	SD		71			51			101			63			
	SE		41			15			15			7			
Females	n					1			61			126			1
Average length						776			791			841			855
	SD								54			42			
	SE								7			4			
Combined	n		3			12			109	1		201			1
Average length			473			585			795	932		858			855
	SD		71			77			78			55			
	SE		41			22			7			4			

<sup>c</sup> Richards et al. *In prep b.*

PANEL F. AVERAGE LENGTH OF CHINOOK SALMON IN ANDREW CREEK IN 2008															
Males	n								11			6			
Average length									684			861			
	SD								79			46			
	SE								24			19			
Females	n								3			19			
Average length									742			823			
	SD								55			38			
	SE								32			9			
Combined	n								14			25			
Average length									696			832			
	SD								76			42			
	SE								20			8			

-continued-

PANEL G. AVERAGE LENGTH OF CHINOOK SALMON IN THE KING SALMON RIVER IN 2008															
BROOD YEAR AND AGE CLASS															
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n								10					4	
Average length									767					883	
	SD								51					86	
	SE								16					43	
Females	n								4					8	
Average length									800					886	
	SD								29					36	
	SE								15					13	
Combined	n								14					12	
Average length									776					885	
	SD								48					53	
	SE								13					15	

PANEL H. AVERAGE LENGTH OF CHINOOK SALMON IN THE TAKU RIVER IN 2008 <sup>d</sup>															
Males	n		92	3	335	2		355	2			46			
Average length			376	377	603	593		758	810			834			
	SD		65	26	73	53		76	7			125			
	SE		7		4			4				18			
Females	n		1		22	2		441	5			82			1
Average length			770		666	765		762	785			818			805
	SD				76	64		50	44			46			
	SE				16			2				5			
Combined	n		93	3	357	4		796	7			128			1
Average length			380	377	607	679		760	792			823			805
	SD		77	26	74	110		63	38			83			
	SE		8		4			2				7			

<sup>d</sup> Jones III et al. (2010).

PANEL I. AVERAGE LENGTH OF CHINOOK SALMON IN THE CHILKAT RIVER IN 2008 <sup>e</sup>															
Males	n		108		129			236				52			525
Average length			351		559			756				894			1,050
	SD		35		79			79				73			
	SE		3		7			5				10			
Females	n				9			183				98			290
Average length					673			785				859			845
	SD				95			48				55			
	SE				32			4				6			
Combined	n		108		138			419				150			815
Average length			351		567			768				871			948
	SD		35		85			65				68			145
	SE		3		7			3				6			103

<sup>e</sup> Chapell *In prep b.*

-continued-

PANEL J. AVERAGE LENGTH OF CHINOOK SALMON IN THE ALSEK RIVER IN 2008															
		Brood year and age class													
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n		2	1		39	1		29	2		30			
Average length			400	475		534	525		710	672		840			
	SD		5			43			101	23		71			
	SE		4			7			19	16		13			
Females	n					4			20	1		30	1		1
Average length						580			722	709		783	794		865
	SD					62			25			21			
	SE					31			6			4			
Combined	n		2	1		43	1		49	3		60	1		1
Average length			400	475		538	525		715	684		811	794		865
	SD		5			46			79	27		59			
	SE		4			7			11	16		8			

PANEL K. AVERAGE LENGTH OF CHINOOK SALMON IN THE SITUK RIVER IN 2008															
Males	n	1			18	3		9	1		2				
Average length		485			609	523		806	815		888				
	SD				45	38		78			18				
	SE				11	22		26			13				
Females	n				9			17			3				
Average length					644			814			858				
	SD				53			41			25				
	SE				18			10			14				
Combined	n	1			27	3		26	1		5				
Average length		485			621	523		811	815		870				
	SD				50	38		55			25				
	SE	1			18	3		9	1		2				

-continued-

SUMMARY. AVERAGE LENGTH OF MALE CHINOOK SALMON SAMPLED IN SOUTHEAST ALASKA IN 2008														
	BROOD YEAR AND AGE CLASS													
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
1. Keta		468		680	655			806			867			
2. Blossom				667	674			779			915			
3. Chickamin		432			640			806			899			
4. Unuk		407			648			772			876			
5. Stikine		348			562			769			878			841
6. Andrew Creek					537			739			829			799
7. King Salmon					613			746			906			
8. Taku		391			595			738			844			
9. Chilkat		356			576			767			886			
10. Alsek					534			740			819			
11. Situk	340			572			798			826				

SUMMARY. AVERAGE LENGTH OF FEMALE CHINOOK SALMON SAMPLED IN SOUTHEAST ALASKA IN 2008														
	BROOD YEAR AND AGE CLASS													
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
1. Keta							800	822		870	894			955
2. Blossom					660		820	832		829	904			940
3. Chickamin					678			807			870			
4. Unuk								802			876			855
5. Stikine					776			791			841			855
6. Andrew Creek								742			823			
7. King Salmon								800			886			
8. Taku		770			666	765		762	785		818			805
9. Chilkat					673			785			859			845
10. Alsek					580			722	709		783	794		865
11. Situk							814			859				

SUMMARY. AVERAGE LENGTH OF CHINOOK SALMON SAMPLED IN SOUTHEAST ALASKA IN 2008 SEXES COMBINED														
	BROOD YEAR AND AGE CLASS													
	2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
	0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
1. Keta		428		585	648		770	816		880	899			955
2. Blossom		495		635	638		804	834		872	923			940
3. Chickamin	275	445			635			797			880			1,000
4. Unuk		393			623			789			883			905
5. Stikine		473			585			795	932		858			855
6. Andrew Creek								696			832			
7. King Salmon								776			885			
8. Taku		380	377		607	679		760	792		823			805
9. Chilkat		351			567			768			871			948
10. Alsek		400	475		538	525		715	684		811	794		865
11. Situk	485			621			811			870				
Averages				426	614	605	602	795	775	803	874	867	794	909

Note: Age classes with fewer than 4 fish sampled were not reported in summary panels.

Appendix A9.–Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.2 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008. Bold numbers indicate probability of <0.01 that they are the same.

PANEL A. DIFFERENCES IN MEAN LENGTHS FOR AGE-1.2 FISH, SEXES COMBINED															
System	Age class	Average length	SE	Difference in mean length											
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk	
1. Keta	1.2	648	10	0	-10	-13	-25	-63				-40	-81	-110	648
2. Blossom	1.2	638	10	10	0	-3	-15	-53				-30	-71	-100	638
3. Chickamin	1.2	635	6	13	3	0	-12	-50				-28	-68	-97	635
4. Unuk	1.2	623	6	25	15	12	0	-38				-15	-56	-85	623
5. Stikine	1.2	585	22	63	53	50	38	0				22	-18	-47	585
6. Andrew Cr	1.2														
7. King Salmon	1.2														
8. Taku	1.2	607	4	40	30	28	15	-22				0	-40	-69	607
9. Chilkat	1.2	567	7	81	71	68	56	18				40	0	-29	567
10. Alsek	1.2	538	7	110	100	97	85	47				69	29	0	538
11. Situk	1.2														

PANEL B. TEST VALUES FOR DIFFERENCES IN MEAN LENGTHS FOR AGE-1.2 FISH, SEXES COMBINED															
System	Age class	Average length	SE	Test statistics for differences in mean length											
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk	
1. Keta	1.2	648	10	0.00	-0.71	-1.08	-2.19	<b>-2.59</b>				<b>-3.79</b>	<b>-6.57</b>	<b>-9.04</b>	
2. Blossom	1.2	638	10	0.71	0.00	-0.21	-1.29	-2.17				<b>-2.81</b>	<b>-5.69</b>	<b>-8.11</b>	
3. Chickamin	1.2	635	6	1.08	0.21	0.00	-1.51	-2.19				<b>-3.88</b>	<b>-7.23</b>	<b>-10.51</b>	
4. Unuk	1.2	623	6	2.19	1.29	1.51	0.00	-1.66				-2.25	<b>-6.06</b>	<b>-9.40</b>	
5. Stikine	1.2	585	22	<b>2.59</b>	2.17	2.19	1.66	0.00				0.99	-0.78	-2.04	
6. Andrew Cr	1.2														
7. King Salmon	1.2														
8. Taku	1.2	607	4	<b>3.79</b>	<b>2.81</b>	<b>3.88</b>	2.25	-0.99				0.00	<b>-4.87</b>	<b>-8.60</b>	
9. Chilkat	1.2	567	7	<b>6.57</b>	<b>5.69</b>	<b>7.23</b>	<b>6.06</b>	0.78				<b>4.87</b>	0.00	<b>-2.88</b>	
10. Alsek	1.2	538	7	<b>9.04</b>	<b>8.11</b>	<b>10.51</b>	<b>9.40</b>	2.04				<b>8.60</b>	<b>2.88</b>	0.00	
11. Situk	1.2														

Appendix A10.–Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.3 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008. Bold numbers indicate probability of <0.01 that they are the same.

PANEL A. DIFFERENCES IN MEAN LENGTHS FOR AGE-1.3 FISH, SEXES COMBINED														
System	Age class	Average length	SE	Difference in mean length										
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk
1. Keta	1.3	816	9	0	18	-19	-27	-21	-120	-40	-56	-48	-101	
2. Blossom	1.3	834	11	-18	0	-37	-45	-39	-138	-57	-74	-66	-119	
3. Chickamin	1.3	797	4	19	37	0	-8	-2	-101	-21	-37	-29	-82	
4. Unuk	1.3	789	5	27	45	8	0	6	-93	-13	-29	-21	-74	
5. Stikine	1.3	795	7	21	39	2	-6	0	-99	-19	-35	-27	-80	
6. Andrew Cr	1.3	696	7	120	138	101	93	99	0	80	64	72	19	
7. King Salmon	1.3	776	13	40	57	21	13	19	-80	0	-16	-8	-61	
8. Taku	1.3	760	2	56	74	37	29	35	-64	16	0	8	-45	
9. Chilkat	1.3	768	3	48	66	29	21	27	-72	8	-8	0	-53	
10. Alsek	1.3	715	11	101	119	82	74	80	-19	61	45	53	0	
11. Situk	1.3													

PANEL B. TEST VALUES FOR DIFFERENCES IN MEAN LENGTHS FOR AGE-1.3 FISH, SEXES COMBINED														
System	Age class	Average length	SE	Test statistics for differences in mean length										
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk
1. Keta	1.3	816	9	0.00	1.25	-2.04	<b>-2.76</b>	<b>-1.92</b>	<b>-10.90</b>	<b>-2.60</b>	<b>-6.37</b>	<b>-5.30</b>	<b>-7.16</b>	
2. Blossom	1.3	834	11	-1.25	0.00	<b>-3.08</b>	<b>-3.64</b>	<b>-2.92</b>	<b>-10.36</b>	<b>-3.38</b>	<b>-6.40</b>	<b>-5.61</b>	<b>-7.44</b>	
3. Chickamin	1.3	797	4	2.04	<b>3.08</b>	0.00	-1.26	-0.25	<b>-12.52</b>	-1.54	<b>-8.06</b>	<b>-5.68</b>	<b>-6.85</b>	
4. Unuk	1.3	789	5	<b>2.76</b>	<b>3.64</b>	1.26	0.00	0.70	<b>-10.88</b>	-0.92	<b>-5.37</b>	<b>-3.60</b>	<b>-6.02</b>	
5. Stikine	1.3	795	7	1.92	<b>2.92</b>	0.25	-0.70	0.00	<b>-9.99</b>	-1.28	<b>-4.75</b>	-3.51	<b>-6.02</b>	
6. Andrew Cr	1.3	696	7	<b>10.90</b>	<b>10.36</b>	<b>12.52</b>	<b>10.88</b>	<b>9.99</b>	0.00	<b>5.54</b>	<b>8.72</b>	<b>9.36</b>	1.43	
7. King Salmon	1.3	776	13	<b>2.60</b>	<b>3.38</b>	1.54	0.92	1.28	<b>-5.54</b>	0.00	-1.26	-0.64	<b>-3.62</b>	
8. Taku	1.3	760	2	<b>6.37</b>	<b>6.40</b>	<b>8.06</b>	<b>5.37</b>	<b>4.75</b>	<b>-8.72</b>	1.26	0.00	<b>2.03</b>	<b>-3.92</b>	
9. Chilkat	1.3	768	3	<b>5.30</b>	<b>5.61</b>	<b>5.68</b>	<b>3.60</b>	<b>3.51</b>	<b>-9.36</b>	0.64	-2.03	0.00	<b>-4.52</b>	
10. Alsek	1.3	715	11	<b>7.16</b>	<b>7.44</b>	<b>6.85</b>	<b>6.02</b>	<b>6.02</b>	-1.43	<b>3.62</b>	<b>3.92</b>	<b>4.52</b>	0.00	
11. Situk	1.3													

Appendix A11.–Differences in mean lengths (Panel A) and test results (Z, Panel B) for statistical differences in mean lengths between age-1.4 Chinook salmon (sexes combined) sampled in 11 rivers in Southeast Alaska in 2008. Bold numbers indicate probability of <0.01 that they are the same.

PANEL A. DIFFERENCES IN MEAN LENGTHS FOR AGE-1.4 FISH, SEXES COMBINED														
System	Age class	Average length	SE	Difference in mean length										
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk
1. Keta	1.4	899	13	0	24	-19	-16	-41	-67	-15	-76	-28	-88	
2. Blossom	1.4	923	21	-24	0	-43	-40	-65	-91	-39	-100	-52	-112	
3. Chickamin	1.4	880	5	19	43	0	3	-22	-48	5	-57	-9	-69	
4. Unuk	1.4	883	4	16	40	-3	0	-25	-50	2	-59	-12	-72	
5. Stikine	1.4	858	4	41	65	22	25	0	-26	27	-35	13	-47	
6. Andrew Cr	1.4	832	4	67	91	48	50	26	0	52	-9	39	-21	
7. King Salmon	1.4	885	15	15	39	-5	-2	-27	-52	0	-61	-14	-74	
8. Taku	1.4	823	7	76	100	57	59	35	9	61	0	48	-12	
9. Chilkat	1.4	871	6	28	52	9	12	-13	-39	14	-48	0	-60	
10. Alsek	1.4	811	8	88	112	69	72	47	21	74	12	60	0	
11. Situk	1.4													

PANEL B. TEST VALUES FOR DIFFERENCES IN MEAN LENGTHS FOR AGE-1.4 FISH, SEXES COMBINED														
System	Age class	Average length	SE	Test statistics for differences in mean length										
				Keta	Blossom	Chickamin	Unuk	Stikine	Andrew Cr	King Salmon	Taku	Chilkat	Alsek	Situk
1. Keta	1.4	899	13	0.00	0.97	-1.44	-1.25	-3.14	<b>-5.09</b>	-0.75	<b>-5.22</b>	-2.07	<b>-6.02</b>	
2. Blossom	1.4	923	21	-0.97	0.00	-1.98	-1.86	<b>-3.01</b>	<b>-4.19</b>	-1.47	<b>-4.42</b>	-2.37	<b>-4.96</b>	
3. Chickamin	1.4	880	5	1.44	1.98	0.00	0.45	<b>-3.44</b>	<b>-7.43</b>	0.28	<b>-6.35</b>	-1.20	<b>-7.57</b>	
4. Unuk	1.4	883	4	1.25	1.86	-0.45	0.00	<b>-4.42</b>	<b>-8.96</b>	0.11	<b>-7.10</b>	-1.74	<b>-8.37</b>	
5. Stikine	1.4	858	4	<b>3.14</b>	<b>3.01</b>	<b>3.44</b>	<b>4.42</b>	0.00	<b>-4.53</b>	1.67	<b>-4.12</b>	1.90	<b>-5.46</b>	
6. Andrew Cr	1.4	832	4	<b>5.09</b>	<b>4.19</b>	<b>7.43</b>	<b>8.96</b>	<b>4.53</b>	0.00	<b>3.28</b>	-1.06	<b>5.64</b>	<b>-2.49</b>	
7. King Sal.	1.4	885	15	0.75	1.47	-0.28	-0.11	-1.67	<b>-3.28</b>	0.00	<b>-3.58</b>	-0.83	<b>-4.29</b>	
8. Taku	1.4	823	7	<b>5.22</b>	<b>4.42</b>	<b>6.35</b>	<b>7.10</b>	<b>4.12</b>	1.06	<b>3.58</b>	0.00	<b>5.15</b>	-1.18	
9. Chilkat	1.4	871	6	2.07	2.37	1.20	1.74	-1.90	<b>-5.64</b>	0.83	<b>-5.15</b>	0.00	<b>-6.37</b>	
10. Alsek	1.4	811	8	<b>6.02</b>	<b>4.96</b>	<b>7.57</b>	<b>8.37</b>	<b>5.46</b>	<b>2.49</b>	<b>4.29</b>	1.18	<b>6.37</b>	0.00	
11. Situk	1.4													

Appendix A12.—Age composition and average length by age and sex of Chinook salmon sampled in the Farragut River, 2008.

AGE COMPOSITION OF MEDIUM AND LARGE CHINOOK SALMON IN THE FARRAGUT RIVER IN 2008																
		BROOD YEAR AND AGE CLASS														
		2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001	2000	2001	2000	Total
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5	
Males	n								3						6	9
	%								21.4						42.9	64.3
	SE of %								11.4						13.7	13.3
	Escapement															
	SE of esc.															
Females	n								1						4	5
	%								7.1						28.6	35.7
	SE of %								7.1						12.5	13.3
	Escapement															
	SE of esc.															
Combined	n								4						10	14
	%								28.6						71.4	100
	SE of %								12.5						12.5	
	Escapement															
	SE of esc.															

AVERAGE LENGTH OF CHINOOK SALMON IN THE FARRAGIT RIVER IN 2008															
		BROOD YEAR AND AGE CLASS													
		2006	2005	2004	2005	2004	2003	2004	2003	2002	2003	2002	2001	2002	2001
		0.1	1.1	2.1	0.2	1.2	2.2	0.3	1.3	2.3	0.4	1.4	2.4	0.5	1.5
Males	n								3						6
Average length									847						917
	SD								38						51
	SE								22						21
Females	n								1						4
Average length									820						876
	SD														50
	SE														25
Combined	n								4						10
Average length									840						901
	SD								34						52
	SE								17						16

Appendix A13.—Numbers of Chinook salmon examined for coded wire tags (CWT) and numbers of tags recovered in rivers in Southeast Alaska and transboundary rivers, 2007–2008. Hatchery CWTs expanded by tag ratio reported in ADF&G Mark, Tag, and Age Laboratory database.

	2008					2007				
	Chinook sampled	Hatchery CWTs	Expanded hatchery CWTs	Non-natal wild CWTs <sup>a</sup>	Natal wild CWTs <sup>b</sup>	Chinook sampled	Hatchery CWTs	Expanded hatchery CWTs <sup>c</sup>	Non-natal wild CWTs	Natal wild CWTs
Situk River	79	0	0	0	0	134	0	0	0	0
Alsek River	465	0	0	0	0	1,056	0	0	0	0
Chilkat River	817	0	0	0	31	648	0	0	0	26
Taku River	3,343	0	0	0	31	2,384	2 <sup>d</sup>	8	0	13
King Salmon R.	26	0	0		0	63	1 <sup>e</sup>	1	0	0
Stikine River	2,279	0	0	1	26	380	0	0	0	4
Andrew Creek	50	0	0	0	0	200	1	13	0	0
Unuk River	996	0	0	0	14	1,823	0	0	0	39
Chickamin R.	459	0	0	0	9	909	0	0	0	30
Blossom River	112	0	0	0	0	62	1	8	0	0
Keta River	157	0	0	0	0	97	0	0	0	0
Farragut River	15	1	1	0	0	30	4	4	0	0
Totals	9,098	1	1	1	111	7,786	9	34	0	108

<sup>a</sup> Non-natal wild CWTs are recoveries in a stream from Chinook smolt that were tagged in another river, i.e. Stikine River had 1 tag from the Taku River in 2008.

<sup>b</sup> Natal CWTs are recoveries of wild Chinook tagged as smolt in that river.

<sup>c</sup> Expanded hatchery numbers are from listed tag ratios in ADF&G Mark, Tag, and Age Laboratory database.

<sup>d</sup> One Little Port Walter and 1 Douglas Island Pink and Chum/Sheep Creek caught at Canyon Island.

<sup>e</sup> One Little Port Walter.

## **APPENDIX B**

The expansion factor provides a means of predicting escapement in years where only an index count of the escapement is available, i.e. no weir counts or mark–recapture experiments were conducted. The expansion factor is the average over several years of the ratio of the escapement estimate (or weir count) to the index count.

SYSTEMS WHERE ESCAPEMENT IS KNOWN

On systems where escapement can be completely enumerated with weirs or other complete counting methods, the expansion factor is an estimate of the expected value of the “population” of annual expansion factors ( $\pi$ ’s) for that system:

$$\bar{\pi} = \frac{\sum_{y=1}^k \pi_y}{k} \quad (1)$$

where  $\pi_y = N_y / C_y$  is the observed expansion factor in year  $y$ ,  $N_y$  is the known escapement in year  $y$ ,  $C_y$  is the index count in year  $y$ , and  $k$  is the number of years for which these data are available to calculate an annual expansion factor.

The estimated variance for expansion of index counts needs to reflect two sources of uncertainty for any predicted value of  $\pi$ , ( $\pi_p$ ). First is an estimate of the process error ( $var(\pi)$ ); the variation across years in the  $\pi$ ’s, reflecting, for example, weather or observer-induced effects on how many fish are counted in a survey for a given escapement. Second is the sampling variance of  $\bar{\pi}$  ( $var(\bar{\pi})$ ), which will decline as we collect more data pairs.

The variance for prediction will be estimated (Neter et al. 1990):

$$\hat{var}(\pi_p) = \hat{var}(\pi) + \hat{var}(\bar{\pi}) \quad (2)$$

where:

$$\hat{var}(\pi) = \frac{\sum_{y=1}^k (\pi_y - \bar{\pi})^2}{k - 1} \quad (3)$$

and:

$$\hat{var}(\bar{\pi}) = \frac{\sum_{y=1}^k (\pi_y - \bar{\pi})^2}{k(k - 1)} \quad (4)$$

such that:

$$\hat{var}(\pi_p) = \frac{\sum_{y=1}^k (\pi_y - \bar{\pi})^2}{k - 1} + \frac{\sum_{y=1}^k (\pi_y - \bar{\pi})^2}{k(k - 1)} \quad (5)$$

-continued-

## SYSTEMS WHERE ESCAPEMENT IS ESTIMATED

On systems where escapement is estimated, the expansion factor is an estimate of the expected value of the “population” of annual expansion factors ( $\pi$ ’s) for that system:

$$\bar{\pi} = \frac{\sum_{y=1}^k \hat{\pi}_y}{k} \quad (6)$$

where  $\hat{\pi}_y = \hat{N}_y / C_y$  is the estimate of the expansion factor in year  $y$ ,  $\hat{N}_y$  is the estimated escapement in year  $y$ , and other terms are as described above.

The variance for prediction will again be estimated:

$$\hat{var}(\pi_p) = \hat{var}(\pi) + \hat{var}(\bar{\pi}) \quad (7)$$

The estimate of  $var(\pi)$  should again reflect only process error. Variation in  $\hat{\pi}$  across years, however, represents process error **plus** measurement error within years (e.g. the mark–recapture induced error in escapement estimation) and is described by the relationship (Mood et al. 1974):

$$V(\hat{\pi}) = V[E(\hat{\pi})] + E[V(\hat{\pi})] \quad (8)$$

This relationship can be rearranged to isolate process error, that is:

$$V[E(\hat{\pi})] = V[\hat{\pi}] - E[V(\hat{\pi})] \quad (9)$$

An estimate of  $var(\pi)$  representing only process error therefore is:

$$\hat{var}(\pi) = \hat{var}(\hat{\pi}) - \frac{\sum_{y=1}^k \hat{var}(\hat{\pi}_y)}{k} \quad (10)$$

where  $\hat{var}(\hat{\pi}_y) = \hat{var}(\hat{N}_y) / C_y^2$  and  $\hat{var}(\hat{N}_y)$  is obtained during the experiment when  $N_y$  is estimated. We can calculate:

$$\hat{var}(\hat{\pi}) = \frac{\sum_{y=1}^k (\hat{\pi}_y - \bar{\pi})^2}{k-1} \quad (11)$$

and we can estimate  $var(\bar{\pi})$  similarly to as we did above:

$$\hat{var}(\bar{\pi}) = \frac{\sum_{y=1}^k (\hat{\pi}_y - \bar{\pi})^2}{k(k-1)} \quad (12)$$

where both process and measurement errors need to be included.

---

-continued-

For large  $k$  ( $k > 30$ ), equations (11) and (12) provide reasonable parameter estimates, however for small  $k$  the estimates are imprecise and may result in negative estimates of variance when the results are applied as in equation (7).

Because  $k$  is typically  $< 10$ , we will estimate  $var(\hat{\pi})$  and  $var(\bar{\pi})$  using parametric bootstrap techniques (Efron and Tibshirani 1993). The sampling distributions for each of the  $\hat{\pi}_y$  are modeled using Normal distributions with means  $\hat{\pi}_y$  and variances  $v\hat{a}r(\hat{\pi}_y)$ . At each bootstrap iteration, a bootstrap value  $\hat{\pi}_{y(b)}$  is drawn from each of these Normal distributions and the bootstrap value  $\hat{\pi}_{(b)}$  is randomly chosen from the  $k$  values of  $\hat{\pi}_{y(b)}$ . Then, a bootstrap sample of size  $k$  is drawn from the  $k$  values of  $\hat{\pi}_{y(b)}$  by sampling with replacement, and the mean of this bootstrap is the bootstrap value  $\bar{\pi}_{(b)}$ . This procedure is repeated  $B = 1,000,000$  times. We can then estimate  $var(\hat{\pi})$  using:

$$v\hat{a}r_B(\hat{\pi}) = \frac{\sum_{b=1}^B (\hat{\pi}_{(b)} - \bar{\pi}_{(b)})^2}{B - 1} \quad (13)$$

where:

$$\bar{\pi}_{(b)} = \frac{\sum_{y=1}^k \hat{\pi}_{y(b)}}{B} \quad (14)$$

and we can calculate  $var_B(\bar{\pi})$  using equations (13) and (14) with appropriate substitutions. The variance for prediction is then estimated:

$$v\hat{a}r(\pi_p) = v\hat{a}r_B(\hat{\pi}) - \frac{\sum_{y=1}^k v\hat{a}r(\hat{\pi}_y)}{k} + v\hat{a}r_B(\bar{\pi}) \quad (15)$$

As the true sampling distributions for the  $\hat{\pi}_y$  are typically skewed right, using a Normal distribution to approximate these distributions in the bootstrap process will result in estimates of  $var(\hat{\pi})$  and  $var(\bar{\pi})$  that are biased slightly high, but simulation studies using values similar to those realized for this application indicated that the bias in equation (15) is  $< 1\%$ .

#### PREDICTING ESCAPEMENT

In years when an index count ( $C_p$ ) is available but escapement ( $N_p$ ) is not known, it can be predicted:

$$\hat{N}_p = \bar{\pi} C_p \quad (16)$$

and:

$$v\hat{a}r(\hat{N}_p) = C_p^2 v\hat{a}r(\pi_p) \quad (17)$$

Appendix B2.–Peak aerial survey counts, estimated total spawning abundance with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Keta River 1975–2008. Statistics in bold come directly from mark–recapture experiments in 1998–2000; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$\nu \hat{N}_L$	CV
1975	203	611	114	388	834	12,921	18.6%
1976	84	253	47	161	345	2,212	18.6%
1977	230	692	129	440	945	16,587	18.6%
1978	392	1,180	220	750	1,610	48,181	18.6%
1979	426	1,283	239	815	1,750	56,901	18.6%
1980	192	578	108	367	789	11,559	18.6%
1981	329	990	184	629	1,352	33,939	18.6%
1982	754	2,270	422	1,442	3,097	178,256	18.6%
1983	822	2,475	460	1,573	3,377	211,858	18.6%
1984	610	1,836	342	1,167	2,506	116,670	18.6%
1985	624	1,879	349	1,194	2,563	122,087	18.6%
1986	690	2,077	386	1,320	2,835	149,279	18.6%
1987	768	2,312	430	1,469	3,155	184,937	18.6%
1988	575	1,731	322	1,100	2,362	103,666	18.6%
1989	1,155	3,477	647	2,210	4,745	418,278	18.6%
1990	606	1,824	339	1,159	2,489	115,145	18.6%
1991	272	819	152	520	1,117	23,197	18.6%
1992	217	653	122	415	891	14,765	18.6%
1993	362	1,090	203	693	1,487	41,088	18.6%
1994	306	921	171	585	1,257	29,359	18.6%
1995	175	527	98	335	719	9,602	18.6%
1996	297	894	166	568	1,220	27,658	18.6%
1997	246	741	138	471	1,011	18,975	18.6%
1998	180	<b>446</b>	<b>50</b>	<b>348</b>	<b>544</b>	<b>2,500</b>	<b>11.2%</b>
1999	276	<b>968</b>	<b>116</b>	<b>741</b>	<b>1,195</b>	<b>13,456</b>	<b>12.0%</b>
2000	300	<b>914</b>	<b>122</b>	<b>675</b>	<b>1,153</b>	<b>14,884</b>	<b>13.3%</b>
2001	343	1,033	192	656	1,409	36,888	18.6%
2002	411	1,237	230	786	1,688	52,965	18.6%
2003	322	969	180	616	1,323	32,510	18.6%
2004	376	1,132	211	719	1,545	44,328	18.6%
2005	497	1,496	278	951	2,042	77,449	18.6%
2006	747	2,248	418	1,429	3,068	174,962	18.6%
2007	311	936	174	595	1,277	30,326	18.6%
2008	363	1,093	203	694	1,491	41,316	18.6%
Averages	425	1,282					
Minimum	84	253					
Maximum	1,155	3,477					
$\bar{\pi}$		3.01					
SE $\bar{\pi}$		0.56					
var $\bar{\pi}$		0.313546					

Appendix B3.—Peak aerial survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Blossom River 1975–2008. Statistics in bold come directly from mark–recapture experiments in; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1975	146	439	151	144	734	22,698	34.3%
1976	68	205	70	67	342	4,924	34.3%
1977	112	337	116	110	563	13,357	34.3%
1978	143	430	148	141	719	21,775	34.3%
1979	54	162	56	53	272	3,105	34.3%
1980	89	268	92	88	448	8,435	34.3%
1981	159	478	164	157	800	26,920	34.3%
1982	345	1,038	356	340	1,735	126,743	34.3%
1983	589	1,772	608	580	2,963	369,418	34.3%
1984	508	1,528	524	501	2,555	274,799	34.3%
1985	709	2,133	732	699	3,567	535,278	34.3%
1986	1,278	3,844	1,319	1,259	6,429	1,739,198	34.3%
1987	1,349	4,058	1,392	1,329	6,786	1,937,810	34.3%
1988	384	1,155	396	378	1,932	157,018	34.3%
1989	344	1,035	355	339	1,730	126,010	34.3%
1990	257	773	265	253	1,293	70,332	34.3%
1991	239	719	247	235	1,202	60,825	34.3%
1992	150	451	155	148	755	23,959	34.3%
1993	303	911	313	299	1,524	97,763	34.3%
1994	161	484	166	159	810	27,602	34.3%
1995	217	653	224	214	1,092	50,143	34.3%
1996	220	662	227	217	1,107	51,539	34.3%
1997	132	397	136	130	664	18,554	34.3%
1998	91	<b>364</b>	<b>77</b>	<b>213</b>	<b>515</b>	<b>5,929</b>	<b>21.2%</b>
1999	212	638	219	209	1,066	47,858	34.3%
2000	231	695	238	228	1,162	56,821	34.3%
2001	204	614	211	201	1,026	44,315	34.3%
2002	224	674	231	221	1,127	53,430	34.3%
2003	203	611	209	200	1,021	43,881	34.3%
2004	333	<b>734</b>	<b>71</b>	<b>609</b>	<b>908</b>	<b>5,073</b>	<b>9.7%</b>
2005	445	<b>926</b>	<b>99</b>	<b>791</b>	<b>1,148</b>	<b>9,801</b>	<b>10.7%</b>
2006	339	<b>1,270</b>	<b>172</b>	<b>933</b>	<b>1,607</b>	<b>29,584</b>	<b>13.5%</b>
2007	135	406	139	133	679	19,378	34.3%
2008	257	773	265	254	1,292	70,227	34.3%
Average	313	935					
Minimum	54	162					
Maximum	1,349	4,058					
$\bar{\pi}$		3.01 <sup>a</sup>					
SE $\bar{\pi}$		1.03					
var $\bar{\pi}$		1.0648					
		47					

<sup>a</sup> Includes 2006 estimate.

Appendix B4.–Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Chickamin River 1975–2008. Statistics in bold come directly from mark–recapture experiments; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1975	370	1,758	259	1,251	2,265	66,965	14.7%
1976	157	745	110	530	959	12,009	14.7%
1977	363	1,722	254	1,225	2,219	64,264	14.7%
1978	308	1,465	216	1,042	1,887	46,484	14.7%
1979	239	1,133	167	806	1,460	27,810	14.7%
1980	445	2,112	311	1,503	2,722	96,655	14.7%
1981	384	1,824	268	1,298	2,350	72,075	14.7%
1982	571	2,712	399	1,930	3,495	159,365	14.7%
1983	599	2,847	419	2,026	3,668	175,597	14.7%
1984	1,102	5,235	770	3,724	6,745	593,586	14.7%
1985	956	4,541	668	3,231	5,851	446,721	14.7%
1986	1,745	8,289	1,220	5,898	10,680	1,488,468	14.7%
1987	975	4,631	682	3,295	5,967	464,654	14.7%
1988	786	3,734	550	2,656	4,811	301,971	14.7%
1989	934	4,437	653	3,157	5,716	426,397	14.7%
1990	564	2,679	394	1,906	3,452	155,481	14.7%
1991	487	2,313	340	1,646	2,981	115,925	14.7%
1992	346	1,644	242	1,169	2,118	58,516	14.7%
1993	389	1,848	272	1,315	2,381	73,964	14.7%
1994	388	1,843	271	1,311	2,375	73,584	14.7%
1995	356	<b>2,309</b>	<b>723</b>	<b>1,388</b>	<b>4,650</b>	<b>522,729</b>	<b>31.3%</b>
1996	422	<b>1,587</b>	<b>199</b>	<b>1,279</b>	<b>2,089</b>	<b>39,601</b>	<b>12.5%</b>
1997	272	1,292	190	919	1,665	36,162	14.7%
1998	391	1,857	273	1,321	2,393	74,726	14.7%
1999	501	2,380	350	1,693	3,066	122,686	14.7%
2000	801	3,805	560	2,707	4,902	313,607	14.7%
2001	1,010	<b>5,177</b>	<b>972</b>	<b>3,780</b>	<b>7,573</b>	<b>944,784</b>	<b>18.8%</b>
2002	1,013	<b>5,007</b>	<b>738</b>	<b>3,892</b>	<b>6,742</b>	<b>544,644</b>	<b>14.7%</b>
2003	964	<b>4,579</b>	<b>592</b>	<b>3,481</b>	<b>5,134</b>	<b>350,464</b>	<b>12.9%</b>
2004	798	<b>4,268</b>	<b>893</b>	<b>2,519</b>	<b>6,018</b>	<b>797,449</b>	<b>20.9%</b>
2005	926	<b>4,257</b>	<b>591</b>	<b>3,099</b>	<b>5,415</b>	<b>349,281</b>	<b>13.9%</b>
2006	1,330	6,318	930	4,495	8,140	864,616	14.7%
2007	893	4,242	624	3,018	5,465	389,783	14.7%
2008	1,111	5,277	777	3,755	6,800	603,321	14.7%
Averages	673	3,231					
Minimum	157	745					
Maximum	1,745	8,289					
$\bar{\pi}$		4.75					
SE $\bar{\pi}$		0.70					
var $\bar{\pi}$		0.48879					

Appendix B5.–Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Unuk River 1977–2008. Statistics in bold come directly from mark–recapture experiments; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$V \hat{N}_L$	CV
1977	974	4,739	584	3,594	5,885	341,523	12.3%
1978	1,106	5,382	664	4,081	6,682	440,365	12.3%
1979	576	2,803	346	2,125	3,480	119,439	12.3%
1980	1,016	4,944	610	3,749	6,139	371,612	12.3%
1981	731	3,557	439	2,697	4,417	192,370	12.3%
1982	1,351	6,574	811	4,985	8,163	657,072	12.3%
1983	1,125	5,474	675	4,151	6,797	455,625	12.3%
1984	1,837	8,939	1,102	6,778	11,099	1,214,845	12.3%
1985	1,184	5,761	710	4,369	7,154	504,668	12.3%
1986	2,126	10,345	1,276	7,845	12,845	1,627,155	12.3%
1987	1,973	9,601	1,184	7,280	11,921	1,401,382	12.3%
1988	1,746	8,496	1,048	6,443	10,549	1,097,466	12.3%
1989	1,149	5,591	689	4,240	6,942	475,272	12.3%
1990	591	2,876	355	2,181	3,571	125,741	12.3%
1991	655	3,187	393	2,417	3,957	154,449	12.3%
1992	874	4,253	524	3,225	5,281	274,995	12.3%
1993	1,068	5,197	641	3,941	6,453	410,625	12.3%
1994	711	<b>4,623</b>	<b>1,266</b>	<b>2,992</b>	<b>9,425</b>	<b>1,602,756</b>	<b>27.4%</b>
1995	772	3,757	463	2,849	4,664	214,554	12.3%
1996	1,167	5,679	700	4,306	7,051	490,280	12.3%
1997	636	<b>2,970</b>	<b>271</b>	<b>2,499</b>	<b>3,636</b>	<b>73,441</b>	<b>9.1%</b>
1998	840	<b>4,132</b>	<b>394</b>	<b>3,433</b>	<b>4,974</b>	<b>155,236</b>	<b>9.5%</b>
1999	680	<b>3,914</b>	<b>480</b>	<b>3,110</b>	<b>5,071</b>	<b>230,400</b>	<b>12.3%</b>
2000	1,341	<b>5,872</b>	<b>644</b>	<b>4,848</b>	<b>7,347</b>	<b>414,736</b>	<b>11.0%</b>
2001	2,019	<b>10,541</b>	<b>1,181</b>	<b>8,705</b>	<b>13,253</b>	<b>1,394,761</b>	<b>11.2%</b>
2002	897	<b>6,988</b>	<b>764</b>	<b>5,759</b>	<b>8,677</b>	<b>583,696</b>	<b>10.9%</b>
2003	1,121	<b>5,546</b>	<b>433</b>	<b>4,814</b>	<b>6,530</b>	<b>187,489</b>	<b>7.8%</b>
2004	1,008	<b>3,963</b>	<b>325</b>	<b>3,406</b>	<b>4,684</b>	<b>105,625</b>	<b>8.2%</b>
2005	929	<b>4,742</b>	<b>396</b>	<b>4,094</b>	<b>5,579</b>	<b>156,816</b>	<b>8.4%</b>
2006	940	<b>5,645</b>	<b>506</b>				
2007	720	<b>5,718</b>	<b>474</b>				
2008	- <sup>a</sup>	<b>3,104</b>	<b>390</b>				
Average	1,092	5,466					
Minimum	576	2,803					
Maximum	2,126	10,541					
$\bar{\pi}$		4.87	Based on 1997–2001 and 2003–2005				
SE $\bar{\pi}$		0.60					
var $\bar{\pi}$		0.355230					

<sup>a</sup> Incomplete surveys in 2008.

Appendix B6.–Peak survey counts, weir counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Stikine River 1975–2008. Statistics in bold come directly from mark–recapture experiments; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	Little Tahltan Weir Counts <sup>a, b</sup>	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1975	700		7,571	1,623				21.4%
1976	400		5,723	933				16.3%
1977	800		11,445	1,865				16.3%
1978	632		6,835	1,465				21.4%
1979	1,166		12,610	2,704				21.4%
1980	2,137		30,573	4,982				16.3%
1981	3,334		36,057	7,731				21.4%
1982	2,830		40,488	6,598				16.3%
1983	594		6,424	1,377				21.4%
1984	1,294		13,995	3,000				21.4%
1985	1,598	3,114	16,691	4,204			17,672,775	25.2%
1986	1,201	2,891	15,496	3,903			15,232,238	25.2%
1987	2,706	4,783	25,637	6,457			41,693,495	25.2%
1988	3,796	7,292	39,085	9,844			96,908,274	25.2%
1989	2,527	4,715	25,272	6,365			40,516,408	25.2%
1990	1,755	4,392	23,541	5,929			35,155,413	25.2%
1991	1,768	4,506	24,152	6,083			37,004,106	25.2%
1992	3,607	6,627	35,521	8,946			80,038,968	25.2%
1993	4,010	11,449	61,367	15,456			238,892,573	25.2%
1994	2,422	6,387	34,234	8,675			75,257,360	25.2%
1995	1,117	3,072	16,466	4,400			19,356,920	25.2%
1996	1,920	4,821	<b>23,886</b>	<b>2,773</b>	NA	NA	<b>3,912,484</b>	11.6%
1997	1,907	5,557	<b>28,185</b>	<b>2,977</b>	NA	NA	<b>8,761,600</b>	10.6%
1998	1,385	4,879	<b>25,968</b>	<b>3,931</b>	NA	NA	<b>15,452,761</b>	15.1%
1999	1,379	4,738	<b>19,947</b>	<b>3,240</b>	NA	NA	<b>10,497,600</b>	16.2%
2000	2,720	6,640	<b>27,531</b>	<b>3,168</b>	<b>22,220</b>	<b>34,565</b>	<b>10,036,224</b>	11.5%
2001	4,158	9,738	<b>63,523</b>	<b>5,853</b>	<b>53,741</b>	<b>75,718</b>	<b>34,257,609</b>	9.2%
2002	no survey	7,490	<b>50,875</b>	<b>5,912</b>	<b>40,675</b>	<b>63,900</b>	<b>34,951,744</b>	11.6%
2003	1,903	6,492	<b>46,824</b>	<b>6,078</b>	<b>34,911</b>	<b>58,738</b>	<b>36,942,084</b>	13.0%
2004	6,014	16,381	<b>48,900</b>	<b>3,896</b>	<b>42,179</b>	<b>58,738</b>	<b>15,178,816</b>	8.0%
2005	2,157	7,253	<b>39,833</b>	<b>2,724</b>	<b>20,052</b>	<b>59,885</b>	<b>6,441,444</b>	6.3%
2006	1,372	3,845	<b>24,400</b>	<b>6,938</b>				
2007		562	<b>16,038</b>	<b>2,778</b>				
2008		2,657	<b>18,843</b>	<b>3,153</b>				
Averages	2,107	5,845	27,175					
Minimum	400	2,891	5,723					
Maximum	6,014	16,381	63,523					
$\bar{\pi}$		5.36 <sup>a</sup>						
SE $\bar{\pi}$		1.35						
var $\bar{\pi}$		1.82250						

<sup>a</sup> Does not include 2006, 2007 or 2008 estimate.

<sup>b</sup> Weir count of large fish only, harvest above weir not removed.

Appendix B7.—Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in Andrew Creek 1975–2008. Statistics in bold come directly from weir counts; all other statistics are expanded from counts based on the relationship between counts and estimates during years with mark–recapture experiments.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1975	260	508	116	280	736	13,555	22.9%
1976		<b>404</b>	0				
1977		<b>456</b>	0				
1978		<b>388</b>	0				
1979	221	<b>327</b>	0				
1980		<b>282</b>	0				
1981	300	<b>536</b>	0				
1982	332	<b>672</b>	0				
1983		<b>366</b>	0				
1984	154	<b>389</b>	0				
1985	320	625	143	344	906	20,533	22.9%
1986	708	1,383	317	762	2,005	100,512	22.9%
1987	788	1,540	353	848	2,231	124,510	22.9%
1988	564	1,102	253	607	1,597	63,784	22.9%
1989	530	1,036	237	571	1,501	56,325	22.9%
1990	664	1,298	297	715	1,880	88,408	22.9%
1991	400	782	179	431	1,133	32,083	22.9%
1992	778	1,520	348	837	2,203	121,370	22.9%
1993	1,060	2,071	475	1,141	3,002	225,302	22.9%
1994	572	1,118	256	616	1,620	65,606	22.9%
1995	343	670	154	369	971	23,591	22.9%
1996	335	655	150	361	949	22,503	22.9%
1997	293	<b>478</b>					
1998	487	952	218	524	1,379	47,557	22.9%
1999	605	1,182	271	651	1,713	73,395	22.9%
2000	690	1,348	309	743	1,954	95,467	22.9%
2001	1,054	2,060	472	1,135	2,985	222,758	22.9%
2002	876	1,712	392	943	2,481	153,873	22.9%
2003	595	1,163	266	640	1,685	70,988	22.9%
2004	1,534	2,998	687	1,651	4,344	471,850	22.9%
2005	1,015	1,979	455	1,093	2,874	206,579	22.9%
2006	1,089	2,124	488	1,168	3,079	237,798	22.9%
2007	890	1,736	399	954	2,517	158,830	23.0%
2008	503	981	225	539	1,422	50,733	23.0%
Averages	619	1,082					
Minimum	154	282					
Maximum	1,534	2,998					
$\bar{\pi}$		1.95					
SE $\bar{\pi}$		0.45					
var $\bar{\pi}$		0.200518					

Appendix B8.—Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in King Salmon River, 1971–2008.

Year	Peak Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$\nu \hat{N}_L$	CV
1971	94	143	25	93	193	644	17.78%
1972	90	137	24	89	184	590	17.78%
1973	211	320	57	209	432	3,245	17.78%
1974	104	158	28	103	213	788	17.78%
1975	42	64	11	42	86	129	17.78%
1976	65	99	18	64	133	308	17.78%
1977	134	204	36	133	274	1,309	17.78%
1978	57	87	15	56	117	237	17.78%
1979	88	134	24	87	180	565	17.78%
1980	70	106	19	69	143	357	17.78%
1981	101	153	27	100	207	744	17.78%
1982	259	393	70	256	530	4,890	17.78%
1983	183	<b>245</b>	0				0.00%
1984	184	<b>265</b>	0				0.00%
1985	105	<b>175</b>	0				0.00%
1986	190	<b>255</b>	0				0.00%
1987	128	<b>196</b>	0				0.00%
1988	94	<b>208</b>	0				0.00%
1989	133	<b>240</b>	0				0.00%
1990	98	<b>179</b>	0				0.00%
1991	91	<b>134</b>	0				0.00%
1992	58	<b>99</b>	0				0.00%
1993	175	266	47	173	358	2,232	17.78%
1994	140	213	38	139	287	1,429	17.78%
1995	97	147	26	96	199	686	17.78%
1996	192	292	52	190	393	2,687	17.78%
1997	238	361	64	236	487	4,129	17.78%
1998	88	134	24	87	180	565	17.78%
1999	200	304	54	198	410	2,916	17.78%
2000	91	138	25	90	186	604	17.78%
2001	98	149	26	97	201	700	17.78%
2002	102	155	28	101	209	758	17.78%
2003	78	118	21	77	160	444	17.78%
2004	89	135	24	88	182	577	17.78%
2005	94	143	25	93	193	644	17.78%
2006	99	150	27	98	203	714	17.78%
2007	119	181	32	118	244	1,032	17.78%
2008	79	120	21	78	162	455	17.78%
Averages	1,210	184					
Minimum	42	64					
Maximum	259	393					
$\bar{\pi}$		1.52					
SE $\bar{\pi}$		0.27					
var $\bar{\pi}$		0.072896					

Appendix B9.—Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Taku River, 1973–2008.

Year	Survey Counts	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1973	2,800	14,560	4,984	4,791	24,329	24,840,256	34.2%
1974	3,079	16,011	5,481	5,269	26,753	30,037,196	34.2%
1975 <sup>a</sup>	2,484	12,917	4,422	4,251	21,583	19,549,839	34.2%
1976	4,726	24,575	8,412	8,087	41,063	70,766,455	34.2%
1977	5,671	29,489	10,094	9,704	49,274	101,896,508	34.2%
1978 <sup>a</sup>	3,292	17,118	5,860	5,633	28,604	34,336,787	34.2%
1979	4,156	21,611	7,398	7,112	36,111	54,725,669	34.2%
1980	7,544	39,229	13,428	12,909	65,548	180,319,778	34.2%
1981	9,528	49,546	16,960	16,304	82,787	287,636,173	34.2%
1982	4,585	23,842	8,161	7,846	39,838	66,606,818	34.2%
1983	1,883	9,792	3,352	3,222	16,361	11,234,161	34.2%
1984 <sup>a</sup>	3,995	20,774	7,111	6,836	34,712	50,567,743	34.2%
1985	6,905	35,906	12,291	11,816	59,996	151,066,223	34.2%
1986	7,327	38,100	13,042	12,538	63,663	170,095,329	34.2%
1987	5,563	28,928	9,902	9,519	48,336	98,052,377	34.2%
1988	8,560	44,512	15,237	14,648	74,376	232,160,074	34.2%
1989	8,986	<b>40,329</b>	<b>5,646</b>	<b>30,936</b>	<b>56,995</b>	<b>29,069,351</b>	<b>14.0%</b>
1990	12,077	<b>52,142</b>	<b>9,326</b>	<b>37,072</b>	<b>80,784</b>	<b>52,507,414</b>	<b>17.9%</b>
1991	9,929	51,645	17,674	16,991	86,271	312,356,844	34.2%
1992	10,745	55,889	19,126	18,387	93,361	365,807,701	34.2%
1993	12,713	66,125	22,629	21,754	110,461	512,077,977	34.2%
1994	9,299	48,368	16,552	15,912	80,797	273,975,987	34.2%
1995	7,971	<b>33,805</b>	<b>5,060</b>	<b>25,455</b>	<b>64,388</b>	<b>22,873,263</b>	<b>15.0%</b>
1996	18,576	<b>79,019</b>	<b>9,048</b>	<b>64,388</b>	<b>99,866</b>	<b>124,224,399</b>	<b>11.5%</b>
1997	13,201	<b>114,938</b>	<b>17,888</b>	<b>88,593</b>	<b>157,717</b>	<b>319,980,544</b>	<b>15.6%</b>
1998	5,969	31,039	10,625	10,214	51,864	112,886,800	34.2%
1999	3,951	<b>19,734</b>	<b>3,957</b>	11,978	27,490	<b>15,657,849</b>	<b>20.1%</b>
2000	5,772	<b>30,529</b>	<b>5,417</b>	19,912	41,146	<b>29,343,889</b>	<b>17.7%</b>
2001	5,040	<b>42,980</b>	<b>6,477</b>	30,285	55,675	<b>41,951,529</b>	<b>15.1%</b>
2002	8,089	<b>52,409</b>	<b>10,958</b>	30,931	73,887	<b>120,077,764</b>	<b>20.9%</b>
2003	5,481	<b>36,435</b>	<b>6,409</b>	23,873	48,997	<b>41,075,281</b>	<b>17.6%</b>
2004	9,138	<b>68,199</b>	<b>9,189</b>	50,189	86,209	<b>84,437,721</b>	<b>13.5%</b>
2005	3,981	<b>38,806</b>	<b>4,528</b>	29,931	47,681	<b>20,502,784</b>	<b>11.7%</b>
2006	5,338	<b>41,831</b>	<b>5,542</b>	31,113	52,837	<b>30,713,764</b>	<b>13.2%</b>
2007		<b>17,516</b>	<b>4,383</b>	8,925	26,107	<b>19,210,689</b>	<b>25.0%</b>
2008	4,817	<b>27,383</b>	<b>2,454</b>	22,573	32,193	<b>73,517,935</b>	<b>9.0%</b>
Preliminary M–R estimates							
Averages	6,948	38,221					
Minimum	1,883	9,792					
Maximum	18,576	114,938					
$\bar{\pi}$		5.20 <sup>b</sup>					
SE $\bar{\pi}$		1.78					
var $\bar{\pi}$		3.168400					

<sup>a</sup> Counts from years when all five index areas were not surveyed are expanded by average proportion of index area missed.

<sup>b</sup> Based on 1989, 1990, 1995–1997 and 2003.

Appendix B10.—Peak survey counts, estimated total spawning abundance  $\hat{N}_L$  with associated SEs and approximate 95% CI's for large Chinook salmon spawning in the Alsek River, 1976–2008.

Year	Weir counts <sup>a</sup>	$\hat{N}_L$	SE ( $\hat{N}_L$ )	Lower 95% CI	Upper 95% CI	$v \hat{N}_L$	CV
1976	1,382	5,765	2,360	1,140	10,389	5,567,461	40.9%
1977	2,517	10,496	4,296	2,076	18,917	18,458,142	40.9%
1978	2,819	11,754	4,811	2,324	21,183	23,146,064	40.9%
1979	4,477	18,670	7,642	3,692	33,648	58,396,875	40.9%
1980	1,937	8,077	3,306	1,597	14,557	10,929,596	40.9%
1981	1,997	8,327	3,408	1,646	15,007	11,615,649	40.9%
1982	2,200	9,174	3,755	1,814	16,534	14,100,518	40.9%
1983	2,645	11,028	4,514	2,181	19,875	20,374,823	40.9%
1984	1,797	7,494	3,068	1,482	13,507	9,410,168	40.9%
1985	1,381	5,758	2,357	1,139	10,378	5,554,894	40.9%
1986	2,394	9,981	4,085	1,974	17,988	16,690,357	40.9%
1987	2,733	11,395	4,664	2,253	20,536	21,752,667	40.9%
1988	1,973	8,227	3,367	1,627	14,827	11,339,073	40.9%
1989	2,183	9,105	3,727	1,800	16,409	13,887,877	40.9%
1990	2,109	8,794	3,599	1,739	15,848	12,955,269	40.9%
1991	3,051	12,722	5,207	2,516	22,928	27,115,966	40.9%
1992	1,323	5,519	2,259	1,091	9,928	5,102,791	40.9%
1993	3,043	12,688	5,193	2,509	22,867	26,970,377	40.9%
1994	2,952	12,312	5,039	2,435	22,189	25,395,683	40.9%
1995	6,072	25,322	10,365	5,007	45,637	107,427,633	40.9%
1996	3,464	14,443	5,912	2,856	26,030	34,949,232	40.9%
1997	3,045	12,697	5,197	2,511	22,883	27,008,922	40.9%
1998	1,131	<b>4,969</b>	<b>1,431</b>	<b>2,164</b>	<b>7,774</b>	<b>3,723,801</b>	<b>28.8%</b>
1999	1,918	<b>11,620</b>	<b>4,427</b>	<b>4,940</b>	<b>22,294</b>	<b>10,719,237</b>	<b>32.5%</b>
2000	1,263	<b>6,835</b>	<b>1,678</b>	<b>3,546</b>	<b>10,124</b>	<b>4,650,300</b>	<b>24.6%</b>
2001	1,679	<b>6,111</b>	<b>805</b>	<b>4,533</b>	<b>7,689</b>	<b>8,210,439</b>	<b>13.2%</b>
2002	2,237	<b>5,396</b>	<b>714</b>	<b>3,997</b>	<b>6,795</b>	<b>14,580,748</b>	<b>13.2%</b>
2003	1,416	<b>4,782</b>	<b>534</b>	<b>3,735</b>	<b>5,829</b>	<b>5,843,285</b>	<b>11.2%</b>
2004	2,481	<b>6,995</b>	<b>556</b>	<b>5,905</b>	<b>8,085</b>	<b>17,926,084</b>	<b>7.9%</b>
2005	1,070	4,462	1,826	882	8,042	3,335,472	40.9%
2006	451	1,881	770	372	3,389	592,574	40.9%
2007	628	2,619	1,072	518	4,720	1,148,971	40.9%
2008	321	1,339	548	265	2,412	300,192	40.9%
Averages	2,243	8,993					
Minimum	451	1,339					
Maximum	6,072	25,322					
$\bar{\pi}$		4.17					
SE $\bar{\pi}$		1.71					
Var $\bar{\pi}$		2.91					

<sup>a</sup> Weir count includes immediate harvest below weir times proportion of large fish at weir.



## **APPENDIX C**

Appendix C1.—Computer files used to complete this report.

File Name	Description
TOTALCHTS.XLS	Excel workbook with tables and charts with annual counts for each index area.
SUMVER08.XLS	Appendix Table A2, with expanded escapement totals for Southeast Alaska.
ESCAP2008.XLS	Table 1. Estimated Chinook escapement in 2008.
GOALS.XLS	Appendix Table A1. Expanded goals for Southeast Alaska.
AGELENGTHSEAK2008.XLS	Appendix Table A4-A7. Length and age summaries for 2008.
PahlkeCWTrecovs_08.xls	Coded wire tag recoveries.
ALSEKESC.XLS	Calculation of historical total escapement of Alsek River Chinook based on latest expansion factor.
Total escs with SE.xls	Calculations of total escapements with standard errors, includes estimates of average EF and variance around them.