

Fishery Management Report No. 03-06

**Fishery Management Report for Sport Fisheries in the
Northwest Alaska Management Area, 2001**

by
Fred DeCicco

March 2003

Alaska Department of Fish and Game

Division of Sport Fish



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used in Division of Sport Fish Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications without definition.

Weights and measures (metric)		General		Mathematics, statistics, fisheries	
centimeter	cm	All commonly accepted abbreviations.	e.g., Mr., Mrs., a.m., p.m., etc.	alternate hypothesis	H_A
deciliter	dL	All commonly accepted professional titles.	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	e
gram	g	and	&	catch per unit effort	CPUE
hectare	ha	at	@	coefficient of variation	CV
kilogram	kg	Compass directions:		common test statistics	F, t, χ^2 , etc.
kilometer	km	east	E	confidence interval	C.I.
liter	L	north	N	correlation coefficient	R (multiple)
meter	m	south	S	correlation coefficient	r (simple)
metric ton	mt	west	W	covariance	cov
milliliter	ml	Copyright	©	degree (angular or temperature)	°
millimeter	mm	Corporate suffixes:		degrees of freedom	df
		Company	Co.	divided by	÷ or / (in equations)
Weights and measures (English)		Corporation	Corp.	equals	=
cubic feet per second	ft ³ /s	Incorporated	Inc.	expected value	E
foot	ft	Limited	Ltd.	fork length	FL
gallon	gal	et alii (and other people)	et al.	greater than	>
inch	in	et cetera (and so forth)	etc.	greater than or equal to	≥
mile	mi	exempli gratia (for example)	e.g.,	harvest per unit effort	HPUE
ounce	oz	id est (that is)	i.e.,	less than	<
pound	lb	latitude or longitude	lat. or long.	less than or equal to	≤
quart	qt	monetary symbols (U.S.)	\$, ¢	logarithm (natural)	ln
yard	yd	months (tables and figures): first three letters	Jan,...,Dec	logarithm (base 10)	log
		number (before a number)	# (e.g., #10)	logarithm (specify base)	log ₂ , etc.
Time and temperature		pounds (after a number)	# (e.g., 10#)	mideye to tail fork	MEF
day	d	registered trademark	®	minute (angular)	'
degrees Celsius	°C	trademark	™	multiplied by	x
degrees Fahrenheit	°F	United States (adjective)	U.S.	not significant	NS
hour (spell out for 24-hour clock)	h	United States of America (noun)	USA	null hypothesis	H_0
minute	min	U.S. state and District of Columbia abbreviations	use two-letter abbreviations (e.g., AK, DC)	percent	%
second	s			probability	P
				probability of a type I error (rejection of the null hypothesis when true)	α
Physics and chemistry				probability of a type II error (acceptance of the null hypothesis when false)	β
all atomic symbols				second (angular)	"
alternating current	AC			standard deviation	SD
ampere	A			standard error	SE
calorie	cal			standard length	SL
direct current	DC			total length	TL
hertz	Hz			variance	var
horsepower	hp				
hydrogen ion activity	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

FISHERY MANAGEMENT REPORT NO. 03-06

**FISHERY MANAGEMENT REPORT FOR SPORT FISHERIES IN THE
NORTHWEST ALASKA MANAGEMENT AREA, 2001**

by

Fred DeCicco,
Division of Sport Fish, Fairbanks

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1599

March 2003

The Fishery Management Reports series was established in 1989 for the publication of an overview of Division of Sport Fish management activities and goals in a specific geographic area. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm> This publication has undergone regional peer review.

Fred DeCicco

*Alaska Department of Fish and Game, Division of Sport Fish, Region III,
1300 College Road, Fairbanks, AK 99701-1599, USA*

This document should be cited as:

DeCicco, F. 2003. Fishery Management Report for sport fisheries in the Northwest Alaska regulatory areas, 2001. Alaska Department of Fish and Game, Fishery Management Report No. 03-06, Anchorage.

The Alaska Department of Fish and Game administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility, or if you desire further information please write to ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; U.S. Fish and Wildlife Service, 4040 N. Fairfield Drive, Suite 300, Arlington, VA 22203 or O.E.O., U.S. Department of the Interior, Washington DC 20240.

For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-4120, (TDD) 907-465-3646, or (FAX) 907-465-2440.

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	iv
LIST OF FIGURES.....	iv
LIST OF APPENDICES.....	v
PREFACE.....	1
INTRODUCTION.....	2
Region III Description.....	2
The Alaska Board of Fisheries.....	4
Advisory Committees.....	4
ADF&G Emergency Order Authority.....	4
Region III Sport Fish Division Research and Management Staffing.....	4
The Statewide Harvest Survey.....	5
SECTION I: NORTHWESTERN MANAGEMENT AREA OVERVIEW.....	5
Management Area Description and Its Fisheries Resources.....	5
Seward Peninsula/Norton Sound Sub-area.....	5
Kotzebue/Chukchi Sea Sub-area.....	12
Rural Alaska Sport Fishing.....	14
AYK Sport Fishing Regulations.....	15
Commercial Fisheries.....	15
Subsistence Fisheries.....	21
Alaska Board of Fisheries Activities.....	29
Established Management Plans and Policies.....	29
Major Issues for the Northwestern Management Area.....	30
Access Program.....	31
SECTION II: SEASON SUMMARY FOR 2002.....	32
Pink Salmon.....	32
Chum Salmon.....	32
Chinook Salmon.....	32
Coho Salmon.....	32
Arctic Grayling.....	33
Sheefish.....	33
Dolly Varden.....	34
Commercial and Subsistence Fisheries.....	34
SECTION III: SPORT FISHING EFFORT IN THE NORTHWESTERN MANAGEMENT AREA.....	35
Sport Angling Effort.....	35
SECTION IV: MAJOR NORTHWESTERN AREA FISHERIES OVERVIEW.....	39
Northwestern Alaska Salmon Fisheries.....	39
Regulatory History.....	39
Unalakleet River Salmon Fisheries.....	42
Fishery Description and Historical Perspective.....	42
Recent Fishery Performance.....	45

TABLE OF CONTENTS (Continued)

	Page
Sport Fishery Management Objectives	46
Current Issues	46
Recent and Ongoing Research and Management Activities	47
Nome Area Roadside Salmon Fisheries	48
Fishery Description and Historical Perspective	48
Recent Fishery Performance	59
Sport Fishery Management Objectives	59
Management History and Recent Board of Fisheries and Management Actions	60
Current Issues	61
Ongoing Research and Management Activities	61
Northwestern Alaska Dolly Varden and Arctic Char	62
Fishery Description and Historical Perspective	62
Recent Fishery Performance	67
Wulik River	68
Fishery Objectives and Management	68
Fishery Outlook	68
Recent Board of Fisheries and Management Actions	68
Current Issues	70
Ongoing Research and Management Activities	70
Northwestern Alaska Arctic Grayling	71
Fishery Description and Historical Perspective	71
Recent Fishery Performance	74
Seward Peninsula/Norton Sound Sub-area	74
Kotzebue Sub-area	74
Fishery Objectives and Management	75
Management Goals	75
Management Objectives for Specific Rivers	75
Fishery Outlook	76
Recent Board of Fisheries and Management Actions	76
Current Issues	76
Ongoing Research and Management Activities	77
Kotzebue Sound Sheefish	77
Fishery Description and Historical Perspective	77
Recent Fishery Performance	79
Fishery Objectives and Management	82
Fishery Outlook	82
Recent Board of Fisheries and Management Actions	82
Current Issues	82
Ongoing Research and Management Activities	83
Northwestern Alaska Northern Pike	83
Fishery Description and Historical Perspective	83
Recent Fishery Performance	85
Fishery Management Objectives	85
Fishery Outlook	87
Recent Board of Fisheries and Management Actions	87
Current Issues	87
Ongoing Research Activities	87
ACKNOWLEDGMENTS	87
LITERATURE CITED	87

TABLE OF CONTENTS (Continued)

	Page
APPENDIX A	93
APPENDIX B.....	95
APPENDIX C LIST OF SPORT FISH EMERGENCY ORDERS ISSUED DURING 2001.....	99

LIST OF TABLES

Table	Page
1. Historic commercial salmon harvests by subdistrict from the Norton Sound district 1980-2001	17
2. Kotzebue district chum salmon commercial harvests and incidental Dolly Varden harvests 1980-2001	22
3. Status of recent chum salmon escapements in Nome subdistrict streams, 1995-2001	23
4. Salmon escapement goals (SEG, BEG, or OEG) for Norton Sound area streams, 2001.....	25
5. Subsistence salmon harvests by subdistrict for the Norton Sound district 1980-2001	26
6. Subsistence salmon harvests for the Port Clarence and Kotzebue districts 1980 - 2001.....	28
7. Sport fishing effort in the A-Y-K region by management sub-areas 1982-2001.....	36
8. Freshwater sport fishing effort in angler-days for major rivers by sub-area in the Northwest Management Area, 1983-2001	38
9. Northwest Management Area historic salmon harvests by sub-area 1977-2001.....	40
10. Sport fish effort, harvest, and catch estimates by species for the Unalakleet River, 1990 - 2001	43
11. Sport fish effort and harvests by species from the Nome River 1983-2001, and catches 1990-2001	49
12. Sport fish effort and harvests by species from the Fish/Niukluk River 1983-2001, and catches 1990-2001	50
13. Sport fish effort and harvests by species from the Pilgrim River 1983-2001, and catches 1990-2001	52
14. Sport fish effort and harvests by species from the Snake River 1983-2001, and catches 1990-2001	53
15. Sport fish effort and harvests by species from the Solomon River 1983-2001, and catches 1990-2001	54
16. Sport fish effort and harvests by species from the Kuzitrin River 1983-2001, and catches 1990-2001	55
17. Sport fish effort and harvests by species from the Penny River 1983-2001, and catches 1990-2001	56
18. Sport fish effort and harvests by species from the Cripple River 1983-2001, and catches 1990-2001	57
19. Sport fish effort and harvests by species from the Sinuk River 1983-2001, and catches 1990-2001	58
20. Documented subsistence harvests of Dolly Varden in Noatak and Kivalina	64
21. Historic Dolly Varden harvests and catches in NWMA by sub-area 1977-2001	65
22. Aerial counts of Dolly Varden spawning in the Noatak River and overwintering in the Wulik and Kivalina rivers, 1968-2001	66
23. Historic Dolly Varden and Arctic char harvests in the NWMA by sub-area and rivers, 1988-2001	69
24. Historic Arctic grayling harvests and catches in Seward Peninsula/Norton Sound waters, 1987-2001	72
25. Historic Arctic grayling harvests and catches in the Kotzebue Sound/Chukchi Sea sub-area 1989-2001	73
26. Reported subsistence sheefish harvests, Kotzebue District, 1966-2001	80
27. Historic sheefish sport fish harvests and catches from northwest Alaska waters, 1977-2001	81
28. Historic northern pike sport fish harvests and catches in NWMA by sub-area, 1977-2001	86

LIST OF FIGURES

Figure	Page
1. Map of the sport fish regions in Alaska and the six Region III management areas.....	3
2. The Northwestern Management Area with lines depicting reporting areas W and X	6
3. The Seward Peninsula/Norton Sound sub-area	7
4. Eastern Norton Sound	9
5. Southern Seward Peninsula with road accessible waters	10
6. National Parks Preserves and Wildlife Refuges in NWMA	11
7. Kotzebue Sound Chukchi Sea sub-area	13
8. Commercial salmon fishing subdistricts in Norton Sound	16
9. Port Clarence commercial fishing district	19
10. Kotzebue commercial salmon fishing district	20
11. Sport fishing effort in angler-days in the NWMA by sub-area, 1978-2000	37
12. Dolly Varden and Arctic char distribution in the NWMA	63
13. Sheefish distribution in the NWMA.....	78
14. Northern pike distribution in the NWMA	84

LIST OF APPENDICES

Appendix	Page
A. National Wild and Scenic Rivers in the NWMA	94
B. Northwestern area sport fishing regulations summary for 2000.....	96
C1. Emergency Order issued at Fairbanks, August 27, 2001.....	100
C2. Emergency Order issued at fairbanks, august 23, 2001.....	102
C3. Emergency order issued at fairbanks, august 20, 2001	104
C4. Emergency order issued at Nome, July 31, 2001	106
C5. Emergency order issued at fairbanks, June 14, 2001.....	108

PREFACE

The goals of the Sport Fish Division of the Alaska Department of Fish and Game (ADF&G) are to conserve wild stocks of sport fish, to provide a diversity of recreational fishing opportunities, and to optimize social and economic benefits from recreational fisheries. In order to implement these goals the Division has in place a fisheries management process.

This report provides information for the Northwestern Alaska Management Area (NWMA) and is one in a series of reports providing an annual update of fisheries management information about important sport fisheries within Region III. The report is written to make that information available to the State Board of Fisheries, Fish and Game Advisory Committees, the general public, and other interested parties. It presents fisheries assessment information and the management strategies that are developed from that information. In addition, the report includes a description of the fisheries regulatory process, the geographic, administrative, and regulatory boundaries, funding sources, and other information concerning Sport Fish Division management programs within this area.

An annual regional Area Review is conducted in mid-winter during which the current status of important fisheries are considered and research needs are identified. Fisheries stock assessment research projects are developed, scheduled, and implemented to meet information needs identified by fisheries managers. Projects are planned within a formal operational planning process. Biological information gathered during the course of these research projects is combined with effort information and input from user groups to assess the need for and to develop fisheries management plans and to propose regulatory strategies.

Sport Fish Division management and research activities are primarily funded by a combination of State of Alaska Fish and Game (F&G) and Federal Aid in Fisheries Restoration (D-J) monies. The F&G funds are from the sale of fishing licenses. The D-J (Dingle-Johnson, named after the congressmen who wrote the Act) funds are from a Federal tax on fishing tackle and equipment. D-J funds are provided to the states at a match of up to three-to-one with the F&G funds. There is also an amendment to the D-J Act (W-B, for Wallop-Breaux) that provides money to states for boating access projects at the same three-to-one match with F&G funds. Funding Source for W-B money is a tax on boat gas and equipment. Other, additional funding sources can include contracts with various government agencies and the private sector.

This report provides fisheries information for 2001 with preliminary information from the 2002 season. Following the introduction, which includes an overview of the region, this report is organized into four major sections. **Section I** provides an overview of the Northwestern Alaska Management Area. Included is a description of the management area and sub-areas, Board of Fish activities, and management information and activities within the area. **Section II** provides a summary of the most recent fishing season. **Section III** provides effort and harvest results for the management area and sub areas. **Section IV** provides more detailed summaries of major fisheries and activities occurring during the reporting period. Included in these summaries are a fishery description; a description of recent performance of the fishery; a description of recent Board of Fishery actions related to the fishery; a discussion of social or biological issues that may be associated with each fishery; and a description of ongoing research and management activities related to each fishery.

INTRODUCTION

REGION III DESCRIPTION

The Alaska Board of Fisheries (BOF) divides the state into ten regulatory areas for the purpose of organizing the sport fishing regulatory system by drainage and fishery. These areas (different from Regional Management Areas) are described in Title 5 of the Alaska Administrative Code (AAC 2001). Sport Fish Division of the Alaska Department of Fish and Game (ADF&G) divides the state into three administrative Regions with boundaries roughly corresponding to groups of the BOF regulatory areas (Figure 1). Region I is Southeast Alaska. Region II covers portions of Southcentral Alaska, Kodiak, Southwestern Alaska, and the Aleutian Islands. Region III includes two and most of a third of the BOF fishery regulatory areas. They are the Upper Copper and Upper Susitna regulatory area, the Arctic-Yukon-Kuskokwim (AYK) regulatory area, and the Tanana River drainage. The lower Yukon and lower Kuskokwim drainage from the Aniak River downstream and Kuskokwim Bay were added to the region in spring 2000.

Region III is the largest region, encompassing the majority of the landmass of the State of Alaska (Figure 1). The region contains over 1,251,300 km² (485,000 mi²) of land, some of the state's largest river systems (the Yukon, portions of the Kuskokwim, the Colville, Noatak, Kobuk and upper Copper River and upper Susitna River drainages), thousands of lakes, and thousands of miles of coastline and streams. Regional coastline boundaries extend from Sheldon Point in the southwest, around all of western, northwestern and northern Alaska to the Canadian border on the Arctic Ocean. Region III as a whole is very sparsely populated, with the most densely populated center located in the Tanana River valley. Fairbanks (population about 31,000) is the largest community. Smaller population centers include Delta, Glennallen, Bethel, Galena, Nome and Kotzebue.

For administrative purposes Sport Fish Division has divided Region III into five fishery management areas. They are:

- (1) The Upper Copper/Upper Susitna Management Area (the Copper River drainage and the Susitna River drainage above the Oshetna River).
- (2) The Upper Tanana River Management Area (The Tanana River drainage upstream from Banner Creek and the Little Delta River).
- (3) The Lower Tanana River Management Area (The Tanana River drainage downstream from Banner Creek and the Little Delta River).
- (4) The Northwestern Management Area (Norton Sound, Seward Peninsula and Kotzebue Sound drainages).
- (5) The AYK Management Area (the North Slope drainages, the Yukon River drainage except the Tanana River drainage, and the Kuskokwim River drainage upstream from the Aniak River).
- (6) The Lower Yukon/Lower Kuskokwim (LYLK) Management Area (the Yukon drainage downstream from Paimute and the Kuskokwim drainage downstream from and including the Aniak River drainage), which was created and added to Region III in 2000.




Figure 1.-Map of the sport fish regions in Alaska and the six Region III management areas.

Area offices for the six management areas are in Glennallen, Delta Junction, Fairbanks, Nome/Fairbanks, and Bethel.

THE ALASKA BOARD OF FISHERIES

The Alaska Board of Fisheries (BOF) is the seven-member board that sets fishery regulations and harvest levels, allocates fishery resources, and approves or mandates fishery conservation plans for the State of Alaska. Board members are appointed by the Governor and must be confirmed by the legislature. Board members are appointed for three years.

Statewide fisheries issues may be considered at any BOF meeting. Under the current operating schedule, the BOF considers fishery issues for regulatory areas or groups of regulatory areas on a 3-year cycle. The BOF meetings are usually in the winter, between early October and late March. Regulation proposals and management plans are received for evaluation by the BOF from ADF&G and the public (any Alaskan can submit a proposal to the BOF), and during its deliberations the BOF receives input and testimony through oral and written reports from staff of the Alaska Department of Fish and Game, members of the general public, representatives of local fish and game Advisory Committees, and special interest groups such as fishermen's associations and clubs.

ADVISORY COMMITTEES

Local Fish and Game Advisory committees have been established throughout the state to assist the Boards of Fish and Game in assessing fisheries and wildlife issues and proposed regulation changes. Advisory committee members are individuals from the local public who are nominated and voted on by all present during an advisory committee meeting. Most active committees in urban areas meet in the fall and winter on a monthly basis; rural committees have generally only one fall and one spring meeting due to the logistics of travel from outlying areas. Advisory meetings allow opportunity for direct public interaction with department staff who answer questions and provide clarification concerning proposed regulatory changes. The Boards Support Section within the Division of Administration provides administrative and logistical support for the BOF and Fish and Game Advisory Committees. During 2001, the department had direct support responsibilities for 56 Advisory committees in the state.

ADF&G EMERGENCY ORDER AUTHORITY

ADF&G has emergency order (EO) authority (AAC 2001) to modify time, area, and bag/possession limit regulations. EOs are implemented to deal with conservation issues that are not adequately controlled by existing regulations. In that scenario, they deal with the issue until it is resolved or the BOF can formally take up the issue. EOs are also the mechanism by which "in-season" management of fisheries is accomplished. In-season management is in accordance with a fisheries management plan approved by the BOF when such plans exist.

REGION III SPORT FISH DIVISION RESEARCH AND MANAGEMENT STAFFING

The Region III Sport Fish Division staff biologists are organized into a research group and a management group. The management group consists of a management supervisor, an area management biologist for each of the six management areas, one or more assistant area management biologists, and two stocked waters biologists. The area biologists evaluate fisheries and propose and implement management strategies through plans and regulations in order to meet division goals. A critical part of these positions is interaction with the BOF, Advisory Committees, other agencies, and the general public. The stocked waters biologists plan and

implement the regional stocking program for recreational fisheries. The research group consists of a research supervisor, research biologists, and various field assistants. The research biologists plan and implement fisheries research projects in order to provide information needed by the management group to meet division goals.

The Statewide Harvest Survey

Recreational angling effort and catch and harvest of important sport fish species in Alaska has been estimated and reported annually since 1977 (Mills 1979-1994, Howe et al. 1995, 1996, 2000a-d; Walker et al. *In prep*; Jennings et al. *In prep*). The Statewide Harvest Survey (SWHS), a questionnaire mailed out to a random selection of sport fish license holders, is the instrument that provides the data analyzed to make these estimates. Estimates for a particular year usually become available in August or September of the following year. Effort, catch, and harvest are estimated on a site-specific basis, but estimates of effort directed toward a single species and the resulting species-specific catch-per-unit-effort (CPUE) information can seldom be derived from the report. Utility of the estimates is strongly dependant on the number of responses for a site (Mills and Howe 1992). Estimates based on 12 or fewer responses are useful only to document that fishing occurred. Twelve to 29 responses produce estimates useful for indicating relative order of magnitude and for assessing long-term trends, and estimates based on 30 or more responses are generally the most useful in tracking harvest trends.

SECTION I: NORTHWESTERN MANAGEMENT AREA OVERVIEW

MANAGEMENT AREA DESCRIPTION AND ITS FISHERIES RESOURCES

The Northwest Alaska sport fish management area (Figure 2) includes all waters north of the Yukon River drainage, in Norton Sound, the Seward Peninsula, Kotzebue Sound including the major drainages of the Kobuk and Noatak rivers, and the eastern Chukchi Sea to Point Hope. The total land area consists of approximately 67,800 sq mi (173,500 km²). The management area is comprised of two sub-areas, the Seward Peninsula/Norton Sound sub-area in the south and the Kotzebue/Chukchi Sea sub-area to the north. Fish species present in the Northwest Management Area include anadromous Dolly Varden *Salvelinus malma*, chinook *Oncorhynchus tshawytscha*, coho *O. kisutch*, chum *O. keta*, sockeye *O. nerka* and pink salmon *O. gorbuscha*; Bering cisco *Coregonus laurettae*, humpback whitefish *Coregonus pidschian*, as well as freshwater resident Arctic grayling *Thymallus arcticus*, Dolly Varden *Salvelinus malma*, Arctic char *Salvelinus alpinus*, northern pike *Esox lucius*, sheefish *Stenodus leucichthys*, round whitefish *Prosopium cylindraceum*, least cisco *C. sardinella*, humpback whitefish *C. pidschian*, broad whitefish *C. nasus*, burbot *Lota lota* and lake trout *Salvelinus namaycush*. Most of these species are harvested in sport, personal use or subsistence fisheries. In addition, marine species such as red king crab *Paralithodes camtschatica*, Pacific herring *Clupea harengus*, rainbow smelt *Osmerus mordax*, saffron cod *Eleginus gracilis*, and starry flounder *Platichthys stellatus* are harvested.

Seward Peninsula/Norton Sound Sub-area

The Seward Peninsula-Norton Sound sub-area (statewide harvest Area W; Figure 3) includes all westerly flowing waters and adjacent marine (salt) waters, north of the Yukon River drainage and south of the Selawik River in the Kotzebue Sound/Chukchi Sea sub-area (ADF&G 1984).




Figure 2.-The Northwestern Management Area with lines depicting reporting areas W and X.




Figure 3.-The Seward Peninsula/Norton Sound sub-area.

Streams in eastern Norton Sound (Figure 4) include the Golsovia, Unalakleet, Egavik, Shaktoolik, Inglutalik, Ungalik and Koyuk rivers. All but the Koyuk, drain the Nulato Hills which separate Norton Sound from the Yukon and Koyukuk River valleys. Of these, the Unalakleet River is the largest and most heavily utilized. The village of Unalakleet is located at the mouth of this river. The Unalakleet River has been designated a National Wild and Scenic River (Appendix A1) and supports anadromous populations of Dolly Varden, chinook, coho, chum and pink salmon and resident populations of Dolly Varden, Arctic grayling and whitefish *Coregonus sp.* Other area streams provide the opportunity for high quality fisheries for the same species, but are not as intensively fished because of their remote nature and difficult access.

Many streams located along the southern half of the Seward Peninsula between Koyuk and Teller, (Figure 5) including the Fish, Niukluk, Bonanza, Eldorado, Nome, Snake, Sinuk, Feather, Tisuk, Pilgrim, and Kuzitrin rivers, are accessible via the Nome road system and offer sport fishing opportunity for Arctic grayling, Dolly Varden, salmon and northern pike (Fish, Pilgrim and Kuzitrin rivers). Small sockeye salmon runs occur in the Pilgrim and Sinuk rivers, and a few remnant late run sockeye are present in most other locations while chinook salmon are present in the Pilgrim and Fish rivers. Trophy Arctic grayling, larger than 1.4 kg (3 lbs), are present in many Seward Peninsula waters and some of Alaska's largest Arctic grayling have been taken there. Of the 134 largest Arctic grayling registered in the ADF&G trophy fish program, 34 were taken from Seward Peninsula waters. Remote streams such as the Koyuk, Tubutulik, Kwiniuk, and Agiapuk rivers are accessible by aircraft or boat from nearby villages. These rivers receive little sport fishing effort but provide opportunity for remote high quality fisheries.

Most of the streams draining the northern half of the Seward Peninsula have never been visited by division personnel but likely have limited sport fishing potential due to relatively small flow volumes and difficult access. Much of the northwestern Seward Peninsula is part of the Bering Land Bridge National Preserve (Figure 6).

Other than thaw lakes on the northern side of the Seward Peninsula, there are few lakes in the sub-area. Unique lake formations include five maar lakes south of Cape Espenberg. These lakes were formed by sub-permafrost steam explosions and contain a combination of Arctic char, least cisco and sticklebacks. The largest inland water body is Imuruk Lake in the north-central portion of the Seward Peninsula. It is approximately 32 km² in area, and drains northward via the Inmachuk River. This lake is very shallow with a heavy load of suspended volcanic ash. It contains a small population of resident Arctic grayling.

Some small alpine lakes in the Kigluaik Mountains northeast of Nome contain lake resident Arctic char, (Kretsinger 1987) while others contain Dolly Varden (Phillips et al. 1999). Glacial Lake in the Sinuk River drainage contains sockeye salmon and round whitefish while Salmon Lake, located about 150 km northeast of Nome in the headwaters of the Pilgrim River, contains sockeye salmon, Arctic grayling, round whitefish, least cisco, slimy sculpin, ninespine stickleback, burbot and Dolly Varden which use it as a migration corridor (DeCicco 1995). Even though this lake can be reached by road, it receives little sport fishing use. However, during the first half of the century it was an important fishing area for gold miners in the area and sockeye were nearly extirpated from the drainage. Subsistence fishing for salmon in Salmon Lake has been prohibited for many years because the sockeye stock was practically eliminated by early fisheries. The sockeye population has recently been showing signs of recovery in part due to lake fertilization in recent years.




Figure 4.-Eastern Norton Sound.




Figure 5.-Southern Seward Peninsula with road accessible waters.




Figure 6.-National Parks Preserves and Wildlife Refuges in NWMA. 1) Cape Krusenstern National Monument, 2) Noatak National Preserve, 3) Kobuk Valley National Park and Preserve, 4) Gates of the Arctic National Park, 5) Selawik National Wildlife Refuge, and 6) Beringland Bridge National Preserve.

Kotzebue/Chukchi Sea Sub-Area

The Kotzebue/Chukchi Sea sub-area, statewide harvest Area X, includes all waters and drainages of the Selawik, Kobuk, Noatak, Wulik, Kivalina and Kukpuk rivers (Figure 7). The area also includes all salt water from the northern half of Eschscholtz Bay, including the Chamisso Island area and the northern half of Kotzebue Sound to and including Point Hope (ADF&G 1984).

The most important streams of Kotzebue/Chukchi Sea sub-area are the Noatak and Kobuk rivers, each of which drains approximately 12,000 sq mi (31,000 km²) of the southern and western slopes of the western Brooks Range. Both rivers are approximately 400 mi (640 km) in length (U.S. Army Corps of Engineers 1967). The third largest drainage is that of the Selawik River, with an approximate area of 4,600 sq mi (11,700 km²). Aquifers provide groundwater that serves to stabilize flow and water temperature fluctuations on the lower main stem of the Noatak River and in tributaries of the Kobuk River. These areas provide important overwintering and spawning habitats for many species of fish.

The Noatak River is a National Wild and Scenic River (Appendix A1) and most of the drainage is included in the Noatak National Preserve (Figure 6). The extreme upper headwaters of both the Noatak and Kobuk rivers are included in the Gates of the Arctic National Park. A portion of the lower Kobuk Valley between the villages of Kiana and Ambler is included in the Kobuk Valley National Park, and the Salmon River tributary, as well as the upper main stem of the Kobuk River are National Wild and Scenic Rivers as is the Selawik River. Much of the Selawik River valley is part of the Selawik National Wildlife Refuge. These three large river systems contain abundant fisheries resources.

The Noatak River produces a large run of chum salmon that contributes to a Kotzebue-based commercial fishery. Many thousands of anadromous Dolly Varden overwinter the lower 300 km of the river and spawn in some of the river's tributary streams. During the commercial salmon fishery in August a significant incidental harvest of adult Dolly Varden is sometimes taken. This system is known for its trophy size Dolly Varden. The fish that held the State of Alaska record for Dolly Varden/Arctic char until 2000 was taken from the Noatak River drainage. Whitefish, Arctic grayling, Dolly Varden, lake trout, Arctic char, burbot and northern pike are resident in the Noatak River drainage. Sheefish use the lower reaches of the river for feeding during the spring of the year, but are not known to spawn there (Alt 1987).

The Kobuk River also supports a large run of chum salmon that contributes to the Kotzebue commercial fishery. Major spawning areas are located in many of the Kobuk's tributary streams and in the upper part of the main stem of the river. The Kobuk River contains the largest spawning population of sheefish in northwestern Alaska. Sheefish migrate upstream 300 to 400 miles to spawn in the upper reaches of the drainage. Hotham Inlet, Selawik Lake and the delta system at the river's mouth serve as winter feeding areas for juvenile and adult sheefish. The Alaska state record sheefish, 24 kg (53 lbs), was taken in 1986 from the upper Kobuk River. Abundant whitefish (*C. sardinella*, *C. nasus*, *C. pidschian*) utilize the river, including Selawik Lake and Hotham Inlet (Kobuk Lake). Whitefish support important subsistence fisheries in villages along the river. Dolly Varden, northern pike, Arctic grayling, burbot, lake trout and Arctic char inhabit various parts of the Kobuk watershed.

The Selawik River also supports a spawning population of sheefish that shares rearing and winter feeding areas with the Kobuk population. Sheefish in both populations are slower growing, but attain a larger size than those in other areas of Alaska (Alt 1987). The Selawik River drainage




Figure 7.-Kotzebue Sound Chukchi Sea sub-area.

and associated wetlands provide abundant habitat for whitefish (*C. sardinella*, *C. nasus*, *C. pidschian*) and northern pike, important subsistence resources to the residents of Selawik.

Other important waters in the sub-area include the Wulik and Kivalina rivers, that drain into the Chukchi Sea near the village of Kivalina. These drainages provide rearing, spawning and winter habitat for diadromous Dolly Varden. All five species of North American Pacific salmon, Arctic grayling, burbot and whitefish also occur in these relatively small drainages, but populations are not large.

Sport fishing effort in the northern part of the NWMA is relatively light compared to most other areas in the state with heaviest use on the Noatak, Kobuk, and Wulik rivers. Many visitors to Gates of the Arctic National Park and Kobuk Valley National Park participate in float trips on the Kobuk River from Walker Lake to Kobuk village (Alt 1984; ADF&G 1986; National Park Service (NPS) 1984, 1985). A small amount of shore based sport fishing for sheefish takes place near Kotzebue in the summer. Guided and unguided anglers and river floaters use the Noatak River as do Kotzebue area residents who boat or fly to different parts of the river to fish or hunt. The most popular fishing area on the Noatak River is the Kelly River, but other tributaries such as the Nimiuktuk and Kugururok rivers are also used occasionally for Arctic grayling, chum salmon and Dolly Varden fishing (Alt 1978). Raft, canoe, and kayak trips are becoming increasingly popular. Arctic grayling, Dolly Varden, northern pike and lake trout are available in the upper Noatak River, and downstream from the Nimiuktuk River, chum salmon also occur. Lake trout occur in Matcharak, Feniak, and Desperation lakes and in some other lakes in the middle and upper Noatak drainage. Some lakes also contain Arctic char or Dolly Varden. Most lakes in the area are accessible during summer months only by floatplane. Thirteen lakes surveyed by Alt (1978) in the upper Noatak River all contained fish. Round whitefish, lake trout and Arctic grayling were the most common species. Least cisco, northern pike, Arctic char, slimy sculpin *Cottus cognatus*, salmon (chum and sockeye), and ninespine stickleback *Pungitius pungitius* were also found. Of six lakes surveyed in 2000, two contained resident populations of Dolly Varden.

The lower floodplains of the Kobuk and Selawik rivers, especially in the vicinity of the Kobuk River delta, and the lower Noatak River contain hundreds of shallow thaw lakes of various sizes. Fisheries resources in this area have been poorly inventoried, but populations of whitefish, and northern pike are known to be seasonally present. Dolly Varden spawn in several Kobuk River tributary streams, including the Squirrel, Salmon, Tutuksuk, Hunt and Ambler rivers. The mountains in the upper Kobuk River drainage contain several relatively large, oligotrophic lakes. Lake trout, Arctic grayling, Arctic char, northern pike and several species of whitefish inhabit Walker, Nutuvukti, and Selby lakes.

Most sport fishing throughout the region is by unguided private individuals. The sport fish guiding industry, while present in many of the region's best fishing waters, is not as large or well developed as in other parts of the state. DeCicco and Barnes (1992) produced a list of guide services by area, species and fishery.

RURAL ALASKA SPORT FISHING

With the exception of the limited road system around Nome, waters of the Northwestern Management Area are not accessible from highways or roads of any kind. Small communities are scattered along the major river systems of the area and along the coast of western Alaska. The communities are invariably located on or near water because of the importance of fish as a

food source to native people historically and today. Native residents harvest substantial amounts of fish and game resources for personal subsistence use. Subsistence fishing is usually conducted using nylon gillnets or seines. Fishing with rod and reel is also practiced to some extent by rural residents, but most often as an extension of subsistence activities and less for recreational purposes. Consequently, harvest estimates of sport caught fish from rural Alaska are generally low, in part because local residents often fish under subsistence regulations and because the small amount of sport fishing done by them is often considered as part of their normal subsistence activities. Since statewide harvest estimates are based upon surveys of licensed sport fishers, rural harvests may not be fully documented.

AYK SPORT FISHING REGULATIONS

Published regulations for the Northwestern Management Area for 2001 are reproduced as Appendix B.

COMMERCIAL FISHERIES

Although small when compared to the major commercial fisheries in southeast and southwest Alaska, the commercial fisheries in northwest Alaska form an economic base for income and employment in many local communities. Commercial harvests for salmon, herring, halibut and crab are much larger than sport harvests for those species. In addition, extremely limited commercial fisheries exist for freshwater species such as sheefish, Dolly Varden and whitefish. Although personal use fisheries are also allowed, there has been no participation in these fisheries in the NWMA largely because all Alaska residents qualify as subsistence users. Subsistence harvests of salmon, Dolly Varden, sheefish, whitefish and crab are very important to the economies of the many small villages in the NWMA, and are much larger than the sport fish harvests which make up the smallest component of overall use in most years.

The Division of Commercial Fisheries Management and Development (CFMD) manages commercial fisheries in the Northwestern Management Area where commercial fisheries for salmon in the Norton Sound management district have been ongoing since 1961. The initial species of interest were chinook and coho, but fisheries have also developed for chum and pink salmon. The district is divided into six subdistricts to facilitate management of individual stocks or stock groups. Subdistricts include: 1) Nome, 2) Golovin, 3) Moses Point, 4) Norton Bay, 5) Shaktoolik, and 6) Unalakleet (Figure 8). Conservation concerns for chum salmon stocks have resulted in very little commercial salmon fishing in the Nome subdistrict since the early 1980s. There has likewise been little recent commercial fishing in the Norton Bay subdistrict, but this has largely been the result of limited markets in this remote area (Brennan et al. 2002). Average commercial harvests over the last five years (1996-2000) in the Norton Sound district have been 5,649 chinook, 37,437 coho, 15,437 chum, and 248,404 pink salmon (Table 1). Commercial harvests in 2001, 213 chinook, 19,492 coho, 11,100 chum, 0 pink and 44 sockeye salmon, were the lowest on record. The Port Clarence District includes all waters from Cape Douglas north to Cape Prince of Wales, including the drainages of the Pilgrim and Kuzitrin rivers (Figure 9). Commercial salmon fishing has been prohibited in this district since 1967. Few stocks are present and their run sizes are relatively small, however, the sockeye run into Salmon Lake that passes through the district has increased to between 5,000 and 10,000 fish in recent years. Because of the existence of important subsistence fisheries on these stocks, commercial fishing has never reopened. The Kotzebue Sound District includes all waters from Cape Prince of Wales to Point Hope (Figure 10) and is the northern most commercial fishing




Figure 8.-Commercial salmon fishing subdistricts in Norton Sound.

Table 1.-Historic commercial salmon harvests by subdistrict from the Norton Sound district 1980-2001.

Year	Nome (Subdistrict 1)					Golovin (Subdistrict 2)					Moses Point (Subdistrict 3)				
	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum
1980	8			10,007	23,937	36	36	328	10,744	52,609	502			1,435	14,755
1981	4		508	3,202	22,380	23	5	13	49,755	58,323	198		5	26,417	29,325
1982	20		1,183	18,512	33,162	78	5	4,281	39,510	51,970	253		318	9,849	40,030
1983	23		261	308	12,283	52	10	295	17,414	48,283	254			17,027	65,776
1984	7		820		4,571	31		2,462	88,588	54,153			5,959	28,035	9,477
1985	21		356		6,596	193	113	1,196	3,019	55,781	816	32	1,803	559	24,466
1986	6		50		8,216	81	8	958	25,425	68,725	600	41	5,847	15,795	20,668
1987	3		577		6,226	166	51	2,203	1,579	44,344	907	15	64	568	17,278
1988	2		54	182	1,866	108	921	2,149	31,599	33,348	663	93	3,974	13,703	18,585
1989	2			123	617			0	0		62			0	167
1990	0	0				52	21	0	0	15,993	202	0	0	501	3,423
1991	0	0				49	1	0	0	14,839	161	0	0	0	804
1992	1	2	693	185	1,762	6	9	2,085	0	1,002	0	0	3,531	0	6
1993	0	2	611	0	745	1	4	2	8,480	2,803	3	0	4,065	0	167
1994	0	1	287	0	354	0	0	3,424	0	111	0	0	5,345	0	414
1995	0	1	369	0	492	0	0	1,616	4,296	1,987	4	44	3,742	2,962	1,171
1996	0	0	9	13	25	0	0	638	0	0	0	0	1,915	68,609	0
1997	0	0	0	0	0	19	2	102	20	8,003	844	0	1,409	0	2,683
1998	0	0	0	0	0	1	0	3	106,761	723	105	0	1,462	145,699	2,311
1999	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2000	0	0	0	0	0	0	0	1,645	17,408	164	10	0	5,182	46,369	535
2001	0	0	0	0	0	0	43	30	0	7,094	7	0	1,696	0	681
96-00 Avg	0	0	2	3	5	4	0	478	24,838	1,778	192	0	1,994	52,135	1,106
91-00 Avg	0	1	219	22	375	8	2	952	13,697	2,963	113	4	2,665	26,364	809

-continued-

Table 1.-Page 2 of 2.

Year	Norton Bay (Subdistrict 4)					Shaktoolik (Subdistrict 5)					Unalakleet (Subdistrict 6)				
	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum
1980	340			47	7,855	1,086		8,001	1,947	27,453	4,339	3	21,512	203,142	64,198
1981	63			177	3,111	1,484	4	1,191	29,695	21,097	6,157	47	29,845	123,233	39,186
1982	96		2,332	2,535	7,128	1,677	3	22,233	17,019	26,240	3,768	2	61,343	142,856	44,520
1983	215		204	3,935	17,157	2,742	4	12,877	12,031	67,310	7,022	13	36,098	26,198	109,220
1984				1,162	3,442	1,613		10,730	1,596	32,309	6,804	6	47,904		43,317
1985	528		384	68	9,948	5,312	29	2,808		13,403	12,621	21	15,421	1	25,111
1986	139	2	1,512	40	1,994	1,075		6,626		16,126	4,494	153	20,580		30,239
1987	544		145	16	3,586	2,214	79	6,193		14,088	3,246	141	15,097	97	17,525
1988	434	2	709	1,749	7,521	671	43	6,096	3,671	21,521	2,218	157	24,232	23,730	25,363
1989						1,241	49	8,066		19,641	4,402	222	36,025		20,825
1990	0	0	0	0	0	2,644	55	4,695		21,748	5,998	358	52,015		23,659
1991	0	0	0	0	0	1,324	56	11,614		31,619	4,534	147	52,033		39,609
1992	27	0	0	0	1,787	1,098	20	14,660		27,867	3,409	229	84,449	6,284	52,547
1993	267	0	0	290	1,678	3,756	8	11,130	106,743	20,864	5,944	251	26,290	42,061	28,156
1994	0	0	0	0	0	885	5	22,065	502,231	5,411	4,400	71	71,019	480,158	12,288
1995	0	0	0	0	0	1,239	1	10,856	37,377	14,775	7,617	78	31,280	37,009	24,843
1996	0	0	0	0	0	1,340		13,444	304,982	3,237	3,644		52,200	113,837	7,369
1997	194	0	0	0	531	2,449		4,694	0	5,747	9,067	159	26,079		17,139
1998	0	0	0	0	0	910		3,624	236,171	7,080	6,413	7	24,534	99,412	6,210
1999	0	0	0	0	0	581	0	2,398	0	2,181	1,927	0	10,264	0	5,700
2000	0	0	0	0	0	160	3	7,779	85,493	2,751	582	11	29,803	17,278	2,700
2001	0	0	0	0	0	90	0	2,664	0	1,813	116	0	15,102	0	1,512
96-00 Avg	39	0	0	0	106	1,088	2	6,388	125,329	4,199	4,327	44	28,576	57,632	7,824
91-00 Avg	49	0	0	29	400	1,374	13	10,226	159,125	12,153	4,754	106	40,795	99,505	19,656




Figure 9.-Port Clarence commercial fishing district.




Figure 10.-Kotzebue commercial salmon fishing district.

district in Alaska. The current commercial fishery opened under state management in 1962, but there are documented sales of salmon in the Kotzebue area dating back to the early 1900s. This is primarily a chum salmon fishery with a few chinook taken annually and an incidental take of Dolly Varden that pass through the fishery in August. Average commercial harvests over the past five years (1996-2000) in the Kotzebue Sound District have been 115,829 chum salmon and about 1,100 Dolly Varden (Table 2). The harvest in 2001 was 211,672 chum salmon and zero Dolly Varden. There is also a directed under ice commercial fishery on sheefish in Hotham Inlet. Documented annual harvests in this fishery have averaged only 250 fish over the past five years, and the harvest quota of 25,000 pounds has never been met. Brennan et al. (2002) documents these fisheries in greater detail.

The CFMD conducts annual assessments of salmon escapements using weirs, counting towers and aerial surveys (Table 3). The status of Norton Sound chum salmon stocks of concern was recently reviewed by the BOF and escapement goals for chum salmon based on aerial surveys, tower or weir counts in Nome Subdistrict streams have been established (Table 4). In addition, Sustainable Escapement Goals (SEGs) have recently been developed salmon stocks that lacked adequate data for the development of more formalized BEGs. Optimal Escapement Goals (OEG) have also been developed for some species/river combinations. An OEG is a specific management objective for escapement that considers biological and allocative factors and may differ from the SEG or BEG. The BOF places an OEG into regulation and the department manages the fisheries to maintain escapements within the bounds of the OEG. A number of these goals are based on tower or weir counts. These include Snake, Nome, Eldorado, and Kwiniuk rivers chum salmon; Nome and North River pink salmon; Kwiniuk and Niukluk rivers coho; and North River chinook. The remainder of the goals are based on aerial surveys (Table 4). Additional goals will be revised using tower or weir counts when sufficient data are available.

SUBSISTENCE FISHERIES

There are approximately 16,000 people in the NWMA. Except for the two larger communities of Nome and Kotzebue, the population is scattered among 26 small villages along the coast and the major area rivers (Alaska Dept. of Labor 1991). Most of the population is comprised of Alaska Natives, many of whom lead a relatively traditional lifestyle. Most area residents rely heavily on the subsistence use of fish and wildlife for their livelihood. Subsistence use of salmon is monitored in village surveys conducted by the Division of Subsistence. Recent subsistence salmon harvests (1995-1999) have averaged about 98,000 fish in the Norton Sound District (Table 5); 11,000 in the Port Clarence District; and about 84,000 in the Kotzebue Sound District (Table 6). In 2001, 71,346 salmon were estimated harvested for subsistence in the Norton Sound District, 8,168 in the Port Clarence District, and 49,844 in the Kotzebue District. In addition to salmon, saffron cod, rainbow smelt, Dolly Varden and whitefish are taken. In the Kotzebue District sheefish are also an important subsistence resource, especially in the villages along the Kobuk River, Kotzebue, and Selawik. In 2001, an estimated 3,838 sheefish were taken for subsistence in the Kobuk River villages (Georgette and Utermohle 2001). The relative importance of whitefish is higher in the Kotzebue Sound District than in many areas of the state. The subsistence harvest of whitefish in 1997 was estimated at 84,851 for the village of Noatak and the five Kobuk River villages combined. In 1998, 1999, and 2000 the whitefish harvests were estimated at 39,754 and 56,326, 70,097 respectively (Brennan et al. 2001; Georgette and

Table 2.-Kotzebue district chum salmon commercial harvests and incidental Dolly Varden harvests 1980-2001.

Year	Chum Salmon	Dolly Varden	
		Sold	Caught
1980	367,284	3,049	
1981	677,239	3	
1982	417,790	3,447	
1983	175,762	190	
1984	320,206	347	1,090
1985	521,406	454	3,600
1986	261,436	5	2,373
1987	109,467	1,261	
1988	352,915	752	
1989	254,617	3,093	
1990	163,263	604	
1991	239,923	6,136	
1992	289,184	1,977	
1993	73,071	76	
1994	153,452	149	
1995	290,730	2,090	
1996	82,110	188	
1997	142,720	3,320	
1998	55,907	349	
1999	138,605	1,502	
2000	159,802	7	
2001	211,662	0	
96-00 Avg	115,829	1,073	
91-00 Avg	162,550	1,579	

Table 3.-Status of recent chum salmon escapements in Nome subdistrict streams, 1995-2001.

Location	Escapement Goal (number of fish)	Year						
		1995 Estimate	1996 Estimate	1997 Estimate	1998 Estimate	1999 Estimate	2000 Estimate	2001 Estimate
Sinuk River								
Aerial Survey	4,000-6,200	3,110	1,815 (early)	2,975	630 (incomplete)	no survey	10 (incomplete)	3,746
Snake River								
Tower	1,600-2,500	4,393	2,772	6,184	11,067	484	1,400 ^a	2,182
Nome River								
Tower/Weir	2,900-4,300	5,092	3,339	5,131	976	1,048	4,051 ^a	2,859
Eldorado River								
Tower	6,000-9,200	39,867	12,655	14,302	13,808	4,218	10,604 ^a	11,635
Flambeau								
Aerial Survey	4,100-6,300	6,455	5,380	905 (incomplete)	2,828 (incomplete)	No survey	819 (incomplete)	3,612
Bonanza River								
Aerial Survey	1,000-1,900	0 (incomplete)	1,980 (incomplete)	881 (incomplete)	no survey	361 (incomplete)	1,130 (incomplete)	1,084
Solomon River								
Aerial Survey	1,100-1,600	315	323	316 (incomplete)	90 (incomplete)	51 (incomplete)	150 (incomplete)	280
Fish River								
Niukluk Tower	(Goal not set)	86,333	80,121	57,304	45,587	35,240	26,724	30,662
Fish R. & Boston Cr.								
Combined Aerial Survey Index	23,200-46,400	43,012	19,077 (incomplete)	40,500 (incomplete)	4,126 (incomplete)	640 (incomplete)	no Survey	6,753 (incomplete)
Kwiniuk River								
Kwiniuk Tower	11,500-23,000	42,703	28,493	20,118	24,248	8,763	12,251 ^a	16,598

-continued-

Table 3.-Page 2 of 2.

Location	Escapement Goal (number of fish)	Year						
		1995 Estimate	1996 Estimate	1997 Estimate	1998 Estimate	1999 Estimate	2000 Estimate	2001 Estimate
Tubutulik River								
Aerial Survey	8,000-16,000	16,518	10,790 (incomplete)	3,105	10,060	no survey	no Survey	863
Unalakleet River								
North Tower	(Goal not set)	no project	9,789 (incomplete)	6,904 (incomplete)	5,421 (incomplete)	5,600 (incomplete)	3,717 ^a	6,515
Unalakleet Test Fish	560 (Index)	1,101	1,424	743	492	956	1,083	
Unalakleet, Old Woman Rivers Combined Aerial Survey Index	2,400-4,800	6,080	296 (incomplete)	4,840 (incomplete)	1,230 (incomplete)	no survey	no survey	no survey

Note: "Incomplete" survey indicates a survey was flown, but not used in the "Goal" assessment due to timing or survey conditions

Table 4.-Salmon escapement goals (SEG, BEG, or OEG) for Norton Sound area streams, 2001.

Location	Type	Chum	Type	Chinook	Type	Sockeye	Type	Coho	Type	Pink
Salmon Lake					Aerial	Combined				
Grand Central River						4,000-8,000				
Pilgrim River										
Glacial Lake					Aerial	800-1,600				
Sinuk River	Aerial	4,000-6,200								
Cripple River										
Penny River										
Snake River	Weir	1,600-2,500								
Nome River	Weir	2,900-4,300							Weir	13,000
Flambeau River		4,100-6,300								
Eldorado River	Weir	6,000-9,200								
Solomon River		1,100-1,600								
Ophir Creek							Aerial	Combined		
Niukluk River								950-1,900	Tower	8,400
Fish River		Combined	Aerial	Combined						
Boston Creek		23,200-46,400		100-250						
Kwiniuk	Tower	11,500-23,000	Tower	300-550			Aerial	650-1,300	Tower	12,500
Tubutulik	Aerial	9,200-18,400								
Inlutalik River										
Ungalik River										
Shaktoolik			Aerial	400-800					Aerial	48,000
Unalakleet River		Combined	Aerial	Combined						
Old Woman River		2,400-4,800		550-1,100						
North River			Tower	1,200-2,400			Aerial	550-1,100	Tower	8,500

Table 5.-Subsistence salmon harvests by subdistrict for the Norton Sound district 1980-2001.

Year	Nome (Subdistrict 1)					Golovin (Subdistrict 2)					Moses Point (Subdistrict 3)				
	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum
1980	129		2,157	22,246	5,983	12		692	10,727	4,057	131		229	4,232	1,393
1981	35	14	1,726	5,584	8,579	8		1,520	5,158	5,543	32		2,345	6,530	2,819
1982	21	6	1,829	19,202	4,831	7		1,289	4,752	1,868	1		1,835	3,785	3,537
1983	74	53	1,911	8,086	7,091										
1984	83	16	1,795	17,182	4,883										
1985	56	114	1,054	2,117	5,667	12	2	430	1,904	9,577	67		1,389	1,212	947
1986	150	107	688	8,720	8,085										
1987	200	107	1,100	1,251	8,394										
1988	63	133	1,076	2,159	5,952										
1989	24	131	469	924	3,399										
1990	58	234	510	2,233	4,246										
1991	83	166	1,279	194	3,715						312		2,153	3,555	2,660
1992	152	163	1,481	7,351	1,684						100		1,281	6,152	1,260
1993	52	80	2,070	873	1,766						368		1,217	1,726	1,635
1994	23	69	983	6,556	1,673	253	168	733	8,410	1,337	322	104	1,180	9,345	3,576
1995	36	211	1,897	486	5,344	165	34	1,649	7,818	10,373	284	17	1,353	2,046	3,774
1996	19	353	1,317	5,802	4,333	86	134	3,014	17,399	2,867	417	52	1,720	9,442	2,319
1997	19	99	534	287	4,996	138	427	555	4,570	4,891	619	50	1,213	1,314	2,064
1998	15	14	1,057	4,797	964	184	37	1,292	13,340	1,893	414	49	1,831	6,891	1,376
1999	11	85	161	58	337	60	48	1,234	469	3,656	424	13	975	1,564	744
2000	7	26	747	2,657	535	169	18	2,335	10,906	1,155	248	46	1,429	5,983	1,173
2001	2	92	425	113	858	89	72	880	1,665	3,291	427	70	1,352	1,390	898
Avg 96-00	14	115	763	2,720	2,233	127	133	1,686	9,337	2,892	424	42	1,434	5,039	1,535
Avg 91-00	42	127	1,153	2,906	2,535	151	124	1,545	8,987	3,739	351	47	1,435	4,802	2,058

-continued-

Table 5.-Page 2 of 2.

Year	Norton Bay (Subdistrict 4)					Shaktoolik (Subdistrict 5)					Unalakleet (Subdistrict 6)				
	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum	Chinook	Sockeye	Coho	Pink	Chum
1980	22		33	4,275	1,132	57		756	3,227	1,827	1,046		4,758	19,071	5,230
1981	7		82	2,314	3,515	8		525	2,225	3,490	869	24	5,808	5,750	4,235
1982	1		484	2,600	2,485	68		2,138	3,865	1,165	913	2	7,037	20,045	4,694
1983											1,868	33	6,888	13,808	4,401
1984											4,650	1	6,675	17,418	3,348
1985						298		1,379	24	298	1,397	3	2,244	55	1,968
1986															
1987															
1988															
1989													4,681	17,500	1,388
1990											2,476				
1991															
1992															
1993															
1994	308	1	370	6,049	4,581	1,175	1	2,777	9,133	1,221	5,294	819	16,081	31,572	12,732
1995	475	46	985	3,514	5,828	1,275	2,480	2,626	7,024	2,480	5,049	807	13,110	17,246	13,460
1996	295	3	676	3,929	4,161	1,114	31	3,615	8,370	4,425	5,324	608	15,963	19,782	16,481
1997	656	54	322	1,795	4,040	1,146	62	2,761	5,779	1,612	6,325	353	9,120	10,804	7,646
1998	684	0	388	2,009	6,192	982	92	1,872	6,270	1,034	3,963	201	7,303	13,173	2,551
1999	327	0	167	1,943	4,153	818	183	1,556	5,092	467	2,691	537	8,140	10,067	3,692
2000	397	2	267	2,255	4,714	440	20	2,799	5,432	2,412	2,429	212	5,878	10,631	3,000
2001	460	14	276	5,203	4,445	963	143	2,090	10,172	1,553	2,810	359	6,270	11,279	2,918
Avg 96-00	472	12	364	2,386	4,652	900	78	2,521	6,189	1,990	4,146	382	9,281	12,891	6,674
Avg 91-00	449	15	454	3,071	4,810	993	410	2,572	6,729	1,950	4,439	505	10,799	16,182	8,509

Table 6.-Subsistence salmon harvests for the Port Clarence and Kotzebue districts 1980-2001.

Year	Port Clarence District					Kobuk River Villages	Kotzebue District Chum Salmon		
	Chinook	Sockeye	Coho	Pink	Chum		Noatak	Kotzebue	Other Villages
1980	7	3,195	5	3,170	1,715	8,494	2,135	2,387	455
1981	8	255	110	765	5,845	9,459	5,465	4,099	1,017
1982	23	405	100	4,345	684	19,648	5,479	347	419
1983	17	261		615	299	5,486	4,035	88	2,140
1984						7,231	6,049	13,494	573
1985						17,411		36,311	
1986						12,901	1,246		
1987						7,067	2,921		
1988						13,723			
1989	28	535	472	395	410	3,894	1,595		
1990						4,353	3,921		
1991						11,103	3,637		
1992						12,260	2,043		
1993						12,160	3,270		
1994	181	1,979	1,692	3,849	2,042	26,612	6,126		3,488
1995	76	4,481	1,739	3,293	6,011	38,867	6,359	50,708	6,947
1996	195	4,558	2,079	2,587	1,264	39,076	10,091	50,573	
1997	158	3,177	829	755	2,099	26,242	5,309	26,355	
1998	287	1,665	1,759	7,812	2,621	21,398	2,614	24,986	
1999	89	2,392	1,030	786	1,936	14,264	1,616	64,768	
2000	72	2,851	935	1,387	1,275	21,538	7,293	37,144	
2001	84	3,692	1,299	1,183	1,910	28,975	2,326	17,713	
Avg 96-00	160	2,929	1,326	2,665	1,839	24,504	5,385	40,765	
Avg 91-00	151	3,015	1,438	2,924	2,464	22,352	4,836	42,422	5,218

Utermohle 2001). In 2001, 30,976 whitefish were estimated to have been harvested (Georgette et al. *In press*).

ALASKA BOARD OF FISHERIES ACTIVITIES

The development of regulations for recreational fisheries in the NWMA occurs within the established Alaska Board of Fisheries (BOF) process. Local fish and game advisory committees have been established throughout Alaska to assist the BOF by bringing local issues to their attention, and proposing or commenting on regulation changes proposed for upcoming meetings. Active committees meet at least once a year and in the fall prior to scheduled BOF meetings in order to provide timely information regarding regulation proposals or concerns that may affect a local area. ADF&G staff is often invited to attend committee meetings, to interact with the public, and to provide information to the committee regarding issues of local concern. Within the NWMA there are eight local advisory committees to serve resource users of the area: Kotzebue, Noatak/Kivalina, Upper Kobuk, Lower Kobuk, Northern Seward Peninsula, Norton Sound, Southern Norton Sound and St. Lawrence Island advisory committees. The current BOF schedule provides for meetings rotated through areas of the state on a 3-year schedule. The last BOF meeting that addressed the NWMA occurred in December 2000. The BOF adopted several regulations that affected sport fisheries in the NWMA during that meeting.

Two regulations dealt with salmon fisheries in the Unalakleet River drainage, and one with sheefish in the Selawik River drainage. One statewide salmon regulation affected salmon fisheries in the NWMA. The statewide chinook salmon regulation set the daily bag and possession limit for chinook salmon less than 20 inches in length (jack kings) at 10 per day. In the Unalakleet River drainage, the daily bag and possession limit for salmon, other than king salmon, was reduced from 10 per day to five per day. To address catch and release mortality of coho salmon released in the estuary, a regulation was passed that prohibited an angler from fishing downstream from the South River after a bag limit of coho had been taken. In the Selawik River drainage, the daily bag and possession limit for sheefish upstream from the Tagagawik River (spawning area) was reduced from 10 fish to 2 fish. This action made the regulations on the Selawik River consistent with those on the Kobuk River. In addition, subsistence fishing for Arctic grayling was closed on the Nome and Solomon rivers, and rod and reel was recognized as a legitimate subsistence method in northern Norton Sound.

The area management biologist has EO authority (5AAC 2001) that allows the in-season modification of time, area, and bag/possession limit regulations as necessary to address conservation concerns on a species, area or fishery basis. EOs issued in the NWMA during the reporting period are summarized in Appendix C, and mentioned in the following sections of this document addressing specific fisheries.

ESTABLISHED MANAGEMENT PLANS AND POLICIES

There are presently no specific BOF adopted management plans that pertain to the NWMA sport fisheries. However, the Division of Sport Fisheries has developed objectives for the region or its constituent areas and has identified them in fishery based management plans. A management plan was recently developed for Nome area roadside Arctic grayling fisheries (DeCicco 2002a). In addition, a series of general divisional criteria that have been prepared to guide the establishment of fishery objectives which are listed below:

1. **Protection and management of existing fish resources.** Divisional activities should strive to manage and protect Alaska's wild stocks of fish resources for future generations.
2. **Public use and benefits of existing fish resources.** Alaska's fishery resources should be made available for public use and benefit on a sustained yield basis.
3. **Rehabilitation of depressed stocks and damaged habitat.** Division activities should strive to restore and maintain fish stocks and habitat damaged by human activities.
4. **Enhancement of natural production or creation of new opportunities.** The Division should pursue creation of new sport fishing opportunities through rehabilitation of natural stocks or creation of new fisheries where these opportunities do not negatively affect other fisheries.

Management plans prepared for specific NWMA fisheries also identify a series of fishery objectives. While in some cases the objectives are different, objectives that recur frequently in the plans include:

1. management of sport fisheries so that harvests do not jeopardize sustained yield of the harvested stocks;
2. management of fisheries to maintain stock abundance and size composition that has been previously documented in given waters;
3. maintenance, and/or improvement of public access to fishing opportunities; and,
4. promote awareness of sport fishing opportunities that exist.

Existing management plans are under review, and a region wide Arctic grayling management plan is being developed.

MAJOR ISSUES FOR THE NORTHWESTERN MANAGEMENT AREA

1. Nome subdistrict salmon. Chum salmon stocks in the Nome subdistrict have been depressed since the mid 1980s. Fisheries divisions of the department have been cooperating in the recovery of these stocks. Efforts, including egg incubation boxes, fishery closures and increased escapement monitoring are ongoing. The Bureau of Land Management is operating a weir to count sockeye as they migrate into Glacial Lake. In addition, a lake fertilization project on Salmon Lake to increase zooplankton production for rearing sockeye has also been undertaken. There is a local initiative to develop a salmon incubation facility located near Nome that would assist in restoring depressed salmon runs. Sport Fish Division has participated in these efforts through the Regional Planning Team (RPT), and through cooperation with other divisions. Chum salmon sport fisheries remain closed.
2. Wulik River Dolly Varden. Development of a world-class zinc deposit at the Red Dog site in the upper Wulik River drainage carries the risk of heavy metal contamination on one of the most important streams in Northwest Alaska for Dolly Varden. There has been concern that heavy metal contamination of Red Dog and Ikalukrok creeks would occur both from natural leaching of the ore body as it was stripped for ore production and from discharge of contaminated waters into the river. A contamination problem in 1989 and 1990 has been controlled with additional waste-water treatment and the construction of a clean water bypass system in Red Dog Creek. Water quality is monitored by the Division of Habitat and mine personnel. Contamination along the road corridor has recently been documented by

- the NPS. The Division of Sport Fish counts Dolly Varden overwintering in the Wulik River annually and collects fish from which tissues are excised for heavy metal analyses.
3. Nome area gold mining. The future development of large-scale lode deposits of gold near Nome has the potential to degrade fish habitat in the Snake, Cripple and Solomon river drainages. Interest in mining is directly related to the world price of gold. In the last few years, development interest has declined with the price of gold, but that is likely to change with the recent increase in gold prices.
 4. Rural resentment of sport fishing and sport anglers. Rural Alaskans often feel resentment toward "outsiders" who come into remote areas traditionally used by local people for subsistence hunting or fishing. They sometimes have a cultural bias against the concept of "sport fishing" and feel that people do not have the right to "play" with food resources. The bias can be particularly strong against catch-and-release practices. This has led to resentment of sport anglers who wish to fish in remote waters of NWMA, and to proposals before the BOF that would eliminate catch and release in some fisheries.
 5. Effects of federal subsistence fisheries management on sport fishing opportunity in the NWMA. During October 1999, the federal government through the USFWS Office of Subsistence Management asserted its authority to manage subsistence fisheries on waters within or adjacent to Federal Conservation units. There is concern that a result of this action will be reduced opportunity for sport fishing throughout much of Alaska. Since there is a large amount of Federal Public land within the NWMA that is used by local residents for subsistence purposes, the potential loss of opportunity in remote areas of the NWMA is of acute concern to anglers and sport fish managers. The ADF&G continues to work with Federal managers and Rural Advisory Councils to address fisheries issues as they arise.

ACCESS PROGRAM

The Sport Fish Access Program was initiated nation-wide in 1984 as a result of the Wallop-Breaux Amendment to the Sport Fish Restoration (Dingell-Johnson or D-J) Act. The Sport Fish Access program is comprised of two parts. The first involves major capital improvement projects, such as boat launches, parking areas, camping areas handicap-accessible public fishing docks, access roads and trails, and the purchase or lease of lands or right-of-ways to ensure public access to fishing sites. The second portion of the program is called the Small Access Site Maintenance Project. This annually funded program involves maintaining and upgrading existing angler access sites. Activities include placing and maintaining signs at lake and river access sites, constructing and maintaining trails, and securing public rights-of way to fishing sites. Portable toilets, picnic tables and trash removal are provided at heavily used roadside sites. At remote sites, this project provides tent platforms and outhouses; it also publishes brochures on fishing and boating opportunities.

To date, few access projects have been proposed for the rural areas of the NWMA, however, a boat launching facility was recently built in the village of Unalakleet using Access funding.

SECTION II: SPORT FISH SEASON SUMMARY FOR 2002

PINK SALMON

Because the fishing season in the Nome subdistrict began with a subsistence salmon closure, sport fisheries for all salmon in the Nome subdistrict were closed by EO no. 3-S-01-02 (Appendix C). Escapements of pink salmon in the Nome subdistrict during 2002 were weak, and sport fishing remained closed for this species until July 16 when sport fisheries were opened in the Nome, Sinuk, Cripple and Penny rivers and in all salt waters west of Cape Nome. Farther east in Norton Sound, the runs were stronger with 636,407 pink salmon counted past the Niukluk River tower and 115,675 counted past the Eldorado River tower. The Unalakleet River had a relatively strong run with 321,756 counted past the tower on the North River. Sport fishing in these areas remained open, with no in-season adjustments to the fisheries.

CHUM SALMON

Chum salmon runs remained depressed throughout the Nome subdistrict in 2002 and only limited fishing was allowed. Due in large part to fishery restrictions, escapement goals were met on most rivers. Sport fisheries remained closed by regulation, no commercial fishing occurred, and subsistence fisheries were regulated under Tier II management. The Nome River with 1,720 chum salmon counted past the weir, failed to make its escapement goal of 2,900 – 4,300 and the Snake River with 2,669 counted past the tower exceeded its goal of 1,600-2,500. The Eldorado River also made its escapement goal range of 6,200 – 9,200 with 10,260 chum salmon counted past the tower. The escapement in the Niukluk River counted past the tower was 33,979 chum salmon. An escapement goal has not yet been set for this river. The North River tower (Unalakleet River) obtained a count of 5,918 chum salmon. The Kwiniuk River with 37,864 chum salmon counted past the tower by August 15, 2002, exceeded its Optimal Escapement Goal of 11,500 – 23,000. Only 600 chum salmon were harvested commercially in Norton Sound in 2002, the lowest on record. Sport fisheries outside the Nome subdistrict remained open without management action in 2002.

CHINOOK SALMON

The chinook salmon fishery in Norton Sound is functionally confined to the Unalakleet River with little effort or harvest elsewhere. By August 5, 2002, 1,484 chinook were counted past the tower on the North River, within the escapement goal range of 1,200-2,400. In 2002, the tower operated without problems. The average count past the tower for all years (1972-1974, 1984-1986, and 1996-1999) has been 2,622 chinook. Data collected during 1997 and 1998 in a radio tagging study indicated that about 40% of the Unalakleet River chinook run migrated past the counting tower on the North River. Therefore 3,710 chinook is a minimum estimate of escapement in the Unalakleet River during 2002. The commercial harvest of chinook in the Unalakleet subdistrict was only five fish in 2002. The sport fishery proceeded without management actions.

COHO SALMON

Coho returns to Nome subdistrict streams were low in 2002. In the Nome River, only 21 were counted past the weir by August 14, and other Northern Norton Sound streams were also recording low escapements. An EO on August 16, (no. 3-SS-01-02), closed sport fishing for coho from Rocky Point to Cape Prince of Wales and in the Pilgrim and Kuzitrin rivers. It also reduced the daily bag limit for coho in eastern Norton Sound from Rocky Point to the Unalakleet

River to one fish. By August 28, 1,561 coho had moved upstream on the Nome River while other rivers continued with lower than average escapements. EO no. 3-SS-02-02, effective August 31, opened the Nome River to sport fishing for coho with a daily bag limit of one fish. To the east, 7,269 coho were counted past the tower on the Niukluk River through September 11 and the daily bag limit remained at one fish through the end of the season. On the Unalakleet River, only 3,020 coho were counted past the tower through August 29, and the daily bag and possession limit remained at one fish throughout the remainder of the 2002 season. This was the second year that the tower on the Kwiniuk River was operated through the coho run; 9,532 coho were counted in 2001, and 6,459 were counted in 2002. A regulation was enacted by the BOF in December 2000 that reduced the “other salmon” daily bag limit in the Unalakleet River from 10 to 5 salmon. A research project was conducted on the Unalakleet River in 2001 to assess whether catch and release mortality of coho varied as fish moved from brackish water to upriver areas. Results of this study found that the mortality of coho caught and released did not change from the mouth of the South River upstream in the Unalakleet River drainage.

ARCTIC GRAYLING

Arctic grayling fisheries in the Northwest Management Area in 2002 progressed without the need for in-season management actions. A stock assessment project was conducted to estimate abundance and size composition of Arctic grayling in an index section of the Snake River during 2001. Abundance within the index area was estimated at about 1,090 fish, within the range of past estimates (565 to 1,235) and composition estimates were similar to what was found during past years. These data indicate that the management objectives of maintaining abundance and size composition for that fishery are being met under the current sport fishing regulatory framework. In 2002, a stock assessment project was conducted in an index area of the Pilgrim River. Data analysis has not yet been completed, but preliminary estimates suggest that the Arctic grayling population in this area is similar in abundance and size composition to that found in 1996. In December 2000, the BOF closed the Nome and Solomon rivers to subsistence fishing for Arctic grayling. In 2002, a restoration project was begun for Arctic grayling in the Nome River. Approximately 4,800 Arctic grayling fry were captured from the Nome River and held in pens in the Banner Creek gravel pit at different densities. Fish in fed and unfed treatment groups were reared through the summer. In September, 698 were released into the Nome River and approximately 1,300 were left in the pond for the winter. During the spring of 2003, surviving fish will be removed from the pond and reintroduced into the Nome River. Survival estimates by treatment group will be estimated at that time and the most successful treatment will be selected for use in 2003.

Kotzebue Sound/Chukchi Sea sub-area Arctic grayling fisheries are geographically dispersed over a wide area. Harvests are typically low, and although grayling abundance has not been estimated anywhere in this region, the dispersed nature of the fishery suggests that no single stock is being heavily exploited.

SHEEFISH

Sheefish fisheries in northwestern Alaska proceeded without the need for in-season management actions in 2002. Spawner abundance was estimated in the Kobuk River during 1996-1998 and ranged from about 32,000 to 43,000. The sport fish sheefish harvest has averaged only a few hundred fish annually and is not considered to have a significant influence on the population. The most recent sport fish harvest estimate for Kobuk River sheefish was 552 fish in 2001.

However, there are social problems associated with this fishery. Upper Kobuk River residents have concerns about increasing use of the upper Kobuk River by sport fishers and other visitors. In addition, catch-and-release fishing is considered disrespectful by many local residents.

DOLLY VARDEN

Dolly Varden are harvested in fisheries throughout the Northwestern Management Area in both the Kotzebue/Chukchi Sea sub-area and the Seward Peninsula/Norton Sound sub-area. Dolly Varden populations in the Kotzebue/Chukchi Sea sub-area are monitored with aerial counts of spawners in selected spawning streams, and counts of overwintering aggregations in the Wulik River. Inclement weather allowed spawning counts on only the Kelly and part of the Kugururok rivers in 2002, where 1,620 and 2,045 were counted respectively. These counts are about half the value of past counts on these rivers. On October 6, 2001 about 92,000 Dolly Varden were counted in the Wulik River. On October 10, 2002 the overwintering count of Dolly Varden was 44,257 fish. Residents of Kivalina did not catch many outmigrating fish this spring because the fish moved to sea very rapidly. However, fall seining on the Wulik River took place in late September aided by unusually high water that allowed easy access to upstream wintering areas. Reports suggest that harvests were higher than in recent years. A new angling record was set in 2002 with a 27 pound 4 ounce Dolly Varden taken from the Wulik River in October. There are no stock concerns regarding Dolly Varden in the sub-area. A project to describe genetic relationships among Dolly Varden stocks north and south of the Bering Strait was begun in 2000, and analysis should be completed in 2003. Another project to assess the spawning stock in Kagvik Creek was begun in 2001. This project was expanded with the construction of a weir in 2002. Inclement weather and high water caused even the weir to fail in 2002, and a decision to terminate the project was made in the fall. These projects were funded by the USFWS Office of Subsistence Management, and are continuing. Incidental commercial harvest in the Kotzebue chum salmon fishery has been negligible for the past three years.

COMMERCIAL AND SUBSISTENCE FISHERIES

The 2002 commercial salmon fishery in Norton Sound was by far the poorest on record for all species. The only commercial harvests occurred in the Unalakleet and Shaktoolik subdistricts. The total harvest was 5 chinook, 1 sockeye, 600 chum, and 1,760 coho. Only 12 permit holders fished. The previous low record salmon harvest for Norton Sound was 23,051 salmon in the 1999 commercial season. No commercial harvest of pink salmon occurred in 2002 because of a lack of market even though there was an abundance of this species in eastern Norton Sound. Subsistence fisheries in eastern Norton Sound were allowed to proceed in what has become a normal manner. All Tier I subsistence salmon fisheries in the Nome subdistrict were closed on June 15. In 1999 the BOF designated the Nome subdistrict as a Tier II chum salmon management area. Permits were issued to 30 qualified fishers for the 2002 season. Regular weekly Tier II fishing periods began on June 25 in marine waters between Cape Nome and Topkok Head. Each Tier II permit holder was allowed up to 100 chum salmon. On July 1, Tier II fisheries were allowed in waters west of the Safety Sound bridge, and in the Eldorado and Flambeau rivers. On July 2, Tier I beach seining and hook and line subsistence fisheries for pink salmon were opened in all Safety Sound drainages with the condition that all chum salmon be released. Additional openings were announced on a stream-by-stream basis as data on escapement became available.

SECTION III: SPORT FISHING EFFORT IN THE NORTHWESTERN MANAGEMENT AREA

SPORT ANGLING EFFORT

Recreational angler effort has been estimated for the Northwest Management Area with a mail survey since 1977 (Mills 1979-1994; Howe et al. 1995, 1996, 2000a,b,c,d; Walker et al. *In prep*; Jennings et al. *In prep*). The results of this survey indicate that effort in the Northwest Management Area has remained more or less stable since 1982, with a slight decline in recent years. Effort over this period has ranged from 20,000 to 30,000 angler days during most years (Table 7; Figure 11). During 2001, the total sport fishing effort for the Northwest Management Area was estimated at 16,859 angler days. The fraction of the entire AYK Region (excluding the Upper Copper/ Upper Susitna Area) effort expended in the Northwest Management Area has declined from over 15% in 1992 to approximately 10% between 1995 and 1999. In 2001, effort was about 11.5% (Table 7) after showing an increase to 14% in 2000. When the Upper Copper/Upper Susitna Area is included in the AYK total, the fraction of effort from the NWMA is about 5%.

The Seward Peninsula and Norton Sound sub-area accounts for most of the sport fishing in the NWMA. Effort in the sub-area has averaged around 18,500 angler days over the past 10 years showing a decline to 13,934 angler days from 1996-1998. The effort increased to about 15,000 angler days in 1999. This trend continued in 2000 with 18,559 angler days of effort, but dropped by over 40% to 11,000 angler days in 2001. Rivers supporting the most sport fishing effort in the NWMA have been the Nome, Unalakleet and Fish/Niukluk rivers. The Nome River has sustained more fishing effort than any other stream in the NWMA for four of the past 10 years. In 1995, the Unalakleet River surpassed the Nome River as the most popular single stream in the management area with 3,832 estimated angler days. This pattern has continued, but estimated effort on the Unalakleet River dropped from 6,257 angler days in 2000 to 2,793 in 2001 (Table 8). The Fish/Niukluk river system surpassed the Unalakleet River in angler effort in 1999 and 2001 with 4,916 and 2,822 angler days estimated respectively. The Nome River has been closed to fishing for Arctic grayling and chum salmon, and it is likely that these closures have contributed to a reduction of fishing effort on this stream. Annual effort in the Nome River has remained around 1,450 angler days for the past three years but dropped to 1,050 days in 2001. The Cripple River was the second most heavily utilized stream in 1998, with an estimated 2,303 angler-days. This was a substantial increase over previous estimates for this stream, 206 and 166 angler days in 1996 and 1997 respectively. Effort in the Cripple River was estimated at 205 angler days in 2001. Most of the sport fishing effort on the Cripple River is generated from a commercially run recreational mining camp that is located at the river's mouth.

In the Kotzebue/Chukchi Sea sub-area, sport fishing effort has been more variable, ranging from 3,700 to 10,200 angler days per year over the past 10 years. After showing a decline to about 3,800 angler days in 1997 and 1998 effort nearly doubled to 6,771 angler days in 1999. In 2000, effort dropped to an estimated 5,225 angler days and in 2001, effort was estimated at 5,900 angler days (Table 8). The large drainages of the Kobuk and Noatak rivers support about half of the freshwater effort in this sub-area during most years while the remainder is dispersed among smaller drainages such as the Wulik, Kivalina and Selawik rivers, and many of the area's lakes. Expense of travel, difficulty of access and small human population likely account for the low levels of sport fishing effort reported in this region.

Table 7.-Sport fishing effort in the A-Y-K region by management sub-areas, 1982-2001.

Year	Arctic-Yukon-Kuskokwim Region ^a												
	Tanana Area		AYK Area						Northwest Alaska				AYK Angler-Days
	Tanana Angler-Days	% AYK	North Slope Angler-Days	% AYK	Yukon Angler-Days	% AYK	Kuskokwim Angler-Days	% AYK	Seward Pen Angler-Days	% AYK	Kotzebue Angler-Days	% AYK	
1982	150,530	75.7	4,879	2.5	11,034	5.6	12,244	6.2	13,198	6.6	6,906	3.5	198,791
1983	144,981	72.8	5,738	2.9	11,070	5.6	12,429	6.2	16,944	8.5	7,963	4.0	199,125
1984	145,142	72.9	8,344	4.2	6,358	3.2	13,970	7.0	17,436	8.8	7,791	3.9	199,041
1985	135,745	72.6	4,490	2.4	8,670	4.6	11,358	6.1	19,919	10.7	6,701	3.6	186,883
1986	144,814	74.4	4,779	2.5	9,381	4.8	11,319	5.8	18,107	9.3	6,313	3.2	194,713
1987	155,346	71.6	5,256	2.4	7,017	3.2	17,856	8.2	21,413	9.9	10,221	4.7	217,109
1988	173,706	74.4	2,541	1.1	8,261	3.5	23,494	10.1	20,278	8.7	5,279	2.3	233,559
1989	185,715	77.5	4,118	1.7	10,712	4.5	16,457	6.9	17,692	7.4	4,932	2.1	239,626
1990	184,887	75.3	3,764	1.5	15,539	6.3	15,858	6.5	21,799	8.9	3,782	1.5	245,629
1991	155,662	70.8	7,291	3.3	10,749	4.9	13,055	5.9	23,622	10.7	9,543	4.3	219,922
1992	120,848	66.5	4,940	2.7	12,831	7.1	14,404	7.9	22,684	12.5	6,145	3.4	181,852
1993	160,117	72.5	5,600	2.5	14,011	6.3	14,505	6.6	18,930	8.6	7,809	3.5	220,972
1994	148,633	70.8	5,407	2.6	12,872	6.1	18,117	8.6	18,922	9.0	6,036	2.9	209,987
1995	201,389	74.5	5,644	2.1	18,677	6.9	16,289	6.0	19,647	7.3	8,495	3.1	270,141
1996	150,227	54.7	4,487	1.6	10,678	3.9	16,420	6.0	13,783	5.0	5,571	2.0	274,566
1997	119,699	65.6	5,278	2.9	12,725	7.0	27,318	15.0	13,850	7.6	3,729	2.0	182,599
1998	112,025	65.5	3,653	2.1	10,127	5.9	27,913	16.3	13,616	8.0	3,801	2.2	171,135
1999	160,427	70.7	5,230	2.3	12,906	5.7	26,563	11.7	15,006	6.6	6,771	3.0	226,903
2000	121,785	66.3	4,739	2.6	11,327	6.2	20,030	10.9	18,559	10.1	7,129	3.9	183,569
2001	91,226	62.6	6,061	4.2	10,531	7.2	21,087	14.5	10,954	7.5	5,900	4.0	145,759
91-00 Avg.	145,081	68	5,227	2	12,690	6	19,461	9	17,862	9	6,503	3	214,165
96-00 Avg.	132,833	65	4,677	2	11,553	6	23,649	12	14,963	7	5,400	3	207,754

a AYK total does not include Copper River drainage.




Figure 11.-Sport fishing effort in angler-days in the NWMA by sub-area, 1978-2000.

Table 8.-Freshwater sport fishing effort in angler-days for major rivers by sub-area in the Northwest Management Area, 1983-2001.

Year	Seward Peninsula/Norton Sound Sub-Area							Kotzebue/Chukchi Sea Sub-Area					MWMA
	Nome	Fish/Niukluk	Unalakleet	Snake	Pilgrim	Others	Total	Noatak	Kobuk	Wulik	Other	Total	Total
1983	3,908	1,999	4,146	119	597	6,175	16,944	1,372	3,609	805	2,177	7,963	24,907
1984	5,714	1,115	2,421	418	732	7,036	17,436	1,805	2,834	677	2,475	7,791	25,227
1985	6,514	889	5,750	361	375	6,030	19,919	3,470	2,297	550	384	6,701	26,620
1986	6,023	1,888	6,457	850	868	2,021	18,107	2,141	2,646	168	1,358	6,313	24,420
1987	1,865	2,473	942		1,159	14,974	21,413	3,584	5,133	303	1,201	10,221	31,634
1988	6,058	2,245	1,219	2,128	4,822	3,806	20,278	2,492	2,100	455	232	5,279	25,557
1989	6,569	2,124	1,701	436	1,678	5,184	17,692	2,552	1,729	107	544	4,932	22,624
1990	7,194	2,059	3,957	775	1,710	6,104	21,799	1,423	1,306	285	768	3,782	25,581
1991	4,646	2,524	5,616	2,384	3,183	5,269	23,622	4,235	2,353	93	2,862	9,543	33,165
1992	6,455	2,742	2,433	2,379	1,184	7,491	22,684	2,611	2,095	469	970	6,145	28,829
1993	3,633	3,962	2,153	1,468	1,195	6,519	18,930	3,013	2,604	350	1,842	7,809	26,739
1994	5,116	3,082	2,349	880	844	6,651	18,922	2,747	1,153	762	1,374	6,036	24,958
1995	3,044	2,603	3,832	1,968	1,253	6,947	19,647	2,504	3,681	647	1,663	8,495	28,142
1996	2,920	2,120	2,539	1,269	840	4,095	13,783	2,039	1,358	274	1,900	5,571	19,354
1997	1,914	3,017	4,393	445	820	3,261	13,850	1,159	825	553	1,192	3,729	17,579
1998	1,371	1,344	3,795	376	546	6,184	13,616	765	2,053	202	781	3,801	17,417
1999	1,463	4,916	4,176	977	433	3,041	15,006	3,142	2,099	737	793	6,771	21,777
2000	1,455	3,451	6,257	377	753	3,385	15,678	1,713	2,298	336	878	5,225	20,903
2001	1,045	2,822	2,793	853	491	1,899	9,903	2,702	925	592	1,275	5,494	15,397
91-00 Avg.	3,202	2,976	3,754	1,252	1,105	5,284	17,574	2,393	2,052	442	1,426	6,313	23,886
96-00 Avg.	1,825	2,970	4,232	689	678	3,993	14,387	1,764	1,727	420	1,109	5,019	19,406

SECTION IV: MAJOR NORTHWESTERN AREA FISHERIES OVERVIEW

NWMA waters offer some of the most remote and diverse angling opportunities available in Alaska. Opportunities to fish for Dolly Varden, sheefish and Arctic grayling in pristine and uncrowded areas are widespread. Angling opportunities for salmon, especially chinook and coho are less common, but can be found in several Norton Sound streams that produce good runs. Marine sport fisheries are practically non-existent. Guided fishing comprises a small amount of the effort in northwestern Alaska. An unpublished survey conducted by the Division of Sport Fish for the 1994 season estimated that only about 1% of the total sport fishing effort in the Seward Peninsula-Norton Sound sub-area was by guided anglers while about 5% of the effort in the Kotzebue-Chukchi Sea sub-area was guided. Through-the-ice jigging for saffron cod, smelt, flounder, sheefish, and other species is common near settlements, but these fisheries generally operate under subsistence fishing regulations. The following sections discuss the major sport fisheries in the NWMA by species and area. Discussion of each fishery will deal with 1) historical perspective, 2) recent fishery performance 3) fishery objectives, 4) fishery outlook, 5) in-season management and recent BOF actions, 6) current issues and 7) recommended management and research programs. Recent fishery performance will focus on data from 2001. Observations regarding the 2002 season may be included for some fisheries, but data on harvest are not yet available. Summaries of historic sport fish harvests and catches are provided by fishery.

NORTHWESTERN ALASKA SALMON FISHERIES

Guided and unguided sport fishing for salmon takes place throughout the management area, however the vast majority of salmon fishing occurs in the Seward Peninsula/Norton Sound sub-area with concentrations near Unalakleet, and in waters accessible from the Nome area road system. Some salmon fishing effort occurs in association with wilderness float trips in Kotzebue Sound drainages, but the amount of sport fishing effort expended toward salmon in the northern part of the management area is light, and consequently, harvests are small.

Total fishing effort estimated for all species of fish in the NWMA has ranged from about 11,000 angler-days in the late 1970s to 33,000 angler-days in 1991 (Table 8). Effort has declined in recent years, and was estimated at 17,000 angler days in 2001. Salmon harvest is estimated to have ranged from 3,800 fish in 1977 to 20,000 fish in 1982 (Table 9). The estimated salmon harvest in 2001 was 8,600 fish. Mean annual harvest of salmon of all species from 1991 to 2000 was about 11,000 fish, with 97% of the harvest reported from Seward Peninsula and Norton Sound, and only about 3.0% from Kotzebue drainages. In 2001, the harvest was from Kotzebue drainages increased to 20% of the NWMA total. Over the past five years (1996-2000), about 50% of the total average harvest has been coho salmon, 35% pink salmon, 9% chum salmon, and 7% chinook salmon. During years of high pink salmon abundance such as 1992, 1994, 1996, and 1998, harvests of this species have comprised about 50% of the total annual salmon harvest. However, during years of low pink salmon abundance such as 1993, 1995, 1997 and 1999 coho salmon have accounted for about 60% of the total salmon harvest (Table 9).

Regulatory History

Prior to 1966, the daily bag limit was 15 fish (of all species) in freshwaters of western Alaska. From 1966 through 1970 an exception was made in the Unalakleet River where the daily bag

Table 9.-Northwest Management Area historic salmon harvests by sub-area, 1977-2001.

Year	Norton Sound/ Seward			NW Management			Norton Sound/ Seward			NW Management		
	Peninsula	Kotzebue	Area	Peninsula	Kotzebue	Area	Peninsula	Kotzebue	Area	Peninsula	Kotzebue	Area
	Effort	Effort	Effort	KS	KS	KS	Coho	Coho	Coho	Chum	Chum	Chum
1977	7,828	3,487	11,315	197	16	213	449	0	449	670	28	698
1978	8,379	4,997	13,376	303	0	303	742	0	742	546	254	800
1979	8,725	2,593	11,318		10			0			27	
1980	7,958	3,841	11,799	52	9	61	1,455	0	1,455	1,601	86	1,687
1981	10,879	5,284	16,163	70	22	92	1,504	0	1,504	1,889	32	1,921
1982	13,198	6,906	20,104	409	0	409	2,986	0	2,986	2,620	346	2,966
1983	16,944	7,963	24,907	687	0	687	3,823	0	3,823	2,042	463	2,505
1984	17,436	7,791	25,227	247	13	260	7,582	0	7,582	1,481	312	1,793
1985	19,919	6,701	26,620	239	0	239	1,177	51	1,228	1,036	310	1,346
1986	18,107	6,313	24,420	1,077	0	1,077	3,926	0	3,926	1,719	749	2,468
1987	20,413	9,288	29,701	615	95	710	2,319	11	2,330	814	402	1,216
1988	20,278	5,279	25,557	400	18	418	5,038	0	5,038	1,583	236	1,819
1989	17,692	4,932	22,624	203	0	203	4,158	0	4,158	1,497	41	1,538
1990	21,799	3,782	25,581	364	0	364	3,305	0	3,305	925	0	925
1991	23,622	9,543	33,165	404	0	404	5,800	0	5,800	1,415	59	1,474
1992	22,684	6,145	28,829	204	8	212	4,671	0	4,671	523	220	743
1993	18,930	7,809	26,739	595	0	595	3,783	9	3,792	691	443	1,134
1994	18,922	6,036	24,958	600	0	600	5,547	0	5,547	536	248	784
1995	19,647	8,495	28,142	438	0	438	3,705	0	3,705	394	321	715
1996	13,783	5,571	19,354	662	0	662	7,289	40	7,329	662	576	1,238
1997	13,850	3,729	17,579	1,106	0	1,106	4,393	0	4,393	278	272	550
1998	13,616	3,801	17,417	590	0	590	4,441	0	4,441	682	177	859
1999	15,006	6,771	21,777	630	20	650	5,582	0	5,582	211	392	603
2000	18,559	7,129	25,688	889	6	895	7,441	10	7,451	1,097	719	1,816
2001	10,955	5,904	16,859	271	0	271	4,802	29	4,831	1,709	1,671	3,380
Avg (91-00)	17,862	6,503	24,365	612	3	615	5,265	6	5,271	649	343	992
Avg (96-00)	14,963	5,400	20,363	775	5	689	5,082	8	5,090	445	348	793

Table 9.-Page 2 of 2.

Year	Norton Sound/ Seward Peninsula			Norton Sound/ Seward Peninsula			Norton Sound/ Seward Peninsula		
	Area	Area	Area	Area	Area	Area	Total	Total	Total
1977	2,402	8	2,410	0	0	0	3,718	52	3,770
1978	7,399	0	7,399	0	0	0	8,990	254	9,244
1979		0			0			37	
1980	7,732	0	7,732	0	0	0	10,840	95	10,935
1981	3,101	0	3,101	0	0	0	6,564	54	6,618
1982	13,742	0	13,742	0	0	0	19,757	346	20,103
1983	4,583	0	4,583	0	0	0	11,135	463	11,598
1984	8,322	0	8,322	351	0	351	17,983	325	18,308
1985	1,138	68	1,206	20	0	20	3,610	429	4,039
1986	3,172	62	3,234	19	0	19	9,913	811	10,724
1987	1,304	0	1,304	924	21	945	5,976	529	6,505
1988	2,912	0	2,912	782	0	782	10,715	254	10,969
1989	3,564	10	3,574	165	0	165	9,587	51	9,638
1990	7,647	0	7,647	198	0	198	12,439	0	12,439
1991	1,738	91	1,829	237	0	237	9,594	150	9,744
1992	6,403	293	6,696	131	0	131	11,932	521	12,453
1993	2,250	0	2,250	10	0	10	7,329	452	7,781
1994	7,051	51	7,102	18	0	18	13,752	299	14,051
1995	928	38	966	104	0	104	5,569	359	5,928
1996	5,972	10	5,982	100	0	100	14,685	626	15,311
1997	1,458	0	1,458	30	0	30	7,265	272	7,537
1998	6,939	13	6,952	16	0	16	12,668	190	12,858
1999	3,039	12	3,051	0	0	0	9,462	424	9,886
2000	2,886	0	2,886	45	0	45	12,358	735	13,093
2001	360	0	360	39	0	39	7,181	1,700	8,881
Avg (91-00)	3,866	51	3,917	69	0	69	10,461	403	10,864
Avg (96-00)	4,059	7	4,066	38	0	38	11,288	449	11,737

limit was 6 salmon (all species). In 1970, the 6 salmon limit on the Unalakleet River was replaced with a daily 15 salmon limit for the entire AYK Region with exceptions outside the NWMA. Salmon Lake and its tributaries were closed to salmon fishing in 1980. The general 15 salmon daily bag limit in the NWMA remained in effect until 1985 when the king salmon limit was set at 5 per day and the “other salmon” limit was set at 10 per day. An exception was made for the Snake and Nome rivers where the “other salmon” limit was 15 per day of which only 5 could be chum or coho. In 1987 emergency regulations were adopted that set the king salmon daily bag limit at 1 fish and the “other salmon” limit at 10 per day. In 1988, the king salmon daily bag limit for the AYK Region outside the Tanana River drainage was set at 3 per day with only 2 over 28 inches and the “other salmon” limit remained at 10 per day, however, in Seward Peninsula waters, the king salmon limit was 1 per day with 10 “other salmon” of which only 3 could be chum or coho. Also in 1988, chum salmon fishing was closed by EO on the Nome River. This closure was extended to all rivers from the Sinuk to the Solomon in 1991. In 1992, this closure was adopted into regulation by the BOF. These regulations remained in effect until 1997 when the BOF broke out the “other salmon” bag limits for northern Norton Sound and adopted daily bag limits of 10 pink, 3 coho, 3 sockeye, and 3 chum salmon. The chum salmon closure in the Nome subdistrict remains in effect. In 2000, the “other salmon” daily bag and possession limit was reduced from 10 salmon to 5 salmon in the Unalakleet River. In addition, a regulation to address the possible high catch and release mortality of coho in the estuary of the Unalakleet River was adopted. This regulation closed the area downstream from the South River for the remainder of that day, to all sport fishing for those anglers that had harvested a daily bag limit of coho salmon.

UNALAKLEET RIVER SALMON FISHERIES

Fishery Description and Historical Perspective

The Unalakleet River has historically supported substantial runs of chinook, chum, coho and pink salmon. Guided and unguided fishing effort is primarily focused on chinook and coho salmon, but chum and pink salmon are also harvested. The City of Unalakleet with a population of about 800 is located on the shore of Norton Sound at the mouth of the river. Daily air service from Anchorage and Nome provides access for nonlocal anglers visiting the Unalakleet area. The U. S. Air Force operated a sport fishing recreational camp on the Unalakleet River during the 1960s, and a commercial sport fishing lodge was constructed there in the late 1960s. This fishing lodge is still being operated, although it has expanded in size and has had several owners. The Unalakleet Native Corporation owned the lodge for several years and contracted operations. The lodge is currently in private ownership and is located about 8 miles upstream from the mouth of the river. Several local residents also guide anglers on the river, and guiding operations from the Yukon River drainage sometimes visit the river during the peak of the chinook and coho salmon runs. The majority of angling, however, is by unguided anglers. An unpublished survey by the Division of Sport Fish estimated that only about 8.5% of salmon anglers on the Unalakleet River were guided. The average annual sport harvest of salmon of all species from the Unalakleet River from 1990 to 1999 has been about 3,682 fish. Coho comprised about 61% of the average harvest while chinook made up about 10% (Mills 1991-1994; Howe et al. 1995, 1996, 2000a-d; Walker et al. *In prep*). The harvest increased to about 4,800 salmon in 1997 and 6,263 salmon in 1999, declining to 5,859 in 2000 and 3,900 in 2001 (Table 10). Coho are estimated to have comprised 43% of the 1999 harvest, 71% of the 2000 harvest, and 64% of the 2001 salmon harvest.

Table 10.-Sport fish effort, harvest, and catch estimates by species for the Unalakleet River, 1990-2001.

	Year									
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	<u>Effort</u>									
Number of Anglers	482	917	685	625	777	1,009	695	941	835	995
Number of Trips	1,642	3,236	2,256	1,398	2,192	3,376	1,886	3,366	3,197	1,726
No. Angler Days	3,974	5,616	2,433	2,153	2,349	3,832	2,539	4,393	3,795	4,176
	<u>Harvest</u>									
Chinook Salmon	276	296	117	382	379	259	384	842	513	415
Coho Salmon	1,826	2,180	1,555	643	2,425	2,033	3,411	2,784	2,742	2,691
Chum Salmon	298	497	379	116	220	207	463	228	447	211
Pink Salmon	1,180	437	779	89	402	222	59	1,055	434	2,946
Dolly Varden	614	1,648	746	602	679	1,061	1,506	936	588	2,384
Arctic Grayling	99	1,909	114	131	353	300	420	210	144	277
Total Fish Harvest	4,293	6,967	3,690	1,963	4,458	4,082	6,243	6,055	4,868	8,924
	<u>Catch</u>									
Chinook Salmon	361	375	476	2,340	517	588	2,059	5,144	1,539	669
Coho Salmon	3,396	2,882	2,802	1,572	2,488	3,086	5,863	4,020	3,213	9,593
Chum Salmon	379	692	1,412	515	561	966	1,589	1,323	2,218	1,916
Pink Salmon	1,513	559	6,503	605	1,020	799	2,594	4,101	4,853	3,475
Dolly Varden	2,222	2,267	1,942	964	1,253	2,732	3,170	4,400	2,336	10,460
Arctic Grayling	448	2,813	1,022	874	1,476	1,332	1,694	4,918	3,256	6,089
Total Fish Catch	8,319	9,588	14,157	6,870	7,315	9,503	16,969	23,906	17,415	32,202

-continued-

Table 10.-Page 2 of 2.

	Year			
	2000	2001	Average (91-00)	Average (96-00)
	<u>Effort</u>			
Number of Anglers	1,002	918	854	894
Number of Trips	3,415	1,365	2,492	2,718
No. Angler Days	6,257	2,793	3,667	4,232
	<u>Harvest</u>			
Chinook Salmon	345	250	380	500
Coho Salmon	4,150	2,766	2,489	3,156
Chum Salmon	403	714	353	350
Pink Salmon	961	188	688	1,091
Dolly Varden	4,462	1,022	1,419	1,975
Arctic Grayling	538	247	422	318
Total Fish Harvest	10,859	5,167	5,752	7,390
	<u>Catch</u>			
Chinook Salmon	1,045	542	1,390	2,091
Coho Salmon	9,287	5,399	4,564	6,395
Chum Salmon	3,652	2,030	1,534	2,140
Pink Salmon	3,982	1,197	2,699	3,801
Dolly Varden	10,293	2,769	3,871	6,132
Arctic Grayling	6,814	2,331	2,965	4,554
Total Fish Catch	35,073	14,268	17,024	25,113

Recent Fishery Performance

During 1997, 1998, and 2000, the Unalakleet River sustained the highest sport fishing effort of any single river in the NWMA. In 1999 and 2001 it ranked second to the Fish/Niukluk river system. In 2000, estimated effort on the Unalakleet River increased by 50% over the previous year, to 6,257 angler days, but in 2001 effort dropped to 2,800 angler days (Table 8). An estimated total of 3,918 salmon were harvested from the Unalakleet River in 2001; 71% (2,766) were coho, and about 6% (249) were chinook, and 18% (714) were chum salmon (Jennings et al. *In prep*). Approximately 92% of the entire NWMA harvest of chinook salmon, and 61% of the coho harvest were taken from the Unalakleet River in 2001.

The chinook salmon run usually begins in mid-June, peaks during the last week of June and continues through mid-July. Anglers access the river by boat from the village of Unalakleet and are composed of a mix of local residents, visitors who rent boats or fish with friends, and visitors who either stay at the Unalakleet Lodge or are guided by local resident guides. Most sport fishing effort occurs in the lower 15 miles of the Unalakleet River and in the lower 5 miles of the North River which enters the Unalakleet about 7 miles upstream from its mouth. The estimated harvest of 842 chinook, from the very strong 1997 run, was the second highest on record. During 1998, the estimated harvest was 513 chinook, dropping to 415 in 1999, 345 in 2000, and 250 in 2001 (Table 10). The total escapements of chinook into the Unalakleet River were estimated for the first time in 1997 and 1998 by expanding the tower estimate from the North River to include the entire drainage based on proportions of radio tagged fish moving up each drainage. In 1997 and 1998, chinook escapement into the Unalakleet River was estimated at 11,204 and 5,220 respectively. A similar expansion would give an estimate of 5,673 chinook for 1999, 2,615 for 2000, and 3,343 for 2001.

Over the past five years (1996-2000) the commercial harvests of chinook salmon in the Unalakleet subdistrict has averaged 5,734 fish (Table 1). Over the course of the fishery since 1961, estimated harvests have ranged from 960 in 1969 to 12,621 in 1985. In 1998, 6,413 chinook were harvested in this fishery, in 1999, 1,927 were taken, in 2000, 582 were taken, and in 2001, only 213 were commercially harvested. Commercial fish managers believe that some of the Unalakleet harvest usually includes some fish bound for Yukon River that mill in Norton Sound before moving into the Yukon. Records of subsistence harvests of chinook salmon in Unalakleet have ranged from 90 fish in 1966 to 6,325 fish in 1997 (Table 5). The recent five-year average (1996-2000) harvest was 4,146 fish. The 2000 subsistence harvest was estimated at 2,429 fish, and the estimated 2001 harvest was 2,810 chinook. The sport fish harvest over the same 5-year period has averaged 380 fish or about 9% of the average Unalakleet subsistence harvest, and 6% of the average total Unalakleet harvest. In 2001, the sport harvest comprised about 9% of the subsistence harvest and also about 9% of the total Unalakleet harvest because of a very low commercial catch due to fishery restrictions.

Coho are the most sought after salmon species in the Unalakleet drainage. The run usually begins around August 1, peaks during mid-August and continues through early September. The sport harvest of coho, from 1996 to 2000, has averaged 2,489 fish, and has remained relatively stable since 1997, however there was a marked increase in harvest to 4,150 coho in 2000 (Table 10). The 1999 estimated harvest of coho in the Unalakleet River was 2,691 fish, and the estimated harvest in 2001 was 2,766 (Table 10). The coho fishery is more consumptive than other Unalakleet salmon fisheries. Approximately 55% of coho caught are harvested while about

27%, 23%, and 25% (1996-2000 average) of chinook, chum and pink salmon caught are harvested.

From 1996-2000 commercial harvests of coho salmon in the Unalakleet subdistrict have averaged 28,576 fish (Table 1). Since 1961, commercial harvests of coho have ranged from 79 in 1964 to 71,019 in 1994 (Table 1). The commercial harvest has trended downward over the past five years. In 1998, 24,534 were harvested, and in 1999, 10,264 were taken. This trend reversed in the 2000 season when 29,803 coho were harvested, but only 15,102 coho were harvested commercially in 2001. Subsistence harvests of coho salmon in the community of Unalakleet averaged 9,281 fish from 1996 to 2000. The 2001 subsistence harvest was estimated at 6,270 coho. The sport fish harvest over the same period has averaged 2,489 fish. In 2001, the sport harvest of coho was about 12% of the total Unalakleet coho harvest. Reliable escapement data for coho salmon in the Unalakleet drainage have not been available, however the tower on the North River has recently been providing data for that portion of the drainage.

Sport Fishery Management Objectives

There have been no specific management objectives identified for salmon fisheries on the Unalakleet River. In comparison to commercial and subsistence salmon harvests, sport harvests have been small. The goal of sport fishery management in the Unalakleet River is to ensure that escapement goals are met and to maintain opportunity for anglers to participate in the fisheries. Emergency actions to restrict harvest are generally not implemented unless other harvests, and escapement monitoring projects, indicate that a particular run is small and escapement goals may not be met, or if restrictions in subsistence fisheries may be necessary in order to allow for sufficient spawning escapement. A Sustainable Escapement Goal (SEG) based on passage past the North River tower has been developed for chinook (1,200-2,400 fish) and pink salmon (8,500), but will not be established for other species until more years of reliable tower count data have been accumulated. When BEGs have been developed, more precise management to attain those goals will become possible.

Current Issues

Although sport fishing has been ongoing in the Unalakleet River drainage for many years, there is some local resentment of visiting anglers because some Unalakleet residents feel that “outsiders” are competing for the local salmon resources. Declines in chum and coho salmon runs throughout western Alaska have impacted the Unalakleet River drainage, although the effect appears to be less dramatic than in Nome subdistrict streams where chum runs have a long history of being depressed. The daily bag limit of “other salmon” was reduced from 10 fish to 5 fish in 2001. While the commercial harvests of chinook in the Unalakleet Subdistrict have remained relatively stable through 1999 but have declined in the past three years. The commercial harvests of coho have declined during the past four years. The 1999 harvest dropped to 10,264 coho. The harvest in 2000 rebounded to nearly 30,000, but dropped again in 2001 to 15,102. In 2002, the commercial harvest of coho in the Unalakleet subdistrict was only 1,079 fish. The upper reaches of the Unalakleet River are a National Wild and Scenic River under federal management. With the recent federal takeover of subsistence management on federal waters, the possibility exists that subsistence management in the upper reaches of the river could affect sport fishing opportunity in the Unalakleet River.

Recent and Ongoing Research and Management Activities

Salmon escapements in the Unalakleet River are monitored using a counting tower in the North River, a test net in the Unalakleet River downstream from the mouth of the North River, and by aerial surveys. The tower is a cooperative project operated by the Unalakleet IRA with funding from the Kawarek Corporation and guidance by the ADF&G, CFMD. Aerial surveys are difficult in the Unalakleet River because of its dark bottom and tannic stained water. These surveys provide a minimum escapement, but are unreliable as a total escapement estimate in this river. Water in the North River is clear, and the tower provides a reliable estimate of escapement into that portion of the river system. A weir site has been located on the mainstem of the Unalakleet River, and this project may be started in the near future in order to estimate passage of salmon in the entire drainage.

A research project was conducted on the Unalakleet River in 1997 and 1998 to estimate the proportions of the chinook salmon escapement that spawn in the North River and the main Unalakleet River upstream from the North River. In 1997, 37.2% of the radio tagged chinook spawned in the North River, and 62.8% spawned in the remainder of the drainage (Wuttig 1998). Proportions estimated in 1998 were similar, 40.1% in the North River and 59.9% in the remainder of the Unalakleet drainage. These relative proportions can now be used to expand the North River tower estimate to allow estimation of the escapement in the entire system. The sport fish staff have frequently assisted and cooperated informally with the Commercial Fisheries Division on projects, including the partial funding of counting towers from which spawning escapements are estimated, surveys for abundance, and observation of spawning concentrations. EOs restricting the harvest of salmon are usually coordinated with the Commercial Fisheries Division. There have been no EOs that have affected Unalakleet River salmon sport fisheries during the reporting period. Because of the low run of coho salmon in eastern Norton Sound in 1999, an EO was issued that lowered the “other salmon” sport fishing bag limit from ten fish per day to three fish per day. The “other salmon” bag limit of 10 fish, while it may be appropriate for pink salmon during years of high abundance, was felt by some to be excessive, especially for coho. The BOF reduced the “other salmon” limit on the Unalakleet River to five per day at their December 2000 meeting. In 1999, about 72% of the coho salmon captured in the Unalakleet River were released, in 2000 about 55% were released, and in 2001 about 49% were released. A department study (Vincent-Lang et al. 1993) found high levels of catch and release mortality with coho captured in estuarine waters of the Little Susitina River. Local residents have expressed concern about mortality of fish captured and released in sport fisheries, and a proposal to eliminate the release of sport caught fish in the Unalakleet River was recently heard by the BOF. The result was a regulation that prohibits sport fishing in estuarine waters of the Unalakleet River downstream from the South River for the remainder of the day after an angler has retained a bag limit of coho. A project was undertaken in 2001 to document relative catch and release mortality of coho in the Unalakleet River from its lower reaches progressively into freshwater upstream areas. This study found that the catch and release mortality of radio-tagged coho salmon was about 15% and that mortality did not differ from the mouth of the South River upstream. It was therefore concluded that the boundary for restrictions downstream of the South River was appropriate, and may be conservative. Another regulation passed in December 2000 made it unlawful to completely remove a salmon from the water if that salmon was to be released.

NOME AREA ROADSIDE SALMON FISHERIES

Fishery Description and Historical Perspective

Nine rivers accessible from the road system near Nome sustain some level of sport fishing effort for salmon. Estimated harvests from these rivers have averaged about 7,800 salmon annually, of which coho and pink salmon have comprised about 85%. Through 1996, the Nome River had sustained more sport fishing effort than any other single water body in northwestern Alaska because of its proximity to Nome and the adjacent road. Sport fishing on the Nome River has accounted for an annual average of 18% of all the fishing effort in the entire northwestern management area since 1990 (Table 8). An average of about 2,000 salmon have been harvested annually from the Nome River over the past 10 years, approximately 79% have been pink salmon (Table 11).

The Fish/Niukluk rivers is also a popular sport fishing location for salmon. Two guiding operations with small lodges are located on the Niukluk River. In addition, Nome based guides fish these rivers as well as other road accessible waters. Many residents of Nome have summer cabins on the Niukluk River at Council or fish camps along the river. Since the construction of the bridge over Safety Sound in 1980, and improvements to the road, access to the Fish/Niukluk has increased, and this area has become a prime destination for the road bound angler. The drainage sustains an average annual effort of about 2,800 angler days. The Fish/Niukluk has been the most popular sport fishing stream in Norton Sound for three of the past 10 years. About 1,300 salmon are harvested annually from the Fish/Niukluk rivers, and with coho accounting for about 74% of the salmon harvest (Table 12). A counting tower on the Niukluk River provides escapement data for salmon that is used for in-season management decisions.

The Pilgrim River, with its headwaters at Salmon Lake is another popular salmon fishery. All five species of North American Pacific salmon occur in the Pilgrim River. Sockeye spawn in Salmon Lake and the runs appear to be responding positively to recently undertaken lake fertilization efforts. There is a Bureau of Land Management campground at the outlet of Salmon Lake, and from there the river can be floated for about 25 river miles to the bridge at mile 65 of the Kougarok Road. Riverboats can be launched at the bridge for access to downstream locations. Over the past five years, the Pilgrim River has sustained an average annual effort of about 700 angler days and about 125 salmon are harvested annually (Table 13). A counting tower was recently installed on the Pilgrim River, and data from that project will be used for in-season management decisions. The Fish/Niukluk and the Pilgrim rivers are rivers where fishing for chum salmon is still allowed, however annual (1996-2000 average) harvests have been very small (Tables 12 and 13).

The mouth of the Snake River is in downtown Nome, and this small stream can also be accessed from a bridge at about mile 8 of the Teller Road and from the Glacier Creek Road. The Snake River sustains an average annual effort of about 700 angler days, with an annual harvest of about 430 salmon, about 43% coho and 57% pink salmon (Table 14).

Other road accessible waters include the Solomon, Kuzitrin, Penny, Cripple and Sinuk rivers (Tables 15–19). The average annual harvests in these rivers combined for the past five years (1996-2000) have averaged 261 coho, 0 chum, and 70 pink salmon. During years of high pink salmon abundance this species has dominated catches and harvests in some Nome roadside streams.

Table 11.—Sport fish effort and harvests by species from the Nome River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	Salmon					All Salmon	Dolly Varden	Arctic	
			King	Coho	Red	Pink	Chum			Grayling	Whitefish
Harvest											
1983	0	3,908	93	204	0	1,782	538	2,617	2,468	464	0
1984	1,288	5,714	13	2,648	0	4,128	325	7,114	935	376	13
1985	1,220	6,514	20	209	0	349	189	767	1,236	528	0
1986	1,087	6,023	0	415	0	491	76	982	1,057	491	0
1987	674	1,865	0	163	0	235	0	398	906	344	0
1988	1,733	6,058	0	1,455	0	528	273	2,256	2,365	946	18
1989	1,231	6,569	19	1,233	0	1,573	495	3,320	3,551	2,032	131
1990	1,625	7,194	39	407	35	2,651	122	3,254	1,078	33	0
1991	1,277	4,646	22	417	0	356	241	1,036	1,220	186	13
1992	1,433	6,455	16	713	0	4,397	0	5,126	573	0	0
1993	1,181	3,633	93	602	0	723	0	1,418	917	0	0
1994	1,025	5,116	0	326	0	4,103	0	4,429	431	16	0
1995	859	3,044	0	143	0	230	0	373	462	0	0
1996	947	2,920	0	598	0	3,280	0	3,878	873	0	0
1997	691	1,914	10	295	0	83	0	388	328	0	0
1998	636	1,371	0	189	0	1,985	0	2,174	302	0	0
1999	564	1,463	0	219	0	0	0	219	791	0	0
2000	450	1,455	0	342	0	578	0	920	340	0	0
2001	312	1,045	0	297	0	0	0	297	43	0	0
Avg 91-00	906	3,202	14	384	0	1,574	24	1,996	624	20	1
Avg 96-00	658	1,825	2	329	0	1,185	0	1,516	527	0	0
Catches											
1990	1,625	7,194	48	896	35	5,483	825	7,287	2,271	613	0
1991	1,277	4,646	22	869	0	894	389	2,174	3,725	1,363	13
1992	1,433	6,455	23	1,466	0	9,810	266	11,565	1,130	90	9
1993	1,181	3,633	121	764	0	1,756	175	2,816	5,153	569	0
1994	1,025	5,116	0	386	0	6,190	36	6,612	631	1,111	0
1995	859	3,044	0	228	0	980	478	1,686	1,474	571	0
1996	947	2,920	21	788	0	5,898	432	7,139	1,311	497	0
1997	691	1,914	20	447	0	190	113	770	873	569	0
1998	636	1,371	19	863	0	3,482	8	4,372	319	207	0
1999	564	1,463	0	231	0	13	0	244	1,486	300	0
2000	450	1,455	0	385	0	876	20	1,281	431	10	0
2001	312	1,045	0	377	0	32	13	422	94	60	0
Avg 91-00	906	3,202	23	643	0	3,009	192	3,866	1,653	529	2
Avg 96-00	658	1,825	12	543	0	2,092	115	2,761	884	317	0

Table 12.—Sport fish effort and harvests by species from the Fish/Niukluk River 1983-2001, and catches 1990-2001.

Year	Num. Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Northern Pike	Whitefish	Burbot
Harvest													
1983	0	1,999	0	1,355	0	631	371	2,357	2,208	5,160	557	0	0
1984	521	1,115	0	1,090	0	78	52	1,220	325	376	13	0	13
1985	466	889	0	40	0	70	10	120	195	945	0	0	35
1986	850	1,888	189	1,359	0	415	0	1,963	1,359	1,114	19	0	0
1987	809	2,473	36	1,032	0	127	72	1,267	1,376	2,119	471	0	0
1988	866	2,245	0	800	0	73	127	1,000	891	1,237	0	0	0
1989	641	2,124	0	728	0	233	107	1,068	734	808	0	70	0
1990	580	2,059	0	267	0	638	216	1,121	348	415	17	0	0
1991	881	2,524	14	977	0	356	272	1,619	1,474	1,320	283	13	35
1992	773	2,742	0	753	0	357	15	1,125	303	158	43	0	0
1993	832	3,962	9	1,185	0	278	514	1,986	1,003	619	75	9	21
1994	766	3,082	10	1,122	0	231	119	1,482	708	644	99	0	0
1995	918	2,603	18	818	0	136	27	999	368	430	0	37	34
1996	692	2,120	11	1,652	0	404	166	2,233	402	313	145	0	24
1997	722	3,017	71	462	0	58	0	591	2,071	734	30	131	148
1998	229	1,344	0	316	0	0	0	316	160	16	0	0	84
1999	819	4,825	44	1,365	0	80	0	1,489	1,952	860	28	0	89
2000	534	3,451	174	1,165	0	51	0	1,390	1,687	442	57	0	0
2001	875	2,822	0	969	0	161	439	1,569	1,197	430	297	129	43
Avg 91-00	717	2,967	35	982	0	195	111	1,323	1,013	554	76	19	44
Avg 96-00	599	2,951	60	992	0	119	33	1,204	1,254	473	52	26	69

-continued-

Table 12.–Page 2 of 2.

Year	Num. Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Northern Pike	Whitefish	Burbot
Catches													
1990	580	2,059	0	1,140	0	2,096	487	3,723	912	2,189	133	0	0
1991	881	2,524	22	1,417	0	579	521	2,539	3,439	7,261	764	39	35
1992	773	2,742	0	1,555	0	1,969	326	3,850	1,041	2,171	256	0	0
1993	832	3,962	9	1,804	0	909	945	3,667	6,130	5,976	75	44	21
1994	766	3,082	29	1,448	0	2,052	1,271	4,800	2,125	2,398	596	56	0
1995	918	2,603	18	1,401	0	300	428	2,147	662	1,169	137	65	34
1996	692	2,120	64	3,348	0	3,512	1,660	8,584	1,872	4,653	513	94	24
1997	722	3,017	125	1,751	106	1,209	714	3,905	9,952	10,452	423	315	277
1998	229	1,344	15	772	0	3,252	822	4,861	1,390	8,159	189	198	84
1999	819	4,825	55	2,151	0	187	265	2,658	5,601	7,245	264	0	101
2000	534	3,451	207	2,952	0	3,989	952	8,100	2,261	1,701	114	210	0
2001	875	2,822	21	1,739	0	279	543	2,582	3,064	3,972	538	416	43
Avg 91-00	717	2,967	54	1,860	11	1,796	790	4,511	3,447	5,119	333	102	58
Avg 96-00	599	2,951	93	2,195	21	2,430	883	5,622	4,215	6,442	301	163	97

Table 13.—Sport fish effort and harvests by species from the Pilgrim River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Northern Pike	Whitefish	Burbot
							<u>Harvest</u>						
1983	0	597	0	37	0	37	111	185	445	761	148	0	0
1984	398	732	65	195	78	104	0	442	195	247	13	0	0
1985	363	375	10	20	20	50	100	200	14	319	0	0	0
1986	238	868	38	76	0	0	113	227	189	227	529	0	0
1987	438	1,159	72	109	435	0	272	888	163	272	199	0	0
1988	836	4,822	55	218	746	36	346	1,401	327	109	91	36	0
1989	1,050	1,678	68	204	78	301	272	923	603	516	415	131	10
1990	862	1,710	19	81	93	208	41	442	498	415	1,194	0	33
1991	1,169	3,183	51	310	124	81	85	651	1,015	459	608	13	0
1992	686	1,184	55	57	66	55	106	339	131	91	231	0	0
1993	570	1,195	28	191	10	0	0	229	730	75	207	0	0
1994	323	844	0	134	9	154	0	297	63	49	108	0	0
1995	531	1,253	19	113	62	0	73	267	74	52	68	18	11
1996	445	840	0	133	84	49	0	266	388	73	75	0	0
1997	456	820	45	0	20	0	0	65	65	81	117	0	0
1998	392	546	32	6	0	0	0	38	14	0	26	0	0
1999	283	433	0	33	0	0	0	33	45	11	94	9	0
2000	177	753	0	179	32	6	0	217	0	57	271	0	0
2001	207	491	0	29	0	0	0	29	270	43	0	0	0
Avg 91-00	503	1,105	23	116	41	35	26	240	253	95	181	4	1
Avg 96-00	351	678	15	70	27	11	0	124	102	44	117	2	0
							<u>Catches</u>						
1990	862	1,710	57	186	198	736	460	1,637	845	1,476	2,918	0	33
1991	1,169	3,183	65	476	374	406	194	1,515	3,155	4,463	1,499	13	0
1992	686	1,184	55	162	90	714	197	1,218	279	526	863	18	0
1993	570	1,195	92	325	106	392	254	1,169	3,038	2,362	959	0	11
1994	323	844	0	436	18	350	146	950	180	266	358	0	0
1995	531	1,253	19	472	163	58	232	944	294	370	656	27	11
1996	445	840	0	265	235	364	133	997	509	785	334	0	0
1997	456	820	90	49	115	0	15	269	254	429	262	0	0
1998	392	546	32	65	145	263	44	549	41	65	77	0	0
1999	283	433	20	77	177	0	0	274	585	694	104	9	0
2000	177	753	0	200	32	109	24	365	0	221	596	0	0
2001	207	491	0	29	14	0	11	54	439	402	0	0	0
Avg 91-00	503	1,105	37	253	146	266	124	825	834	1,018	571	7	2
Avg 96-00	351	678	28	131	141	147	43	491	278	439	275	2	0

Table 14.—Sport fish effort and harvests by species from the Snake River 1983-2000, and catches 1990-2001.

Year	Number of Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Whitefish
Harvest											
1983	0	119	19	0	0	37	0	56	223	278	0
1984	245	418	0	273	0	143	0	416	65	26	0
1985	129	361	0	120	0	0	0	120	0	139	0
1986	136	850	0	94	0	378	94	566	57	378	0
1987											
1988	340	2,128	0	800	0	546	437	1,783	218	709	0
1989	148	436	0	10	0	291	97	398	44	101	0
1990	298	775	10	47	0	111	41	209	66	116	0
1991	647	2,384	7	798	62	71	93	1,031	1,252	402	0
1992	461	2,379	8	510	0	183	0	701	115	16	0
1993	622	1,468	9	248	0	151	0	408	331	467	0
1994	341	880	0	145	0	452	7	604	117	32	0
1995	640	1,968	0	85	0	19	0	104	131	18	9
1996	433	1,269	0	426	0	659	0	1,085	97	121	0
1997	235	445	0	98	0	0	0	98	81	0	0
1998	164	376	0	0	0	463	0	765	0	8	0
1999	313	977	0	209	0	0	0	209	44	113	0
2000	185	377	0	209	0	103	0	312	199	16	0
2001	329	853	0	175	0	0	0	175	108	63	0
Avg 91-00	404	4,252	2	273	6	210	10	502	237	119	1
Avg 96-00	266	689	0	188	0	245	0	433	84	52	0
Catches											
1990	298	775	29	58	0	222	122	431	315	199	0
1991	647	2,384	14	1,798	73	234	109	2,228	3,471	2,096	0
1992	461	2,379	8	640	0	1,182	0	1,830	180	158	0
1993	622	1,468	9	306	0	429	37	781	1,003	1,614	0
1994	341	880	60	235	0	648	37	980	420	377	0
1995	640	1,968	0	245	0	300	189	734	507	887	9
1996	433	1,269	0	530	0	967	111	1,608	255	1,055	0
1997	235	445	0	118	0	0	9	127	243	123	135
1998	164	376	0	64	0	463	0	527	0	218	0
1999	313	977	0	606	0	0	0	606	257	723	0
2000	185	377	0	209	0	103	0	312	199	449	0
2001	329	853	0	214	0	21	78	313	108	1,385	0
Avg 91-00	404	1,252	9	475	7	433	49	973	654	770	14
Avg 96-00	266	689	0	305	0	307	24	636	191	514	27

Table 15.—Sport fish effort and harvests by species from the Solomon River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Whitefish
Harvests											
1983	0	30	0	0	0	0	0	0	37	0	0
1984	215	766	0	299	0	0	0	299	221	0	0
1985	391	2,667	0	80	0	120	219	419	611	0	0
1986	102	102	0	0	0	38	0	38	0	0	0
1987	270	272	0	109	0	0	72	181	1,576	91	0
1988	155	309	0	18	0	0	0	18	36	127	0
1989	263	492	10	136	0	243	49	438	745	152	0
1990	216	458	0	12	0	361	14	387	182	17	0
1991	593	1,057	7	83	0	173	0	263	2,219	158	0
1992	685	962	0	316	0	210	0	526	131	0	0
1993	317	1,404	28	420	0	259	0	707	893	0	61
1994	328	1,193	0	235	0	256	0	491	269	0	19
1995	426	781	0	38	0	87	0	125	366	0	0
1996	230	335	0	142	0	0	0	142	49	0	0
1997	250	434	0	10	0	15	0	25	186	0	26
1998	245	340	0	0	16	154	0	170	383	0	0
1999	193	438	0	22	0	0	0	22	154	0	0
2000	88	242	0	32	0	113	0	145	0	0	0
2001	254	615	0	39	0	0	0	39	162	0	0
Avg 91-00	336	719	4	130	2	127	0	262	465	16	11
Avg 95-99	201	358	0	41	3	56	0	101	154	0	5
Catches											
1990	216	458	0	12	58	736	108	914	415	33	0
1991	593	1,057	7	83	0	620	47	757	4,549	602	0
1992	685	962	0	316	0	998	91	1,405	197	38	0
1993	317	1,404	47	650	0	633	0	1,330	1,725	140	61
1994	328	1,193	0	255	0	784	7	1,046	520	212	19
1995	426	781	0	208	0	190	22	420	734	200	0
1996	230	335	0	237	0	39	0	276	49	97	0
1997	250	434	0	39	0	74	0	113	415	703	26
1998	245	340	17	59	64	433	0	573	410	0	0
1999	193	438	0	185	0	13	0	198	573	21	0
2000	88	242	57	53	119	288	278	795	1,537	853	0
2001	254	615	0	39	0	407	0	446	399	0	0
Avg 91-00	336	719	13	209	18	407	45	691	1,071	287	11
Avg 96-00	201	358	15	115	37	169	56	391	597	335	5

Table 16.—Sport fish effort and harvests by species from the Kuzitrin River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	King Salmon	Coho Salmon	Red Salmon	Pink Salmon	Chum Salmon	All Salmon	Dolly Varden	Arctic Grayling	Northern Pike	Whitefish	Burbot
Harvest													
1983	0	179	0	0	0	0	0	0	0	371	0	0	0
1984	153	279	0	0	0	325	0	325	260	195	156	0	0
1985	103	84	0	0	0	0	0	0	0	195	14	0	0
1986	204	318	0	0	0	0	0	0	38	189	151	0	0
1987	135	1,392	0	0	0	0	0	0	91	181	127	0	0
1988	217	1,037	0	0	36	55	54	145	109	1,255	437	36	0
1989	115	313	0	0	0	0	0	0	0	283	233	0	0
1990	282	572	0	0	0	28	14	42	0	133	746	0	0
1991	414	836	0	0	10	10	0	20	222	286	481	0	0
1992	287	469	0	8	0	46	0	54	8	0	128	0	0
1993	293	463	0	0	0	0	0	0	146	101	209	0	0
1994	267	643	0	109	0	0	0	109	0	98	169	0	0
1995	214	413	0	0	0	0	40	40	22	44	137	0	0
1996	230	483	0	0	0	39	0	39	85	230	497	0	0
1997	236	440	0	0	0	0	0	0	0	108	216	0	0
1998	122	122	0	0	0	0	0	0	0	8	38	0	0
1999	197	355	0	0	0	0	0	0	55	23	233	0	0
2000	111	373	0	0	0	0	0	0	0	64	63	36	0
2001	186	297	0	0	0	0	0	0	21	51	52	646	7
Avg 91-00	237	460	0	12	1	10	4	26	54	96	217	4	0
Avg 96-00	179	355	0	0	0	8	0	8	28	87	209	7	0
Catches													
1990	282	572	0	0	0	194	27	221	0	298	1,094	0	0
1991	414	836	0	0	10	41	0	51	333	1,349	1,937	0	0
1992	287	469	0	89	0	82	0	171	8	481	1,956	37	0
1993	293	463	0	0	0	0	0	0	263	288	751	9	0
1994	267	643	0	109	0	16	0	125	0	351	722	0	0
1995	214	413	0	0	0	0	40	40	54	192	1,005	0	11
1996	230	483	0	0	0	49	11	60	85	388	2,015	0	0
1997	236	440	0	0	64	0	0	64	85	1,068	1,503	0	0
1998	122	122	0	0	0	0	0	0	8	0	38	0	0
1999	197	355	0	7	7	0	0	14	176	158	1,840	0	0
2000	111	373	0	0	0	103	0	103	0	859	578	36	0
2001	186	297	0	0	0	0	0	0	22	760	2,738	646	7
Avg 91-00	237	460	0	21	8	29	5	63	101	513	1,235	8	1
Avg 96-00	179	355	0	1	14	30	2	48	71	495	1,195	7	0

Table 17.—Sport fish effort and harvests by species from the Penny River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	Salmon					All Salmon	Dolly Varden	Arctic Grayling
			King	Coho	Red	Pink	Chum			
Harvest										
1983	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1984	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1985	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1986	306	396	0	113	0	396	189	698	189	189
1987	34	34	0	0	0	0	0	0	0	0
1988	31	91	0	73	0	0	0	73	73	0
1989	33	34	0	10	0	0	0	10	99	30
1990	66	343	0	35	0	416	0	451	0	0
1991	36	61	0	0	0	0	0	0	95	0
1992	37	37	0	16	0	0	0	16	0	0
1993	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1994	78	101	0	0	9	34	0	43	46	0
1995	42	43	0	0	0	0	0	0	0	0
1996	40	80	0	95	0	0	0	95	12	0
1997	15	15	0	0	0	0	0	0	0	0
1998	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1999	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
2000	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
2001	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Avg 91-00	41	56	0	19	2	6	0	26	26	0
Avg 96-00	28	48	0	48	0	0	0	48	6	0
Catches										
1990	66	343	0	35	0	416	0	451	0	0
1991	36	61	0	0	0	0	0	0	95	0
1992	37	37	0	16	0	0	0	16	0	0
1993	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1994	78	101	0	20	87	341	0	448	84	0
1995	42	43	0	0	0	0	0	0	0	0
1996	40	80	0	95	0	0	0	95	61	73
1997	15	15	0	0	0	0	0	0	0	20
1998	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
1999	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
2000	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd
2001	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Avg 91-00	41	56	0	22	15	57	0	93	40	16
Avg 96-00	28	48	0	48	0	0	0	48	31	47

Table 18.—Sport fish effort and harvests by species from the Cripple River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	Salmon					All Salmon	Dolly Varden	Arctic Grayling
			King	Coho	Red	Pink	Chum			
<u>Harvest</u>										
1983	0	179	0	37	0	93	0	130	111	0
1984	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
1985	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
1986	102	170	0	113	0	189	189	491	57	0
1987	101	181	0	0	0	0	0	0	0	0
1988	155	509	0	528	0	0	0	528	127	127
1989	66	257	0	78	0	165	39	282	0	0
1990	66	813	0	140	0	1,138	41	1,319	663	0
1991	108	221	0	24	0	0	0	24	0	0
1992	37	149	0	32	0	92	0	124	0	0
1993	125	1,121	0	57	0	212	0	269	0	0
1994	91	320	0	122	0	85	0	207	27	0
1995	78	272	0	10	0	84	0	94	38	0
1996	164	307	0	134	0	156	0	290	24	0
1997	75	166	0	11	0	11	0	22	0	0
1998	513	2,303	0	366	0	1,141	0	1,507	272	0
1999	64	66	0	0	0	0	0	0	452	0
2000	120	614	0	149	0	177	0	326	100	0
2001	44	205	0	117	0	0	0	117	32	0
Avg 91-00	138	554	0	91	0	196	0	286	91	0
Avg 96-00	187	691	0	132	0	297	0	429	170	0
<u>Catches</u>										
1990	66	813	0	314	0	2,665	162	3,141	1,492	0
1991	108	221	0	24	0	0	0	24	0	0
1992	37	149	0	49	0	1,053	0	1,102	0	0
1993	125	1,121	0	57	0	403	0	460	44	0
1994	91	320	0	268	0	426	0	694	27	0
1995	78	272	0	10	0	225	22	257	38	0
1996	164	307	0	219	0	435	318	972	110	0
1997	75	166	0	11	0	22	290	323	0	20
1998	513	2,303	0	4,178	0	6,808	1,331	12,317	570	52
1999	64	66	0	72	0	0	0	72	965	0
2000	120	614	0	239	0	711	66	1,016	100	0
2001	44	205	0	117	0	34	0	151	32	0
Avg 91-00	138	554	0	513	0	1,008	203	1,724	185	7
Avg 96-00	187	691	0	944	0	1,595	401	2,940	349	14

Table 19.—Sport fish effort and harvests by species from the Sinuk River 1983-2001, and catches 1990-2001.

Year	Number of Anglers	Days Fished	Salmon					All Salmon	Dolly Varden	Arctic Grayling	Whitefish
			King	Coho	Red	Pink	Chum				
<u>Harvest</u>											
1983	0	477	0	0	0	0	19	19	1,132	130	0
1984	306	366	0	234	26	1,272	143	1,675	844	428	0
1985	311	806	0	10	0	120	0	130	292	0	0
1986	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
1987	405	5,198	0	0	380	91	72	543	652	724	0
1988	464	1,055	73	91	0	946	146	1,256	146	73	0
1989	230	906	0	10	0	0	10	20	175	51	104
1990	116	343	0	12	0	0	14	26	17	0	0
1991	557	885	0	71	41	51	47	210	729	129	0
1992	436	1,504	0	40	0	293	0	333	139	0	0
1993	463	874	9	96	0	115	0	220	536	37	0
1994	463	1,132	0	109	0	145	0	254	305	8	0
1995	485	1,295	0	19	21	28	0	68	158	18	0
1996	376	553	0	189	8	285	0	482	485	97	0
1997	235	443	0	0	0	54	0	54	346	0	0
1998	75	123	0	0	0	0	0	0	311	8	0
1999	78	244	0	0	0	0	0	0	88	11	0
2000	155	294	0	11	0	10	0	21	59	0	0
2001	240	490	0	62	39	0	0	101	86	43	0
Avg 91-00	332	735	1	54	7	98	5	164	316	31	0
Avg 96-00	184	331	0	40	2	70	0	111	258	23	0
<u>Catches</u>											
1990	116	343	0	12	0	0	406	418	66	232	0
1991	557	885	0	167	41	224	186	618	2,584	1,291	0
1992	436	1,504	0	65	0	1,429	15	1,509	770	300	0
1993	463	874	9	143	10	547	28	737	1,179	879	0
1994	463	1,132	0	172	0	348	22	542	830	417	0
1995	485	1,295	0	113	66	125	44	348	723	498	9
1996	376	553	0	246	8	736	200	1,190	618	339	0
1997	235	443	0	196	10	76	160	442	1,249	1,464	0
1998	75	123	0	0	0	0	0	0	311	25	0
1999	78	244	0	0	0	0	0	0	198	22	0
2000	155	294	0	21	0	21	12	54	95	26	0
2001	240	490	0	96	39	11	0	146	108	218	0
Avg 91-00	332	735	1	112	14	351	67	544	856	526	1
Avg 96-00	184	331	0	93	4	167	74	337	494	375	0

Recent Fishery Performance

The alternate year strong pink salmon run in Norton Sound has strongly influenced the salmon sport fishery on road accessible streams. However, a production failure of the strong pink salmon run in 1998 has resulted in relatively weak even year returns of pink salmon to Nome subdistrict streams in 2000 and 2002. Salmon harvests from the Nome River increased 10 fold between 1997 and 1998, and dropped 10 fold in 1999, reflecting the strong even-year pink salmon run in spite of a reduced amount of angler effort. Weak even year returns since 1998 have resulted in a change in this pattern. Effort on the Nome River has dropped continuously from a high of 7,200 angler days in 1990 to about 1,460 angler days in 1999 and 2000, and 1,000 angler days in 2001. The pink salmon harvest of about 1,985 fish in 1998 may also have been influenced by reduced subsistence opportunity on depressed chum salmon stocks that likely focused local sport fishing effort on the abundant pink salmon, in part to meet the local need for salmon. The 2000 harvest of coho salmon in the Nome River was about 342 fish, and the 2001 harvest was 297 fish, near the recent five-year average harvest of about 329 coho (Table 11). Chum salmon fishing has been closed for many years because of depressed stocks and harvests of sockeye and chinook salmon in the Nome River are negligible. A similar declining trend in sport fishing effort had been seen in the Fish/Niukluk River system from a high of about 4,000 angler days in 1993 to 1,344 angler days in 1998 (Table 12). However, in 1999, and 2000 effort increased to 4,825 angler days and 3451 angler days respectively, suggesting a shift in effort away from the Nome subdistrict. This trend continued in 2001 with 2,822 angler days estimated, although there was a general decline in sport fishing effort throughout the area. Sport fishing for chum salmon is allowed in the Fish/Niukluk, and the estimated harvest of chum salmon increased to 439 in 2001. The coho harvest in 2001 was about 970 fish from an estimated catch of 1,740 fish, and the estimated chinook harvest was negligible. The harvest of pink salmon was estimated at 161 fish in 2001, and has been trending upward over the past four years, but is still very small in spite of abundant runs of this species with over 1,500,000 counted past the tower in 1998, 962,000 past the tower in 2000, and 46,600 in 2001. The Pilgrim River is the other road accessible water where chum salmon fishing is still allowed. Effort there in 2001 was estimated at 490 angler days near the recent five-year average of 687 angler days (Table 13). Recent harvests of salmon have been very small with only about 30 salmon estimated to have been harvested in 2001.

Sport Fishery Management Objectives

There have been no specific management objectives identified for salmon fisheries on the in Nome roadside streams. The goal of sport fishery management in the in these waters is to assure that escapement goals are met and to maintain opportunity for anglers to participate in the fisheries. Sport fishery harvests are small, and emergency actions to restrict harvest are generally not contemplated unless escapement-monitoring projects indicate that the particular run is small and that restrictions in subsistence fisheries may be necessary thereby mandating restrictions in the sport fisheries in order for escapements to reach designated goals. Biological Escapement Goals (BEGs) based aerial surveys are in place, and goals based on tower or weir estimates (Snake and Nome rivers) are used to assess the runs. Goals have not been established for species other than chum salmon in these streams, and pink salmon in the Nome River, but they will be developed as more years of reliable counts from escapement assessment projects have been accumulated.

Management History and Recent Board of Fisheries and Management Actions

Chum salmon stocks in the Nome area are depressed, and an effort to restore these runs is ongoing. Salmon sport fisheries in northwestern Alaska are managed in cooperation with the Division of Commercial Fisheries. Subsistence uses are given priority. Since the availability of salmon resources is limited and local chum salmon populations are depressed, particularly in the Nome area where sport fishing effort is greatest, the Nome area has most often required restrictive management measures. Chum salmon sport and commercial fisheries in the Nome subdistrict have been closed. Subsistence fisheries are managed to achieve escapement goals.

In 1984 the BOF reduced the bag and possession limits in the Nome and Snake rivers to 15 salmon other than king salmon, only 5 of which could be chum and coho in combination, and in 1985 all but the lower 2 miles of the Nome River was closed to all sport fishing for salmon by EO.

In 1987, additional regulations were adopted in Seward Peninsula drainages (Cape Prince of Wales to Cape Darby), which reduced the bag and possession limit for salmon other than chinook to 10 per day, 10 in possession, only 3 of which could be chum or coho salmon in combination. The limit for chinook salmon was set at 1 per day and in possession.

The Nome River was closed in July 1990 to the taking of chum salmon on sport fishing gear, and the following year, another EO closed Nome area waters to the retention of both chum and pink salmon. The area affected by this action was more widespread than in previous years and included all waters from the Sinuk River in the west to the Solomon River in the east.

Due to continued low escapements of chum salmon in Nome area streams, a proposal to close sport fishing for chum salmon in the rivers addressed in the 1991 EO was brought before the BOF and enacted into regulation. This regulation is intended to protect chum salmon stocks and will remain in effect until stocks recover and surpluses above the escapement goals are available for harvest by sport anglers.

Two EOs were issued addressing salmon in the Nome area in 1992. The first closed the Tubutulik and Kwiniuk rivers to sport fishing for chum salmon. The second, because of near record pink salmon runs, increased the bag and possession limits for pink salmon from 10 per day to 20 per day in Nome area streams.

During the December 1997 meeting the BOF adopted two salmon regulation changes for the NWMA. The first established a uniform daily bag and possession limit for chinook salmon in the entire NWMA at one fish. This replaced a three fish (only one over 28 inches) daily bag and possession except for the Unalakleet River where a one fish limit was already in place. Since the Unalakleet River has the strongest chinook run in the NWMA, having a more liberal limit in other parts of the area where chinook runs were small made little sense. The other change was for northern Norton Sound that includes the streams accessible from the Nome road system. The "other salmon" aggregate limit of 10 fish per day only 3 of which could be chum or coho was split out to provide separate daily bag and possession limits by species. The new limits were set at 10 pink, 3 coho, 3 chum, and 3 sockeye. The new regulation will provide for more precise management by species, and is less ambiguous for the inexperienced angler. Nome subdistrict streams still remain closed to fishing for chum salmon.

Because of weak coho runs throughout northern Norton Sound, an EO was issued on August 15, 1997 that mandated the release of all coho caught while sport fishing in northern Norton Sound

waters, in addition, bait was eliminated as a legal gear while fishing for coho. This action remained in effect until September 30, 1997.

The BOF scheduled a special meeting in Nome for March 1998 to discuss the chum salmon situation in the Nome subdistrict. The BOF directed Nome area residents to create a Subsistence Salmon Working Group to address the issue of the chronic chum salmon shortage and consider the possibility of Tier-II management. The group did not recommend Tier-II management, however in the March 1999 meeting, the BOF directed the department to implement Tier II subsistence management for chum salmon in the Nome subdistrict.

In 2000 three EOs affecting sport fishing for salmon were issued in northern Norton Sound. One closed all waters to sport fishing for salmon in the Nome subdistrict so that subsistence closures could proceed. A later EO, issued after the earlier closure had expired, reduced the coho salmon daily bag limit from three to one. A later EO closed coho salmon fishing on the Nome River, which was later reopened by a third EO after returns strengthened.

In 2001 five EOs affecting sport fishing for salmon were issued in northern Norton Sound (Appendix C). EO No 3-S-01-01 closed all waters to sport fishing for salmon in the Nome subdistrict so that subsistence closures could precede the runs and open on a river by river basis as the runs progressed. EO No 3-PS-01-01 effective on July 31 closed pink salmon sport fisheries in the Nome subdistrict. EO No 3-SS-01-01 on August 20, reduced the coho daily bag limit in Nome subdistrict streams and the Pilgrim River from 3 to 1 coho. On August 23, EO 3-SS-02-01 closed coho sport fishing on the Nome River, and on August 27, EO 3-ss-03-01 reopened coho fishing on the Nome River with a daily bag limit of one coho.

Current Issues

Chum salmon stocks have steadily declined on the Seward Peninsula since the early 1980s, as evidenced by failure to achieve desired spawning escapements during recent years in many key streams where spawners are enumerated. This has created the need for increasingly restrictive sport, commercial and even subsistence fishing regulations. It is anticipated that until chum salmon populations recover, there will be a need to continue with very restrictive measures to protect local stocks. All rivers in northern Norton Sound from the Sinuk in the west to Topkok in the east are closed to sport fishing for chum salmon, and will remain closed until runs rebuild. In addition, restrictions to the sport harvest of coho salmon in the Nome area have been necessary during recent years due to weak or late runs. Increased effort is being directed toward the enumeration of coho salmon escapements in Nome area streams using tower and weir projects. Data from these projects have allowed more precise management of coho sport fisheries in the area.

Ongoing Research and Management Activities

Current research and management activities on Nome roadside salmon populations are primarily conducted by the CFMD in conjunction with Kawarek Corporation's fisheries office. These groups cooperatively staff and manage escapement enumeration projects on the Niukluk, Eldorado, Pilgrim and Snake rivers. All projects are counting towers except the Nome River where a weir is run by CFMD throughout the salmon runs. The weir obstructs the movement of all fish, and fish are counted as they are permitted to pass through an opening in the weir several times each day. Weirs may replace towers in the Eldorado, Snake and Pilgrim rivers. In addition, the BLM has operated a weir at the outlet of Glacial Lake in 2001 and 2002 to

enumerate sockeye migrating into the lake. A potential weir site has been located in the mainstem of the Unalakleet River and a structure may be placed there in the near future.

NORTHWESTERN ALASKA DOLLY VARDEN AND ARCTIC CHAR

Fishery Description and Historical Perspective

In the Northwestern Management Area, Arctic char occur in lakes in the Kigluaik Mountains and in some headwater lakes in the Kobuk and Noatak River drainages, while Dolly Varden are common inhabitants of most coastal streams and large rivers (Figure 12). Although the department groups Dolly Varden and Arctic char for bag limits and record keeping, the two species are separate with Arctic char present only as lake resident populations, while Dolly Varden may be present as lake resident, stream resident, dwarf, or anadromous populations. Arctic char distribution is very limited in the NWMA and the vast majority of char fisheries are directed toward Dolly Varden. There have been no recent directed commercial fisheries on char in northwestern Alaska.

Many northwestern Alaska residents maintain a traditional lifestyle, and are dependent to some degree on locally harvested fish resources. Dolly Varden comprise an important part of this traditional harvest, and in Kivalina they outrank salmon and whitefish in importance to the subsistence economy. The number of Dolly Varden harvested for subsistence purposes in northwestern Alaska vastly exceeds the number taken by sport anglers. Intermittent community subsistence harvest estimates dating to 1959 for Kivalina and Noatak (Table 20) and personal observation of the area biologist suggest that 15,000 to 25,000 Dolly Varden are harvested annually in this area, however, the actual magnitude of the annual harvests throughout the NWMA are not known. Fish are captured with gill nets or beach seines during open water periods, and with hook and line during winter. Dolly Varden are also an important subsistence resource in Norton Sound, however their relative importance is minor compared to salmon.

Estimated harvests of Dolly Varden by sport anglers in the Seward Peninsula/Norton Sound sub-area have averaged about 5,000 fish over the past 10 years (Table 21). Observations and aerial surveys suggest that Dolly Varden spawner abundance is low in most rivers, however, spawning occurs in almost all drainages of Norton Sound, some northern Seward Peninsula rivers, and the major drainages of Kotzebue Sound and the Chukchi Sea. Aerial surveys of spawning Dolly Varden conducted during the mid 1980s indicated that about 12,000-15,000 spawn annually in the Noatak drainage (Table 22). Total abundance of spawning Dolly Varden in northwestern Alaska is unknown. Recent surveys and reports from local residents suggest that spawner abundance in Kotzebue area streams has been lower in the past two years than was indicated in earlier surveys.

Drainages of Kotzebue Sound and the Chukchi Sea are known for the large size of anadromous Dolly Varden available to the sport angler. Since the inception of ADF&G's Trophy Fish Program in 1967, out of 179 qualifying fish in the Dolly Varden/Arctic char category, 108 (60%) have come from the NWMA, and in the past 10 years (1990-1999) 88 out of 96 (92%) have come from northwestern Alaska. In addition, the current Alaska sport fish angling record for Arctic char/Dolly Varden (20 lbs. 12 oz.) is a Dolly Varden taken from the Wulik River in 2000. In October 2002, a new record Dolly Varden weighing more than 27 pounds was taken from the same river.

During summer, spawning Dolly Varden are targeted in some northwestern Alaskan streams, however, most sport fisheries for char target overwintering populations of Dolly Varden either in




Figure 12.-Dolly Varden and Arctic char distribution in the NWMA.

Table 20.-Documented subsistence harvests of Dolly Varden in Noatak and Kivalina.

Year	Kivalina		Noatak Number
	Number	Pounds	
1959	34,240	85,600 ^a	
1960	49,720	124,300 ^a	
1962			27,623 ^b
1963			4,130
1964		93,995	
1965		28,140	
1968	49,512	120,214	
1969	64,970	152,750	32,350
1970	33,820	79,420	3,700
1971	29,281	68,518	5,320
1972	48,807	114,637	1,492
1979 ^c	14,600		
1980			9,060
1981 ^c	15-18,000		7,220
1982 ^c	18,438	69,059	3,056
1983 ^c	16,270	68,467	2,676
1984 ^c	12,000		4,545
1985 ^c	10,500		2,542
1986 ^c	7,436		
1991			4,814 ^d
1992			4,395 ^d
1993			4,275 ^d
1995			5,762 ^d
1996			5,031 ^d
1997			4,763 ^d
1998			3,872 ^d
1999			
2000			3,315 ^d
2001			2,702 ^e

^a Sarrio and Kessel 1966

^b Foote and Williamson 1966

^c Data from Sport Fish Division surveys.

^d Data from ADF&G Subsistence Division household surveys Georgette and Utermohle 1998-2001.

^e Data from ADF&G Subsistence Division household surveys Georgette et al. *In press*.

Table 21.—Historic Dolly Varden harvests and catches in NWMA by sub-area, 1977-2001.

Year	Seward Peninsula/Norton Sound					Kotzebue/Chukchi Sea				
	Number of Anglers	Effort Angler Days	Dolly Varden Harvest	Dolly Varden Catch	% Harvested	Number of Anglers	Effort Angler Days	Dolly Varden Harvest	Dolly Varden Catch	% Harvested
1977		7,828	1,621				3,487	469		
1978		8,379	1,690				4,997	199		
1979		8,725					2,593	1,772		
1980		7,958	5,811				3,841	301		
1981		10,879	3,981				5,284	1,177		
1982		13,198	6,498				6,906	1,531		
1983		16,944	9,779				7,963	2,192		
1984	1,597	17,436	4,260			696	7,791	3,804		
1985	2,854	19,919	5,695			1,788	6,701	1,557		
1986	2,872	18,107	5,381			1,570	6,313	1,300		
1987	2,528	20,413	5,506			2,090	9,288	1,072		
1988	2,661	20,278	4,437			959	5,279	983		
1989	2,560	17,692	7,003			1,028	4,932	999		
1990	2,686	21,799	3,765	9,118	41	991	3,782	806	3,747	22
1991	3,236	23,622	10,365	25,425	41	1,606	9,543	1,149	1,658	69
1992	3,540	22,684	2,382	6,012	40	1,421	6,145	582	7,054	8
1993	3,134	18,930	5,907	22,166	27	1,575	7,809	914	7,190	13
1994	3,016	18,922	3,071	7,344	42	1,100	6,036	2,365	10,733	22
1995	3,719	19,647	2,908	7,921	37	1,957	8,495	939	7,804	12
1996	2,958	13,783	4,285	8,427	51	1,407	5,571	913	5,376	17
1997	2,773	13,850	4,467	17,988	25	824	3,729	598	7,346	8
1998	3,206	13,616	2,240	5,711	39	1,089	3,801	440	8,606	5
1999	3,124	15,006	6,708	21,428	31	1,313	6,771	796	8,259	10
2000	2,713	18,559	7,952	16,348	49	1,387	7,129	1,599	8,031	20
2001	2,371	10,955	3,174	7,395	43	1,177	5,904	1,693	4,766	36
Avg (91-00)	3,142	17,862	5,029	13,877	38	1,368	6,503	1,030	7,206	18
Avg (96-00)	2,955	14,963	5,130	13,980	39	1,204	5,400	869	7,524	12

Table 22.—Aerial counts of Dolly Varden spawning in the Noatak River and overwintering in the Wulik and Kivalina rivers, 1968-2001.

Year	Spawners		Nonspawners	
	Noatak River	Wulik River	Kivalina River	
1968		90,286	27,640	
1969		297,257	nd	
1976		68,300	12,600	
1979		55,030	15,744	
1980		113,553	39,692	
1981	7,922	101,826	45,355	
1982	8,275	65,581	10,932	
1984	9,290	30,923	5,474	
1985	11,073	nd	nd	
1986	nd	5,590	5,030	
1988	nd	80,000	nd	
1989	nd	56,384	nd	
1990	7,261	nd	nd	
1991	9,605	126,985	35,275	
1992	nd	135,135	nd	
1993	9,560	144,138	16,534	
1994	nd	66,752	nd	
1995	6,500	128,705	28,870	
1996	12,184	61,005	nd	
1997	nd	95,412	nd	
1998	nd	104,043	nd	
1999	9,636	nd	nd	
2000	nd	92,614	nd	
2001	nd	92,614	nd	

^a Only Kelly River and part of Kugururok River counted.

nd = no data

the fall as they enter freshwater from the sea, or in the spring as they move toward the sea for feeding. Since overwintering populations are composed of mixed stocks, potentially from a wide geographic area, harvests in the few rivers with good angler access have been sustainable. In some streams along the Nome road system, if such harvests were directed towards a single stock they would likely not be sustainable. Movements of Norton Sound Dolly Varden are tied to those of salmon, and Dolly Varden are often present in streams during late summer to feed on salmon eggs, especially during years of high pink salmon abundance. They are likely to remain longer in streams during the spring following a large pink salmon run in order to feed on outmigrating fry. The timing of the fall movement of Dolly Varden into Seward Peninsula streams has varied widely over the past 10 years resulting in annual changes in the availability of Dolly Varden to the fall fishery. Consequently, fisheries and harvests in this area follow these patterns of availability. In 1988, the BOF adopted the bag limit of 10 Dolly Varden/Arctic char per day with 10 in possession with exceptions for the Noatak, Wulik, and Kivalina rivers where only 2 of the 10 fish could be over 20 inches in length. In 1994, the BOF adopted the current daily bag and possession limits for char in the entire AYK region with 10 fish per day, only 2 over 20 inches allowed in marine or flowing waters; and 2 fish per day allowed in lakes. Due to habitat preferences, these regulations allow a liberal limit for Dolly Varden while protecting spawning fish, and a conservative limit for Arctic char without the need for anglers to differentiate between the two species.

Recent Fishery Performance

In the Kotzebue/Chukchi Sea sub-area, sport harvests have averaged about 1,000 Dolly Varden annually. Estimated mean annual catch (which includes fish that are kept and those released) since 1990 has been 14,000 Dolly Varden in the Seward Peninsula/Norton Sound area, and 7,200 in the Kotzebue/Chukchi Sea area (Table 21). The data suggest that about 60% of all Dolly Varden captured in the Seward Peninsula/Norton Sound area are released while about 82% in the Kotzebue area are released. The higher harvest rate in the Seward Peninsula/Norton Sound area is likely because local residents have good road access to fishing areas where fish taken on rod and reel are used for food. In the Kotzebue area, fishing sites are accessed by air and much of the effort is from outside the local area by anglers seeking a quality fishing experience. While effort levels in both the Seward Peninsula/Norton Sound area and the Kotzebue area were lower in 1997 and 1998, effort increased in 1999. Catches of Dolly Varden in the Seward Peninsula/Norton Sound sub-area increased dramatically from 5,700 in 1998 to 21,000 in 1999. They declined to about 7,400 in 2001. Catches have remained more stable in the Kotzebue/Chukchi Sea sub-area, but have dropped to about 4,700 fish in 2001. The catch of Dolly Varden per angler day has averaged much higher in the Kotzebue area than in the Seward Peninsula area. This is likely because much of the effort on the Seward Peninsula is directed at other species, while most of the Kotzebue area effort is directed at Dolly Varden. The Kotzebue area seems to be attracting more visiting anglers seeking to catch large northern form Dolly Varden.

Dolly Varden harvests in the Seward Peninsula/Norton Sound sub-area are distributed among most of the sampled rivers with highest harvests coming from the Nome, Unalakleet, Solomon and Fish/Niukluk rivers (Table 23). In the Kotzebue/Chukchi Sea sub-area, highest harvests are from the Noatak and the “other rivers” category that includes the Wulik and Kivalina rivers.

Wulik River

The Wulik River is located about 90 miles north of Kotzebue and is well known as a “char” fishing destination (Figure 7). The river is about 90 miles long and enters the Chukchi Sea through Kivalina Lagoon near the village of Kivalina. The river is heavily used by the residents of Kivalina for subsistence and the majority of the village harvest of Dolly Varden comes from the Wulik River (Table 21). During the most recent five-year period (1996-2001), estimated sport fishing effort has averaged about 500 angler-days (Howe et al. 1996, 2000a-d; Walker et al. *In prep*; Jennings et al. *In prep*). Sport fishing occurs throughout the open water period, but the majority of effort and harvest occurs during late August and September when Dolly Varden return from the sea to winter in the river.

The average harvest of Dolly Varden from the Wulik River (1996-2000) was 248 fish. The estimated sport harvest during 2001 was 427 fish (Walker et al. *In press*) with the 2001 catch estimated at 2,925 Dolly Varden. Approximately 49% of the catch was of fish greater than 20 inches in length. Effort on the Wulik River in 2001 was estimated at 591 angler days. Effort more than doubled between 1998 and 1999 to almost 750 angler days, but dropped to 336 angler days in 2000. Although the river is well known for its large run of anadromous Dolly Varden, effort has remained relatively low because of the river’s remote location and difficulty of access. However, this may change with publicity regarding the recent record sized fish having been caught there in 2000 and in 2002.

Fishery Objectives and Management

Management of Dolly Varden in Norton Sound streams is structured to maintain opportunity and allow a relatively liberal bag on mixed stock population aggregations. In the Kotzebue sub-area, the intent is to maintain a high quality fishery with the opportunity to harvest a small number of large sized char under a conservative bag that protects the spawning component of the population, minimizes conflicts with subsistence users, and does not adversely affect the population structure. Because of the differential size structure of the population groups north and south of the Bering Strait, it is felt that these objectives are being addressed with the general bag and possession limit regulation of 10 fish per day with only 2 over 20 inches in length.

Fishery Outlook

Dolly Varden in Norton Sound are widespread, they spawn in most rivers and overwinter in all major drainages. The outlook is for fisheries to continue into the future without further management actions unless participation in the fisheries increases dramatically. In the Kotzebue area, the fishery is likely to grow in popularity as more anglers experience these high quality-fishing opportunities and publicity regarding the availability of large size fish increases. Until these fisheries grow to the point that harvests are thought to affect spawner abundance, spawner success, or population structure, it is unlikely that additional management actions will be necessary.

Recent Board of Fisheries and Management Actions

Population assessments conducted on the Nome and Solomon rivers in 1991 and 1992 suggested that the number of fish overwintering in these drainages could not sustain harvest levels occurring at that time. Consequently, the daily bag limit was reduced by EO to two fish. Subsequent studies showed that these populations were comprised of mixed stocks and that at least 20% of the fish overwintering in a given river could be expected to overwinter in a different river the next year. Tag recoveries showed that fish ranged over a wide geographic area. Since

Table 23.—Historic Dolly Varden and Arctic char harvests in the NWMA by sub-area and river, 1988-2001.

Areas	Year															Average	
	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	(91-00)	(96-00)
<u>Seward Peninsula/Norton Sound Historic Dolly Varden and Arctic Char Harvests</u>																	
Salt Water	0	418	55	183	0	204	205	90	0	12	189	0	330	1,069	1,66	210	320
Nome River		2,001	3,551	1,078	1,220	557	917	431	462	873	328	302	791	340	43	622	527
Pilgrim River		327	603	166	856	131	448	63	74	388	65	14	45	0	270	2,208	102
Unalakleet R.		891	570	614	1,474	746	427	410	976	1,506	936	588	2,384	4,462	1,002	1,391	1,975
Fish-Niukluk R.		0	734	348	1,474	270	1,003	699	346	402	2,071	160	1,952	1,687	1,197	1,006	1,254
Sinuk R.					729	139	536	305	158	485	346	311	88	59	86	316	258
Snake R.					1,252	115	331	117	131	97	81	0	44	199	108	237	84
Solomon R.					2,219	131	893	197	366	49	186	383	154	0	162	458	154
Other Streams	5,506	1,218	1,545	1,227	1,141	89	1,050	759	395	473	265	482	920	136	140	571	455
Lakes ^a		0	0	332	0	0	97	0	0	0	0	0	0	0	0	10	0
Freshwater																	
Total	5,506	4,437	7,003	3,765	10,365	2,178	5,702	2,981	2,908	4,273	4,278	2,240	6,378	6,883	3,008	4,819	4,810
Grand Total	5,506	4,855	7,058	3,948	10,365	2,382	5,907	3,071	2,908	4,285	4,467	2,240	6,708	7,952	3,174	5,029	5,130
<u>Kotzebue/Chukchi Sea Historic Dolly Varden and Arctic Char Harvests</u>																	
Salt Water	148	0	0	0	199	0	0	27	22	0	28	0	0	281	108	56	62
Kobuk R.	127		23	34	170	99	9	132	28	172	82	49	49	47	79	84	80
Noatak R.	844		651	386	936	197	325	786	124	632	139	175	255	763	1,026	433	393
Other Streams	401	965	302	302	412	279	533	1,402	676	97	347	216	181	467	430	461	262
Lakes ^a	0	18	23	471	0	107	47	18	0	12	0	0	22	41	50	25	15
Freshwater																	
Total	1,372	983	999	1,193	1,518	682	914	2,338	828	913	568	440	507	1,318	1,585	1,003	749
Grand Total	1,520	983	999	1,193	1,717	682	914	2,365	850	913	596	440	507	1,599	1,693	1,058	811

^a Lake totals are for Arctic char.

exploitation occurs primarily on mixed stocks in only a few locations and many of the represented stocks sustain no other exploitation, harvest levels were thought to be sustainable. Long-term harvest data supported this assumption and the reduced bag limit was rescinded. In the November 1994 meeting, the BOF adopted regulations that created a 10 fish with only 2 over 20 inches daily bag and possession limit for Dolly Varden/Arctic char in flowing and marine waters for the entire AYK Region. A separate daily bag limit of 2 fish (no size limit) was also created for lakes. The effects of these new bag limits are to have a fairly liberal limit for resident and migratory Dolly Varden that protects spawning sized fish, while maintaining a conservative limit for lake resident Arctic char without requiring anglers to differentiate between the two species. This bag limit has not needed to be adjusted, and no recent EOs have been issued regarding char in NWMA.

Current Issues

With over 100,000 anadromous Dolly Varden overwintering annually, the Wulik River is probably the most important Dolly Varden stream in northwestern Alaska (Table 22). Fish from this river are also very important as a subsistence food to the residents of Kivalina who harvest 15,000 to 20,000 annually (Table 20). The Red Dog Mine is located in the headwaters of this drainage and poses a potential threat to these fish and the water quality of the river. Water quality near the mine is systematically monitored and except for a pollution event in 1989-1990, the mine has operated in an environmentally sensitive manner. The Red Dog Mine funds a program run by the Division of Habitat to monitor heavy metals concentrations in receiving waters and in fish tissues. Fish tissues are sampled for heavy metals in the spring and the fall each year on a continuing basis in cooperation with the Division of Habitat. The recent discovery of additional new ore bodies will add new challenges to mineral development in this important drainage.

The question of how great an impact Dolly Varden have on salmon, especially chum salmon populations that have been depressed in Norton Sound for several years, has been raised by Nome residents in a number of public meetings. The department has no data concerning the possible effects of Dolly Varden egg predation on salmon numbers, however Dolly Varden have not been found to be significant predators on juvenile chum salmon in published predation studies.

Ongoing Research and Management Activities

The ADF&G began an effort to assess Dolly Varden populations in waters of the Seward Peninsula in 1991. Abundances and size compositions have been estimated for Dolly Varden overwintering in the Nome River in 1991 and 1992, and in the Solomon River in 1991. In addition, the movement of marked fish from the Nome River in 1991 to other rivers in 1992 was estimated (DeCicco 1993). These data in combination with harvest estimates and observed changes in abundances have been used to guide ADF&G management activities. It has been learned that Dolly Varden that overwinter in a particular stream may overwinter in other streams during subsequent years. Hence, a restrictive bag limit in one stream does not necessarily protect a single stock because fish range widely and stocks mix over a broad geographic area. Periodic assessment of Dolly Varden populations will continue as needed. During the winter of 2000/2001 Dolly Varden were radio tagged in the Nome and Solomon rivers in order to document the critical wintering areas in these rivers (DeCicco 2001). Fish remained in the Nome River throughout the winter, but about half those with radio tags in the Solomon River moved to the nearby Bonanza River during the course of the winter.

Studies in the Kotzebue area have continued intermittently since 1967, but in recent years have been limited to counting spawning Dolly Varden in Noatak River tributary streams with the assistance of the National Park Service (NPS), and counting Dolly Varden overwintering in the Wulik River with the assistance of the Habitat Division of ADF&G. Data on the abundance of Dolly Varden spawning in the Noatak River system and overwintering in the Wulik River will continue to be collected in cooperation with the NPS and the Habitat Division. A genetics study funded through the USFWS Office of Subsistence Management to determine the relationships among stocks north and south of the Bering Strait is being conducted. Samples have been collected and are being analyzed by the USFWS Conservation Genetics Lab in Anchorage. A spawning stock study on Kagvik Creek, a tributary to the Kugururok River in the Noatak River system has been unsuccessful for two years due to high water, and has been curtailed. The effects of multiple catch and release on spawning success should be investigated.

NORTHWESTERN ALASKA ARCTIC GRAYLING

Fishery Description and Historical Perspective

Arctic grayling are the most numerous species harvested in the Kotzebue/Chukchi Sea sub-area and the third or fourth most commonly harvested species in the Seward Peninsula/Norton Sound sub-area. In general, the sport fisheries for grayling in the northwestern area are small with average estimated annual harvests of 1,500 in the Seward Peninsula/Norton Sound sub-area and 1,400 in the Kotzebue/Chukchi Sea sub-area (Tables 24 and 25).

The Seward Peninsula has long been known for its production of large Arctic grayling with approximately 25% of all trophy grayling registered with the department's trophy fish program coming from this area. However, most populations are quite small and since they are resident in separate, often small streams, they must be managed as independent units with regulations tailored to the individual populations or groups of similarly structured populations. A management plan for Nome area roadside Arctic grayling fisheries has recently been developed (DeCicco 2002a).

Since 1989, the stock status of grayling populations in several rivers where sport fishing occurs on the Seward Peninsula has been investigated (DeCicco 1990-1999, 2002b, DeCicco and Wallendorf 2000; Gryska *In prep*). The Nome River stock was found to be overexploited, while the Niukluk, Fish, Pilgrim, Snake and Sinuk rivers populations are believed to be sustaining current levels of harvest. The Solomon River was found to have a very small Arctic grayling population.

Grayling densities in most Seward Peninsula rivers are low. They ranged from about 40 to 60 grayling per mile in the Nome and Sinuk rivers, to about 200 grayling per mile in the Pilgrim River. Densities in the Niukluk and Fish rivers were higher at about 375 and 185 grayling per mile respectively in 1991. More recent data have shown density in the Niukluk River had increased to about 470 grayling per mile in 1998 while density in the Fish River was estimated at about 500 grayling per mile in 1999. In contrast, interior Alaskan populations often exceed 500 fish per mile. Arctic grayling from Seward Peninsula rivers are generally older and larger when they first spawn than grayling in interior Alaska streams. Since they can live for more than 20 years, some grayling survive to grow very large, particularly in rivers where fishing effort is light. A grayling from the Eldorado River was recently aged at 31 years. For example in the lightly exploited Sinuk River almost 70% of the 1991 sample was age-8 or older and the average total length of all fish sampled was almost 19 inches. However, the density of fish was low,

Table 24.—Historic Arctic grayling harvests and catches in Seward Peninsula/Norton Sound waters, 1987-2001.

Areas	Year															Average	
	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	(91-00)	(96-00)
<u>Harvests</u>																	
Salt Water	0	55	0	0	0	0	0	131	0	0	0	0	0	0	0	13	16
Nome River		891	2,032	33	186	0	0	16	0	0	0	0	0	0	0	20	2
Pilgrim River		109	516	415	445	91	75	49	52	73	81	0	11	58	43	94	50
Unalakleet R			142	99	1,708	98	131	353	291	420	210	144	277	538	247	417	296
Fish-Niukluk R.		1,237	748	415	1,320	128	585	506	404	313	734	16	1,029	442	430	548	504
Sinuk R.					129	0	37	8	18	97	0	8	11	0	43	31	22
Snake R.					402	16	467	32	18	121	0	8	113	16	63	119	97
Solomon R.					158	0	0	0	0	0	0	0	0	0	0	16	0
Other Streams	4,600	2,636	767	416	773	159	289	236	254	461	236	122	159	149	168	284	238
Lakes		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Freshwater																	
Total	4,600	4,873	4,205	1,378	5,121	492	1,584	1,200	1,037	1,485	1,261	298	1,600	1,203	994	1,528	1,209
Grand Total	4,600	4,928	4,205	1,378	5,121	492	1,584	1,331	1,037	1,485	1,261	298	1,600	1,203	994	1,541	1,225
<u>Catches</u>																	
Salt Water				0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nome River				613	1,363	90	569	1,111	571	497	569	207	300	10	60	529	479
Pilgrim River				1,476	4,463	526	2,362	266	370	821	429	65	694	221	403	1,022	654
Unalakleet R				448	4,104	1,459	874	1,639	1,471	1,694	4,918	3,256	6,089	6,814	2,331	3,232	3,344
Fish-Niukluk R.				2,189	7,261	2,171	5,976	2,389	1,169	4,653	10,452	8,159	7,414	1,701	3,972	5,135	5,239
Sinuk R.				232	1,291	300	879	417	498	339	1,464	25	22	29	218	526	459
Snake R.				199	2,096	158	1,614	377	887	1,055	123	218	723	449	1,385	770	681
Solomon R.				33	602	38	140	212	200	97	703	0	21	853	0	287	278
Other Streams				929	1,980	1,030	809	670	622	1,250	1,529	1,570	869	992	1,098	1,132	1,039
Lakes				0	0	0	0	0	0	0	0	0	0	0	0	0	0
Freshwater																	
Total				6,119	23,160	5,772	13,223	7,081	5,788	10,406	20,187	13,500	16,133	11,069	9,467	12,632	12,173
Grand Total				6,119	23,160	5,772	13,223	7,081	5,788	10,406	20,187	13,500	16,133	11,069	9,467	12,632	12,173

Table 25.—Historic Arctic grayling harvests and catches in the Kotzebue Sound/Chukchi Sea sub-area, 1989-2001.

Area	Year													Average	
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	(91-00)	(96-00)
<u>Harvests</u>															
Salt Water	0	0	0	0	0	10	0	0	0	0	0	0	0	1	0
Kobuk R.	268	67	446	255	305	178	383	513	476	1,729	672	836	355	579	845
Noatak R.	912	269	817	105	322	407	185	1,136	872	42	412	223	620	452	537
Other Streams	150	286	631	248	234	186	263	393	555	0	97	45	111	265	218
Lakes	85	0	87	360	55	33	79	94	0	17	66	129	158	92	61
Freshwater															
Total	1,415	622	1,981	968	916	804	910	2,136	1,903	1,788	1,247	1,233	1,244	1,389	1,661
Grand Total	1,415	622	1,981	968	916	814	910	2,136	1,903	1,788	1,247	1,233	1,244	1,390	1,661
<u>Catches</u>															
Salt Water		0	0	0	0	10	0	0	0	0	0	0	0	1	0
Kobuk R.		790	1,535	1,593	1,717	1,593	5,146	2,469	2,815	5,280	6,680	5,753	4,103	3,458	4,599
Noatak R.		1,462	2,402	1,112	1,718	842	1,114	3,886	2,179	964	3,621	1,668	2,123	1,951	2,464
Other Streams		1,076	1,264	738	3,151	2,653	7,921	3,516	3,182	548	5,114	1,934	975	3,002	2,859
Lakes		0	174	1,548	642	374	1,560	1,306	216	404	66	376	171	667	474
Freshwater															
Total		3,328	5,375	4,991	7,228	5,462	15,741	11,177	8,392	7,196	15,481	9,731	7,372	9,077	10,395
Grand Total		3,328	5,375	4,991	7,228	5,472	15,741	11,177	8,392	7,196	15,481	9,731	7,372	9,078	10,395

approaching that of the Nome River, which has been the most heavily fished stream in the area and has a depressed grayling population.

Populations of grayling in the Kotzebue area are inaccessible by road and are lightly exploited. Grayling occur in almost all streams of the area, and in many of the lakes as well. Most grayling in this area are captured in association with wilderness float trips or as an alternate species in trips directed toward fishing for Dolly Varden or sheefish. Over the past five years the estimated harvest has been about 1,650 fish or about 16% of those captured (Table 25).

Prior to 1988, the daily bag limit for Arctic grayling in the NWMA was 15 with only 2 over 20 inches. In 1988, the BOF established a separate daily bag and possession limit for Arctic grayling in Northern Norton Sound of 5 per day, with only one over 15 inches. The effect of this change is reflected in harvest estimates that averaged about 4,300 grayling annually from 1980-1988, but dropped to about 1,550 from 1991-2001. Measurable increases in populations in the Fish and Niukluk rivers were likely due to this regulatory change. Both populations nearly doubled in abundance when compared to abundance estimates from the early 1990s.

Recent Fishery Performance

Seward Peninsula/Norton Sound Sub-area

Estimated harvests of Arctic grayling by sport anglers in the Seward Peninsula/Norton Sound area have been declining since a high of 5,121 reached in 1991. Since then, harvests have averaged about 1,100 per year (Table 24). Estimated harvests have ranged from about 300 fish about 1,600 fish. The harvest increased to 1,600 Arctic grayling in 1999 and was estimated at about 1,000 fish in 2001. The estimated catch of Arctic grayling tripled from 6,342 in 1996 to 20,117 in 1997, dropping in 1998 to 12,408 fish and rebounding to 16,000 in 1999. It was estimated at 11,000 in 2000, and 9,500 in 2001. It appears that catch and release practices are increasing in these grayling fisheries. The percentage of captured grayling that were harvested averaged about 17% from 1993 through 1996 and dropped to about 6% in 1997 and 2.4% during 1998. However, it has stabilized at about 10% over the past three years.

Current exploitation rates on most northwestern Alaska grayling populations are unknown, but since most fisheries are in remote areas with little effort, exploitation is believed to be light. Some estimates of exploitation in Nome area roadside streams are available by combining harvest data with abundance data. Using these data for years with abundance estimates, exploitation rates of Arctic grayling have been estimated to range from 10 to 20% in some streams during the early 1990s. More recent estimates for the Niukluk and Fish rivers suggest that exploitation rates in these streams are nearer 1%. These data suggest a change in angler motivation toward enjoyment of a quality fishing experience, away from harvest as a primary reason for Arctic grayling fishing.

Kotzebue Sub-area

In the Kotzebue/Chukchi Sea sub-area, harvests over the past five years have ranged between 900 and 2100 fish (Howe et al. 1996, 2000a-d; Walker et al. *In press*). Catches over the same period have ranged quite widely from about 16,000 in 1995 to about 7,200 in 1998. The percentage of catch harvested has ranged from about 37% in 1992 to 6% in 1995. It was about 8% in 1999, 13% in 2000 and 15% in 2001. Most grayling from this area are harvested in association with float trips or for variety, while fishing for other species. It is likely that harvests will remain relatively stable until participation in this area increases significantly.

Fishery Objectives and Management

Research on the status of resident Arctic grayling populations in the rivers accessible by the road system in northern Norton Sound has been ongoing for about 12 years. Arctic grayling in northwestern Alaska may live for more than 20 years and attain a large size. They spawn in the spring and the summer is spent feeding to recover condition in order to be able to spawn the next year. Data on population abundance, age, and size composition by river throughout this period has allowed the development of regulations tailored to individual rivers or groups of rivers that share population characteristics. Overall management objectives for these Arctic grayling populations are to maintain a minimum number of spawning sized fish (>15 inches) in populations where size composition has been documented, and to allow for population recovery in systems that have been stressed by over exploitation. Maintaining the opportunity to participate in high quality Arctic grayling fisheries is also an objective of management. The background daily bag and possession limits are 5 fish per day with only 1 over 15 inches. This bag limit is appropriate for drainages with Arctic grayling populations that have characteristics of lightly exploited populations. These characteristics include large average size and a high proportion of sexually mature fish that are seven years of age or older in the population. Abundance is related more to the river's size and flow characteristics, therefore, both abundance and population density vary by river. Rivers that share these characteristics and regulations include the Fish/Niukluk River system, the Eldorado, Kuzitrin and Sinuk rivers. On the other extreme are over exploited populations where abundance is very low. Rivers like the Nome and Solomon are in this category. These rivers are closed to fishing for Arctic grayling. Populations intermediate between these two categories include those in the Pilgrim and Snake rivers. These populations contain a smaller proportion of sexually mature fish, have been impacted somewhat by harvest, but Arctic grayling are still relatively abundant. In these rivers the regulations allow harvest of 2 Arctic grayling per day with only 1 over 15 inches.

MANAGEMENT GOALS

The ADF&G intent for the management objectives are to regulate Nome roadside Arctic grayling fisheries to maintain populations with characteristics that users presently consider to be producing a high quality sport fishery and maintain minimum spawning stock abundances. Each population will be managed to maintain a minimum number of Arctic grayling >15 inches in length (sexually mature fish).

MANAGEMENT OBJECTIVES FOR SPECIFIC RIVERS

Maintain a population of Arctic grayling > 15 inches in length in index sections of the following rivers at these levels:

<u>River</u>	<u>Section</u>	<u>Section Length</u>	<u>#>15 inches</u>
1. Niukluk River (Council to Casadepage River)		14 mi	3,500
2. Fish River (Cache Cr to lower end of canyon)		16 mi	4,500
3. Pilgrim River (7.5 mile section below bridge)		7.5 mi	350
4. Snake River (Boulder Creek to bridge)		12 mi	600
5. Sinuk River		25 mi	1,000
6. Nome River (Hobson Creek to weir)		26 mi	2,000

The numbers of fish by river section in rivers were determined from past estimates of abundance of Arctic grayling within the size ranges and sections noted (numbers 1-5), where it has been shown that populations are supporting existing levels of effort and harvest. For the Nome River, the number is a desired number of fish >15 inches that is believed would sustain a modest level of harvest based on the river's accessibility and the amount of Arctic grayling habitat available.

It is likely that levels of fishing effort will increase at some time in the future. Effort level is something that managers cannot control, and as this occurs, harvests and populations will continue to be monitored to ensure that the grayling stocks are not adversely impacted. Reduction of daily bag limits is a tool that may be needed in the future to decrease the risk of negatively impacting the size structure and abundance of Arctic grayling populations if increased effort levels result in harvests that may reduce populations below these threshold levels.

Providing diversity of opportunity in the form of the various harvest options for Arctic grayling is built into current management practices.

Management objectives have not been developed for remote Arctic grayling waters of the remainder of the Seward Peninsula or the Kotzebue sub-area. These waters are rarely visited by anglers, and populations are in pristine condition. The general regulations for these waters provide for a liberal daily bag and possession limit of 10 fish with no size limits. Until effort and harvests increase, it is likely that regulations will remain unchanged. A statewide Arctic grayling management plan is being developed by ADF&G that may address background regulations.

Fishery Outlook

Northwestern Alaska, particularly Seward Peninsula waters provide some of the best opportunities in the state to capture large sized Arctic grayling because many populations are dominated by large sexually mature fish. Populations are managed to maintain the reproductive potential of the stock by limiting the harvest of large fish. The result is a quality Arctic grayling fishing opportunity. The outlook for these fisheries to be maintained at current levels of participation and exploitation is favorable. Populations in the Fish and Niukluk rivers have recovered from relatively low levels of abundance in the early 1990s, and the outlook in these rivers is good.

Recent Board of Fisheries and Management Actions

In 1992 the daily bag and possession limit for Arctic grayling in the Pilgrim River was reduced to 2 per day with only 1 over 15 inches, and the Nome and Solomon rivers were closed to fishing for Arctic grayling by EO. In 1993, the daily bag and possession limit in the Snake River was made the same as that in the Pilgrim River. In the 1994 meeting, the BOF adopted these bag limit changes for the Snake and Pilgrim rivers into regulation. After a population assessment in the Nome River in 1997 found that the population had not increased after five years of emergency closure, the BOF adopted regulations closing the Nome and Solomon rivers to fishing for Arctic grayling. In 2000 the BOF also closed these rivers to subsistence fishing for Arctic grayling.

Current Issues

There is concern on the part of the public and ADF&G staff that populations of grayling in the vicinity of Nome that are road accessible, especially the Nome and Solomon rivers, have been over harvested and may not recover for many years. The Nome River population showed no increase over the past five years. An experimental restoration project in 1998 to increase survival of young-of-the-year Arctic grayling by rearing them in a gravel pit failed. Additional

restoration efforts are underway using a different rearing pond. Other road accessible populations would be vulnerable to over exploitation if fishing practices and motivations were to change, however, at this time other populations appear to be healthy, and able to sustain the current levels of harvest.

Ongoing Research and Management Activities

The ADF&G began an ongoing active effort to assess Arctic grayling populations in waters of the Seward Peninsula in 1989. Abundance and age and size compositions have been estimated for Arctic grayling in the Fish, Niukluk, Nome, Pilgrim, Snake and Sinuk rivers. These data in combination with harvest estimates and observed changes in abundance or size compositions have been used to guide ADF&G management activities. Special regulations in some streams and the closure of both the Solomon and Nome rivers to grayling have resulted. Assessments of the Nome River grayling population in 1997 and 2000 found that it had not recovered even with years of closure to sport fishing. This project has shown the need to close these rivers to all harvest of Arctic grayling. Stock assessment in the Fish and Niukluk Rivers in 1998 and 1999 found that grayling populations had increased significantly since the early 1990s. These changes are likely a delayed effect of regulation changes made in 1988. A management plan has been developed that addressed Nome roadside Arctic grayling fisheries (DeCicco 2002a), and threshold levels of abundance will be used as guidelines in the management of these fisheries.

KOTZEBUE SOUND SHEEFISH

Fishery Description and Historical Perspective

Within the NWMA, except for a small population of sheefish that resides in the Koyuk River of Norton Bay, spawning stocks of sheefish occur only in the Kobuk and Selawik rivers (Alt 1975).

The drainages of Kotzebue Sound are known for the large size of sheefish that are available to the sport angler. These remote high quality sport fisheries, and are considered by many to be some of the "crown jewels" of Alaskan freshwater sport fishing. Since the inception of ADF&G's Trophy Fish Program in 1967, all but one of the qualifying sheefish has come from the Kobuk River.

Kotzebue Sound sheefish are distributed throughout the nearshore estuarine areas of Kotzebue Sound. The major concentration is in Hotham Inlet but a few fish occur in the Sheshalik and Krusenstern areas as well as in southern Kotzebue Sound, especially in summer (Figure 13). Nearly all sheefish occupying the estuarine environment during summer are immature or nonspawning adults, while adult prespawning fish move upstream on the Kobuk and Selawik rivers to spawn just before freeze-up in the fall. The Kobuk River stock with 32,000 to 43,000 spawning in 1995-1996 (Taube 1997, Taube and Wuttig 1998) is the largest and most heavily utilized. They spawn upstream from the village of Kobuk, with the greatest observed concentrations between the Maneluk River and Beaver River. After spawning is complete in late September, fish disperse to downstream overwintering areas. Abundance of sheefish spawning in the Selawik River was estimated at about 5,200 fish during 1995 and 1996 (Underwood et al. 1998). Tag recoveries showed that these stocks mixed in Hotham Inlet winter habitats, but maintained fidelity to their spawning areas.

Sport fisheries for sheefish are managed by the Sport Fish Division of ADF&G. Subsistence fisheries are given priority. The commercial fishery and much of the subsistence harvest takes place through the ice while sport fisheries are mainly summer and fall activities. The same populations contribute to all harvests. The annual commercial sales of sheefish in Kotzebue




Figure 13.-Sheefish distribution in the NWMA.

have ranged from 200 to 850 fish since 1991 (Brennan et al. 1998, 1999, 2001). The magnitude of the subsistence harvest in the villages of the Kobuk River was estimated at about 7,000 in 1996, 9,800 in 1997, 5,350 in 1998 and 8,526 in 1999, and 7,446 in 2000 (Georgette and Utermohle 1997-2001, *In press*). In 2001, the estimated sheefish harvest was 3,838 fish (Table 26). Since subsistence practices have not changed appreciably in recent years, it is likely that Kobuk River subsistence harvests have been relatively stable at, or near, these levels. Winter gill net harvests from the fishery near Kotzebue were estimated at about 15,000 in 1995- 1996, and about 14,000 in 1996-1997 (Taube 1997, Taube and Wuttig 1998). During the winter of 2000/2001, a complete census of participants in the winter fishery documented the harvest at 14,533 (Savereide 2002). Sheefish are also taken by jigging lures under the ice in Hotham Inlet and Selawik Lake, but harvests in this fishery are undocumented.

The Sport Fish Division of ADF&G conducted studies of the ecology, movements, and growth of sheefish between 1966 and 1979. Much of this work was conducted in northwestern Alaska and was summarized by Alt (1987). After some familiarization work in 1994, ADF&G Division of Sport Fish in cooperation with the National Park Service (NPS), began a project to estimate abundance of sheefish spawning in the Kobuk River. This project continued through 1997 and established base line estimates on spawner abundance, age, size and sex composition of the spawning population. Tag recovery data indicated that, although some sheefish spawned in consecutive years, most spawned every other year. The abundance of sheefish spawning in the Selawik River was estimated at 5,200 and 5,150 in 1995 and 1996 by the U. S. Fish and Wildlife Service (Underwood et al. 1998).

Most sheefish sport fishing effort occurs on the Kobuk River spawning population. Most of the area-wide subsistence harvest and the entire commercial harvest of sheefish occur on the entire (spawners and nonspawners) population. When taken in isolation, recent sport harvests of about 800 fish annually are easily sustainable. Although spawner abundances have recently been estimated, the total size of the area wide population is not known, and the sport harvest must be viewed in relation to other ongoing harvests. It was always assumed that subsistence harvests were much greater than either commercial or sport harvests, and recent data support this assumption. Taube and Wuttig (1998) estimated the under ice gill net harvest in Hotham Inlet near Kotzebue at 13,704 fish in the winter of 1997/98. Savereide (2002) estimated the winter 2000/01 harvest at 14,533 sheefish. In order to ensure sustained yields from these population(s), a management approach involving the subsistence and commercial fisheries for sheefish is recommended. Sheefish are very fecund fish with some large females containing over 400,000 eggs. Such populations may be subject to episodic recruitment events depending on environmental conditions. If spawner abundances are maintained above some threshold level, intermittent years of good recruitment should carry the population through years when environmental conditions are less favorable.

Recent Fishery Performance

Estimated annual sport harvests of sheefish by anglers in northwestern Alaska since 1977 have fluctuated from a high of about 1,900 to a low of about 151 with an average annual harvest of about 900 fish over the past 10 years (Table 27). The harvest in 2001 was estimated at 1,300 sheefish, and the most recent five year (1995-1999) average harvest has been about 730 sheefish. In addition to harvests, catches have been estimated through the SWHS since 1990. Estimates of sheefish catch (which includes fish that are kept and those released) for the past five years was about 3,000 fish, indicating that about 75% of all sheefish captured in northwestern Alaska by

Table 26.—Reported subsistence sheefish harvests, Kotzebue District, 1966-2001^a.

Year	Number of Fishermen Interviewed	Reported Harvest	Average Catch Per Fisherman	Hotham Inlet Winter Harvest
1966-67	135	22,400	166	
1967-78	146	31,293	214	
1968-69	144	11,872	82	
1970	168	13,928	83	
1971	155	13,583	88	
1972	79	3,832	49	
1973	65	4,883	75	
1974	58	1,062	18	
1975	69	1,637	24	
1976	57	966	17	
1977	95	1,810	19	
1978	95	1,810	19	
1979	75	3,985	53	
1980	74	3,117	42	
1981	62	6,651	107	
5/82-4/83 ^b	430	4,704	36	
5/83-4/84 ^b	27	764	28	
5/84-9/84 ^b	30	2,803	93	
1985 ^c	2	60	30	
1986 ^{b,c}	72	721	10	
1987 ^c	46	276	6	
1988 ^{c,d}				
1989 ^d				
1990 ^d				
1991	40	2,180	55	
1992	43	2,821	66	
1993 ^d				
1994	171	3,181 ^e	84	
1995	314	9,465 ^e	24.6	15,161 ^f
1996	389	6,465 ^e	18	13,704 ^f
1997	338	9,805 ^e	24.6	
1998	435	5,350 ^e	13.6	
1999	191	8,256 ^e	18.6	
2000	237	7,446 ^e	16.6	14,533 ^f
2001	257	3,838 ^e	10.6	

^a Due to limited survey effort during many years, total catch and effort are minimums and are not comparable among years. Data from Brennan et al. 1999.

^b Summer harvests only.

^c Data from fall subsistence salmon surveys may include summer and winter harvests.

^d Subsistence sheefish harvests not documented.

^e Reported harvests from Kobuk River villages only.

^f Data from Sport Fish Division harvest estimates.

Table 27.—Historic sheefish sport fish harvests and catches from northwest Alaska waters, 1977-2001.

Year	Kotzebue/Chukchi Sea Sub-area								
	Sheefish			Kobuk River			Selawik River		
	Harvest	Catch	% Harvested	Harvest	Catch	% Harvested	Harvest	Catch	% Harvested
1977	656								
1978	506								
1979	709								
1980	1,713								
1981	1,263			1,015					
1982	2,222			1,886					
1983	2,079			1,448					
1984	3,050								
1985	1,645			1,330					
1986	3,363			1,590					
1987	1,836			865					
1988	964			964					
1989	629			131					
1990	151	403	37	151	336	44.9	0	0	0
1991	603	1,616	37	579	1,568	36.9	24	48	50.0
1992	1,904	3,678	52	627	2,034	30.8	411	411	100.0
1993	1,029	2,273	45	395	1,074	36.8	111	111	100.0
1994	564	958	59	135	386	35.0	95	95	100.0
1995	1,142	3,270	35	748	2,669	28.0	38	47	80.9
1996	485	3,183	15	360	2,850	12.6	94	271	34.7
1997	906	2,341	39	318	1,334	23.8	108	108	100.0
1998	414	924	45	145	617	23.5	148	186	79.6
1999	635	5,134	12	621	5,070	12.2	nd	nd	nd
2000	1,201	3,372	36	362	2,338	15.5	0	0	0
2001	1,305	5,146	25.4	552	4,105	13.4	0	0	0.0
91-00 Avg	888	2,675	37.5	429	1,994	25.5	114	142	71.7
96-00 Avg	728	2,991	29.3	361	2,442	17.5	88	141	53.6

nd = no data

sport anglers are released. In a 1997 hook and release study, the mortality of sheefish caught and released on sport fishing gear was found to be low, 3.3% for treble hook lures, and 1.7% for single hook lures (Stuby and Taube 1998). Overall mortality was 2.4%. Approximately 67% of all sheefish caught on the Kobuk River over the past ten years have been released. The Kobuk River is probably the most popular sheefish destination in North America, and people from the world over come there to fish for this unique species. In spite of the world-wide reputation of this destination, the level of fishing effort is still quite low. An estimated 286 anglers fished for a total of 924 angler-days on the Kobuk River during 2001. This was about 16% of the overall freshwater sport fishing effort in the Kotzebue sub-area (5,900 angler-days) in 2001.

Fishery Objectives and Management

The Kobuk River sheefish fishery is managed to maintain opportunity to participate in this unique high-quality sport fishery while keeping harvests from spawning areas low. In order to accommodate local use of this resource downstream from major spawning areas, the daily bag limit is generous downstream from the Mauneluk River (10 sheefish per day). In the spawning area upstream from the Mauneluk River, only two fish per day are allowed to be harvested or in possession. The majority of anglers visiting the Kobuk River for sheefish use the area upstream from the Mauneluk River while the majority of local use is closer to villages downstream from the Mauneluk River.

Fishery Outlook

The outlook for sheefish fisheries in northwestern Alaska is good for the immediate future. Although overall harvest levels are substantial, populations appear to be healthy, spawner abundances are high and sport harvests are low.

Recent Board of Fisheries and Management Actions

During 1988, the Board of Fisheries adopted the current regulations for sheefish in the waters of northwestern Alaska: 10 fish per day and 10 in possession, with an exception for the Kobuk River upstream of the Mauneluk River where only 2 sheefish may be caught per day or possessed. The ADF&G believes that these regulations are sufficient to allow ample opportunity for sport fishing, yet keeping harvests low. The 10 fish limit in the lower Kobuk River and the remainder of the management area is liberal enough to allow local fishermen who choose to catch sheefish or food on sport fishing tackle the opportunity to take sheefish without the need to fish with nets. During the December 2000 meeting, the BOF reduced the sheefish daily bag and possession limit in spawning areas on the Selawik River to two sheefish. This made regulations on the Selawik River consistent with those already in place on the Kobuk. Another sheefish proposal that would have eliminated catch and release of sheefish was not supported.

Current Issues

Local native residents of Kobuk River villages have expressed concern over some practices of sport anglers on the upper Kobuk River in the vicinity of the sheefish spawning grounds. Catch and release fishing is considered by some local residents to be disrespectful and damaging to the fish. The discarding of filleted carcasses in the water is thought to drive other sheefish away from the area. The ADF&G Division of Subsistence investigated local concerns in the upper Kobuk River in 1986 and determined that some concerns could be addressed if sport fishers were more aware of local customs and culture. Catch and release fishing is viewed as a conservation tool by ADF&G and by many anglers and although sheefish may be sensitive to rough handling, the department has demonstrated that they can be released without significant mortality. An

educational brochure explaining proper hook and release techniques for sheefish and concerns of local residents has been developed in association with the NPS. This brochure has been made available to those fishing on the upper Kobuk River. It is hoped that with proper handling, impacts of catch and release fishing to the spawning population can be minimized.

Because the subsistence component of the harvest is high, and some sheefish spawning areas are located within federally managed lands, the possibility exists that federal subsistence management in these areas may affect sport fishing opportunity there. This issue may be addressed in the future as the USFWS Office of Subsistence Management becomes more involved with active management of fisheries in remote areas of Alaska.

Ongoing Research and Management Activities

The department believes that recent research conducted cooperatively with the USFWS and the NPS has provided substantial background data on spawner abundance for the two stocks comprising the Kobuk-Selawik sheefish population. These data will be used as a base line to which future population assessments can be compared. Additional studies to monitor harvests may be planned. Current low levels of sport fishing harvest are unlikely to affect sustained yields of this species in northwestern Alaska. A study to document the winter subsistence harvest of sheefish in Kotzebue Sound was completed in 2001 (Savereide 2002). This study was funded through the USFWS Office of Subsistence Management.

NORTHWESTERN ALASKA NORTHERN PIKE

Fishery Description and Historical Perspective

Northern pike are present throughout the northern regions of the world. They are primarily a freshwater resident species, but are known to enter weakly brackish waters in the Baltic and in some other areas. The known distribution of northern pike in northwestern Alaska is shown in Figure 14.

Northern pike occur in most of the lakes and flowing waters of the Noatak and Kobuk rivers and are particularly common in wetlands of the lower reaches, delta areas, and lakes in lowland areas adjacent to these rivers. Northern pike are also common residents of the waters along the western shores of Hotham Inlet, Selawik Lake and the entire Selawik lowland area. They occur in the lower portions of the Buckland River drainage, and may be present in some lakes and streams on the northern Seward Peninsula. On the remainder of the Seward Peninsula, northern pike are common residents of Imuruk Basin and the middle and lower reaches of the Pilgrim and Kuzitrin rivers. These two adjacent drainages form a large interconnected wetland area (approximately 380 km²) in their lower reaches. In addition there is another large wetland area (approximately 650 km²) farther upstream in the Kuzitrin River drainage. Northern pike are also found in the Fish River drainage and have even been observed in the fast clear waters of the Niukluk River downstream from Council. They occur in the Koyuk River and may be present in the Kwik River near Moses Point, but are not known to be present in other Norton Sound drainages.

The majority of northern pike harvested in northwestern Alaska are taken for subsistence. Few community harvest estimates are available, however, in 1986, 5,750 northern pike were estimated to have been harvested by the community of Kotzebue. During the mid 1980s a commercial freshwater fishery occurred near Selawik. In 1985, the USFWS estimated that the spring subsistence/commercial harvest (only some of the fish were sold) of northern pike was between 5,671 and 9,138 fish. Currently, without the commercial fishery, the annual harvest at




Figure 14.-Northern pike distribution in the NWMA.

Selawik is still likely several thousand northern pike. Northern pike are also harvested by residents of the lower Kobuk River villages of Noorvik and Kiana, and the residents of Teller who fish in Imuruk Basin drainages. Additional harvests of northern pike may take place near other area villages. The total annual northwestern Alaska northern pike subsistence harvest is likely 10,000 to 15,000 fish.

Sport fisheries for northern pike occur around Kotzebue in lakes in the lower Noatak and Kobuk river drainages, but participation and harvests are low. About 195 northern pike were estimated to have been harvested in the Kotzebue sub-area of NWMA during 1998, 193 in 1999, 357 in 2000, and 471 in 2001 (Table 28). In the Seward Peninsula/Norton Sound sub-area, virtually the entire harvest of northern pike comes from Imuruk Basin drainages, and most of that from the Kuzitrin or lower Pilgrim rivers. Estimated northern pike harvests have been 75 in 1998, 355 in 1999, 420 in 2000, and in 2001, 349 northern pike were estimated harvested in this fishery.

Recent Fishery Performance

Estimated harvests of northern pike by sport anglers on the Seward Peninsula have averaged about 498 fish over the past 10-years (1991-2000). The largest annual harvest was estimated at nearly 2,000 in 1990 (Table 28). The average annual harvest for the past five years was 388 fish, and the 2001 estimated harvest was 349 fish. Estimates of catch (which includes fish that are kept and those released) since 1991 indicate that about 78% of all pike caught in the past ten years have been released. It is assumed that anglers are selectively retaining larger sized northern pike. Most of the harvest of pike on the Seward Peninsula takes place in the Pilgrim or Kuzitrin River drainages. During 1992 and 1993 the abundance of northern pike in the lower Pilgrim and Kuzitrin rivers was estimated at about 10,000 fish over 300 mm (12 in) in length for the portion of the population inhabiting these rivers from the road crossings downstream to their confluence (Burkholder 1993, 1994). Maximum sustained yield (MSY) for northern pike populations has been estimated to be between 12% and 24% for interior Alaskan lakes (Pearse and Hansen 1993), and annual harvests of about 15% are thought to be sustainable. Current exploitation of the Pilgrim-Kuzitrin population appears to be less than 5%, which is well within what are thought to be sustainable levels.

Estimated sport harvests of northern pike in the Noatak-Kobuk-Selawik area of northwestern Alaska have averaged about 283 fish over the last 10-years (Table 28). Estimated harvests reached a high of 2,752 fish in 1986, and a low of 64 fish in 1989. The harvest in 2000 was about 357 fish, the highest harvest since 1993, and the 2001 harvest was estimated at 122 northern pike. The average annual harvest for the past five years has also been about 200 fish. Since assessment of northern pike populations has not been carried out in this area of northwestern Alaska, the health of populations and relative influence of harvests can only be inferred by comparing the area to other parts of Alaska. The amount of suitable northern pike habitat in the Noatak-Kobuk-Selawik area is much greater (by approximately 50 times) than that available to northern pike in the Pilgrim-Kuzitrin area, and the sport harvests are lower. It is unlikely that sport fisheries are adversely impacting northern pike populations, even when taken in addition to subsistence harvests.

Fishery Management Objectives

There are no specific management objectives for northern pike fisheries in NWMA. Regulations are liberal and management is structured to encourage participation. Liberal regulations provide the opportunity for rural residents to harvest northern pike with rod and reel within the sport fishing regulatory framework. Baseline data exist for the Pilgrim/Kuzitrin portion of the Imuruk

Table 28.—Historic northern pike sport fish harvests and catches in the NWMA by sub-area, 1977-2001.

Year	Seward Peninsula/Norton Sound sub-area					Kotzebue/Chukchi Sea sub-area				
	Number	Effort	Northern Pike		%	Number	Effort	Northern Pike		%
	of	Angler	Harvest	Catch		of	Angler	Harvest	Catch	
	Anglers	Days			Anglers	Days	Harvest			
1977		7,828	302			3,487	147			
1978		8,379	389			4,997	389			
1979		8,725				2,593	527			
1980		7,958	284			3,841	852			
1981		10,879	303			5,284	465			
1982		13,198	210			6,906	454			
1983		16,944	798			7,963	1,262			
1984	1,597	17,436	208			696	7,791	312		
1985	2,854	19,919	56			1,788	6,701	383		
1986	2,872	18,107	699			1,570	6,313	2,752		
1987	2,528	20,413	906			2,090	9,288	813		
1988	2,661	20,278	564			959	5,279	1,565		
1989	2,560	17,692	648			1,028	4,932	64		
1990	2,686	21,799	1,957	4,145	47	991	3,782	320	1,730	18
1991	3,236	23,622	1,429	4,257	34	1,606	9,543	394	1,879	21
1992	3,540	22,684	479	3,742	13	1,421	6,145	333	1,666	20
1993	3,134	18,930	537	2,117	25	1,575	7,809	559	2,209	25
1994	3,016	18,922	376	1,731	22	1,100	6,036	287	1,488	19
1995	3,719	19,647	215	1,856	12	1,957	8,495	256	1,421	18
1996	2,958	13,783	728	3,239	22	1,407	5,571	112	1,423	8
1997	2,773	13,850	363	2,188	17	824	3,729	145	657	22
1998	3,206	13,616	75	452	17	1,089	3,801	195	1,104	18
1999	3,124	15,006	355	2,217	16	1,313	6,771	193	1,869	10
2000	2,713	18,559	420	1,317	32	1,387	7,129	357	1,249	29
2001	2,371	10,955	349	3,276	11	1,177	5,904	122	337	36
Avg (91-00)	3,142	17,862	498	2,312	21	1,368	6,503	283	1,497	19
Avg (96-00)	2,955	14,963	388	1,883	21	1,204	5,400	200	1,260	17

Basin population complex. In northern Norton Sound, rod and reel has been recognized by the BOF as a subsistence method, but regulations are the same as those for sport fishing. Because of the proximity of Nome, it is likely that this population will be the first in the NWMA to require more restrictive regulations as the human population in the Nome area grows and participation in the fishery increases. Harvests are monitored through the SWHS. If large changes in harvest or participation occur, additional stock assessment work will be conducted.

Fishery Outlook

Changes in fishing regulations for northern pike in NWMA are not expected in the near future. Pike populations are largely unexploited, and both participation and harvest are low.

Recent Board of Fisheries and Management Actions

There have been no recent BOF or management actions concerning northern pike. The current daily bag and possession limit in the NWMA is 10 fish with no size limit.

Current Issues

There are no current issues regarding northern pike in the NWMA. Harvest level will continue to be monitored through the SWHS. If harvests increase dramatically, additional research may be undertaken.

Ongoing Research Activities

There are no current research activities associated with northern pike in the NWMA.

ACKNOWLEDGMENTS

I thank Sara Case, Region III publications technician for assistance in final report preparation.

LITERATURE CITED

- AAC (Alaska Administrative Code). 2001. 5AAC 75.003. In Alaska Fish and Game Laws and Regulations Annotated 2001-2002 Edition. Lexis Publishing, NY.
- Alaska Department of Fish and Game (ADF&G). 1984. Sport Fish Survey. Booklet published by Alaska Department of Fish and Game, Division of Sport Fisheries. Juneau, Alaska.
- Alaska Department of Fish and Game (ADF&G). 1986. Alaska Habitat Management Guide, Arctic Region. Vol. II: Distribution, abundance, and human use of fish and wildlife. Division of Habitat, Alaska Department of Fish and Game, Juneau.
- Alaska Department of Labor (ADL). 1991. Alaska Population Overview 1990 Census and Estimates. Demographic Unit, Research and Analysis. Juneau, Alaska. 144 pp.
- Alt, K. T. 1975. A life history study of sheefish and whitefish in Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration. Annual Performance Report. 1974-1975, Project F-9-7, 16(R-II): 19 pp.
- Alt, K. T. 1978. Inventory and cataloging of sport fish and sport fish waters of western Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Volume 19, Study G-I-P, Annual Performance Report 1977-1978.
- Alt, K. T. 1984. Inventory and cataloging of sport fish and sport fish waters of western Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, F-9-16, Volume 25, Study G-I, Annual Performance Report 1983-1984.
- Alt, K. T. 1987. Review of sheefish (*Stenodus leucichthys*) studies in Alaska. Alaska Department of Fish and Game, Fishery Manuscript No. 2, Juneau.

LITERATURE CITED (Continued)

- Brennan, E. L., F. J. Bue, C. F. Lean and T. L. Lingnau. 1998. Annual Management Report 1997, Norton Sound-Port Clarence-Kotzebue. Regional Information Report No. 3A98-28. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Anchorage, Alaska.
- Brennan, E. L., C. F. Lean, F. J. Bue and T. Kohler. 1999. Annual Management Report 1998, Norton Sound-Port Clarence-Kotzebue. Regional Information Report No. 3A99-32. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Anchorage, Alaska.
- Brennan, E. L., C. F. Lean, F. J. Bue and T. Kohler. 2001. Annual Management Report 1999, Norton Sound-Port Clarence-Kotzebue. Regional Information Report 3A01-05. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage, Alaska.
- Brennan, E. L., F. J. Bue, J. Menard and T. Kohler. 2002. Annual Management Report 2000, Norton Sound-Port Clarence-Kotzebue. Regional Information Report 3A02-02. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage, Alaska.
- Burkholder, A. 1993. Abundance and length-at-age composition of northern pike near the confluence of the Pilgrim and Kuzitrin rivers, 1992. Alaska Department of Fish and Game, Fishery Data Series No. 93-16, Anchorage.
- Burkholder, A. 1994. Abundance and length composition of northern pike near the confluence of the Pilgrim and Kuzitrin rivers, 1992-1993. Alaska Department of Fish and Game, Fishery Data Series No. 94-20, Anchorage.
- DeCicco, A. L. 1990. Seward Peninsula Arctic grayling study 1989. Alaska Department of Fish and Game, Fishery Data Series No. 90-11, Anchorage.
- DeCicco, A. L. 1991. Seward Peninsula Arctic grayling study 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-24, Anchorage.
- DeCicco, A. L. 1992. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, Alaska during 1991, Alaska Department of Fish and Game, Fishery Data Series No. 92-13, Anchorage.
- DeCicco, A. L. 1992. Assessment of Dolly Varden overwintering in selected streams of the Seward Peninsula, Alaska, during 1991. Alaska Department of Fish and Game, Fishery Data Series No. 92-11, Anchorage.
- DeCicco, A. L. 1993. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, Alaska during 1992., Alaska Department of Fish and Game, Fishery Data Series No. 93-36, Anchorage.
- DeCicco, A. L. 1993. Assessment of Dolly Varden overwintering in selected streams of the Seward Peninsula, Alaska, during 1992. Alaska Department of Fish and Game, Fishery Data Series No. 93-20, Anchorage.
- DeCicco, A. L. 1994. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, Alaska during 1993. Alaska Department of Fish and Game, Fishery Data Series No. 94-12, Anchorage.
- DeCicco, A. L. 1995. Assessment of selected stocks of Arctic grayling in streams and a survey of Salmon Lake, Seward Peninsula, 1994. Alaska Department of Fish and Game, Fishery Data Series No. 95-19, Anchorage.
- DeCicco, A. L. 1996. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, 1995. Alaska Department of Fish and Game, Fishery Data Series No. 96-21, Anchorage.
- DeCicco, A. L. 1997. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-15, Anchorage.
- DeCicco, A. L. 1998. Assessment of selected stocks of Arctic grayling in streams of the Seward Peninsula, 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-19, Anchorage.
- DeCicco, A. L. 1999. Niukluk River Arctic grayling stock assessment, Seward Peninsula, 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-23, Anchorage.
- DeCicco, A. L. 2001. Over-wintering areas used by Dolly Varden in the Nome and Solomon, and Bonanza rivers, Seward Peninsula, Alaska 2000/2001. Alaska Department of Fish and Game, Fishery Data Series No. 01-25, Anchorage.

LITERATURE CITED (Continued)

- DeCicco, A. L. 2002a. Fishery Management Plan for Arctic grayling sport fisheries along the Nome Road System, 2001-2004. Alaska Department of Fish and Game, Fishery Management Report No. 02-03, Anchorage.
- DeCicco, A. L. 2002b. Stock assessment of Arctic grayling in the Nome River and age validation of Arctic grayling in the Eldorado River, Seward Peninsula, Alaska 2000. Alaska Department of Fish and Game, Fishery Data Series No. 02-01, Anchorage.
- DeCicco, A. L. and M. J. Wallendorf. 2000. Stock assessment of Arctic grayling in the Fish River, Seward Peninsula, Alaska 1999. Alaska Department of Fish and Game, Fishery Data Series No. 00-29, Anchorage.
- DeCicco, A. L. and R. M. Barnes. 1992. Listing of guiding services for recreational fishing in the Arctic-Yukon-Kuskokwim (AYK) Region. Alaska Department of Fish and Game, Special Publication No. 92-3, Anchorage.
- Foote, D. C. and H. A. Williamson. 1966. A human geographical study. Pages 1041-1111 in Environment of the Cape Thompson region, Alaska. N. Wilimovsky and J. Wolfe, eds. U. S. Atomic Energy Commission.
- Georgette, S. and C. Utermohle. 1997. Subsistence salmon harvest summary, northwest Alaska 1996. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Georgette, S. and C. Utermohle. 1998. Subsistence salmon harvest summary, northwest Alaska 1997. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Georgette, S. and C. Utermohle. 1999. Subsistence salmon harvest summary, northwest Alaska 1998. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Georgette, S. and C. Utermohle. 2000. Subsistence salmon harvest summary, northwest Alaska 1999. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Georgette, S. and C. Utermohle. 2001. Subsistence salmon harvest summary, northwest Alaska 2000. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Georgette, S., D. Caylor, and S. Tahbone. *In press*. Subsistence salmon harvest summary, northwest Alaska 2001. Alaska Department of Fish and Game, Division of Subsistence, Nome.
- Gryska, A. *In prep*. Abundance and age, and length composition of Arctic grayling in the Snake River 2001. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Kretsinger, C. 1987. Fishery inventory of lakes and streams in the Kigluaik mountains and Imuruk Basin watershed (Seward Peninsula). Unpublished paper, U.S. Bureau of Land Management. Kobuk District Office, Fairbanks.
- Howe A. H., G. Fidler and M. Mills. 1995. Harvest, catch and participation in Alaska sport fisheries during 1994. Alaska Department of Fish and Game. Fishery Data Series No. 95-24. 212 pp.
- Howe, A. L., G. Fidler, A. E. Bingham, and M. J. Mills. 1996. Harvest, catch, and participation in Alaska sport fisheries during 1995. Alaska Department of Fish and Game, Fishery Data Series No. 96-32, Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001a. Revised Edition: Harvest, catch, and participation in Alaska sport fisheries during 1996. Alaska Department of Fish and Game, Fishery Data Series 97-29 (revised), Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001b. Revised Edition: Harvest, catch, and participation in Alaska sport fisheries during 1997. Alaska Department of Fish and Game, Fishery Data Series 98-25 (revised), Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001c. Revised Edition: Participation, catch, and harvest in Alaska sport fisheries during 1998. Alaska Department of Fish and Game, Fishery Data Series 99-41 (revised), Anchorage.
- Howe, A. L., G. Fidler, C. Olness, A. E. Bingham, and M. J. Mills. 2001d. Participation, catch, and harvest in Alaska sport fisheries during 1999. Alaska Department of Fish and Game, Fishery Data Series 01-8, Anchorage.
- Jennings, G. B., K. Sundet, A. E. Bingham, and H.K. Sigurdsson. *In prep*. Participation, catch, and harvest in Alaska sport fisheries during 2001. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.

LITERATURE CITED (Continued)

- Mills, M. J. 1979. Alaska statewide sport fish harvest studies. 1977 data. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1977-1978. Project F-9-11, 20 (SW-1): 112 pp.
- Mills, M. J. 1980. Alaska statewide sport fish harvest studies. 1978 data. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1979-1980. Project F-9-12, 21 (SW-1): 65 pp.
- Mills, M. J. 1981. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1980-1981. Project F-9-13, 22 (SW-1): 78 pp.
- Mills, M. J. 1982. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1981-1982. Project F-9-13, 23 (SW-1): 115 pp.
- Mills, M. J. 1983. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1982-1983. Project F-9-14, 24 (SW-1): 118 pp.
- Mills, M. J. 1984. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1983-1984. Project F-9-16, 25 (SW-1): 122 pp.
- Mills, M. J. 1985. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1984-1985. Project F-9-17, 26 (SW-1): 88 pp.
- Mills, M. J. 1986. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1985-1986. Project F-9-18, 27 (SW-1): 137 pp.
- Mills, M. J. 1987. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1986-1987. Project F-9-19, 28 (SW-1): 91 pp.
- Mills, M. J. 1988. Alaska statewide sport fisheries harvest report 1987. Alaska Department of Fish and Game, Fishery Data Series No. 52, Juneau.
- Mills, M. J. 1989. Alaska statewide sport fisheries harvest report 1988. Alaska Department of Fish and Game, Fishery Data Series No. 122, Juneau.
- Mills, M. J. 1990. Harvest and participation in Alaska sport fisheries during 1989. Alaska Department of Fish and Game, Fishery Data Series No. 90-44, Anchorage.
- Mills, M. J. 1991. Harvest, catch and participation in Alaska sport fisheries during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-58, Anchorage.
- Mills, M. J. 1992. Harvest, catch, and participation in Alaska sport fisheries during 1991. Alaska Department of Fish and Game, Fishery Data Series Number 92-40, Anchorage.
- Mills, M. J. 1993. Harvest, catch, and participation in Alaska sport fisheries during 1992. Alaska Department of Fish and Game, Fishery Data Series Number 93-42, Anchorage.
- Mills, M. J. 1994. Harvest, catch, and participation in Alaska sport fisheries during 1993. Alaska Department of Fish and Game, Fishery Data Series Number 94-28, Anchorage.
- Mills, M. J. and A. L. Howe. 1992. An evaluation of estimates of sport fish harvest from the Alaska statewide mail survey. Alaska Department of Fish and Game, Special Publication No. 92-2, Anchorage, Alaska.
- National Park Service (NPS). 1984. Kobuk Valley National Park draft statement for management. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- National Park Service (NPS). 1985. Gates of the Arctic National Park General Management Plan. USDI: NPS,
- Pearse, G. A. and P. A. Hansen. 1993. Estimates of sustainable yield for the northern pike populations in George, Volkmar, T, and Harding lakes. Alaska Department of Fish and Game, Fishery Manuscript No. 93-1, Anchorage.
- Phillips, R. B., L. I. Gaudex, K. M. Westrich and A. L. DeCicco. 1999. Combined phylogenetic analysis of ribosomal ITS1 sequences and new chromosome data supports three subgroups of Dolly Varden char (*Salvelinus malma*). Canadian Journal of Fisheries and Aquatic Sciences 56:1504-1511.

LITERATURE CITED (Continued)

- Sarrio, R. and B. Kessel. 1966. Human ecological investigations at Kivalina. Pages 969-1040 in Environment of the Cape Thompson region, Alaska. N. Wilimovsky and J. Wolfe, eds. U. S. Atomic Energy Commission.
- Savereide, J. W. 2002. Under-ice gill net harvest of sheefish in Hotham Inlet in 2000-2001. Alaska Department of Fish and Game, Fishery Data Series No. 02-04, Anchorage.
- Stuby, L. and T. Taube 1998. Mortality of sheefish captured and released on sport fishing gear in the Kobuk River, 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-15. Anchorage.
- Taube, T. T. 1997. Abundance and composition of sheefish in the Kobuk River, 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-1, Anchorage.
- Taube, T. T. and K. Wuttig. 1998. Abundance and composition of sheefish in the Kobuk River, 1997. Alaska Department of Fish and Game, Fishery Manuscript Report No. 98-3, Anchorage.
- Underwood, T., K. Whitten, and K. Secor. 1998. Population characteristics of spawning Inconnu (sheefish) in the Selawik River, Alaska, 1993-1996, Final Report. Alaska Fisheries Technical Report Number 49, U.S. Fish and Wildlife Service, Fairbanks Fishery Resource Office, Fairbanks, Alaska.
- U.S. Army Corps of Engineers, Alaska District. 1967. Harbors and Rivers in Alaska. Interim Report No. 6. Northwestern Alaska.
- Vincent-Lang, D., M. Alexandersdottir, and D. McBride. 1993. Mortality of coho salmon caught and released using sport tackle in the Little Susitna River, Alaska. Fisheries Research 15: 339-356
- Walker, R. J., C. Olnes, K. Sundet, A. L. Howe, and A. E. Bingham. *In prep.* Participation, catch, and harvest in Alaska sport fisheries during 2000. Alaska Department of Fish and Game, Fishery Data Series, Anchorage.
- Wild and Scenic Rivers Act. 1968. U.S. Public Law 90-542. Amended December, 1980 with Public Law 96-603-605.
- Wuttig, K. G. 1998. Escapement of chinook salmon in the Unalakleet River in 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-8, Anchorage.

APPENDIX A

Appendix A.—National Wild and Scenic Rivers in the NWMA.

- Streams Within The National Park System

Kobuk River. The portion within the Gates of the Arctic National Park and Preserve.

Noatak River. The river from its source in the Gates of the Arctic National Park to its confluence with the Kelly River in the Noatak National Preserve.

Salmon River. The portion within the Kobuk Valley National Park.

- Streams Within The National Wildlife Refuge System

Selawik River. The portion from a fork of the headwaters in township 12N, Range 10E, Kateel River meridian to the confluence of the Kugarak River; within the Selawik National Wildlife Refuge.

- Streams Located Outside National Parks and Refuges

Unalakleet River. The segment of the main stem from the headwaters in township 12S, Range 3W, Kateel River meridian extending downstream approximately 65 miles to the western boundary of township 18S, range 8W.

APPENDIX B

Appendix B.–Northwestern Area sport fishing regulations summary for 2000.

NORTHWESTERN ALASKA

SEASONS

Entire year for all species except halibut. Halibut season is February 1-December 31.

BAG, POSSESSION, AND SIZE LIMITS GENERAL REGULATIONS

The general regulations for all waters of the Northwestern Area are listed below..Special regulations for individual water bodies appear afterward (at bottom of page).

Species	Daily Bag	Possession & Size Limit
King salmon 20 “ or greater	1	
Less than 20 “	10	
Other salmon	10	(no size limit)
Arctic char/Dolly Varden (all lakes)	2	(no size limit)
Flowing and salt water	10	(only 2 over 20 inches)
Lake trout	4	(no size limit)
Arctic grayling	10	(no size limit)
Sheefish	10	(no size limit)
Northern pike	10	(no size limit)
Burbot	15	(no size limit)
Halibut	2 per day	4 in possession, no size limit
Outlet stream (Pilgrim River) 300 feet downstream from - Salmon Lake		Closed to salmon fishing
Shellfish		

-continued-

SPECIAL REGULATIONS

KOBUK RIVER DRAINAGE (upstream of the mouth of the Mauneluk River):

- 1. Sheefish:** Daily bag and possession limit is 2 fish, no size limit

NORTHERN NORTON SOUND (all waters draining into Norton Sound from Cape Darby to Cape Prince of Wales (see map):

- 1. Chum salmon:** Daily bag and possession limit is 3 fish, not size limit
All freshwater drainages and marine waters between the west bank of the Sinuk River and Topkok Head (see map) are *closed to chum salmon fishing.*
- 2. Coho salmon:** Daily bag and possession limit is 3 fish, no size limit
- 3. Sockeye salmon:** Daily bag and possession limit is 3 fish, not size limit
- 4. Pink salmon:** Daily bag and possession limit is 10 fish, no size limit
- 5. Arctic grayling:** Daily bag and possession limit is 5 fish, **only 1 over 15 inches**
- 6. Salmon Lake, its tributaries, and the lake outlet are:** *Closed to salmon fishing.*
- 7. Nome River:** *Closed to fishing for Arctic grayling*
- 8. Pilgrim River drainage:** Arctic grayling daily bag and possession limit is 2 fish, **only 1 over 15 inches.**
- 9. Snake Rive drainage:** Arctic grayling daily bag and possession limit is 2 fish, **only 1 over 15 inches.**
- 10. Solomon River:** *Closed to fishing for Arctic grayling*

UNALAKLEET RIVER DRAINAGE:

- 1. Arctic grayling:** Daily bag and possession limit is 5 fish, **only 1 over 15 inches**
- 2. Other salmon:** Daily bag and possession limit is 5 fish, no size limit.
After taking a bag limit of coho salmon from the Unalakleet River drainage, a person may not sport fish for any species of fish downstream from the South River for the remainder of that same day.
- 3. All salmon:** Any salmon completely removed from the water shall be retained and becomes part of the bag limit of the person originally hooking it. A person may not completely remove a salmon from the water before releasing it.
-

APPENDIX C

Appendix C1.-Emergency Order issued at Fairbanks, August 27, 2001.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH & GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-SS-03-01 Issued at Fairbanks, August 27, 2001

Effective date: 12:01 A.M. August 28, 2001

Expiration Date: October 31, 2001
Unless superseded by subsequent
Emergency Order.

EXPLANATION:

This Emergency Order opens the Nome River to sport fishing for coho salmon effective at 12:01 AM, Tuesday August 28, 2001, with a daily bag and possession limit of one coho per day.

REGULATION:

This emergency order rescinds E.O. No. 3-SS-02-01, thereby opening the Nome River to sport fishing for coho salmon with a daily bag and possession limit of one coho.

Frank Rue
Commissioner

by delegation to: _____
Alfred L. DeCicco
Northwest Area
Management Biologist

-continued-

Appendix C1.-Page 2 of 2.

Emergency Order Number: 3-SS-03-01

**August 27, 2001
Page 2**

JUSTIFICATION:

The "special provisions" of 5AAC 70.022 (b) (6) were superseded by Emergency Order No. 3-SS-02-01. Under that Emergency Order, the following special provision were effective at 12:01 PM on Friday August 24, 2001: the Nome River was closed to fishing for coho salmon. The Nome River weir count as of August 23 indicated that only 251 coho had migrated upriver past the structure. There has been a push of coho counted past the weir since that time, and over 1,100 coho had moved upstream as August 26. It appears that escapement numbers will exceed recent odd-year (weak year) returns. Therefore, the department feels that it can reopen the Nome River to the harvest of one coho per day without adversely impacting the reproductive potential of the escapement.

The Department will continue to evaluate in-season run strength and may reduce or expand fishing opportunity as the return of coho nears its completion.

DISTRIBUTION:

Office of the Governor; Lt. Governor; Commissioner, Department of Fish and Game; Director, Division of Sport Fish; Regional Supervisors, Division of Sport and Commercial Fisheries; Members of Board of Fisheries; Director, Fish and Wildlife Protection; Detachment Commander and Area Officer, Board of Fisheries Members; Local Fish and Game Advisory Chairman; Anchorage, Cook Inlet, Delta, Fairbanks and Palmer Area Biologists, Division of Sport Fish and Commercial Fisheries; Juneau, Region I; Anchorage, Region II; Fairbanks, Region III; and selected area newspapers, radio and television stations.

Appendix C2.-Emergency Order issued at Fairbanks, August 23, 2001.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH & GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-SS-02-01 Issued at Fairbanks, August 23, 2001

Effective date: 12:01 A.M. August 24, 2001

Expiration Date: October 31, 2001

Unless superseded by subsequent
Emergency Order.

EXPLANATION:

This Emergency Order closes the Nome River to sport fishing for coho salmon effective at noon, Friday August 24, 2001. Any coho salmon caught while fishing for other species must be immediately released.

REGULATION:

The "special provisions" of 5AAC 70.022 (b) (6) are superseded by this Emergency Order. Under this Emergency Order, the following special provision is effective at 12:01 PM on Friday August 24, 2001: the Nome River is closed to fishing for coho salmon.

Frank Rue
Commissioner

by delegation to: _____

Alfred L. DeCicco
Northwest Area
Management Biologist

-continued-

Emergency Order Number: 3-SS-02-01

August 23, 2001
Page 2

JUSTIFICATION:

Based on weir counts, the coho salmon run in the Nome river is lagging behind recent weak-year estimates and fish are not moving upstream in adequate numbers. The Nome River weir count as of August 23 indicated that only 251 coho had migrated upriver past the structure. In the past four days, no fish have moved upstream past the weir. This ranks the 2001 coho run in the Nome river behind the very weak runs in 1997 and 1999 as of this date. It is likely that other fisheries for coho salmon on the Nome River will also be further restricted to allow the few fish in the river to migrate upstream to spawning areas.

The Department will continue to evaluate in-season run strength and may reduce or expand fishing opportunity as the return of coho nears its completion.

DISTRIBUTION:

Office of the Governor; Lt. Governor; Commissioner, Department of Fish and Game; Director, Division of Sport Fish; Regional Supervisors, Division of Sport and Commercial Fisheries; Members of Board of Fisheries; Director, Fish and Wildlife Protection; Detachment Commander and Area Officer, Board of Fisheries Members; Local Fish and Game Advisory Chairman; Anchorage, Cook Inlet, Delta, Fairbanks and Palmer Area Biologists, Division of Sport Fish and Commercial Fisheries; Juneau, Region I; Anchorage, Region II; Fairbanks, Region III; and selected area newspapers, radio and television stations.

Appendix C3.- Emergency Order issued at Fairbanks, August 20, 2001.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH & GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-SS-01-01 Issued at Fairbanks, August 20, 2001

Effective Date: 12:01 A.M. August 21, 2001

Expiration Date: October 30, 2001
Unless superseded by subsequent
Emergency Order.

EXPLANATION:

This Emergency Order reduces the sport daily bag and possession limit of coho salmon from 3 fish per day to 1 coho salmon per day in all waters of the Nome subdistrict and the Pilgrim River effective 12:01 P.M., Tuesday August 21, 2001.

The waters impacted by this action include all salt and fresh waters of Norton Sound between and including the Sinuk River in the west and the Solomon River in the East. In addition, this action also includes the Pilgrim River. Specific freshwater rivers affected by this Emergency Order are the Sinuk, Cripple, Penny, Snake, Nome, Flambeau, Eldorado, Bonanza, Solomon and Pilgrim rivers.

REGULATION:

The provisions of 5AAC 70.022 (b) (6,7,8,10, and 11) are superseded by this Emergency Order. Under this Emergency Order, the following provisions are effective: In all waters of the Nome subdistrict from and inclusive of the Sinuk River in the west and the Solomon River in the east and the Pilgrim River 12:01 P.M. Tuesday August 21, 2001 the sport daily bag and possession limit for coho salmon is reduced to fish per day.

Frank Rue
Commissioner

by delegation to: _____
Alfred L. DeCicco
Northwest Area
Management Biologist

-continued

Appendix C3.-Page 2 of 2.

Emergency Order Number: 3-SS-01-01

**August 20, 2001
Page 2**

JUSTIFICATION:

The coho salmon run in the Nome Subdistrict appears to be below average. Tower and weir counts as of August 20 indicate that the return is somewhat stronger than the very weak run in 1997 when all subsistence and sport fisheries were closed and only slightly stronger than the 1999 and 2000 returns. Subsistence fishing in the lower reaches of the Nome River is being further restricted to allow more fish to migrate to spawning areas.

The Department will continue to evaluate in-season run strength and may reduce or expand fishing opportunity as the return of coho nears its completion. At the present time, it is prudent to reduce the harvest potential of the sport fishery.

DISTRIBUTION:

Office of the Governor; Lt. Governor; Commissioner, Department of Fish and Game; Director, Division of Sport Fish; Regional Supervisors, Division of Sport and Commercial Fisheries; Members of Board of Fisheries; Director, Fish and Wildlife Protection; Detachment Commander and Area Officer, Board of Fisheries Members; Local Fish and Game Advisory Chairman; Anchorage, Cook Inlet, Delta, Fairbanks and Palmer Area Biologists, Division of Sport Fish and Commercial Fisheries; Juneau, Region I; Anchorage, Region II; Fairbanks, Region III; and selected area newspapers, radio and television stations.

Appendix C4.-Emergency Order issued at Nome, July 31, 2001.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH & GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-PS-01-01

Issued at: Nome, July 31, 2001

Effective Date: 12:01 A.M. August 1, 2001

Expiration Date: October 31, 2001

EXPLANATION:

This emergency order closes sport fishing for pink salmon in all fresh and salt waters of the Nome Subdistrict, effective 12:01 A.M., Wednesday August 1, 2001, until October 31, 2001. This emergency order prohibits the taking of pink salmon in the sport fishery in all fresh and salt waters from Cape Rodney in the west to Topkok Head in the east. All pink salmon incidentally caught while sport fishing for other species must be immediately released.

REGULATION:

5AAC 01.175 WATERS CLOSED TO SPORT FISHING is superseded by adding:

(d) the following waters are closed to sport fishing for pink salmon:

(1) All fresh and salt waters from Cape Rodney to Topkok Head

Frank Rue
Commissioner

by delegation to: _____

Alfred L. DeCicco
Northwest Area Management Biologist
Division of Sport Fish
Fairbanks

-continued-

JUSTIFICATION:

All available data indicates extremely low abundance of pink salmon in northern Norton Sound streams, particularly those in the Nome Subdistrict. As of July 29, only 465 pink salmon had moved upstream past the Nome River weir and fewer than 200 had been estimated to have passed the Snake River counting tower. Historically, approximately half the pink salmon run has returned to Nome Subdistrict streams by August 1.

Because of the weak runs, sport fishing for pink salmon in the Nome Subdistrict will remain closed for the rest of the open water season. Pink salmon caught while angling for other species must be immediately released.

DISTRIBUTION:

Office of the Governor; Lt. Governor; Commissioner, Department of Fish and Game; Director, Division of Sport Fish; Regional Supervisors, Division of Sport and Commercial Fisheries; Members of Board of Fisheries; Director, Fish and Wildlife Protection; Detachment Commander and Area Officer, Board of Fisheries Members; Local Fish and Game Advisory Chairman; Anchorage, Cook Inlet, Delta, Fairbanks and Palmer Area Biologists, Division of Sport Fish and Commercial Fisheries; Juneau, Region I; Anchorage, Region II; Fairbanks, Region III; and selected area newspapers, radio and television stations.

Appendix C5.-Emergency Order issued at Fairbanks, June 14, 2001.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH & GAME

Under Authority of AS 16.05.050

Emergency Order No. Emergency Order No. 3-S-01-01 Issued at: Fairbanks, June 14, 2001

Effective 12:01 A.M. June 15, 2001
Date:

Expiration Date: July 31, 2001
Unless superseded by subsequent
Emergency Order.

EXPLANATION:

This emergency order closes fresh and salt waters in the Nome Subdistrict (from Cape Rodney in the west to Topkok Head in the east) to sport fishing for all salmon. Effective Thursday, June 14, until further notice, all salmon caught while sport fishing for any other species in fresh and salt waters from the Cape Rodney to Topkok Head must be immediately released.

This Emergency Order has no bearing on sport fishing for other fish species.

REGULATION:

5AAC 01.175. WATERS CLOSED TO SPORT FISHING is superseded by adding:

- (d) the following waters are closed to sport fishing for salmon:
(1) All fresh and salt waters from Cape Rodney to Topkok Head

Frank Rue
Commissioner

by delegation to: _____
Alfred L. DeCicco
Northwest Area
Management Biologist

-continued-

Appendix C5.-Page 2 of 2.

JUSTIFICATION:

Based on normal odd-year weak pink salmon runs, and the exceptionally weak run in 1999, the Department anticipates another very weak pink salmon return to northern Norton Sound in 2001. The waters of the Nome Sub-district will be closed to subsistence fishing for all salmon until such time that run strength can be assessed. Before subsistence fisheries can be closed, other fisheries must also be closed. Therefore, in order to comply with the subsistence priority, the sport fisheries for salmon will be closed to allow for the preseason subsistence salmon closure. This closure is being done in coordination with the Commercial Fisheries Division that manages Norton Sound subsistence fisheries. These fisheries may reopen if runs appear strong enough to support both subsistence harvests and sport fisheries.

DISTRIBUTION:

Office of the Governor; Lt. Governor; Commissioner, Department of Fish and Game; Director, Division of Sport Fish; Regional Supervisors, Division of Sport and Commercial Fisheries; Members of Board of Fisheries; Director, Fish and Wildlife Protection; Detachment Commander and Area Officer, Board of Fisheries Members; Local Fish and Game Advisory Chairman; Anchorage, Cook Inlet, Delta, Fairbanks and Palmer Area Biologists, Division of Sport Fish and Commercial Fisheries; Juneau, Region I; Anchorage, Region II; Fairbanks, Region III; and selected area newspapers, and radio stations.

List is available at ADF&G, Sport Fish Division, 1300 College Road, Fairbanks 99701

