

# Kobuk River Test Fishing Project, 1999

By

Tom Kohler



Regional Information Report<sup>1</sup> No. 3A00-07

Alaska Department of Fish and Game  
Commercial Fisheries Management and Development Division, AYK Region  
333 Raspberry Road  
Anchorage, Alaska 99518-1599

January 2000

<sup>1</sup> The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished divisional reports. These reports frequently serve diverse ad hoc informational purposes or archive basic uninterpreted data. To accommodate needs for up-to-date information, reports in this series may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Commercial Fisheries Division.

## TABLE OF CONTENTS

	Page
LIST OF TABLES .....	iii
LIST OF FIGURES .....	iv
INTRODUCTION .....	1
METHODS .....	1
Site Description .....	1
Test Fishing .....	2
Standardized Catches.....	2
RESULTS .....	3
CONCLUSIONS .....	4
LITERATURE CITED.....	5

## LIST OF TABLES

Table	Page
1. Kobuk River chum salmon drift test fish CPUE by day, drift and site, 1999.....	7
2. Kobuk River chum salmon drift test fish mean daily and cumulative CPUE, 1993-1999.....	8
3. Kobuk River chum salmon drift test fish daily and cumulative CPUE proportions, 1993-1999.	10
4. Kobuk River chum salmon drift test fish CPUE indices, mean CPUE and percent by drift and site, 1999.....	12
5. Kobuk River chum salmon drift test fish diurnal and spatial distribution expressed as mean CPUE by drift period and by site, 1993-1999.....	13
6. Kobuk River chum salmon drift test fish CPUE and cumulative CPUE by drift, 1993-1999.....	14
7. Kobuk River chum salmon drift test fish catch age and sex composition, 1999.....	18
8. Comparison of chum salmon age and sex composition and mean length from the Kobuk and Noatak River drift test fish catch, and the Kotzebue District commercial catch, 1999..	19

## LIST OF FIGURES

Figure	Page
1. Kotzebue Sound commercial fishing districts, villages and subsistence fishing areas.....	20
2. Lower Kobuk River drift test fishing sites.....	21
3. Kobuk River chum salmon drift test fish cumulative CPUE, 1993-1999.....	22

## INTRODUCTION

The Kobuk River originates on the south side of the Brooks Range in the Arrigetch Mountains inside the Gates of the Arctic National Park. The river flows roughly 500 river miles west where it terminates at Hotham Inlet. The lower two-thirds of the river is stained by tannin primarily from the Pah River, an upper river tributary. Five villages are located on the Kobuk River and all depend on chum salmon for subsistence use. Residents of Kotzebue also depend on Kobuk River chum salmon as a subsistence resource. The Kobuk River is also thought to support up to 60% of the commercial catch of chum salmon in the Kotzebue District.

This was the sixth consecutive year a drift gillnet test fishing project operated in the lower Kobuk River (Lingnau, 1993; Lingnau, 1994; Lingnau, 1995; Lingnau, 1996; Lingnau, 1997; Lingnau, 1998; Kohler, 2000). Because of the Kobuk River's tannic stain, test fishing is less susceptible to net avoidance by salmon than in clear water systems. The only previous salmon project in the Kobuk River drainage was a counting tower site on the Squirrel River, which was too distant to provide timely information for fisheries management. This report presents the results of the seventh year of the Kobuk River drift test-fishing project.

Management of the Kotzebue District commercial salmon fishery, particularly during the month of July, is dependent primarily on comparing commercial fishing period and cumulative season catch statistics to those of prior years. Because of the change in market demand in recent years, these comparisons are no longer reliable. The drift test-fishing project was initiated because of the need for an inseason index of run timing and abundance for Kobuk River chum salmon stocks, which largely support the first portion of the salmon migration into the Kotzebue District. While test fishing is a relatively low cost approach, it can also be susceptible to inter-annual variability in catch rates which typically requires the data be interpreted in a somewhat qualitative way as an abundance index if calibration is not possible between years. The objectives of the test fishing project for 1999 were:

1. To evaluate chum salmon abundance migrating into the Kobuk River drainage using a comparison of systematic drift gill net catches.
2. To assess, in a qualitative way, the impact of the Kotzebue District commercial salmon fishery on chum salmon abundance in the lower Kobuk River for fisheries management purposes.
3. Describe the migratory timing for chum salmon in the lower Kobuk River.
4. Sample for age, sex and length.

## METHODS

### Site Description

The site is approximately 70 river miles from the eastern boundary of the commercial salmon

fishing district (Figure 1). This is the furthest downstream site where the river runs through a single channel and is below all tributaries which support spawning chum salmon. The test-fishing site was also selected because of its desirable stream characteristics. The site consists of roughly a 1-mile river section located approximately 3 miles downstream from Kiana. The width of the river was approximately 300 meters and was divided into two sites (Figure 2). Site N is the north side of the river (right bank), which is the cut bank side of the river with the swiftest current. Site S is located on the south side of the river (left bank). Site S is located downstream from a major sandbar and has a gradual gradient. It is also the site with the slowest current. A bottom profile at the test fish site in 1997 revealed a near uniform bottom with a maximum depth of 6 meters. The deepest portion of the river was in the first quartile from the right bank.

### **Test Fishing**

Fishing was scheduled to sample salmon passage during three different segments of the day at each of the two sites; morning (0800), mid-day (1500), and late evening (2200). A two-person crew conducted drifts, six days per week. During the peak of the run, drifts were conducted every day of the week.

All test-fishing drifts were made from a 20-foot open outboard motorboat for approximately 20 minutes with a 50-fathom gill net. If catch rates were high, fishing time was reduced in order to control mortality. The net was composed of 6-inch (15.2 cm) stretched mesh multifilament webbing, 40 meshes deep, and hung at a ratio of 2:1. Age-sex-length data were collected from up to 80 chum salmon per day. Mortalities were primarily given to elders but some were given to other individuals for subsistence purposes. The availability of chum salmon was announced over the CB radio.

### **Standardized Catches**

Actual catches were converted to catch per unit of effort (CPUE) by considering fishing time and the length of net used. Each CPUE index was the number of fish, which would have been caught if 100 fathoms of net had been fished for 60 minutes. The index ( $I$ ) was calculated as follows:

$$I = \frac{6,000 (c)}{(l)(t)}$$

Where:  $c$  = number of chum salmon caught

$l$  = length of net in fathoms

$t$  = mean fishing time in minutes

Mean fishing time ( $t$ ) was defined as the amount of time the entire net was fishing plus half the time it took to deploy and retrieve the net. Mean daily drift CPUE indices were calculated using the sum of the total time fished and total fish caught for each day. The mean daily indices were summed to produce total seasonal CPUE indices for the period of data collection. Cumulative proportions of

seasonal total test fish CPUE indices were also calculated.

Catch rate for each time period and site was determined by using the fishing time and number of fish caught for those specific time periods and sites. Seasonal abundance by site and time period was indexed by summing CPUE indices for each of the daily sites and time periods. Temporal distribution was depicted as a percent calculated by dividing each time period total by the total CPUE indices. Spatial distribution was described as a percent by dividing each site's CPUE seasonal total by the total of both site's CPUE indices. Temporal and spatial distribution are described as a percent since the number of drifts made at each site and the amount of time fished varied (Lingnau 1998).

## RESULTS

Drifting began on 11 July and continued through 13 August. CPUE indices were calculated for each drift and site (Table 1). There were 1,373 chum salmon caught in a total of 176 drifts (88 drift time periods) producing 3,359 chum salmon drift period CPUE index points (Table 6). The peak catch and CPUE occurred on 1 August with a catch of 171 salmon, which was a mean daily CPUE of 145.02 (4.3% of the seasonal CPUE index). Totals of 44.4, 34.5, and 21.2 percent of the seasonal CPUE indices were caught at 0800, 1500, and 2200 hours (Table 4). Totals of 17.0 and 83.0 percent of the total seasonal CPUE indices were caught at sites N and S. The CPUE was highest for period 1 at 49.7 followed by period 2 at 38.6 and period 3 at 25.4 in 1999 (Table 5).

There were 913 aged chum salmon scales from test net samples. Enough scale samples were taken to stratify age and sex composition into four periods (Table 7). The age composition was 0.4% age-0.2, 92.0% age-0.3, 5.9% age-0.4, 1.4% age-0.5 and 0.2% age-0.6 (Table 7). The age composition of the 1999 Kotzebue commercial and Noatak River drift test fish catch is shown for comparison in Table 8. Length by age comparison (nsc) indicates that males from the Kotzebue commercial catch and Noatak River test fishing samples were larger than the Kobuk River. Females from the Noatak River sample were larger than the other two sample sources. Chum salmon samples were caught with similar mesh size gear. Samples from the Kobuk and Noatak Rivers were from 6-inch mesh drift gillnet catches while commercial gear is typically 5-7/8 or 6 inch mesh set gillnet.

The test fishing methods for the Kobuk River project were the same as they had been in the prior five years. The test fishing gear was intended to match the gear typically used in the commercial fishery. One day of test fishing were missed due to a regular day off. Ten drifts were missed due to mechanical problems and one drift was missed due to weather. Seasonal test fishing data for 1993-1999 is presented in Tables 2, 3, 5 and 6. Figure 3 shows test fishing cumulative CPUE by day for 1993-1999.

The test fishing CPUE indices generated (number of salmon caught) can be influenced considerably by normal commercial fishing activity in Kotzebue Sound, as well as by the number of drifts conducted and their timing compared to commercial periods. In addition, local salmon migration patterns can be greatly influenced by weather conditions. For these reasons, no interpolations were made for missing data points since the accuracy of such estimates may not be reliable.

## CONCLUSIONS

The Kobuk River test-fishing project was once again successful in 1999. Due to the high water levels and turbidity in 1999 the Kobuk test fishery was the only indication of escapement in the Kotzebue district during the commercial season. During periods of low water at the beginning of the season, tannic staining of the river prevented net avoidance by migrating salmon. This contributed to stable catch efficiency throughout the season. The tannic stain provides concealment of the gillnets so that the ability to catch fish throughout the run remains relatively constant. This allows comparability within and between years.

This year's chum salmon passage by time of day was higher during the first two drift periods. When looking at the historical catch information, in most years, there has been very little difference in salmon passage during different times of the day. Just as consistent is the catch rate by site. In all but one year, roughly seventy percent of the salmon CPUE occurred on the south side of the channel. The one year that was different (1994), was a 50 year flood event.

This year's peak catches occurred between 25 July and 7 August. Fluctuations in the test fishery data due to commercial openings indicated that the migration timing was 5 days at the beginning of the run and 4 days at the end. Previous information from local residents and the department indicate that the migration time is 5 to 6 days. Local subsistence fishers were interviewed throughout the season by the test fishing crew. Catch rates from the test fishery seemed to track with subsistence catches throughout the season. The test fishery is most likely catching mixed stocks of fish. Kiana residents are thought to harvest predominantly Squirrel River stocks. With the Kobuk River test fishing project providing fish to the community, pressure of subsistence harvests on Squirrel River stocks is most likely reduced.

The project was run as long as the budget would allow. Its six week duration is believed to have covered most of the migration. However, catches at the end of test fishing indicated the latest portion of the run was missed. Test fishing on the Kobuk River at the current drift gillnet site near Kiana is feasible and provides management with usable escapement indexing information in a cost effective manner. This project was used this year, and will be used in the future, as a management tool, to index escapements into the Kobuk River.

## LITERATURE CITED

- Kohler, T. 2000. Kobuk River Test Fishing Project, 1998. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 3A00-06, Anchorage.
- Lingnau, T. L. 1993. Kobuk River Test Fishing Project, 1994. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 3A93-20, Anchorage.
- Lingnau, T. L. 1994. Kobuk River Test Fishing Project, 1994. Alaska Department of Fish and Game, Commercial Fisheries Division, Regional Information Report 3A94-35, Anchorage.
- Lingnau, T. L. 1995. Kobuk River Test Fishing Project, 1995. Alaska Department of Fish and Game, Commercial Fisheries Division, Regional Information Report 3A95-30, Anchorage.
- Lingnau, T. L. 1996. Kobuk River Test Fishing Project, 1996. Alaska Department of Fish and Game, Commercial Fisheries Division, Regional Information Report 3A96-33, Anchorage.
- Lingnau, T. L. 1997. Kobuk River Test Fishing Project, 1997. Alaska Department of Fish and Game, Commercial Fisheries Division, Regional Information Report 3A97-38, Anchorage.

## OEO / ADA STATEMENT

The Alaska Department of Fish and Game administers all programs and activities free from discrimination on the basis of sex, color, race, religion, national origin, age, marital status, pregnancy, parenthood or disability. For information on alternative formats available for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-4120, (TDD) 907-465-3646. Any person who believes s/he has been discriminated against should write to:

ADF&G  
P.O. Box 25526  
Juneau, AK 99802-5526

or

O.E.O.  
U.S. Department of Interior  
Washington, D.C. 20240

Table 1. Kobuk River chum salmon drift test fish CPUE by day, drift and site, 1999. <sup>a</sup>

Date	CPUE by Drift <sup>b</sup>			CPUE by Site <sup>c</sup>		Daily CPUE	Cum. CPUE
	#1	#2	#3	N	S		
11-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00
12-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00
13-Jul <sup>d</sup>	0.00	0.00	0.00	0.00	0.00	0.00	0.00
14-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00
15-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00
16-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00
17-Jul	0.00	10.00	2.58	0.00	8.45	4.26	4.26
18-Jul	25.26	0.00	0.00	0.00	16.78	8.48	12.74
19-Jul	2.53	5.16	9.90	0.00	11.75	5.89	18.63
20-Jul	2.55	12.77	0.00	0.00	10.07	5.11	23.74
21-Jul	15.16	48.51	7.66	3.43	43.64	23.75	47.49
22-Jul	12.77	2.55	20.43	0.00	23.50	11.91	59.40
23-Jul	0.00	13.04	5.22	1.75	10.36	6.09	65.49
24-Jul	33.19	33.91	7.74	3.45	46.29	24.95	90.44
25-Jul	17.87	45.00	22.98	20.57	36.67	28.73	119.17
26-Jul	52.50	7.74	58.11	1.74	75.62	39.72	158.89
27-Jul	107.66	50.00	<sup>e</sup>	35.37	120.00	80.39	239.28
28-Jul	<sup>e</sup>						239.28
29-Jul	<sup>e</sup>	<sup>e</sup>	55.00	31.30	76.80	55.00	294.28
30-Jul	37.50	80.00	30.32	28.94	69.26	49.66	343.94
31-Jul	206.61	13.33	weather	51.82	312.38	160.53	504.47
1-Aug	47.50	224.00	158.05	45.88	236.73	145.02	649.49
2-Aug	50.00	42.50	32.50	15.43	66.49	41.67	691.16
3-Aug	33.19	<sup>e</sup>	<sup>e</sup>	10.43	55.00	33.19	724.35
4-Aug	<sup>e</sup>	52.50	95.51	25.81	118.81	74.23	798.58
5-Aug	152.94	135.00	30.97	27.83	180.39	108.04	906.62
6-Aug	97.96	54.43	96.52	50.35	115.00	82.79	989.41
7-Aug	107.18	127.35	7.83	15.43	144.31	82.73	1,072.14
8-Aug <sup>d</sup>							1,072.14
9-Aug	58.11	86.60	20.65	10.29	99.31	55.58	1,127.72
10-Aug	60.00	59.35	15.32	24.35	64.68	44.73	1,172.45
11-Aug	125.83	20.65	20.65	12.09	100.80	58.13	1,230.58
12-Aug	99.80	28.39	12.90	12.17	82.15	48.50	1,279.08
13-Aug	144.47	5.16 <sup>f</sup>		30.64	122.35	78.37	1,357.45

<sup>a</sup> Catch per unit effort is calculated in catch/100fm/hour.

<sup>b</sup> Drift 1 begins at 0800, Drift 2 at 1500, Drift 3 at 2200.

<sup>c</sup> Site N is the North Bank (right bank), Site S is the South Bank (left bank).

<sup>d</sup> Regular Day Off

<sup>e</sup> Breakdown

<sup>f</sup> End of the season, no further drifts were conducted.

Table 2. Kobuk River chum salmon drift test fish mean daily and cumulative CPUE, 1993-1999.

Date	1993		1994		1995		1996		1997		1998		1999	
	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.
09-Jul							12.77	12.77	5.85	5.85				
10-Jul							15.00	27.77	0.00	5.85	5.22	5.22		
11-Jul							98.38	126.15	5.31	11.16	0.85	6.07	0.00	0.00
12-Jul	11.18	11.18			0.00	0.00	45.54	171.69	7.19	18.35		6.07	0.00	0.00
13-Jul	14.22	25.40	0.00	0.00	0.93	0.93	74.29	245.98	a	18.35	15.89	21.96	0.00	0.00
14-Jul	20.57	45.97	2.68	2.68	2.80	3.73	a	245.98	6.25	24.60	7.53	29.49	0.00	0.00
15-Jul	35.08	81.05	2.58	5.26	2.77	6.50	83.75	329.73	3.65	28.25	14.07	43.56	0.00	0.00
16-Jul	13.19	94.24	11.35	16.61	a	6.50	71.35	401.08	14.28	42.53	17.33	60.89	0.00	0.00
17-Jul	17.27	111.51	a	16.61	0.00	6.50	55.49	456.57	15.17	57.70	5.07	65.96	4.26	4.26
18-Jul	a	111.51	7.16	23.77	1.81	8.31	89.86	546.43	16.12	73.82	9.02	74.98	8.48	12.74
19-Jul	10.71	122.22	12.40	36.17	9.89	18.20	54.74	601.17	17.98	91.80		74.98	5.89	18.63
20-Jul	2.76	124.98	+ 3.65	39.82	16.30	34.50	63.70	664.87	a	91.80	18.66	93.64	5.11	23.74
21-Jul	3.20	128.18	7.30	47.12	38.54	73.04	52.12	716.99	18.53	110.33	11.87	105.51	23.75	47.49
22-Jul	5.52	133.70	3.56	50.68	21.18	94.22	50.97	767.96	13.28	123.61	0.00	105.51	11.91	59.40
23-Jul	27.15	160.85	16.49	67.17	50.58	144.80	91.36	859.32	10.79	134.40	29.58	135.09	6.09	65.49
24-Jul	9.06	169.91	a	67.17	28.46	173.26	91.89	951.21	22.86	157.26	27.33	162.42	24.95	90.44
25-Jul	a	169.91	14.38	81.55	40.16	213.42	76.80	1,028.01	21.57	178.83	24.68	187.10	28.73	119.17
26-Jul	15.22	185.13	47.65	129.20	35.15	248.57	55.68	1,083.69	14.66	193.49		187.10	39.72	158.89
27-Jul	8.06	193.19	40.66	169.86	63.94	312.51	29.79	1,113.48	18.46	211.95	23.91	211.01	80.39	239.28
28-Jul	16.36	209.55	57.83	227.69	62.49	375.00	49.06	1,162.54	30.53	242.48	51.91	262.92		239.28
29-Jul	0.93	210.48	33.62	261.31	46.11	421.11	70.13	1,232.67	28.13	270.61	34.16	297.08	55.00	294.28
30-Jul	0.92	211.40	69.21	330.52	57.86	478.97	35.29	1,267.96	22.33	292.94	24.59	321.67	49.66	343.94
31-Jul	12.58	223.98	a	330.52	29.89	508.86	82.27	1,350.23	32.57	325.51	15.69	337.36	160.53	504.47
01-Aug	a	223.98	82.16	412.68	72.91	581.77	167.7	1,517.90	41.41	366.92	25.44	362.80	145.02	649.49
02-Aug	6.74	230.72	65.12	477.80	48.71	630.48	62.02	1,579.92	22.41	389.33		362.80	41.67	691.16
03-Aug	54.49	285.21	71.79	549.59	48.40	678.88	48.7	1,628.62	35.21	424.54	26.67	389.47	33.19	724.35
04-Aug	44.23	329.44	108.98	658.57	53.00	731.88	65.93	1,694.55	26.67	451.21	42.35	431.82	74.23	798.58
05-Aug	89.30	418.74	59.74	718.31	49.95	781.83	60.33	1,754.88	24.47	475.68	8.57	440.39	108.04	906.62

(continued)

Table 2. (Page 2 of 2)

Date	1993		1994		1995		1996		1997		1998		1999	
	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.
06-Aug	18.60	437.34	102.56	820.87	a	781.83	80.47	1,835.35	42.25	517.93	6.00	446.39	82.79	989.41
07-Aug	20.52	457.86	a	820.87	48.39	828.22	90.99	1,926.34	36.00	553.93	5.11	451.50	82.73	1,072.14
08-Aug	a	457.86	62.75	883.62	44.02	872.24	146.9	2,073.28	45.07	599.00	16.40	467.90		1,072.14
09-Aug	1.84	459.70	96.86	980.48	68.22	940.46	106.1	2,179.39	55.14	654.14	17.20	485.10	55.58	1,127.72
10-Aug	12.63	472.33	45.83	1,026.31	56.33	996.79	56.95	2,236.34	a	654.14	9.46	494.56	44.73	1,172.45
11-Aug	18.11	490.44	57.02	1,083.33	37.95	1,034.74	a	2,236.34	43.45	697.59	10.29	504.85	58.13	1,230.58
12-Aug	3.74	494.18	90.54	1,173.87	63.92	1,098.66	72.29	2,308.63	37.36	734.95	19.44	524.29	48.5	1,279.08
13-Aug			11.36	1,185.23	a	1,098.66	114.6	2,423.26	45.93	780.88	10.21	534.50	78.37	1,357.45
14-Aug			a	1,185.23	29.35	1,128.01	158.1	2,581.39	16.01	796.89	3.85	538.35		
15-Aug			5.13	1,190.36	25.26	1,153.27					0	538.35		
16-Aug			16.23	1,206.59	35.04	1,188.31								
17-Aug			0.00	1,206.59										
18-Aug			0.00	1,206.59										
19-Aug			3.12	1,209.71										
20-Aug			0.00	1,209.71										
21-Aug			a	1,209.71										
22-Aug			0.00	1,209.71										
23-Aug			0.00	1,209.71										
24-Aug			0.00	1,209.71										
25-Aug			0.91	1,210.62										
26-Aug			5.56	1,216.18										
27-Aug			1.86	1,218.04										
28-Aug			0.93	1,218.97										
29-Aug			0.00	1,218.97										
30-Aug			0.00	1,218.97										
31-Aug														

a Regular day off.

Table 3. Kobuk River chum salmon drift test fish mean daily and cumulative CPUE proportions, 1993-1999.

Date	1993		1994		1995		1996		1997		1998		1999	
	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.
09-Jul							0.005	0.005	0.007	0.007				
10-Jul							0.006	0.011	0.000	0.007	0.010	0.01		
11-Jul							0.038	0.049	0.007	0.014	0.002	0.011	0.00	0.00
12-Jul	0.023	0.023			0.000	0.000	0.018	0.067	0.009	0.023	0.000	0.011	0.00	0.00
13-Jul	0.029	0.051	0.000	0.000	0.001	0.001	0.029	0.095	0.000	0.023	0.030	0.041	0.00	0.00
14-Jul	0.042	0.093	0.002	0.002	0.002	0.003	<sup>a</sup> 0.095	0.095	0.008	0.031	0.014	0.055	0.00	0.00
15-Jul	0.071	0.164	0.002	0.004	0.002	0.005	0.032	0.128	0.005	0.035	0.026	0.081	0.00	0.00
16-Jul	0.027	0.191	0.009	0.014	<sup>a</sup> 0.000	0.005	0.028	0.155	0.018	0.053	0.032	0.113	0.00	0.00
17-Jul	0.035	0.226	<sup>a</sup> 0.006	0.014	0.000	0.005	0.021	0.177	0.019	0.072	0.009	0.123	0.003	0.00
18-Jul	<sup>a</sup> 0.022	0.226	0.006	0.020	0.002	0.007	0.035	0.212	0.020	0.093	0.017	0.139	0.006	0.01
19-Jul	0.022	0.247	0.010	0.030	0.008	0.015	0.021	0.233	0.023	0.115	0.000	0.139	0.004	0.01
20-Jul	0.006	0.253	0.003	0.033	0.014	0.029	0.025	0.258	0.000	0.115	0.035	0.174	0.004	0.02
21-Jul	0.006	0.259	0.006	0.039	0.032	0.061	0.020	0.278	0.023	0.138	0.022	0.196	0.017	0.03
22-Jul	0.011	0.271	0.003	0.042	0.018	0.079	0.020	0.297	0.017	0.155	0.000	0.196	0.009	0.04
23-Jul	0.055	0.325	0.014	0.055	0.043	0.122	0.035	0.333	0.014	0.169	0.055	0.251	0.004	0.05
24-Jul	0.018	0.344	<sup>a</sup> 0.012	0.055	0.024	0.146	0.036	0.368	0.029	0.197	0.051	0.302	0.018	0.07
25-Jul	<sup>a</sup> 0.031	0.344	0.012	0.067	0.034	0.180	0.030	0.398	0.027	0.224	0.046	0.348	0.021	0.09
26-Jul	0.031	0.375	0.039	0.106	0.030	0.209	0.022	0.420	0.018	0.243	0.000	0.348	0.029	0.12
27-Jul	0.016	0.391	0.033	0.139	0.054	0.263	0.012	0.431	0.023	0.266	0.044	0.392	0.059	0.17
28-Jul	0.033	0.424	0.047	0.187	0.053	0.316	0.019	0.450	0.038	0.304	0.096	0.488	0.000	0.17
29-Jul	0.002	0.426	0.028	0.214	0.039	0.354	0.027	0.478	0.035	0.340	0.063	0.552	0.041	0.22
30-Jul	0.002	0.428	0.057	0.271	0.049	0.403	0.014	0.491	0.028	0.368	0.046	0.598	0.037	0.25
31-Jul	0.025	0.453	<sup>a</sup> 0.067	0.271	0.025	0.428	0.032	0.523	0.041	0.408	0.029	0.627	0.118	0.37
01-Aug	<sup>a</sup> 0.014	0.453	0.067	0.339	0.061	0.490	0.065	0.588	0.052	0.460	0.047	0.674	0.107	0.46
02-Aug	0.014	0.467	0.053	0.392	0.041	0.531	0.024	0.612	0.028	0.489	0.000	0.674	0.031	0.51
03-Aug	0.110	0.577	0.059	0.451	0.041	0.571	0.019	0.631	0.044	0.533	0.050	0.723	0.024	0.53
04-Aug	0.090	0.667	0.089	0.540	0.045	0.616	0.026	0.656	0.033	0.566	0.079	0.802	0.055	0.59
05-Aug	0.181	0.847	0.049	0.589	0.042	0.658	0.023	0.680	0.031	0.597	0.016	0.818	0.080	0.67

(continued)

Table 3. (Page 2 of 2)

Date	1993		1994		1995		1996		1997		1998		1999	
	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.	Daily	Cum.
06-Aug	0.038	0.885	0.084	0.673	<sup>a</sup>	0.658	0.031	0.711	0.053	0.650	0.011	0.829	0.061	0.73
07-Aug	0.042	0.927	<sup>a</sup>	0.673	0.039	0.697	0.035	0.746	0.045	0.695	0.009	0.839	0.061	0.79
08-Aug	<sup>a</sup>	0.927	0.051	0.725	0.037	0.734	0.057	0.803	0.057	0.752	0.030	0.869	0.000 <sup>a</sup>	0.79
09-Aug	0.004	0.930	0.079	0.804	0.057	0.791	0.041	0.844	0.069	0.821	0.032	0.901	0.041	0.83
10-Aug	0.026	0.956	0.038	0.842	0.047	0.839	0.022	0.866	0.000	0.821	0.018	0.919	0.033	0.86
11-Aug	0.037	0.992	0.047	0.889	0.032	0.871	0.000	0.866	0.055	0.875	0.019	0.938	0.043	0.91
12-Aug	0.008	1.000	0.074	0.963	0.054	0.925	0.028	0.894	0.047	0.922	0.036	0.974	0.036	0.94
13-Aug			0.009	0.972	<sup>a</sup>	0.925	0.044	0.939	0.058	0.980	0.019	0.993	0.058	1.00
14-Aug			<sup>a</sup>	0.972	0.025	0.949	0.061	1.000	0.020	1.000	0.007	1.000		
15-Aug			0.004	0.977	0.021	0.971					0.000	1.000		
16-Aug			0.013	0.990	0.029	1.000								
17-Aug			0.000	0.990										
18-Aug			0.000	0.990										
19-Aug			0.003	0.992										
20-Aug			0.000	0.992										
21-Aug			<sup>a</sup>	0.992										
22-Aug			0.000	0.992										
23-Aug			0.000	0.992										
24-Aug			0.000	0.992										
25-Aug			0.001	0.993										
26-Aug			0.005	0.998										
27-Aug			0.002	0.999										
28-Aug			0.001	1.000										
29-Aug			0.000	1.000										
30-Aug			0.000	1.000										

<sup>a</sup> Regular day off

Table 4. Kobuk River chum salmon drift test fish CPUE indices, mean CPUE and percent by drift and site, 1999.

Drift Period	Season CPUE Indices	No. of Period Drifts	Season Mean CPUE	Percent	Station	Season CPUE Indices	No. of Site Drifts	Season Mean CPUE	Percent
1 0800 hr.	1,490.6	30	49.7	44.4	N North Bank	459.1	32	14.3	17.0
2 1500 hr.	1,157.9	30	38.6	34.5	S South Bank	2,247.6	32	70.2	83.0
3 2200 hr.	710.8	28	25.4	21.2					
Total	3,359.4	88	38.2	100.0		2,706.7	64	42.3	100.0

Table 5. Kobuk River chum salmon drift test fish diurnal and spatial distribution expressed as mean CPUE by drift period and by site, 1993-1999. <sup>a</sup>

Year	Mean CPUE by Drift Period			Yearly Mean CPUE	Percent Mean CPUE by Drift Period			Mean CPUE by Site		Yearly Mean CPUE	Percent Mean CPUE by Site	
	1	2	3		1	2	3	N	S		N	S
1993	13.0	21.3	15.9	16.8	25.4	43.4	31.1	10.0	24.9	17.4	28.7	71.3
1994	25.8	33.2	23.7	27.5	31.7	39.8	28.5	4.9	53.5	29.2	8.4	91.6
1995	29.4	37.6	38.7	35.0	29.6	34.7	35.7	25.2	48.2	36.7	34.3	65.7
1996	73.2	81.7	66.5	73.8	32.4	37.2	30.3	40.7	108.1	74.4	27.3	72.7
1997	23.9	23.3	23.6	23.6	33.1	33.2	33.7	12.7	33.8	23.3	27.3	72.7
1998	18.6	20.1	14.0	17.6	35.9	38.9	25.9	7.4	22.9	15.1	24.4	75.6
1999	49.7	38.6	25.4	38.2	44.4	34.5	21.2	14.3	70.2	42.3	17.0	83.0

<sup>a</sup> Drift 1 begins at 0800, Drift 2 at 1500, Drift 3 at 2200. Site N is the North Bank (right bank), Site S is the South Bank (left bank).

Table 6. Kobuk River chum salmon drift test fishing CPUE and cumulative CPUE by drift, 1993-1999.

Date	1993			1994			1995			1996			1997			1998			1999		
	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.									
9-Jul										1			1								
										2	7.7	7.7	2	0.0	0.0						
										3	17.9	25.5	3	10.7	10.7						
10-Jul										1	5.2	30.7	1	0.0	10.7	1					
										2	21.1	51.8	2	0.0	10.7	2					
										3	19.1	70.9	3	0.0	10.7	3	5.2	5.2			
11-Jul										1	78.1	149.0	1	0.0	10.7	1	0.0	5.2	1	0.0	0.0
										2	103.3	252.3	2	5.3	15.9	2	2.5	7.8	2	0.0	0.0
										3	122.1	374.4	3	10.8	26.7	3	0.0	7.8	3	0.0	0.0
12-Jul	1	15.5	15.5				1	0.0	0.0	1	88.4	462.8	1	0.0	26.7	1 <sup>a</sup>		7.8	1	0.0	0.0
	2	2.5	18.0				2	0.0	0.0	2	32.2	495.0	2	16.0	42.7	2		7.8	2	0.0	0.0
	3	16.0	34.0				3	0.0	0.0	3	38.4	533.4	3	5.3	48.1	3		7.8	3	0.0	0.0
13-Jul	1	5.4	39.4	1	0.0	0.0	1	0.0	0.0	1	61.9	595.4	1 <sup>a</sup>		48.1	1	25.0	32.8	1	0.0	0.0
	2	15.5	54.9	2	0.0	0.0	2	2.9	2.9	2	97.2	692.6	2		48.1	2	12.8	45.5	2	0.0	0.0
	3	25.4	80.3	3	0.0	0.0	3	0.0	2.9	3	66.0	758.6	3		48.1	3	9.9	55.4	3	0.0	0.0
14-Jul	1	13.2	93.5	1	0.0	0.0	1	2.8	5.7	1 <sup>a</sup>		758.6	1	0.0	48.1	1	12.6	68.1	1	0.0	0.0
	2	0.0	93.5	2	5.3	5.3	2	5.5	11.2	2		758.6	2	7.9	56.0	2	9.9	78.0	2	0.0	0.0
	3	46.1	139.5	3	2.6	7.9	3	0.0	11.2	3		758.6	3	10.8	66.8	3	0.0	78.0	3	0.0	0.0
15-Jul	1	20.6	160.1	1	5.0	12.8	1	5.6	16.8	1	100.7	859.2	1	2.8	69.5	1	22.3	100.2	1	0.0	0.0
	2	33.9	194.0	2	2.6	15.4	2	0.0	16.8	2	52.9	912.2	2	8.2	77.7	2	12.4	112.5	2	0.0	0.0
	3	46.5	240.5	3	0.0	15.4	3	2.8	19.5	3	100.7	1,012.8	3	0.0	77.7	3	7.5	120.1	3	0.0	0.0
16-Jul	1	2.7	243.2	1	5.1	20.6	1 <sup>a</sup>		19.5	1	50.2	1,063.0	1	10.7	88.4	1	18.1	138.2	1	0.0	0.0
	2	32.5	275.7	2	10.4	31.0	2		19.5	2	82.3	1,145.3	2	11.3	99.7	2	12.8	150.9	2	0.0	0.0
	3	2.7	278.5	3	18.9	49.9	3		19.5	3	85.0	1,230.3	3	20.9	120.5	3	21.3	172.3	3	0.0	0.0
17-Jul	1	23.5	302.0	1 <sup>a</sup>		49.9	1	0.0	19.5	1	93.7	1,323.9	1	21.3	141.9	1	10.0	182.3	1	0.0	0.0
	2	28.7	330.7	2		49.9	2	0.0	19.5	2	34.3	1,358.2	2	8.3	150.2	2	5.1	187.3	2	10.0	10.0
	3	0.0	330.7	3		49.9	3	0.0	19.5	3	56.7	1,414.9	3	15.7	165.8	3	0.0	187.3	3	2.6	12.6
18-Jul	1 <sup>a</sup>		330.7	1	2.6	52.5	1	2.8	22.3	1	59.2	1,474.1	1	16.0	181.8	1	25.5	212.8	1	25.3	37.8
	2		330.7	2	0.0	52.5	2	2.7	25.0	2	98.3	1,572.4	2	10.9	192.7	2	5.1	217.9	2	0.0	37.8
	3		330.7	3	18.5	71.0	3	0.0	25.0	3	117.8	1,690.2	3	21.3	214.0	3	25.3	240.2	3	0.0	37.8
19-Jul	1	5.5	336.1	1	23.7	94.7	1	0.0	25.0	1	69.8	1,760.1	1	8.1	222.1	1	10.2	253.4	1	2.5	40.4
	2	2.7	338.8	2	10.3	105.0	2	12.9	37.9	2	61.2	1,821.2	2	18.7	240.8	2	17.7	271.1	2	5.2	45.5
	3	23.5	362.3	3	2.8	107.8	3	16.2	54.1	3	36.9	1,858.2	3	27.3	268.1	3	7.7	278.8	3	9.9	55.4
20-Jul	1	2.8	365.1	1	2.9	110.6	1	10.8	64.8	1	70.3	1,928.5	1 <sup>a</sup>		268.1	1 <sup>a</sup>		278.8	1	2.6	58.0
	2	5.4	370.5	2	8.1	118.7	2	16.4	81.2	2	69.9	1,998.3	2		268.1	2		278.8	2	12.8	70.8
	3	0.0	370.5	3	0.0	118.7	3	21.8	103.0	3	48.7	2,047.0	3		268.1	3		278.8	3	0.0	70.8

(continued)

Table 6. (Page 2 of 4)

Date	1993			1994			1995			1996			1997			1998			1999		
	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.
21-Jul	1	2.8	373.2	1	10.8	129.5	1	39.1	142.2	1	66.7	2,113.7	1	11.0	279.1	1	40.9	319.7	1	15.2	85.9
	2	5.5	378.7	2	11.0	140.6	2	27.0	169.1	2	45.7	2,159.4	2	34.3	313.4	2	12.8	332.4	2	48.5	134.4
	3	1.9	380.6	3	0.0	140.6	3	49.0	218.2	3	47.4	2,206.8	3	8.3	321.7	3	35.0	367.4	3	7.7	142.1
22-Jul	1	2.8	383.4	1	5.5	146.0	1	20.7	238.8	1	27.6	2,234.4	1	8.1	329.8	1	38.3	405.7	1	12.8	154.9
	2	0.0	383.4	2	2.6	148.6	2	24.0	262.8	2	72.3	2,306.7	2	0.0	329.8	2	10.3	416.0	2	2.6	157.4
	3	13.2	396.6	3	2.7	151.3	3	18.9	281.7	3	58.2	2,364.9	3	31.3	361.1	3	33.2	449.2	3	20.4	177.8
23-Jul	1	2.7	399.3	1	24.8	176.1	1	53.1	334.7	1	53.0	2,417.9	1	18.5	379.5	1	15.3	464.6	1	0.0	177.8
	2	26.1	425.4	2	13.5	189.6	2	59.2	394.0	2	142.9	2,560.8	2	10.8	390.3	2	38.3	502.9	2	13.0	190.9
	3	51.6	477.0	3	11.2	200.8	3	37.7	431.7	3	105.3	2,666.0	3	2.8	393.1	3	20.4	523.3	3	5.2	196.1
24-Jul	1	8.2	485.2	1 <sup>a</sup>		200.8	1	39.1	470.7	1	62.8	2,728.8	1	16.4	409.4	1		523.3	1	33.2	229.3
	2	8.1	493.3	2		200.8	2	36.5	507.2	2	100.3	2,829.1	2	20.1	429.5	2		523.3	2	33.9	283.2
	3	10.9	504.2	3		200.8	3	10.9	518.1	3	122.8	2,951.9	3	31.0	460.5	3		523.3	3	7.7	270.9
25-Jul	1 <sup>a</sup>		504.2	1	24.3	225.0	1	16.2	534.3	1	30.0	2,981.9	1	13.8	474.3	1	15.3	538.6	1	17.9	288.8
	2		504.2	2	13.5	238.5	2	10.9	545.2	2	157.7	3,139.6	2	24.3	498.6	2	40.4	579.0	2	45.0	333.8
	3		504.2	3	5.4	243.9	3	109.4	654.6	3	16.8	3,156.4	3	26.4	524.9	3	15.7	594.7	3	23.0	356.8
26-Jul	1	10.9	515.1	1	32.7	276.6	1	20.6	675.2	1	113.2	3,269.6	1	21.8	546.7	1	37.9	632.5	1	52.5	409.3
	2	8.1	523.2	2	63.7	340.3	2	35.5	710.6	2	5.2	3,274.9	2	11.0	557.8	2	48.5	681.1	2	7.7	417.0
	3	26.4	549.6	3	44.7	384.9	3	47.4	758.0	3	27.7	3,302.6	3	11.0	568.8	3	69.7	750.7	3	58.1	475.1
27-Jul	1	15.5	565.1	1	31.3	406.3	1	50.2	808.3	1	15.2	3,317.8	1	26.4	595.2	1	35.4	786.1	1	107.7	582.8
	2	8.1	573.1	2	59.4	465.6	2	34.7	842.9	2	19.6	3,337.4	2	15.8	611.0	2	51.7	837.8	2	50.0	632.8
	3	0.0	573.1	3 <sup>b</sup>		465.6	3	102.9	945.8	3	72.7	3,410.1	3	13.2	624.2	3	15.5	853.3	3 <sup>b</sup>		632.8
28-Jul	1	11.2	584.3	1 <sup>b</sup>		465.6	1	39.4	985.2	1	52.0	3,462.1	1	29.3	653.5	1	15.0	868.3	1 <sup>b</sup>		632.8
	2	16.2	600.5	2 <sup>b</sup>		465.6	2	88.2	1,073.4	2	83.8	3,545.9	2	28.1	681.6	2	43.4	911.7	2 <sup>b</sup>		632.8
	3	21.6	622.1	3	57.8	523.5	3	67.9	1,141.3	3	8.3	3,554.2	3	33.9	715.5	3	15.5	927.1	3 <sup>b</sup>		632.8
29-Jul	1	2.7	624.8	1	34.3	557.7	1	48.8	1,190.0	1	110.0	3,664.2	1	34.3	749.8	1	20.4	947.6	1 <sup>b</sup>		632.8
	2	0.0	624.8	2	52.5	610.2	2	8.4	1,198.4	2	77.3	3,741.5	2	33.6	783.4	2	17.9	965.4	2 <sup>b</sup>		632.8
	3	0.0	624.8	3	19.3	629.6	3	85.1	1,283.5	3	20.4	3,761.9	3	16.2	799.6	3	5.7	971.1	3	55.0	687.8
30-Jul	1	0.0	624.8	1	83.1	712.6	1	67.1	1,350.5	1	51.1	3,813.0	1	13.3	812.9	1	40.4	1,011.5	1	37.5	725.3
	2	0.0	624.8	2	38.5	751.2	2	59.2	1,409.7	2	36.0	3,849.0	2	21.3	834.2	2	25.3	1,036.8	2	80.0	805.3
	3	2.8	627.5	3	82.0	833.1	3	48.6	1,458.3	3	22.9	3,871.8	3	33.0	867.2	3	10.3	1,047.1	3	30.3	835.6
31-Jul	1	15.2	643.7	1 <sup>a</sup>		833.1	1	43.0	1,507.4	1	71.3	3,943.1	1	24.6	891.8	1 <sup>a</sup>		1,047.1	1	206.6	1042.2
	2	16.2	659.9	2		833.1	2	20.9	1,528.2	2	120.0	4,063.1	2	30.0	921.8	2		1,047.1	2	13.3	1055.6
	3	5.4	665.3	3		833.1	3	19.1	1,547.3	3	59.1	4,122.2	3	42.5	964.3	3		1,047.1	3 <sup>b</sup>		1055.6
1-Aug	1 <sup>a</sup>		665.3	1	51.4	884.5	1	61.5	1,608.8	1	122.2	4,244.4	1	55.6	1,019.9	1	43.9	1,091.0	1	47.5	1103.1
	2		665.3	2	124.7	1,009.2	2	81.0	1,689.8	2	252.2	4,496.6	2	32.0	1,051.9	2	20.4	1,111.4	2	224.0	1327.1
	3		665.3	3	67.2	1,076.4	3	76.9	1,766.8	3	80.0	4,576.6	3	37.6	1,089.5	3	15.7	1,127.1	3	158.1	1485.1

(continued)

Table 6. (Page 3 of 4)

Date	1993			1994			1995			1996			1997			1998			1999		
	Drift	Daily	Cum.	Drift	Daily	Cum.															
2-Aug	1 <sup>b</sup>		665.3	1	27.0	1,103.4	1	45.0	1,811.8	1	120.0	4,696.6	1	20.9	1,110.3	1	43.87	1,170.9	1	50.0	1535.1
	2	0.0	665.3	2	74.6	1,178.0	2	66.2	1,878.0	2	30.6	4,727.2	2	28.4	1,138.8	2	40.85	1,211.8	2	42.5	1577.6
	3	13.3	678.6	3	92.8	1,270.8	3	35.5	1,913.4	3	28.5	4,755.7	3	18.9	1,157.6	3		1,211.8	3	32.5	1610.1
3-Aug	1	42.2	720.8	1	62.3	1,333.1	1	53.7	1,967.1	1	76.7	4,832.3	1	33.9	1,191.5	1	5.11	1,216.9	1	33.2	1643.3
	2	71.5	792.3	2	93.9	1,427.0	2	74.4	2,041.4	2	60.9	4,893.2	2	36.3	1,227.8	2	15.32	1,232.2	2 <sup>b</sup>		1643.3
	3 <sup>b</sup>		792.3	3	51.7	1,478.7	3	22.1	2,063.5	3	3.8	4,896.9	3	35.5	1,263.3	3	5.22	1,237.4	3 <sup>b</sup>		1643.3
4-Aug	1	16.7	809.1	1	124.9	1,603.6	1	45.3	2,108.8	1	52.0	4,948.9	1	29.6	1,292.9	1	0.00	1,237.4	1 <sup>b</sup>		1643.3
	2	60.0	869.1	2	120.0	1,723.6	2	60.0	2,168.8	2	26.0	4,974.9	2	31.4	1,324.3	2	15.16	1,252.6	2	52.5	1695.8
	3	51.3	920.3	3	82.4	1,806.0	3	53.8	2,222.6	3	145.0	5,119.9	3	19.3	1,343.6	3	2.61	1,255.2	3	95.5	1791.3
5-Aug	1	40.9	961.2	1	78.9	1,884.9	1	55.1	2,277.8	1	53.8	5,173.7	1	23.0	1,366.6	1	5.11	1,260.3	1	152.9	1944.2
	2	191.6	1,152.8	2	14.1	1,899.0	2	38.8	2,316.6	2	40.8	5,214.4	2	30.6	1,397.3	2	5.00	1,265.3	2	135.0	2079.2
	3	2.7	1,155.5	3	78.3	1,977.3	3	56.7	2,373.3	3	80.0	5,294.4	3	19.1	1,416.4	3	5.22	1,270.5	3	31.0	2110.2
6-Aug	1	12.8	1,168.3	1	116.1	2,093.5	1 <sup>a</sup>		2,373.3	1	44.1	5,338.5	1	41.9	1,458.2	1	17.87	1,288.4	1	98.0	2208.2
	2	13.8	1,182.1	2	93.3	2,186.8	2		2,373.3	2	43.3	5,381.8	2	39.6	1,497.8	2	15.65	1,304.0	2	54.4	2262.6
	3	29.3	1,211.4	3	92.9	2,279.7	3		2,373.3	3	148.0	5,529.8	3	45.7	1,543.5	3	15.65	1,319.7	3	96.5	2359.1
7-Aug	1	47.5	1,258.9	1 <sup>a</sup>		2,279.7	1	55.8	2,429.1	1	136.3	5,666.1	1	35.1	1,578.6	1	27.79	1,347.5	1	107.2	2466.3
	2	2.8	1,261.6	2		2,279.7	2	68.1	2,497.2	2	57.6	5,723.7	2	34.7	1,613.4	2	5.22	1,352.7	2	127.4	2593.7
	3	8.4	1,270.0	3		2,279.7	3	19.8	2,516.9	3	51.8	5,775.4	3	38.1	1,651.4	3	18.26	1,371.0	3	7.8	2601.5
8-Aug	1 <sup>a</sup>		1,270.0	1	77.7	2,357.3	1	21.6	2,538.5	1	94.6	5,870.0	1	69.0	1,720.5	1	2.52	1,373.5	1 <sup>a</sup>		2601.5
	2		1,270.0	2	64.8	2,422.1	2	74.4	2,612.9	2	221.8	6,091.8	2	49.9	1,770.3	2	18.26	1,391.7	2 <sup>a</sup>		2601.5
	3		1,270.0	3	49.7	2,471.8	3	41.7	2,654.6	3	98.8	6,190.6	3	11.4	1,781.8	3	7.83	1,399.6	3 <sup>a</sup>		2601.5
9-Aug	1	5.5	1,275.5	1	85.2	2,556.9	1	38.9	2,693.5	1	120.0	6,310.6	1	45.5	1,827.2	1	15.32	1,414.9	1	58.1	2659.6
	2	0.0	1,275.5	2	125.7	2,682.6	2	58.1	2,751.6	2	133.3	6,443.9	2	37.2	1,864.4	2	10.21	1,425.1	2	86.6	2746.2
	3	0.0	1,275.5	3	74.8	2,757.4	3	114.1	2,865.7	3	66.5	6,510.4	3	94.3	1,958.7	3	5.22	1,430.3	3	20.7	2766.8
10-Aug	1	0.0	1,275.5	1	9.5	2,766.9	1	73.2	2,938.9	1	32.5	6,542.9	1 <sup>b</sup>		1,958.7	1	0.00	1,430.3	1	60.0	2826.8
	2	8.1	1,283.6	2	54.9	2,821.8	2	29.6	2,968.5	2	98.6	6,641.5	2		1,958.7	2	50.00	1,480.3	2	59.4	2886.2
	3	29.3	1,313.0	3	86.0	2,907.8	3	71.3	3,039.8	3	42.6	6,684.1	3		1,958.7	3	7.66	1,488.0	3	15.3	2901.5
11-Aug	1	11.3	1,324.2	1	105.8	3,013.6	1	56.8	3,096.6	1 <sup>a</sup>		6,684.1	1	43.6	2,002.4	1	2.55	1,490.5	1	125.8	3027.3
	2	40.4	1,364.7	2	50.7	3,064.3	2	20.0	3,117.5	2		6,684.1	2	32.7	2,035.1	2	17.00	1,500.2	2	20.7	3048.0
	3	0.0	1,364.7	3	9.4	3,073.7	3	34.3	3,151.8	3		6,684.1	3	56.0	2,091.1	3	10.32	1,518.5	3	20.7	3068.6
12-Aug	1	11.3	1,376.0	1	17.9	3,091.6	1	31.3	3,183.1	1	123.3	6,807.4	1	36.7	2,127.8	1	7.58	1,526.1	1	99.8	3168.4
	2	0.0	1,376.0	2	183.2	3,274.8	2	105.5	3,288.5	2	39.1	6,846.5	2	54.0	2,181.8	2	0.00	1,526.1	2	28.4	3196.8
	3	0.0	1,376.0	3	0.0	3,274.8	3	56.3	3,344.8	3	28.2	6,874.7	3	16.0	2,197.8	3 <sup>b</sup>		1,526.1	3	12.9	3209.7

(continued)

Table 6. (Page 4 of 4)

Date	1993			1994			1995			1996			1997			1998			1999		
	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.	Drift	Daily	Cum.
13-Aug				1	23.5	3,298.3	1 <sup>a</sup>		3,344.8	1	105.2	6,979.9	1	41.1	2,238.9	1 <sup>b</sup>		1,526.1	1	144.5	3,354.2
				2	10.0	3,308.3	2		3,344.8	2	136.6	7,116.5	2	55.4	2,294.3	2	0.00	1,526.1	2	5.2	3,359.4
				3	3.4	3,311.7	3		3,344.8	3	102.9	7,219.4	3	39.3	2,333.6	3	0.00	1,526.1	3		3,359.4
14-Aug				1 <sup>a</sup>		3,311.7	1	8.1	3,352.9	1	77.3	7,296.7	1	35.2	2,368.8						
				2		3,311.7	2	54.4	3,407.3	2	197.3	7,493.9	2	13.0	2,381.8						
				3		3,311.7	3	23.5	3,430.8	3	181.5	7,675.4	3								
15-Aug				1	7.0	3,318.7	1	25.5	3,456.2												
				2	8.1	3,326.8	2	18.5	3,474.7												
				3	0.0	3,326.8	3	32.0	3,506.7												
16-Aug				1	3.3	3,330.1	1	22.9	3,529.5												
				2	33.8	3,363.9	2	45.4	3,574.9												
				3	11.3	3,375.1															
17-Aug				1	0.0	3,375.1															
				2	0.0	3,375.1															
				3	0.0	3,375.1															
18-Aug				1	0.0	3,375.1															
				2	0.0	3,375.1															
				3	0.0	3,375.1															
19-Aug				1	8.3	3,383.4															
				2	0.0	3,383.4															
				3	3.0	3,386.4															
20-Aug				1	0.0	3,386.4															
				2	0.0	3,386.4															
				3	0.0	3,386.4															
21-Aug				1 <sup>a</sup>		3,386.4															
				2		3,386.4															
				3		3,386.4															
22-Aug				1	0.0	3,386.4															
				2	0.0	3,386.4															
				3	0.0	3,386.4															
23-Aug				1	0.0	3,386.4															
				2	0.0	3,386.4															
				3	0.0	3,386.4															
24-Aug				1	0.0	3,386.4															
				2	0.0	3,386.4															
				3	0.0	3,386.4															

<sup>a</sup> Regular day off.

<sup>b</sup> No drift conducted because of mechanical problems or bad weather.

Table 7. Kobuk River chum salmon drift test fish catch age and sex composition, 1999.

		Brood Year and (Age Group)					Total
		1996 (0.2)	1995 (0.3)	1994 (0.4)	1993 (0.5)	1992 (0.6)	
Stratum Dates:	7/17-7/24						
Sampling Dates:	7/17-7/24						
Sample Size:	88						
Male	Percent of Catch	0.0	51.1	1.1	2.3	0.0	54.5
	Number in Catch	0	45	1	2	0	48
Female	Percent of Catch	0.0	42.1	2.3	0.0	1.1	45.5
	Number in Catch	0	37	2	0	1	40
Total	Percent of Catch	0.0	93.2	3.4	2.3	1.1	100.0
	Number in Catch	0	82	3	2	1	88
Stratum Dates:	7/25-7/31						
Sampling Dates:	7/25-7/31						
Sample Size:	253						
Male	Percent of Catch	0.4	59.3	5.5	2.0	0.4	67.6
	Number in Catch	1	150	14	5	1	171
Female	Percent of Catch	0.0	29.0	2.8	0.4	0.0	32.4
	Number in Catch	0	74	7	1	0	52
Total	Percent of Catch	0.4	88.5	8.3	2.4	0.4	100.0
	Number in Catch	1	224	21	6	1	253
Stratum Dates:	8/1-8/7						
Sampling Dates:	8/2-8/15						
Sample Size:	356						
Male	Percent of Catch	0.0	52.8	3.1	0.3	0.0	56.2
	Number in Catch	0	188	11	1	0	200
Female	Percent of Catch	0.0	41.0	2.8	0.0	0.0	43.8
	Number in Catch	0	146	10	0	0	156
Total	Percent of Catch	0.0	93.8	5.9	0.3	0.0	100.0
	Number in Catch	0	334	21	1	0	356
Stratum Dates:	8/9-8/13						
Sampling Dates:	8/9-8/13						
Sample Size:	216						
Male	Percent of Catch	1.4	53.2	2.3	1.4	0.0	58.3
	Number in Catch	3	115	5	3	0	126
Female	Percent of Catch	0.0	39.4	1.9	0.5	0.0	41.7
	Number in Catch	0	85	4	1	0	90
Total	Percent of Catch	3.0	92.6	4.2	1.9	0.0	100.0
	Number in Catch	1	200	9	4	0	216
Stratum Dates:	7/17-8/13						
Sampling Dates:	7/17-8/13	Season Total					
Sample Size:	913						
Male	Percent of Catch	0.4	54.5	3.4	1.2	0.1	59.7
	Number in Catch	4	498	31	11	1	545
Female	Percent of Catch	0.0	37.5	2.5	0.2	0.1	40.3
	Number in Catch	0	342	23	2	1	368
Total	Percent of Catch	0.4	92.0	5.9	1.4	0.2	100.0
	Number in Catch	4	840	54	13	2	913

Table 8. Comparison of chum salmon age and sex composition and mean length from the Kobuk and Noatak River drift test fish catch, and the Kotzebue District commercial catch, 1999.

		Brood Year and (Age Group)					
		1996 (0.2)	1995 (0.3)	1994 (0.4)	1993 (0.5)	1992 (0.6)	Total
Stratum Dates:		7/17-8/13					
Sample Size:		913					
		Kobuk River					
Male	Percent of Catch	0.4	54.5	3.4	1.2	0.1	59.7
	Number in Catch	4	498	31	11	1	545
	Mean Length (mm)	573.0	608.0	607.0	609.0	590.0	
Female	Percent of Catch	0.0	37.5	2.5	0.2	0.1	40.3
	Number in Catch	0	342	23	2	1	368
	Mean Length (mm)		592.0	587.0	580.0	610.0	
Total	Percent of Catch	0.4	92.0	5.9	1.4	0.2	100.0
	Number in Catch	4	840	54	13	2	913
Stratum Dates:		8/14 - 8/29					
Sample Size:		140					
		Noatak River					
Male	Percent of Catch	0.7	40.0	10.7	0.7	0.7	52.9
	Number in Catch	1	56	15	1	1	74
	Mean Length (mm)	627	613	625	628	623	
Female	Percent of Catch	0.0	40.7	6.4	0.0	0.0	47.1
	Number in Catch	0	57	9	0	0	66
	Mean Length (mm)	560	606	621	634		
Total	Percent of Catch	0.7	80.7	17.1	0.7	0.7	100
	Number in Catch	1	113	24	1	1	140
Stratum Dates:		7/13-8/27					
Sample Size:		3,288					
		Kotzebue Commercial Catch					
Male	Percent of Sample	0.5	44.9	5.4	0.5	0.1	51.3
	Number in Catch	687	62,439	7,496	660	111	71,369
	Mean Length (mm)	583	609	636	626	636	
Female	Percent of Sample	0.4	42.6	5.2	0.4	0.1	48.7
	Number in Catch	556	59,310	7,213	534	653	67,751
	Mean Length (mm)	554	589	606	613	621	
Total	Percent of Sample	0.9	87.5	10.6	0.9	0.2	100.0
	Number in Catch	1,189	121,749	14,709	1,194	280	139,120




Figure 1. Kotzebue Sound commercial fishing districts, villages and subsistence fishing areas.




Figure 2. Lower Kobuk River drift test fishing sites.

Figure 3. Kobuk River chum salmon drift test fish cumulative CPUE, 1993-1999.

