

KODIAK MANAGEMENT AREA
ANNUAL HERRING MANAGEMENT REPORT, 1996

By

Dennis Gretsch,
Dave Prokopowich,
and
Kevin Brennan

Regional Information Report¹ No. 4K97-37

Alaska Department of Fish and Game
Commercial Fisheries Management and Development Division
211 Mission Road
Kodiak, Alaska 99615

May 1997

¹The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished division reports. These reports frequently serve diverse ad hoc informational purposes or archive basic uninterpreted data. To accommodate timely reporting of recently collected information, reports in this series undergo only limited internal review and may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Commercial Fisheries Management and Development Division.

ACKNOWLEDGEMENTS

The author extends his appreciation to the industry personnel, buyers/processors, spotters and permit holders, whose cooperation, observations, and comments continue to support the harvest strategies under which the Kodiak herring fisheries are managed. A special thanks goes to the ADF&G herring field crew personnel, Morris Lambdin, Ed Hajdys, Tom Emerson, Mike Anderson, Donna Higginbotham, Jon Becker, Keith Wible, Paul Kuriscak, and Forrest Bowers for their inseason catch reporting and fishery data collection. Thanks also goes to other ADF&G employees Joan Brodie, Joanne Shaker for AWL analysis, and Lucinda Neel for publication assistance.

TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES	i
LIST OF FIGURES	ii
LIST OF APPENDICES	iv
SAC ROE HERRING FISHERY	1
Area Description	1
Historical Perspective	1
New Regulations	2
Fishery Characteristics	3
Fishery Monitoring	4
Fishing Seasons and Weekly Fishing Periods	5
Districts and Management Units	5
Guideline Harvest Levels	6
Inseason Fishery Management	6
Fish Ticket Data	7
Biomass Estimates	7
Commercial Catch Sampling	7
1996 Season Summary	8
District Summaries	9
West Afognak District	9
North Afognak District	9
South Afognak District	10
Uganik District	10
Uyak District	10
Alitak District	10

TABLE OF CONTENTS (Cont.)

	<u>Page</u>
Eastside District	11
Northeast District	11
Inner Marmot District	12
North Mainland District.....	12
Mid-Mainland District	12
South Mainland District.....	12
Sturgeon/Halibut District.....	12
Age Composition, Weights, and Lengths	12
Spawning Biomass	13
Enforcement Issues	13
1997 Management Plans and Issues.....	14
HERRING FOOD/BAIT FISHERY	14
Historical Perspective.....	14
Fishery Characteristics	15
Management Plan History.....	15
1996-97 Season Summary.....	16
1997-98 Management Plans and Issues	17
HERRING SUBSISTENCE/PERSONAL USE FISHERY	18
Fishery Characteristics	18
1996 Season Summary	18
LITERATURE CITED	19
TABLES	20
FIGURES	32
APPENDIX	48

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Historical harvest and effort level for the sac roe herring fishery for the Kodiak Management Area, 1964-1996.....	20
2. Summary of Commercial Fisheries Entry Commission status of sac roe herring limited entry permits, Kodiak Management Area, 1990-1996	21
3. Sac roe herring fishery summary of season length, guideline harvest level (GHL), harvest data by gear type, percentage of harvest by gear type, number of landings, and exvessel earnings, Kodiak Management Area , 1979-1996	22
4. Sac roe herring fishery guideline harvest level by management unit, harvest by gear type, and the date each management unit was closed, Kodiak Management Area, 1996	23
5. Age composition of sac roe herring harvests by management unit, Kodiak Management Area, 1996	27
6. Comparison of age composition, weight, and length (AWL) data from the sac roe herring fishery, Kodiak Management Area, 1989-1996.....	28
7. Historical food/bait herring harvest for the Kodiak Management Area, 1912-1996	29
8. Commercial food/bait herring age-weight-length summary of harvest by gear type for the Uganik Food/Bait Management District F/B 4, Kodiak Management Area, 1996	30
9. Commercial food/bait herring age-weight-length summary of the purse seine harvest from the Alitak Food/Bait Management District F/B 7, Kodiak Management Area, 1996	31
10. Commercial food/bait herring age-weight-length summary of the purse seine harvest from the Eastside Food/Bait Management District F/B 8, Kodiak Management Area, 1996	31

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Map of southwestern Alaska emphasizing the Kodiak Management Area and its relationship to surrounding management areas	32
2. Historical sac roe herring harvest for the Kodiak Management Area, 1964-1996	33
3. Number of units of each gear type which made landings in the Kodiak Management Area sac roe herring fishery, 1979-1996.....	33
4. Map of the Kodiak Management Area illustrating the sac roe herring fishery districts and management units which were closed to fishing for the 1996 season.....	34
5. Comparison of the projected guideline harvest level (GHL) to the actual sac roe herring harvest in the Kodiak Management Area, 1979-1996	35
6. Percent of sac roe herring harvest by gear type for the Kodiak Management Area, 1979-1996	35
7. Sac roe herring harvest by district and gear type for the Kodiak Management Area, 1996.....	36
8. Statistical chart showing the North Afognak, West Afognak, South Afognak, and Inner Marmot Districts and Sections.....	37
9. Statistical chart showing the Uganik and Uyak Districts and Sections.....	38
10. Statistical chart showing the Alitak District and Sections.....	39
11. Statistical chart showing the Eastside District and Sections	40
12. Statistical chart showing the Northeast District and Sections	41
13. Statistical chart showing the North Mainland District and Sections.....	42
14. Statistical chart showing the Mid Mainland and Sturgeon-Halibut Districts and Sections.....	43
15. Statistical chart showing the South Mainland District and Sections.....	44

LIST OF FIGURES (Cont.)

<u>Figure</u>	<u>Page</u>
16. Summary of sac roe herring average length and weight at age, and percent of each age class in commercial catch samples for the Kodiak Management Area, 1996	45
17. Age frequency in percent by management unit for the sac roe herring fishery, Kodiak Management Area, 1996	46
18. Historic food/bait herring harvest for the Kodiak Management Area, 1912-1996	47
19. Historic food/bait herring harvest for the Kodiak Management Area, 1965-1996	47

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
A. Summary of emergency order abstracts issued for the sac roe herring fishery, Kodiak Management Area, 1996	49

SAC ROE HERRING FISHERY

Area Description

The Kodiak Management Area (KMA) comprises the entire Kodiak archipelago and that portion of the Alaska Peninsula which drains into Shelikof Strait between Cape Douglas and Kilokak Rocks at Imuya Bay. The archipelago is approximately 150 miles long and extends from Shuyak Island south to the Trinity Islands. The Alaska Peninsula portion is about 160 miles long and is separated from the archipelago by the Shelikof Strait which averages 30 miles in width (Figure 1).

Historical Perspective

The Pacific herring *Clupea pallasii*, sac roe fishery began in Kodiak in 1964. From 1964-1996 sac roe herring harvests have averaged 1,909 short tons (Table 1; Figure 2). Prior to 1974, the fishery was unregulated regarding harvest quotas, gear types, seasons, and fishing periods. From 1964-1977 purse seine gear was used exclusively, with an average annual harvest of 898 tons and up to ten vessels participating in the fishery. Starting in 1974 purse seine gear was restricted to 150 fathoms in length and 1,000 meshes in depth. Between 1974 and 1978 the season ran from March 1 through June 30 with a harvest quota of 3,400 tons. Annual harvests, along with effort levels, fish abundance, prices and processor interest, fluctuated greatly from 1964 through 1977. Improved market conditions in 1978 prompted increased effort in this fishery with 28 purse seiners and 7 gillnetters participating. It was during this time period that spotter aircraft and tenders became involved in the fishery.

Between 1977 and 1982 the regulatory and management strategy went through a rapid developmental phase. Regulatory changes focused on gear efficiency, gear conflicts between seiners and gillnetters, gear level restrictions (exclusive registration and limited entry) and closed waters. In 1979 the sac roe season was reduced to May 1 through June 30, and the overall Guideline Harvest Level (GHL) was reduced to 2,400 tons distributed throughout the management area. A limit of 300 fathoms was also placed on the maximum length of gillnets and fishing periods were first limited to 48 hours open followed by 24 hour closures.

The maximum lengths for gillnets and purse seines were reduced again in 1981 to 150 fathoms and 100 fathoms, respectively. In addition, trawl and beach seine gear were eliminated as legal gear during the sac roe season. Fishing periods were established by emergency order in 1981, in which 24 hour fishing periods were followed by 24 hour closures. Beginning in 1982 the starting date for the season was changed from May 1, to April 15. In 1985 the fixed overall GHL of 2,400 tons was replaced by the current harvest strategy where GHL's are set annually on a stock by stock basis.

With the implementation of limited entry following the 1981 sac roe season, new entry into the fishery was restricted to past participants until permanent transferable permits could be awarded. Since 1982, gear levels have been relatively constant with 90 to 144 units of gear fished annually.

Transferrable permits for both gear types are still increasing as the Limited Entry Commission continues its determinations of participants who may qualify for a transferable permit (Table 2).

The overall regulatory effect during the developmental phase (1977-1982) has been the emergence of a relatively stable commercial sac roe herring fishery through 1991. During the last four years the herring stocks have dramatically increased with record to near record harvests occurring in 1992 through 1995, averaging 4,927 tons.

New Regulations

With the increase in herring stock biomass and effort levels, competition in this fishery has dramatically increased and changes in the fishing schedule were made inseason in 1995. The fishing schedule change was effective from April 21 through May 2 and limited fishing time to ten hour periods from 12:00 noon until 10:00 p.m. on odd numbered days of the month. This change in the fishing schedule was prompted by: 1) ADF&G's inability to monitor the purse seine fleet during the night, 2) near record numbers of purse seine gear participating in the fishery, and 3) low roe recovery standards which were set by some processors, (Gretsch 1995). With the changes in the fishing schedule coupled with a diminishing percentage of the catch going to the gillnet fleet, tension between the two gears types escalated. During the Fall of 1995 the Kodiak Fish and Game Advisory Committee established a work group comprised of gillnet and seine fishers along with ADF&G staff to work on a resolution to better manage the sac roe herring fishery which was to be presented to the Alaska Board of Fisheries (BOF).

The BOF met in Kodiak in December 1995 and after lengthy discussions adopted new regulations concerning gear specifications for seine and gillnets, fishing periods, and a harvest strategy which guides management of the sac roe herring fishery. The new gillnet specifications regulation (Regulation 5 AAC 27.520 c) limits gillnets to a maximum of 230 meshes in depth, the previous regulation had no depth requirements. Additionally the time limit allowed for gillnetters to remove their gear from the water after an emergency closure was increased from two hours to four hours (Regulation 5 AAC 27.520 d) (ADF&G 1996).

The new seine specifications regulation (5 AAC 27.525 a) limits the depth of seines, whereby no purse seine shall exceed 20 fathoms stretch measure in depth or 100 fathoms of length. The depth shall be determined by using a stretch measure of the web from the corkline to the bottom of the net including any lines that hang below the lead line. The previous regulation only specified a seine depth limit of 1,025 meshes with no limits on the size of mesh. This regulation allowed fishers to use larger mesh in part of their seines to increase the depth and efficiency of the gear.

The new fishing periods regulation (5 AAC 27.510) (1) establishes fishing periods from April 15 through May 4 for purse seines to run from 12:00 noon until 10:00 p.m. on odd-numbered days, and from 9:00 a.m. until 12:00 noon on even-numbered days if a harvestable surplus is available; from May 5 through June 30, the fishing period for purse seines is 12:00 noon on odd-numbered days until 12:00 noon on even numbered days; the commissioner may open or close, by emergency order, the season before or after May 5 depending on the assessment of effort, harvest rate, and

ADF&G's ability to monitor the fishery. This new regulation eliminates the majority of the night fishing for purse seiners during the time period when gear levels are at the highest. For gillnetters (2) from April 15 through June 30, fishing periods will run from 12:00 noon on odd-numbered days until 12:00 noon on even-numbered days. Gillnet fishing periods remain unchanged from when first established in 1981.

Lastly, the BOF put into regulation a harvest strategy (5 AAC 27.535 e) which in essence has been used by ADF&G staff to manage the sac roe fishery since the early 1980's. The harvest strategy outlines how ADF&G shall manage the sac roe fishery as follows; the commissioner shall issue emergency orders as necessary: 1) depending on the ADF&G's assessment of effort levels, harvest rates, and its ability to monitor the fishery; the length of the fishing periods for either or both gear groups may be increased or reduced; 2) the ADF&G shall establish guideline harvest levels (GHL's) by section based on information such as historical data, current and past fishery performance, sampling of commercial catch, and aerial surveys; 3) an individual section shall be closed to fishing once its GHL has been achieved; the GHL for an individual section may be adjusted up or down inseason depending on observed stock abundance; the ADF&G may make adjustments in the management of a section based on inseason changes in fish behavior or harvest patterns; these changes may result in an adjacent section closure; 4) a section that does not have an extensive history of sac roe production may be designated as exploratory with no specified GHL; 5) a section that has been closed because its GHL has been achieved may be reopened, if ADF&G determines that additional herring arriving on the grounds have increased the available biomass to the point that the initial exploitation rate has dropped below 10 percent for that stock; any reopening will require information indicating that juvenile herring, post spawners, or other forage fish are not present and that the ADF&G has the ability to monitor and regulate the reopening on the grounds; ADF&G shall give at least 24 hours advance notice before a section is reopened; 6) the KMA sac roe herring fishery is intended to occur in an orderly fashion, with minimal waste of the resource and within conservation limits as determined by ADF&G without regard to roe recovery standards.

Fishery Characteristics

The current KMA sac roe herring fishery occurs from April 15 through June 30 in 40-50 bays and coastal locations. The fishery opens at 12:00 Noon on April 15, with the entire management area opened at one time, excluding those areas where local stocks require protection. A unique characteristic of this fishery is that it typically commences prior to any major build-up of herring. This allows for a greater distribution of effort, which should reduce harvest rates within individual bays. Both gear types fish the same areas during the same time periods, except for April 15 to approximately May 4 time period when purse seine fishing is closed from 10:00 p.m. to 9:00 a.m. during an opening.

The overall trend in harvest during 1979-1991 was a relatively stable catch level, averaging 2,101 tons per year (Table 3). With the increase in the herring stock's biomass starting in 1992 catches also increased and averaged 3,891 tons for the years 1992 through 1996. Prior to 1978, the entire sac roe herring harvest was taken by seine gear. In 1978 seven units of gillnet gear accounted for 3% of the total harvest. From 1979 to 1995 the percentage of the total harvest by seine gear ranged

from a high of 85% to a low of 60% and averaged 76%. Gillnet percentage of total harvest peaked in 1988 at 40%, and averaged 24% from 1979-1995. For the 1996 fishery 69% of the harvest was from seine gear and 31% was from gillnet gear.

To reduce operational costs and to cover more areas, most purse seiners form combines of two to ten vessels. These combines include one or several tenders and spotter aircraft. By 1979 the use of small, single engine, float equipped airplanes became more prevalent. Airplanes are the most productive way to find and direct seiners to harvestable herring. In 1986, several seiners started using side scanning sonar to locate schools of herring. This technology enabled fishermen to work during any time of the day or night and in adverse weather conditions which were unworkable for airplanes. Sonar technology continues to improve and most seiners are now equipped with scanning sonar equipment.

Gillnet vessels generally work independently and usually rely on processors to provide tenders to deliver their fish to the processing location. A few gillnetters are equipped with scanning sonar but the majority of these fishermen rely on color down-sounding sonar to locate herring schools, or fish areas where seiners are making sets. During the last five years the use of mechanical shakers has increased. The shaker is a common tool and greatly reduces the time and effort needed to remove herring from the net and greatly increases gear efficiency.

From 1978-1983 herring were harvested at or near their spawning area. As fishermen's knowledge increased in identifying these areas, gillnet gear has been fished in deeper waters, (15-25 fathoms) further from the spawning destination. Fishing deeper waters and nets has increased the amount of herring harvested with low roe recovery. In most cases this fish is dumped. If ADF&G field crews document this, then the poundage is subtracted from the management unit GHL. In some cases the low quality herring is sold as bait, which is also subtracted from the management unit GHL.

The ADF&G relies on the fishing industry to establish roe recovery standards. Generally, tenders will have a processor representative onboard to ensure that marketable quality herring are harvested. Competition among shore-based and floating processors results in this fishery having one of the highest exvessel values per ton in the state. The quality of Kodiak sac roe herring is generally high, due to inseason handling of relatively small amounts of herring, over a long time period.

Fishery Monitoring

The ADF&G Commercial Fisheries Management and Development Division (CFMDD) manages this fishery from its Kodiak Office. From 1974-1993 ADF&G used one state vessel to monitor this fishery. In 1994 and 1995 a second state vessel was utilized to monitor the early portion of the fishery. The ADF&G vessel R/V Resolution was used in 1994 and the Fish and Wildlife Protection (FWP) vessel M/V Trooper assisted in 1995, with an ADF&G biologist aboard. For the 1996 season both the R/V Resolution and the M/V Trooper with biologist aboard monitored the fishery. From 1979-1995, in conjunction with the state vessels, several two person ADF&G field crews were also utilized to monitor this fishery.

Field and vessel crews are stationed in management units which have historically produced the major harvests for a district. These crews are positioned in remote bays by chartered float planes or vessel and are equipped with an inflatable boat or aluminum skiff. Daily contact with fishermen, spotters, and tender operators is maintained to acquire fishery data. Current harvest, effort levels, and fleet movements are reported via single side band (SSB) radio at least three times per day. The use of field crews has been a key element in preventing an excessive harvest from occurring and exceeding the GHL. Field crews also identify herring spawning areas and collect age-weight-length (AWL) samples from the commercial harvest. The ADF&G aerial surveillance of the entire area supplements, and often directs, placement of fishery monitoring field crews. The ADF&G office staff tally field crew, processor, and tender reports to assess herring harvests and decide which management units need to be closed to fishing. Industry spotter reports are also used to provide information concerning all aspects of the fishery. A "Kodiak Sac Roe Herring Harvest Strategy" is distributed annually, which describes in detail the guideline harvest levels, regulatory changes, and expected fishing periods (Gretsch et al. 1996).

Fishing Seasons and Weekly Fishing Periods

The KMA sac roe herring fishery opens by regulation on April 15 and closes by regulation on June 30 (ADF&G 1996). Fishing periods are established by emergency order. For the previous 15 years fishing periods have started at 12:00 noon on odd numbered days and closed at 12:00 noon on even numbered days of the month, for both gear types. The 1996 fishery was conducted with the new BOF regulations in effect, which may restrict fishing time from April 15 through May 4 for purse seiners from (10:00 pm to 9:00 am) during an opening. Gillnet fishing periods remain unchanged. Staggered days of fishing provide clearly defined closed periods, which allows ADF&G staff time to assess, summarize, and update all harvest data from previous fishing periods.

For the 1996 season, purse seine periods were allowed from 12:00 noon to 10:00 pm on odd-numbered days and from 9:00 am until 12:00 noon on even-numbered days through April 30. During the early portion of the season as many as 79 seiners were involved in the fishery and effort levels diminished by late April to approximately 15 vessels, with many vessels leaving the KMA for the Togiak herring fishery. With the reduction in seine effort beginning on May 1 fishing periods were increased to 24 hours in duration for the remainder of the season.

Districts and Management Units

The KMA is divided into 13 districts which define geographical areas used in managing the sac roe and food\bait herring fisheries (Figure 4). For the sac roe fishery, each district is then divided into management units which are intended to define the spawning area used by a stock of herring or may be used to define a geographical area. The descriptions of districts and management units is based upon the 1983 datum baseline on current NOAA marine charts and all coordinates are listed in decimal degrees and minutes. There are a total of 81 management units.

Guideline Harvest Levels

Preseason guideline harvest levels (GHL's) are established for all management units which have produced consistent herring harvests in previous seasons. These GHL's reflect the status of a particular stock of herring by management unit or district. Criteria for establishing the 1996 GHL include: 1) 1995 expected biomass vs. actual biomass estimates, 2) trends in age composition, 3) level of recruitment primarily (age-3), 4) proportion of the spawning population age-5 and younger, 5) proportion of age-2 fish in the spawning biomass (indicator of future recruitment), and 6) spawn observations (extent, frequency, amount deposited). This information is supplemented by fishery performance information, i.e. the expected vs. actual harvest timing, duration, and level. Some management units are designated "exploratory" and are assigned no GHL because these areas have had sporadic or no harvest of herring in past years. Inseason closures in these exploratory areas are used to ensure that excessive harvests are minimized. If at any time during the season it appears that preseason expectations are incorrect, GHL's can be adjusted above or below preseason levels.

From 1979-1982 the KMA GHL was fixed at 2,400 tons. From 1983-1996 the GHL has varied each year (based on the criteria listed above) from a low of 1,640 in 1987 and a high of 4,550 in 1994. The preseason GHL has accurately reflected the actual harvests (Figure 5). These preseason harvest projections aid fishermen and processors in planning prior to the start of each season.

Inseason Fishery Management

Inseason management of the sac roe fishery relies primarily on ADF&G field crews stationed in management units where harvests are anticipated. Mobility of field crews to cover management units has improved in recent years with the use of three, 21' skiffs and state vessels. Presently two skiffs are used by field crews and a third skiff works from the ADF&G vessel R/V K-Hi-C. These skiffs allow field crews to monitor more management units and under rougher sea conditions than the 12 foot inflatable boats. Two crews still utilize inflatable boats which permit the crews to be rapidly moved by aircraft. The R/V K-Hi-C acts as mobile field station along with providing logistical support to field crews.

Generally, once the preseason GHL has been achieved for a management unit, it is closed for the season. Due to the rapid pace at which some fisheries occur, in-period closures are frequent. In management units which have an ADF&G field crew present, in-period closures may occur with as little as 15 minutes advance notice. In management units which do not have field crews present in-period closures may occur by: 1) announcement on single side band frequency 4.125 Mhz following the marine weather forecast at 8:00 a.m. or 6:00 p.m. daily and at 11:00 P.M. by ADF&G announcement, or 2) field announcement with the arrival of an ADF&G representative.

Processors and independent tender operators are required to provide daily tallies of herring deliveries by management unit and accurate estimates of herring onboard tenders that have not yet delivered to the cannery. Timely and accurate harvest reports from ADF&G field crews, fishermen, spotters, and processors are critical for assessing herring harvests which guide the management of the fishery. To date industry cooperation has been excellent in support of this fishery.

Actual fishery performance is used to evaluate the health of a particular fishery. Key components include: 1) duration of fishing time to harvest the management unit GHL, 2) catch per unit of effort, and 3) quality of herring harvested (including roe recovery, weight, and age composition).

Fish Ticket Data

Commercial catch data is compiled by ADF&G, CFMDD personnel. Actual dock weights of delivered herring are used inseason to verify initial harvest estimates. All final data are compiled post season from sales receipts (fish tickets) received from processors of purchased tonnages of herring. Fish ticket data is then compiled, and a summary of the herring harvest is generated. The ADF&G staff edits this summary for errors and lost fish tickets.

Biomass Estimates

The ADF&G has attempted in previous years to conduct aerial surveys to assess the total KMA herring biomass. The results of aerial assessments provided only a limited evaluation of the biomass and did not give a true representation of the total. Problems associated with aerial surveys in the KMA include: 1) herring tend to spawn in the evening, night, and early morning hours, limiting the time fish are visible in shallow water, 2) most management units have many distinct schools of herring which will spawn from April through June, 3) large numbers of juvenile herring, spawning herring, spawned out herring, and other fishes such as capelin *Mallotus villosus* can be found in sac roe herring fishery areas (fish may stay within an area for the duration of the sac roe season or may move, so that aerial biomass estimates may be duplicated or be incomplete), 4) the large geographical area for the KMA (57 management units which have identified spawning stocks), and 5) adverse weather conditions. Industry spotters have helped greatly in past seasons by providing biomass estimates, spawn observations, fleet movements, and harvest estimates. These spotters are very experienced, many having been involved for several seasons in the KMA and other statewide herring fisheries. Biomass estimates are compiled for each district from surveys flown by industry and ADF&G spotters. It has been estimated by both ADF&G and industry spotters that only 25% to 50% of the actual biomass is observed for the KMA herring stocks. There appears to be a significant amount of subtidal spawning occurring in waters 10-20 fathoms in depth. These fish and spawning activity may not be detected from aerial surveys. Previous attempts to assess this subtidal spawning with divers were not successful.

Commercial Catch Sampling

Commercial catch samples are taken from purse seine harvests, except when a management unit has only a gillnet harvest. Seine caught herring are preferred for samples, since this gear type is less size selective than gillnet gear. Field crews collect samples from multiple seine sets within a management unit to obtain a representative sample of all age classes in the catch. Samples are also obtained from tenders and/or fishing boats delivering to the processor if it is known that the catch

being delivered came from a single management unit. Catch samples are frozen upon arrival in Kodiak and are analyzed by the end of the sac roe season. Commercial catch samples are thawed and analyzed for age, weight, length, sex, and sexual maturity.

A single scale is removed from the preferred area, located on the left side of the fish, three rows below the lateral line and three scales posterior to the center of the opercular plate (J. Brodie, ADF&G, Kodiak, personal communication 1994). The scale is visually analyzed with the aid of a microscope to estimate the age of the fish.

Standard length measurements are taken on all herring sampled. This length is the straight line distance from the anterior most part of the fish, including the lower jaw with the mouth closed, to the end of the vertebra (hypural plate). Lengths are taken on all samples using a herring measuring board to the nearest millimeter (mm).

Weight measurements are taken on a Mettler balance to the nearest gram (g) of all fish within a sample.

The sex and sexual maturity of all sampled herring are recorded. Each fish is slit open and visually inspected for relative gonad maturity; relative maturity is broken down into a scale of key characteristics ranging from virgin herring through spawned out herring, with eight levels of maturity identifying gonad key characteristics.

1996 Season Summary

The 1996 KMA sac roe herring season was 69 days in duration and a total of 3,386 tons of herring were harvested, which was 19 percent lower than the preseason GHF of 4,180 tons (Table 3). This was the fifth largest harvest for the fishery, exceeded by the 1992-1995 seasons which averaged 4,927 tons annually. Seine caught herring totaled 2,322 tons, (69% of the total harvest), while gillnet gear accounted for 1,064 tons or 31%. Harvest by gear type from 1979-1995 has averaged 76% (2,140 tons) and 24% (626 tons) for seine and gillnet gear, respectively (Figure 6; Table 3). The 1996 roe recovery averaged 10.9% for seine caught fish which is about a 1% increase from the 1995 fishery and 10.6% for gillnet caught fish which is slightly down from the 1995 fishery. The average price per ton paid for 10% roe recovery herring was approximately \$2,000 per ton. This was the highest price per ton paid in the history of the fishery. The average exvessel earnings for purse seine fishers was a record \$81,500 and for gillnetters a record \$28,800. The total exvessel value of the fishery was estimated at a record \$6.8 million, which is over twice the 1986-1995 average value of the fishery at \$2.8 million. (Table 3).

A total of 78 seiners and 74 gillnetters participated in the 1996 fishery, approximately 21 seiners did not make a delivery during the season. The 1996 seine effort was the highest experienced since 1981. A total of 102 tenders registered to transport herring within the KMA in 1996. There were 11 floating processors and seven shorebased plants, representing 15 companies registered to process herring within the KMA.

The increase in seine and floating processor effort can mainly be attributed to the closure of the Prince William Sound Management Area sac roe herring fishery. The relatively high preseason GHL for the KMA attracted seiners and processors to Kodiak. The floating processors operated in the vicinity of Port Bailey in Kupreanof Strait, near the major harvest locations of west Afognak Island and the Uganik District. The floating processors departed the KMA at the end of April to participate in the Togiak herring fishery.

District Summaries

The majority of the 1996 harvest was taken in the West Afognak, Uganik, Alitak, and Eastside Districts (Figure 7). Of the 81 management units in the KMA, herring were harvested from 43 units (Table 4), 20 of these units were closed inseason by emergency order (Appendix A), and 20 units were closed prior to the start of the season due to low stock abundance.

West Afognak District. The bays along the westside of Afognak Island are among the earliest areas in which herring are harvested in the KMA. Fishery monitoring crews were stationed in management units which were anticipated to have the earliest harvests. A field crew monitored the Raspberry Strait management unit, the R/V Resolution monitored the Paramanof and Foul Bay units, and the M/V Trooper monitored Malina Bay (Figure 8). For the first fishing period on April 15, a large portion of the seine and gillnet fleet fished the Paramanof Bay management unit and harvested approximately 460 tons. Although the initial harvest was near the 500 ton GHL the fishery performance along with biomass observations indicated that the stock status for this unit was better than expected and the fishery was allowed to reopen on April 17. The second opening lasted 21 minutes and closed at 12:21 PM, with approximately 235 tons harvested. The seasonal total harvest was 695.4 tons. Fishery activity was slow in the remaining management units of this district through the first week of the season and most of the effort moved to the Uganik District after the Paramanof Bay closure. A total of 118.3 tons (GHL of 200 tons) was harvested from Malina Bay and poor fishery performance and low biomass estimates prompted the closure of this unit on April 26. The M/V K-Hi-C monitored the Foul Bay management unit in late April to early May, which had a harvest of 123.9 tons (GHL of 150 tons) and this unit was closed on May 4.

The Raspberry Strait management unit for the second year in a row experienced poor fishery performance with only 7 tons harvested during the last two seasons combined. The 1994 biomass observations and fishery harvests indicated a strong stock status for this unit. There is no clear explanation for the dramatic change in the Raspberry Strait stock status and the decline in the Malina Bay stock.

Of the six management units in this district which have GHL's, three were closed with the assistance of ADF&G field crews and the remainder were open through June 30. The total GHL for the district was 1,180 tons and a total of 944 tons were harvested. Purse seiners accounted for 81% of the harvest and gillnetters accounted for 19%.

North Afognak District. There are five management units within this district and four were closed prior to the start of the 1996 season (Figure 8). Herring stocks in these sections have declined in

recent years, and a closure of these units was initiated in 1995 in an attempt to rebuild these stocks. There was no herring harvest from the one open management unit, the Shuyak Island Section.

South Afognak District. There are six management units within this district and all were closed prior to the start of the 1996 season (Figure 8). The herring stocks in this district have also declined and all units were closed starting in 1995 to help rebuild these stocks.

Uganik District. A field crew was stationed in the Terror Bay management unit and the R/V K-Hi-C was stationed at the Village Islands management unit to monitor the fishery within the Uganik District (Figure 9). On April 15 a large concentration of herring were present within the Village Island management unit, however the herring were in deep waters and were difficult to catch with 35 tons harvested through the first opening. With the closure of Paramanof Bay on April 17 this fleet moved to the Village Islands where the effort levels grew to approximately 75 seiners, 30 gillnetters, and 40 tenders. Fishery activity intensified near darkness with many sets being made at the 10:00 p.m. seine period closure time. After tallying the harvest, the Village Islands management unit was closed at 11:00 p.m. with a total catch of 332.6 tons, (GHL 250 tons). The fleet next dispersed to the Terror Bay, East Arm Uganik, and South Arm Uganik management units. The East Arm Uganik Bay management unit was closed on April 20 with a harvest of 112.7 tons (GHL 125 tons). The Viekoda Bay unit was closed on April 26 with a harvest of 70.9 tons, (GHL 50 tons). The R/V K-Hi-C was then moved to the Foul Bay management unit in late April while the field crew continued to monitor Terror Bay through May 11. The South Arm Uganik management unit was closed on May 4 with a harvest of 283.3 tons, (GHL 180 tons). Catches slowly accumulated in the West Uganik Passage and this unit was closed on May 10 with a harvest of 36.2 tons, (GHL of 40 tons). Fishery performance for the Terror Bay management unit was below expectations for the season with 116.6 tons harvested, (GHL of 250 tons), however fisher and spotter observations of Age-2 and -3 herring within this unit indicated good recruitment. The Northeast Arm Uganik Bay management unit remained open through June 30 and experienced a harvest of 10.3 tons, (GHL 30 tons). Of the eight management units in the Uganik District with GHL's, five were closed with the assistance of ADF&G field crews and the remainder were open through June 30. A district total of 963 tons were harvested from a district GHL of 935 tons, 74% of this harvest was from purse seine gear and 26% by gillnet.

Uyak District. The Uyak District was the largest herring producing district within the KMA through the 1980's (Figure 9). Since 1991 fishery performance and spotter observations have indicated a decline in abundance of herring in this district. The ADF&G responded to this decline by reducing the GHL's for these management units from 1992-1994. The entire district was closed during the 1995 and 1996 herring seasons as a further step to promote the recovery of these stocks.

Alitak District. The Alitak District is comprised of ten management units, two are exploratory areas, and eight have GHL's (Figure 10). Fishing activity started within this district at the end of April and the majority of the seine fleet departed Alitak in early May to participate in the Togiak herring fishery. The Sulua Bay management unit has been the most productive unit within the Alitak District during the last two years with 288 tons harvested in 1996, (GHL of 240 tons) and this unit was closed on May 4. An ADF&G skiff equipped fieldcrew was stationed in the Inner Deadman Bay management unit on May 11. The Geese/Twoheaded management unit was closed

on May 10 with a harvest of 33.2 tons, (GHL 15 tons). The Portage Bay section, (GHL 75 tons) was closed inseason on May 22 with a harvest of 75.9 tons, (GHL of 75 tons). All remaining management units of the Alitak District remained open through June 30. The fishery performance for the Inner and Outer Deadman Bay management units was poor and the harvest totaled 13 tons (GHL 40 tons) and 23.8 tons (GHL 95 tons) respectively. The ADF&G crew moved from Inner Deadman Bay to Olga Bay on May 30 to monitor the herring fishery and to begin installation of salmon weir projects. The Olga Bay herring stock has declined over the last five years and the GHL has been adjusted lower to reflect this changing stock status. A total of 10.2 tons were harvested this season and the GHL was 40 tons. The Inner Alitak exploratory unit had a harvest of 3 tons and the Lower Olga/ Moser unit had a harvest of 2.2 tons. The two remaining management units within this district had no harvest. The Alitak District total GHL was 530 tons with 449.3 tons harvested. Seine gear accounted for 57% of the harvest and gillnet gear 43%. Of the eight management units in the Alitak District with GHL's, one was closed with the assistance of the ADF&G field crew, two were closed by ADF&G office staff, and the remainder were open through June 30.

Eastside District. In general the Barling Bay and the East and West Sitkalidak Strait management units experience early herring harvests, however this season these fisheries appeared to have developed later as catches for the early openings were small. The R/V Resolution deployed a two person ADF&G field crew at Ameer Bay within the East Sitkalidak management unit (Figure 11). This crew was equipped with a 21 foot skiff which enabled them to monitor seven management units along the eastside of Kodiak Island. A raft equipped field crew was also flown into Kiliuda Bay. On April 21 the Barling Bay management unit (GHL of 50 tons) was closed with a harvest of 54.3 tons. The Shearwater Bay management unit was closed prior to reaching the GHL due to an error the fieldcrew made in assessing the harvest, with a closure occurring on May 2 with a harvest of 53.5 tons, (GHL 90 tons). The Outer Kiliuda management unit was closed on May 7 with a harvest of 103.8 tons, (GHL of 90 tons). The Outer Ugak Bay management unit was closed on May 8 with a harvest of 54.2 tons, (GHL of 60 tons). The Three Saints Bay unit was closed on May 9 with a harvest of 47.3 tons, (GHL of 40 tons). The Inner Kiliuda Bay management unit was closed on May 11, with a harvest of 79.9 tons, (GHL of 90 tons), and the field crew was moved to the Inner Ugak Bay management unit. Fishery activity remained relatively slow in the East and West Sitkalidak Strait management units and the fieldcrew was moved to the Alitak District on May 11. On May 13 the Inner Ugak Bay management unit was closed with a harvest of 94.1 tons, (GHL of 90 tons) and the fieldcrew returned to Kodiak. Of the 14 management units with GHL's in the Eastside District, six units were closed with assistance of ADF&G field crews, one unit was closed by the ADF&G office staff, and the remaining units were open until June 30. The total GHL for the Eastside District was 1,190 tons. A total of 825.5 tons were actually harvested, with 49% of the harvest from purse seine gear and 51% from gillnet gear.

Northeast District. There are five management units in the Northeast District and four have GHL's (Figure 12). The Woman's Bay management unit was closed on June 4, with a harvest of 49.9 tons, (GHL of 50 tons). Harvests also occurred in the Kalsin Bay and Middle Bay management units of 4.4 tons (GHL of 10 tons) and 0.2 tons (GHL of 10 tons) respectively. Four management units were open through June 30.

Inner Marmot District. There are five management units within the Inner Marmot District and only one unit has a GHL (Figure 8). The Anton Larsen Bay, Sharatin Bay, and Kizhuyak Bay management units were closed prior to the start of the season due to low stock abundance. The Spruce Island management unit had the only harvest 0.5 tons, (GHL of 10 tons). Seine gear and gillnet gear accounted for 40% and 60%, respectively, of the total district harvest, both units were open through June 30.

North Mainland District. The North Mainland District is comprised of four management units (Figure 13). One unit has a GHL, two units are exploratory, and one unit is offshore. The offshore unit is not expected to produce a sac roe herring harvest. The three Mainland Districts experience more extreme weather conditions than the other districts around Kodiak and Afognak Islands. Sea conditions encountered while crossing the Shelikof Strait to reach these districts greatly reduces the mobility of vessels fishing this district. The Mainland Districts frequently experience high winds, low cloud ceilings, and limited visibility, greatly limiting the effectiveness of spotters. Fishing effort in these three districts generally involves only one or two seine combines and 5-10 gillnet vessels annually. No field crews are stationed in these districts due to the high expense of placing and supplying crews in this remote area. The weather conditions, combined with the small number of vessels which fish these units reduces the likelihood that excessive harvests will occur. A total of 11.9 tons were harvested by gillnetters from the Inner Kukak Bay management unit (GHL 50 tons). There were no other harvest from this district.

Mid-Mainland District. The Mid-Mainland District is comprised of six management units. Two units have a GHL, two units are exploratory, and two units are offshore. The offshore units are not expected to produce a sac roe herring harvest (Figure 14). The Inner Katmai management unit had a harvest of 63.3 tons, (GHL 50 tons) and was closed on May 13. The Alinchak management unit had a harvest of 20.2 tons (GHL 40 tons) and remained open through June 30. There were no other harvests from this district and all harvests were from purse seine gear.

South Mainland District. The South Mainland District consists of two management units, one has a GHL and the second is an exploratory unit (Figure 15). These two management units on the southern part of this district are the farthest units from the port of Kodiak. A total of 54.6 tons were harvested with seine gear from the Wide Bay management unit (GHL 95 tons).

Sturgeon/Halibut District. The Sturgeon/Halibut District on the southwest portion of Kodiak Island has no management units or GHL and consists mostly of offshore areas that are not likely to produce a sac roe herring harvest (Figure 14).

Age Composition, Weights, and Lengths

During the 1996 season, age-3 herring were the dominant age class (39%) found in the purse seine harvest (Figure 16). The remaining age classes represented the following percentages of the harvest: age-2 (1%), age-4 (6%), age-5 (12%), age-6 (5%), age-7 (2%), age-8 (23%), age-9 (10%) and Age-10+ (2%). In general, the West Afognak and Uganik Districts had a dominance of age-8

and age-3 herring while the Eastside District management units had a dominance of age-3 and age-9 herring (Figure 17 and Table 5).

Age-3 herring are considered "recruit herring", entering into the commercial fishery and spawning for the first time. When compared to the previous two brood years, age-3 herring showed a slight decrease in average weight in the 1996 harvest at 93 grams (Table 6). The 1991 age-3 harvest of 81 gram average weight herring coincided with the strong age class which contributed to four years of record harvests. The 1996 lower age-3 weights combined with a high percentage of age-3 herring caught in the fishery may indicate a strong recruitment of age-3 herring into the KMA. Age-5 herring caught in the 1996 fishery were the largest by weight for the years 1989-1995. All other age classes had comparable growth to the past seven years.

Spawning Biomass

In 1996 the spawning biomass index for that portion of the KMA fished, after the 1996 fishery, was estimated to range from 26,000 to 36,000 tons, as determined by industry spotter and ADF&G surveys. The 1996 biomass is similar to the 1992 estimated biomass of 32,000 tons, a year which had a harvest of 4,283 tons. The indexed biomass has increased dramatically from 5,500 tons to 15,500 tons in 1990 and steadily increased to a high which ranged from 42,500 to 56,000 tons in 1993. The estimated biomass has declined during the last three years. A breakdown by district where data is available includes; 7,000-10,000 tons in the West Afognak District, 8,000-10,000 tons in the Eastside District, 8,000-10,000 tons in the Uganik District, 2,000-4,000 tons in the Alitak District, and 1,000-2,000 tons in the Mainland Districts. The 1996 sac roe herring harvest of 3,386 tons represents a total indexed exploitation rate ranging from 9% to 13% of the spawning biomass.

These exploitation rates should be qualified, since surveys represent an unknown and undoubtedly highly variable proportion of the actual biomass. These exploitation rates can be used for trend evaluation, but should not be compared to the spawning biomass indices achieved by ADF&G in Prince William Sound, Cook Inlet, and Bristol Bay. These areas have a relatively large biomass available for aerial indexing and the observed biomass is annually less variable, so there is more opportunity for observing a greater and more consistent proportion of the actual total biomass. The exploitation rates achieved in these fisheries would be more comparable between areas.

Enforcement Issues

The Alaska Department of Public Safety, Fish and Wildlife Protection (FWP) substantially increased their enforcement coverage of the KMA herring fishery during the last two seasons. As previously mentioned, the FWP vessel M/V Trooper worked jointly with ADF&G in monitoring the fishery and conducting enforcement work, with an ADF&G biologist onboard. Additionally, the FWP vessel M/V Spiridon and a FWP float equipped Cessna 185 provided surveillance of the fishery.

The presence of FWP greatly reduced the burden on ADF&G field crews, especially during openings, closures, and emergency closures. With the new seine depth regulation in effect for the

1996 season many purse seiners voluntarily had their nets checked for legal depth in Kodiak by FWP. During the fishery the majority of the enforcement activity occurred on late (after the closure time) purse seine sets. There were some problems with fishers determining management unit boundaries within the Uganik and West Afognak Districts, which will need to be corrected during the next BOF meeting. It is hoped that FWP will continue this level of enforcement activity which contributed to a more orderly fishery.

1997 Management Plans and Issues

Based on the age class data collected, fishery performance, spawn observations, and juvenile herring observations in 1996, the preliminary GHF for the KMA in 1997 is expected to be approximately 3,500 tons, down from the 1996 GHF of 4,180 tons. The 1997 harvest is expected to be comprised of 40-50% age-3 and age-4 herring along with 20-30% age-9 and age-10 herring. The observations of age-2 and-3 herring during the 1996 season indicate a good level of recruitment for KMA, which suggests that the fishery will support harvests of 3,000-3,500 tons over the next few seasons.

The ADF&G will continue to rely greatly on industry spotter pilots, processors, and fishermen to provide information to help manage this fishery. The ADF&G will be conducting an Age Structured Analysis Study for the West Afognak District in 1997 to better evaluate the status of the herring stocks and aid in the setting of GHF's for this district. The ADF&G will also be conducting a test fish program to generate funds for herring research in 1997. Research activity will involve an increase in aerial surveys of closed management units to assess trends in the stock status. Further, to help evaluate the stock status in the Uyak District a purse seine vessel may be contracted to conduct hydroacoustic surveys and collect AWL samples in 1997.

The KMA harvest strategy has been tested with record harvests and high gear levels during the last four years. The competition between fishers is intense and gear conflicts between fishers occasionally occur. A report which reviews the results of the 1996 fishery and evaluates the BOF approved harvest strategy was published in October 1996, "Kodiak Management Area 1996 Commercial Sac Roe Herring Fishery Season Summary" (Gretsch and Brodie 1996).

HERRING FOOD/BAIT FISHERY

Historical Perspective

The earliest recorded food/bait herring harvest for the KMA occurred in 1912 (Table 7). In the early 1920's the fishery expanded and large herring were sought for food products, such as salted and pickled herring which were much in demand after World War I. By the late 1920's the demand for herring food products had declined and the fishery switched from a food product to reduction products, such as fish meal and oil. During the peak years of the reduction fishery (1934-1950) the average harvest was 31,600 tons which vastly surpasses the current food/bait herring harvests

(Figure 18). During the reduction fishery the major harvest areas were located in eastern Shelikof Strait and adjacent bays and straits along the west side of Kodiak and Afognak Islands. Quotas and harvest weights were measured by barrels (where 250 lbs. of herring equals one barrel) until 1956 when the unit of measure was changed to short tons. Historically large, approximately 70 foot, "sardine seiner" type vessels were used in conjunction with holding pounds to supply herring to five major reduction plants. In addition, small seine and gillnet vessels participated in a portion of the food fishery and delivered to floating and shore based salting and pickling operations.

From the early 1960's to 1973, there were no harvest quotas or closed seasons. Beginning in 1974, an open fishing season was established between August 1 through February 28 which remains in effect; however no regulatory GHL's were in effect until 1979. In 1979 and 1980, the GHL was 12,600 tons for the food and bait season. As a result of the rapidly developing sac roe fishery, the GHL for the food/bait season was reduced to 1,000 tons in 1981 and remained at that level through 1987. Regulatory GHL's for the food/bait herring fishery were replaced with a regulatory harvest strategy in 1988.

Gear used in this fishery includes trawl, gillnet, and seine. Gear for this fishery was first regulated for the 1986/87 season when seine gear was restricted to 100 fathoms in length and 1,025 meshes in depth. Gillnet gear was also restricted to 150 fathoms in length with no depth requirements. For the 1993/94 season, purse seine specifications were increased to 150 fathoms in length and 1,625 meshes in depth. There are no trawl restrictions. The entire KMA is open to continuous fishing on August 1 for all legal gear types and there are no exclusive gear areas.

Fishery Characteristics

The current food/bait herring fishery can be characterized as a secondary commercial fishery on herring concentrations located in KMA waters. It is primarily a bait fishery providing a frozen product for longline and crab/cod pot fishers. Effort and harvest levels are at historical lows for the food/bait fishery, which is converse to the sac roe fishery. The food/bait fishery is an open-to-entry fishery, while the sac roe fishery has been limited-to-entry since 1981. Existing regulations designate priority status to the sac roe fishery.

Management Plan History

During the Fall and Winter months of the early 1980's major concentrations of herring were observed in eastern Shelikof Strait and adjacent bays along the west side of Kodiak and Afognak Islands. The biomass exceeded that of known KMA spawning stocks. These herring were targeted by food/bait fishers and questions arose concerning the stock of origin of these fish. In 1986, a stock identification study based on scale pattern analysis was performed on herring harvested from a large biomass located in the east part of the Shelikof Strait (Johnson et al. 1988). The study suggested that at least 80% of the East Shelikof herring sampled were of Kamishak Bay spawning stock origins, which is within the Lower Cook Inlet Management Area.

In 1988, the BOF allocated not more than two percent of the previous season's total available spawning biomass from Kamishak to be harvested during Kodiak's food/bait herring fishery. For local Kodiak spawning stocks, which are exploited during the sac roe fishery, the food/bait GHL on those same stocks is 10% of the previous seasons sac roe herring harvest.

Problems arose from this management plan because it was difficult to assign harvest from the intermixed stocks to Kodiak or Kamishak. This plan was in effect through the 1992/93 season.

In 1992, the BOF approved the Kamishak Bay District Herring Management Plan (5AAC27.465) which outlines criteria for the management of the Kamishak Bay sac roe herring and the Shelikof Strait food/bait fishery (ADF&G 1996). This plan defines allocations to these fisheries based on biomass estimates.

In 1993 the BOF placed into regulation a harvest strategy defining the criteria for managing the Kodiak food/bait herring fishery (5AAC 27.535). This strategy combines the Kamishak stock GHL with the Kodiak stock GHL for food/bait management units FB 1, FB 4, and FB 5 (Figures 8 and 9). When this combined GHL is achieved, the Shelikof Strait food and bait management units are closed collectively. This plan alleviates the problem of identifying the spawning stock of a harvest in areas where intermixing may occur.

The allocation of Kamishak Bay herring stocks to the Shelikof Strait food and bait fishery is based on the spawning biomass of age-5 and older herring and not on the biomass of juveniles. The quantity of herring stocks aged four years and younger caught during the food/bait fishery are adjusted to approximate the biomass of a similar number of age-5 herring. Age-4 and younger herring were selected because in the Kamishak spawning stocks, herring are not considered to have attained complete recruitment into the spawning biomass until they have reached age-5.

1996-97 Season Summary

The 1996/97 allocation for Kamishak herring stocks over wintering in Shelikof Strait was 380 tons. In addition, the GHL for Kodiak stocks in food/bait management units FB 1 and FB 4 was 190 tons for a combined GHL for both stocks affected by (5 AAC 27.465) of 570 tons. A "KMA Commercial Food/Bait Herring Fishery Harvest Strategy" is distributed annually, which describes in detail the guideline harvest levels, regulatory changes, and registration requirements (Gretsch et al. 1996).

The Kodiak spawning stocks total GHL was 333 tons, which represents 10% of the previous spring's sac roe herring harvest on a stock by stock basis. The total GHL for the KMA was 713 tons (380 tons Kamishak allocation and 333 tons Kodiak stocks).

The KMA food/bait herring season began August 1, 1996 and remained open until February 28, 1997. Fishing periods were 24 hours per day and seven days a week. Three emergency orders (E.O.'s) were issued during the season which closed districts after the GHL was harvested. The East Afognak (Food/Bait unit #3), North Afognak (Food/Bait unit #2), Uyak (Food/Bait unit #5),

and the Inner Marmot (Food/Bait unit #10) were closed prior to the start of the season. The sac roe herring stocks within these districts have declined during the last five years so these units were closed to prevent further exploitation during the food/bait fishery.

All permit holders and buyers are required to register at the Kodiak ADF&G office prior to fishing or purchasing herring. At that time, management plans are issued and catch reporting procedures and current regulations are reviewed. Each landing is sampled for age, weight, length (AWL) information and skipper interviews are conducted to evaluate which sac roe stocks are being impacted. For the 1996/97 season a total 651 tons were harvested in the KMA. From 1965 through 1995 the food/bait harvest has averaged 205 tons (Figure 19). Eight vessels and five buyers/processors registered for this fishery. Purse seine gear accounted for 68% of the total harvest and trawl gear 32%. The total exvessel value of this fishery was approximately \$260,000 dollars.

The food/bait management districts FB1, FB4, FB11, and FB12 which are affected by the combined harvest of Kodiak-Kamishak herring stocks, were closed on September 16 with a harvest of 511 tons, (GHL of 580 tons). As stated in the Kamishak Bay District Herring Management Plan (5 AAC 27.465) the quantity of age-4 and younger herring harvested is adjusted to a similar number of age-5 herring. This estimated adjusted harvest totaled 626 tons. During this fishery purse seine gear harvested primarily age-3 and age-4 herring while trawl gear harvested a mix of age classes, primarily age-6 and age-8 (Table 8).

The Alitak food/bait management district FB7 was closed on October 30 with a harvest of 53.3 tons (GHL of 44 tons). This harvest by purse seine gear was primarily of age-2 and age-3 herring (Table 9).

The Eastside food/bait management district FB8 was closed on November 3 with a harvest of 86.6 tons (GHL of 80 tons). This harvest by purse seine gear was primarily of age-8 and age-9 herring (Table 10).

No biomass surveys were conducted by the ADF&G in 1996 on overwintering herring concentrations.

1997/98 Management Plans and Issues

The success of purse seine gear in this fishery during the last four years will likely encourage other seiners to participate next season. On-grounds monitoring by the ADF&G of the fishery may be necessary as gear levels escalate. Improvements in catch reporting will be needed with increasing effort levels to help limit harvests to guideline levels.

HERRING SUBSISTENCE/PERSONAL USE FISHERY

Fishery Characteristics

The subsistence and personal use fishery for herring is regulated only during the sac roe herring fishery season, April 15 through June 30. During this time period, a permit is required for individuals who are not sac roe commercial fishermen to harvest herring. Sac roe commercial fishermen may retain herring from their lawfully taken commercial catch to fulfill their subsistence or personal use needs. Most of the herring caught during this time period are used for bait in commercial longline fisheries. However small amounts are used for food, sport fishing bait, and fertilizer.

1996 Season Summary

A total of 22 permits were issued in 1996 and 3 permits were returned with harvest data. The total harvest was 550 pounds from the South Afognak, Inner Marmot, and Northeast Districts. Generally, most permits are returned during the spring months and the actual harvest will likely be higher than reported at this time.

LITERATURE CITED

- ADF&G (Alaska Department Fish and Game). 1996-97 Commercial herring fishing regulations, 1996. Alaska Department of Fish and Game, Division of Commercial Fisheries, Juneau.
- Gretsch, D., 1995. Kodiak management area sac roe herring fishery briefing document. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K95-41, Kodiak.
- Gretsch, D., D. Prokopowich, and K. Brennan. 1996. Kodiak management area sac roe herring harvest strategy. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K96-17, Kodiak.
- Gretsch, D. and J. Brodie 1996. Kodiak management area 1996 commercial sac roe herring fishery season summary. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K96-49, Kodiak.
- Gretsch, D., D. Prokopowich, and K. Brennan. 1996. Kodiak management area commercial food/bait herring fishery harvest strategy 1996/97. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K96-43, Kodiak.
- Johnson, B.A., C. Burkey, and D. Gaudet. (Draft manuscript 1988). Stock identification of Pacific herring in the bait fishery in Shelikof Strait, Alaska, 1985/86. Alaska Department of Fish and Game, Division of Commercial Fisheries. Juneau.

Table 1. Historical commercial harvest data and effort levels for the sac roe herring fishery Kodiak Management Area, 1964-1996.

Year	Total Harvest (Tons)	Harvest by Gear Type		Number of Vessels Making Landings			Number of Processors
		Seine (Tons)	Gillnet (Tons)	Seine	Gillnet	Total	
1964	568	568	0	5	0	5	2
1965	657	657	0	8	0	8	2
1966	2,769	2,769	0	11	0	11	4
1967	1,662	1,662	0	5	0	5	4
1968	2,001	2,001	0	10	0	10	4
1969	1,130	1,130	0	21	0	21	9
1970	342	342	0	13	0	13	5
1971	284	284	0	4	0	4	2
1972	215	215	0	4	0	4	1
1973	831	831	0	11	0	11	4
1974	868	868	0	26	0	26	4
1975	8	8	0	2	0	2	3
1976	5	5	0	1	0	1	1
1977	338	338	0	11	0	11	3
1978	904	881	23	28	7	35	7
1979	1,735	1,457	278	57	125	182	8
1980	2,383	2,009	374	92	109	201	9
1981	2,065	1,596	469	79	114	193	9
1982	1,771	1,447	324	45	67	112	6
1983	2,318	1,797	521	41	64	105	7
1984	2,163	1,691	472	39	69	108	7
1985	1,968	1,244	724	34	81	115	7
1986	1,558	1,111	447	31	71	102	8
1987	2,146	1,591	555	29	62	91	8
1988	2,171	1,304	867	33	76	109	6
1989	2,249	1,513	736	37	83	120	6
1990	2,347	1,644	703	27	63	90	6
1991	2,432	1,697	735	32	64	96	6
1992	4,283	3,260	1,023	40	74	114	6
1993	4,929	4,203	726	41	86	127	6
1994	5,893	4,976	917	66	57	123	15
1995	4,604	3,837	767	73	71	144	17
1996	3,386	2,322	1,064	57 ^a	74	131	15
Average 1964-1996	1,909	1,553	355	30	43	74	6
Average 1991-1995	4,428	3,595	834	50	70	121	10
Average 1986-1995	3,261	2,514	748	41	71	112	8

^a Approximately 20 vessels participated in the fishery but made no deliveries.

Table 2. Summary of Commercial Fisheries Entry Commission status of sac roe herring fishery limited entry permits, Kodiak Management Area, 1990-1996.

Gear Type	Year						
	1990	1991	1992	1993	1994	1995 ^a	1996 ^b
Gillnet							
Transferable	72	74	97	95	99	94	94
Non-Transferable	27	28	11	8	8	5	9
Total Gillnet Permits	99	102	108	103	107	99	103
Purse Seine							
Transferable	47	48	59	66	69	65	68
Non-Transferable	25	22	13	12	14	13	13
Total Seine Permits	72	70	72	78	83	78	81
Combination (Seine or Gillnet)							
Total Transferable	2	2	2	2	2	2	2
Combined Total							
Transferable	121	124	158	163	170	161	164
Non-Transferable	52	50	24	20	22	18	22
Total Permits	173	174	182	183	192	179	186

^a There were 11 gillnet permits and 1 seine permit which were not renewed in 1995, which are not included in this table.

^b There were 20 gillnet permits and 6 seine permits which were not renewed in 1996, which are not included in this table.

Table 3. Sac roe herring fishery summary of season length, guideline harvest level (GHL), harvest data by gear type, percentage of harvest by gear type, number of landings, and exvessel earnings, Kodiak Management Area, 1979-1996.

Year	Season Length (Days)	G-H-L (Tons)	Total Harvest (Tons)	Harvest by Gear Type		Percent Harvest by Gear Type		Number of Landings by Gear Type		Units of Gear Making Landings		Average Catch by Gear Type		Average Earnings		Price per Ton (\$)	Exvessel Total Value (\$)
				Seine (Tons)	Gillnet (Tons)	Seine	Gillnet	Seine	Gillnet	Seine	Gillnet	Seine (Tons)	Gillnet (Tons)	Seine (\$)	Gillnet (\$)		
1979	36	2,400	1,735	1,457	278	84%	16%	-	-	57	125	26	2	\$ 38,342	\$ 3,336	\$ 1,500	\$ 2,602,500
1980	35	2,400	2,383	2,009	374	84%	16%	-	-	92	109	22	3	\$ 15,068	\$ 2,368	\$ 690	\$ 1,844,270
1981	48	2,400	2,065	1,596	469	77%	23%	207	406	79	114	20	4	\$ 14,647	\$ 2,983	\$ 725	\$ 1,497,125
1982	59	2,400	1,771	1,447	324	82%	18%	138	191	45	67	32	5	\$ 17,686	\$ 2,660	\$ 550	\$ 974,050
1983	51	2,400	2,319	1,797	522	77%	23%	164	284	41	64	44	8	\$ 35,063	\$ 6,525	\$ 800	\$ 1,855,200
1984	54	2,400	2,163	1,691	472	78%	22%	138	212	39	69	43	7	\$ 34,687	\$ 5,472	\$ 800	\$ 1,730,400
1985	59	2,000	1,968	1,244	724	63%	37%	118	348	34	81	37	9	\$ 32,929	\$ 8,044	\$ 900	\$ 1,771,200
1986	61	1,690	1,558	1,110	448	71%	29%	132	385	31	71	36	6	\$ 34,016	\$ 5,994	\$ 950	\$ 1,480,100
1987	61	1,840	2,146	1,591	554	74%	26%	122	411	29	62	55	9	\$ 54,862	\$ 8,935	\$ 1,000	\$ 2,146,000
1988	59	2,065	2,171	1,304	867	60%	40%	169	555	33	76	40	11	\$ 51,370	\$ 14,830	\$ 1,300	\$ 2,822,300
1989	76	2,415	2,249	1,513	736	67%	33%	171	627	37	83	41	9	\$ 34,749	\$ 7,535	\$ 850	\$ 1,911,149
1990	75	2,375	2,347	1,644	703	70%	30%	156	544	27	63	61	11	\$ 51,756	\$ 9,485	\$ 850	\$ 1,994,950
1991	83	2,510	2,432	1,697	735	70%	30%	169	587	32	64	53	11	\$ 45,077	\$ 9,762	\$ 850	\$ 2,087,200
1992	77	2,720	4,283	3,260	1,023	76%	24%	185	706	40	74	82	14	\$ 40,750	\$ 6,912	\$ 500	\$ 2,141,500
1993	77	3,525	4,929	4,203	726	85%	15%	237	294	41	86	103	8	\$ 56,382	\$ 4,643	\$ 550	\$ 2,710,950
1994	71	4,550	5,893	4,978	917	84%	16%	285	485	66	57	75	16	\$ 60,315	\$ 12,870	\$ 800	\$ 4,714,400
1995	73	4,480	4,604	3,637	768	83%	17%	280	642	73	71	53	11	\$ 66,859	\$ 13,759	\$ 1,272	\$ 5,856,288
1996	69	4,180	3,386	2,322	1,064	69%	31%	202	690	57	74	41	14	\$ 61,474	\$ 28,757	\$ 2,000	\$ 6,772,000
Averages																	
1979-1995	62	2,610	2,766	2,140	626	76%	24%	157	393	47	70	48	9	\$40,268	\$7,415	\$876	\$2,348,211
1979-1991	68	2,228	2,101	1,546	544	74%	26%	130	350	44	81	38	7	\$35,404	\$6,764	\$905	\$1,884,342
1992-1996	73	3,891	4,619	3,720	900	80%	20%	238	603	55	72	71	13	\$61,156	\$13,366	\$1,024	\$4,439,028

Table 4. Sac roe herring fishery guideline harvest level by management unit, harvest by gear type, and the date each management unit was closed, Kodiak Management Area, 1996.

STAT. AREA	MGMT. UNIT	GUIDELINE HARVEST LEVEL (Tons)	SEINE HARVEST (Tons)	GILLNET HARVEST (Tons)	TOTAL HARVEST (Tons)	DATE UNIT CLOSED
NORTH AFOGNAK DISTRICT						
NA10	Shuyak Is.	20	0	0	0	6/30
NA20	Delphin Bay	Closed	0	0	0	-
NA30	Perenosa Bay	Closed	0	0	0	-
NA40	Seal Bay	Closed	0	0	0	-
NA50	Tonki Bay	Closed	0	0	0	-
District Total		20	0	0	0	-
WEST AFOGNAK DISTRICT						
WA10	Raspberry St.	300	0	3.8	3.8	6/30
WA20	Malina Bay	200	116.1	2.2	118.3	4/26
WA31	Paramanof Bay	500	538	157.4	695.4	4/17
WA32	Foul Bay	150	108.9	15	123.9	5/4
WA40	Devils Inlet	10	0	0	0	6/30
WA40	Blue Fox	20	0	2.6	2.6	6/30
WA50	Offshore W. Afog. ^a	-	0	0	0	6/30
District Total		1,180	763.0	181.0	944.0	-
SOUTH AFOGNAK DISTRICT						
SA10	Izhut Bay	Closed	0	0	0	-
SA20	Kitoy Bay	Closed	0	0	0	-
SA30	MacDonalds Lagoon	Closed	0	0	0	-
SA40	Danger Bay	Closed	0	0	0	-
SA50	Litnik	Closed	0	0	0	-
SA60	Duck Bay	Closed	0	0	0	-
District Total		Closed	0	0	0	-
UGANIK DISTRICT						
UG10	Kupreanof	10	0	0.1	0.1	6/30
UG20	Viekoda	50	69.2	1.7	70.9	4/26
UG21	Terror	250	32.7	83.9	116.6	6/30
UG30	Village Island	250	305.6	27	332.6	4/17
UG31	W. Uganik Pass	40	24.9	11.3	36.2	5/10
UG32	NE Arm Uganik	30	10.3	0	10.3	6/30
UG33	E. Arm Uganik	125	82	30.7	112.7	4/20
UG34	S. Arm Uganik	180	191.4	91.9	283.3	5/4
UG40	Offshore Uganik ^a	-	0	0	0	6/30
District Total		935	716.1	246.6	962.7	-

-Continued-

Table 4. (page 2 of 4)

STAT. AREA	MGMT. UNIT	GUIDELINE HARVEST LEVEL (Tons)	SEINE HARVEST (Tons)	GILLNET HARVEST (Tons)	TOTAL HARVEST (Tons)	DATE UNIT CLOSED
UYAK DISTRICT						
UY10	Offshore Uyak ^a	Closed	0	0	0	-
UY20	Harvester Island	Closed	0	0	0	-
UY30	Inner Uyak	Closed	0	0	0	-
UY32	Browns Lagoon	Closed	0	0	0	-
UY31	Larsen Bay	Closed	0	0	0	-
UY40	Zachar Bay	Closed	0	0	0	-
UY50	Spiridon Bay	Closed	0	0	0	-
District Total		Closed	0	0	0	-
ALITAK DISTRICT						
AL10	Outer Alitak	(Exploration)	0	0	0	6/30
AL20	Inner Alitak	(Exploration)	0	3	3	6/30
AL21	In. Deadman Bay	40	12.9	0.1	13	6/30
AL22	Outer Deadman Bay	95	22.8	1	23.8	6/30
AL30	Sulua Bay	240	107.2	180.8	288	5/4
AL31	Portage Bay	75	69.8	6.1	75.9	5/22
AL40	Lower Olga/Moser	15	0	2.2	2.2	6/30
AL41	No. Upper Olga Bay	10	0	0	0	6/30
AL50	Upper Olga Bay	40	10.2	0	10.2	6/30
AL60	Geese/Twoheaded	15	33.2	0	33.2	5/10
District Total		530	256.1	193.2	449.3	-
STURGEON/HALIBUT DISTRICT						
SH10	Sturg./Halibut	(Exploration)	0	0	0	6/30
EASTSIDE DISTRICT						
EA10	Kaiugnak	20	0	0	0	6/30
EA20	SW. Sitkalidak	20	0	2.3	2.3	6/30
EA21	Three Saints Bay	40	41.3	6	47.3	5/9
EA22	Newman Bay	20	9	2.9	11.9	6/30
EA23	W. Sitkalidak St.	275	38.2	34.3	72.5	6/30
EA24	Barling Bay	50	25.6	28.7	54.3	4/21
EA30	E. Sitkalidak St.	300	94.9	133.7	228.6	6/30
EA31	Tanginak Anchorage	15	0	1.4	1.4	6/30
EA40	Outer Sitka.	(Exploration)	0	0	0	6/30
EA41	Boulder Bay	(Exploration)	15.4	6.2	21.6	6/30
EA42	Shearwater Bay	90	0	53.5	53.5	5/2
EA43	Outer Kiliuda Bay	90	84.9	18.9	103.8	5/7

-Continued-

Table 4. (page 3 of 4)

STAT. AREA	MGMT. UNIT	GUIDELINE HARVEST LEVEL (Tons)	SEINE HARVEST (Tons)	GILLNET HARVEST (Tons)	TOTAL HARVEST (Tons)	DATE UNIT CLOSED
EASTSIDE DISTRICT (Cont.)						
EA44	Inner Kiliuda Bay	90	77.5	2.4	79.9	5/11
EA50	Outer Ugak Bay	60	9.9	44.3	54.2	5/8
EA51	Inner Ugak Bay	90	7.5	86.6	94.1	5/13
EA52	Pasagshak	30	0	0.1	0.1	6/30
District Total		1,190	404.2	421.3	825.5	-
NORTHEAST DISTRICT						
NE10	Womens Bay	50	44.6	5.3	49.9	6/4
NE20	Kalsin Bay	10	0	4.4	4.4	6/30
NE30	Middle Bay	10	0	0.2	0.2	6/30
NE40	Inshore Chiniak	10	0	0	0	6/30
NE50	Off. Chiniak	(Exploration)	0	0	0	6/30
District Total		80	44.6	9.9	54.5	-
INNER MARMOT DISTRICT						
IM10	Monashka Bay	(Exploration)	0	0	0	6/30
IM20	Anton Larsen Bay	Closed	0	0	0	-
IM30	Sharatin Bay	Closed	0	0	0	-
IM40	Kizhuyak Bay	Closed	0	0	0	-
IM50	Spruce Island	10	0.2	0.3	0.5	6/30
District Total		10	0.2	0.3	0.5	-
NORTH MAINLAND DISTRICT						
NM10	Hallo Bay	(Exploration)	0	0	0	6/30
NM20	Inner Kukak	50	0	11.9	11.9	6/30
NM30	Outer Kukak	-	0	0	0	6/30
NM40	Missak Bay	(Exploration)	0	0	0	6/30
District Total		50	0	11.9	11.9	-
MID MAINLAND DISTRICT						
MM10	Inner Katmai	50	63.3	0	63.3	5/13
MM20	Outer Katmai	-	0	0	0	6/30
MM30	Alinchak	40	20.2	0	20.2	6/30
MM40	Puale Bay	(Exploration)	0	0	0	6/30
MM50	Portage Bay	(Exploration)	0	0	0	6/30
MM60	Outer Portage ^a	-	0	0	0	6/30
District Total		90	83.5	0	83.5	-

-Continued-

Table 4. (page 4 of 4)

STAT. AREA	MGMT. UNIT	GUIDELINE HARVEST LEVEL (Tons)	SEINE HARVEST (Tons)	GILLNET HARVEST (Tons)	TOTAL HARVEST (Tons)	DATE UNIT CLOSED
SOUTH MAINLAND DISTRICT						
SM10	Wide Bay	95	54.6	0	54.6	6/30
SM20	Low. Shelikof	(Exploration)	0	0	0	6/30
District Total		95	54.6	0	54.6	-
GRAND TOTAL		4,180	2,322.3	1,064.2	3,386.5	

^a These are offshore management units which are not expected to yield herring of sac roe quality. These units are more applicable to the food/bait fishery. (See Herring Food/Bait Fishery Management Plan.)

Table 5. Age composition of sac roe herring harvests by management unit, Kodiak Management Area, 1996.^{a, b}

Management Unit	Harvest (tons)	Percent at Age										N
		Age-2	3	4	5	6	7	8	9	10	11+	
Raspberry Strait	4	-	-	6.8	15.9	13.6	11.3	47.7	4.5	-	-	44
Malina Bay	118	-	15.9	7.0	10.5	9.4	1.6	52.2	2.1	-	1.0	371
Paramanof Bay	695	0.3	22.7	7.2	3.2	5.4	2.7	52.7	3.6	-	1.8	610
Foul Bay	124	-	38.2	14.3	13.2	9.6	1.8	21.4	0.4	-	0.7	424
Viekoda Bay	71	-	87.3	7.8	1.3	0.8	-	2.6	-	-	-	229
Terror Bay	117	2.6	65.9	10.9	3.5	4.9	1.4	7.8	0.4	0.4	1.8	485
Village Islands	333	-	1.4	3.1	8.6	11.1	1.9	69.0	3.6	-	0.9	414
W. Uganik Passage	36	-	23.9	6.2	7.2	5.2	2.0	25.0	11.4	4.1	14.5	96
N.E. Arm Uganik	10	3.7	38.8	6.7	11.1	1.4	2.2	4.4	7.4	1.4	22.3	134
E. Arm Uganik	113	0.5	6.5	5.7	6.8	10.2	1.8	63.6	3.9	-	0.7	382
S. Arm Uganik	283	0.3	6.0	4.6	10.7	10.3	4.0	57.5	4.3	-	2.0	299
Zachar Bay ^c	-	-	-	8.5	40.4	8.5	8.5	23.4	8.5	-	2.1	47
Inner Deadman Bay	13	-	7.1	10.0	44.2	8.5	4.2	21.4	4.2	-	-	70
Sulua Bay	288	-	1.0	6.2	31.2	14.5	1.0	43.7	2.0	-	-	96
Portage Bay	76	-	10.3	2.7	35.3	-	-	10.8	30.4	1.6	8.6	184
Upper Olga Bay	10	5.4	42.8	7.1	15.9	1.6	9.3	3.8	2.7	9.3	1.6	182
Geese-Twoheaded	33	-	41.0	12.5	37.5	1.7	-	3.5	3.5	-	-	56
Three Saints Bay	47	1.7	79.4	0.5	2.2	-	-	0.5	14.2	-	1.1	175
W. Sitkalidak	73	-	66.1	6.1	17.9	1.5	-	2.0	5.6	-	0.5	195
Barling Bay	54	-	27.4	8.6	27.9	5.9	0.5	13.9	14.5	1.0	-	186
E. Sitkalidak	229	-	37.7	3.4	16.0	5.3	0.3	11.8	20.6	0.7	3.8	262
Outer Kiliuda Bay	104	-	9.6	2.3	18.4	2.3	1.8	4.1	60.3	0.4	0.4	217
Inner Kiliuda Bay	80	1.7	34.2	1.9	12.1	1.9	1.3	3.5	42.8	-	-	502
Shearwater Bay	54	-	-	4.1	27.7	-	-	4.1	62.5	1.3	-	72
Inner Ugak Bay	94	-	88.2	-	1.3	-	1.1	0.2	7.8	0.4	0.6	444
Womans Bay	50	0.3	97.8	0.1	0.3	0.3	0.1	-	0.3	-	0.1	502
Inner Katmai Bay	63	-	23.7	21.7	34.6	1.9	8.9	8.9	-	-	-	101
Wide Bay	55	-	16.2	36.9	32.4	7.2	1.8	5.4	-	-	-	111
Total 28 Mgmt. Units	3,227	0.6	38.6	6.2	11.4	5.0	1.8	23.3	10.4	0.5	1.7	6,890

^a Of the 41 stocks exploited in 1996, samples were collected from 27 (68%). These 27 stocks yielded 3,227 tons or 95% of the management area's total harvest of 3386 tons.

^b The above samples were collected by ADF&G employees from commercial herring seine catches with the exception of a gillnet caught sample from Raspberry Straits, a combined commercial gillnet and seine caught sample from Shearwater, and the gillnet sample caught by ADF&G employees in Zachar Bay.

^c Zachar Bay was closed to commercial herring fishing for the 1996 season.

Table 6. Comparison of age composition, weight, and length (AWL) data from the sac roe herring fishery, Kodiak Management Area, 1989-1996.

YEAR	Age Class										Average	N
	2	3	4	5	6	7	8	9	10	11+		
<i>Percent at Age</i>												
1989	0.3	7.1	6.2	42.0	19.3	1.0	10.0	4.5	8.5	1.1		3,026
1990	0.7	52.0	3.7	3.3	20.4	8.5	0.6	3.0	2.6	5.2		7,672
1991	<0.1	26.3	49.4	2.6	1.4	8.7	5.2	0.6	2.7	3.0		5,498
1992	0.5	2.9	62.3	26.9	2.0	0.7	2.0	1.2	<0.1	1.4		9,325
1993	0.3	8.7	4.5	63.5	18.2	1.2	0.2	1.6	0.8	1.0		7,396
1994	0.1	11.5	11.6	3.8	42.4	24.5	1.9	0.7	2.0	1.5		6,965
1995	0.3	3.2	16.0	8.9	2.2	37.4	28.3	0.7	0.4	2.3		3,775
1996	0.6	38.6	6.2	11.4	5.0	1.8	23.3	10.4	0.5	1.7		6,890
<i>Average Weights (Grams)</i>												
1989	60	85	130	176	224	251	269	271	276	284	199	2,569
1990	52	95	141	167	191	248	229	280	283	290	152	4,885
1991	53	81	137	169	191	221	255	261	301	292	153	4,239
1992	40	87	114	176	201	226	247	284	283	303	140	8,139
1993	58	94	134	144	203	226	226	250	281	310	156	6,852
1994	78	99	129	166	182	251	251	256	282	316	188	6,836
1995	49	109	146	166	200	208	264	270	270	310	209	3,763
1996	52	93	132	187	193	227	230	294	284	305	171	6,768
<i>Average Lengths (Millimeters)</i>												
1989	168	185	210	227	247	254	259	259	260	263	235	3,026
1990	158	188	214	226	232	250	245	261	258	260	212	7,671
1991	166	184	215	225	235	244	251	257	266	263	215	5,497
1992	162	194	202	231	240	250	254	264	265	268	214	9,323
1993	161	191	211	216	240	247	244	254	261	264	219	7,394
1994	175	194	211	227	231	253	252	253	259	267	231	6,965
1995	163	201	220	228	241	243	261	264	265	272	242	3,774
1996	158	192	212	235	236	248	247	266	261	268	223	6,803

Table 7. Historical food/bait herring harvest for the Kodiak Management Area, 1912-1996.

YEAR	TONS	YEAR	TONS	YEAR	TONS
1912	20.0	1941	40,083.5	1970	7.5
1913	0.0	1942	16,791.0	1971	44.2
1914	0.0	1943	35,352.0	1972	49.8
1915	0.0	1944	26,835.0	1973	178.0
1916	70.0	1945	31,114.0	1974	40.1
1917	137.9	1946	47,505.9	1975	5.2
1918	118.4	1947	50,743.0	1976	No Data
1919	259.7	1948	46,428.0	1977	No Data
1920	45.9	1949	0.0	1978	398.9
1921	944.9	1950	44,132.5	1979	124.8
1922	1,482.6	1951	4,299.0	1980	380.7
1923	321.5	1952	1,389.0	1981	18.0
1924	4,823.0	1953	725.0	1982	326.0
1925	9,997.0	1954	0.0	1983	33.4
1926	2,680.9	1955	0.0	1984	123.0
1927	2,592.9	1956	13,524.0	1985	102.0
1928	625.0	1957	21,218.5	1986	213.0
1929	No Data	1958	1,711.0	1987	217.1
1930	622.0	1959	3,831.0	1988	340.2
1931	1,000.0	1960	0.0	1989	344.6
1932	3,594.0	1961	0.0	1990	312.6
1933	2,312.5	1962	0.0	1991	215.3
1934	60,000.0	1963	0.0	1992	311.5
1935	No Data	1964	309.8	1993	837.0
1936	24,748.0	1965	35.0	1994	677.0
1937	27,659.3	1966	198.0	1995	507.0
1938	24,522.0	1967	300.3	1996	651.0
1939	38,600.5	1968	15.4		
1940	22,677.0	1969	11.0		

Table 8. Commercial food/bait herring age-weight-length summary of the harvest by gear type for the Uganik Food/Bait Management District F/B 4, Kodiak Management Area, 1996.

Age (years)	Sex			Total	Percent of Total	Weight			Std. Length			
	Male	Female	Unknown			Mean (gm)	Std. Dev.	Number Weighed	Mean (mm)	Std. Dev.	Number Measured	
Seine Gear												
0	-	-	-	-	-	-	-	-	-	-	-	-
1	-	-	-	-	-	-	-	-	-	-	-	-
2	6	3	-	9	4.5	91	14.8	8	187	11.3	9	
3	51	45	-	96	48.2	137	18.0	89	209	9.3	96	
4	36	35	-	71	35.7	168	20.4	64	223	8.9	71	
5	4	3	-	7	3.5	219	27.4	7	239	11.3	7	
6	5	5	-	10	5.0	225	40.6	9	239	11.6	10	
7	-	1	-	1	0.5	288	-	1	269	-	1	
8	4	1	-	5	2.5	248	19.1	4	251	8.1	5	
9	-	-	-	-	-	-	-	-	-	-	-	
10	-	-	-	-	-	-	-	-	-	-	-	
11+	-	-	-	-	-	-	-	-	-	-	-	
Total	106	93	-	199	100.0	157	38.9	182	217	16.0	199	
Trawl Gear												
0	-	-	-	-	-	-	-	-	-	-	-	
1	-	-	-	-	-	-	-	-	-	-	-	
2	-	-	-	-	-	-	-	-	-	-	-	
3	11	9	-	20	10.4	143	30.9	20	208	10.9	20	
4	8	11	-	19	9.9	181	21.9	19	223	8.8	19	
5	4	6	-	10	5.2	208	26.0	10	234	8.0	10	
6	27	16	-	43	22.4	236	31.3	43	241	10.0	43	
7	10	12	-	22	11.5	253	23.6	22	246	6.6	22	
8	35	26	-	61	31.8	265	33.6	61	251	9.1	61	
9	2	1	-	3	1.6	279	61.0	3	256	14.2	3	
10	1	1	-	2	1.0	247	29.7	2	253	17.7	2	
11+	8	4	-	12	6.3	305	46.7	12	265	11.8	12	
Total	106	86	-	192	100.0	235	53.3	192	241	17.6	192	

Table 9. Commercial food/bait herring age-weight-length summary of the purse seine harvest from the Alitak Food/Bait Management district F/B 7, Kodiak Management Area, 1996.

Age (years)	Sex			Total	Percent of Total	Weight			Std. Length		
	Male	Female	Unknown			Mean (gm)	Std. Dev.	Number Weighed	Mean (mm)	Std. Dev.	Number Measured
1	0	1	0	1	0.7	57	0	1	157	0.0	1
2	16	17	0	33	23.7	112	13	33	196	6.7	33
3	26	20	0	46	33.0	153	19.6	46	216	9.4	46
4	3	5	0	8	5.7	164	17.7	8	219	9.2	8
5	12	9	0	21	15.1	237	27.2	21	246	7.0	21
6	2	5	0	7	5.0	248	33.3	7	248	8.3	7
7	2	3	0	5	3.5	251	50.7	5	250	10.2	5
8	3	2	0	5	3.5	301	28.5	5	264	8.9	5
9	4	3	0	7	5.0	308	28.6	7	262	7.4	7
10	1	0	0	1	0.7	324	0	1	263	0.0	1
11+	3	2	0	5	3.5	320	33.2	5	275	8.6	5
Total	72	67	0	39	100.0	185	71.3	139	225	25.9	139

Table 10. Commercial food/bait herring age-weight-length summary of the purse seine harvest from the Eastside Food/Bait Management district F/B 8, Kodiak Management Area, 1996.

Age (years)	Sex			Total	Percent of Total	Weight			Std. Length		
	Male	Female	Unknown			Mean (gm)	Std. Dev.	Number Weighed	Mean (mm)	Std. Dev.	Number Measured
2	0	2	0	2	0.8	108	14.1	2	203	7.8	2
3	15	9	0	24	10.1	167	17.1	23	224	6.0	24
4	6	3	0	9	3.7	226	40.2	9	245	16.3	9
5	17	23	0	40	16.8	256	31.8	40	255	11.0	40
6	12	5	0	17	7.1	276	25.2	17	259	6.8	17
7	5	2	0	7	2.9	305	23.6	7	267	7.6	7
8	42	25	0	67	28.2	326	32.6	67	269	9.2	67
9	37	29	0	66	27.8	352	34.3	66	277	8.4	66
11+	3	2	0	5	2.1	346	12.2	5	277	5.3	0
Total	137	100	0	237	100.0	297	67.3	236	262	18.8	232




Figure 1. Map of southwestern Alaska emphasizing the Kodiak Management Area and its relationship to surrounding management areas.




Figure 2. Historical sac roe herring harvest for the Kodiak Management Area, 1964-1996.




Figure 3. Number of units of each gear type which made landings in the Kodiak Management Area sac roe herring fishery, 1979-1996.




Figure 4. Map of the Kodiak Management Area illustrating the sac roe herring fishery districts and management units which were closed to fishing for the 1996 season.




Figure 5. Comparison of the projected guideline harvest level (GHL) to the actual sac roe herring harvest in the Kodiak Management Area, 1979 - 1996.




Figure 6. Percent of sac roe herring harvest by gear type for the Kodiak Management Area, 1979 - 1996.




Figure 7. Sac roe herring harvest by district and gear type for the Kodiak Management Area, 1996.




Figure 8. Statistical chart showing the North Afognak, West Afognak, South Afognak, and Inner Marmot Districts and Sections.




Figure 9. Statistical chart showing the Uganik and Uyak Districts and Sections.




Figure 10. Statistical chart showing the Alitak District and Sections.




Figure 11. Statistical chart showing the Eastside District and Sections.




Figure 12. Statistical chart showing the Northeast District and Sections.




Figure 13. Statistical chart showing the North Mainland District and Sections.




Figure 14. Statistical chart showing the Mid Mainland and Sturgeon-Halibut Bay Districts and Sections.






Figure 15. Statistical chart showing the South Mainland District and Sections.



N	47	2666	431	786	347	130	1608	717	36	122	6890
% BY AGE	0.6	38.6	6.2	11.4	5.0	1.8	23.3	10.4	0.5	1.7	100%
N	39	2619	423	778	339	129	1574	710	36	121	6768
AVG. WEIGHT (gm.)	52	93	132	187	193	227	230	294	284	305	171
N	46	2631	427	785	341	130	1574	712	36	121	6803
AVG. LENGTH (mm.)	158	192	212	235	236	248	247	266	261	268	223

Figure 16. Summary of sac roe herring average length and weight at age, and percent of each age class in commercial catch samples for the Kodiak Management Area, 1996.



1/ All samples were collected by ADF&G personnel from commercial seine catches with the exception of a commercial gillnet caught sample from Raspberry Straits, a partial gillnet caught sample from Shearwater Bay and the ADF&G variable mesh caught test sample from Zachar Bay.

Figure 17. Age frequency in percent (%) by management unit for the sac roe herring fishery Kodiak Management Area, 1996.^{1/}




Figure 18. Historic food/bait herring harvest for the Kodiak Management Area, 1912-1996.




Figure 19. Historic food/bait herring harvest for the Kodiak Management Area, 1965-1996.

APPENDIX

Appendix A. Summary of herring emergency orders issued in the Kodiak Management Area, 1996.

E.O.#	Issued	Effective	Action Taken				
1			This EO pertains to the 1995/96 food/bait herring fishery				
2	10:00 AM 4/11/96	12:00 noon 4/15/96	<p><u>Opening</u> beginning 12:00 noon 4/15 as described in the 1996 Kodiak Herring Sac Roe Harvest Strategy for <u>gillnetters</u> fishing time will be for 24 hour open periods beginning at 12:00 noon on the odd numbered days of the month followed by 24 hour closures beginning at 12:00 noon on the even numbered days of the month</p> <p>for <u>purse seiners</u> fishing time will be for 10 hour periods running from 12:00 noon to 10:00 PM on the odd numbered days of the month followed by 3 hour open periods running from 9:00 AM to 12:00 noon on the even numbered days of the month. There will be a 24 hour closure for both gear types beginning at 12:00 noon on the even numbered days of the month.</p> <p>NOTE: The following management units will remain closed for the entire 1996 herring season:</p> <table border="0"> <tr> <td style="vertical-align: top;"> <p><u>South Afognak District</u> (SA10) Izhut Bay (SA20) Kitoi Bay (SA30) MacDonalds Lagoon (SA40) Danger Bay (SA50) Litnik (SA60) Inshore Marmot (Duck Bay)</p> </td> <td style="vertical-align: top;"> <p><u>Uyak District</u> (UY10) Offshore Uyak (UY20) Harvester Island (UY30) Inner Uyak Bay (UY31) Larsen Bay (UY32) Browns Lagoon (UY40) Zachar Bay (UY50) Spiridon Bay</p> </td> </tr> <tr> <td style="vertical-align: top;"> <p><u>North Afognak District</u> (NA20) Delphin Bay (NA30) Perenosa Bay (NA40) Seal Bay (NA50) Tonki Bay</p> </td> <td style="vertical-align: top;"> <p><u>Inner Marmot District</u> (IM20) Anton Larson Bay (IM30) Sharatin Bay (IM40) Kizhuyak Bay</p> </td> </tr> </table>	<p><u>South Afognak District</u> (SA10) Izhut Bay (SA20) Kitoi Bay (SA30) MacDonalds Lagoon (SA40) Danger Bay (SA50) Litnik (SA60) Inshore Marmot (Duck Bay)</p>	<p><u>Uyak District</u> (UY10) Offshore Uyak (UY20) Harvester Island (UY30) Inner Uyak Bay (UY31) Larsen Bay (UY32) Browns Lagoon (UY40) Zachar Bay (UY50) Spiridon Bay</p>	<p><u>North Afognak District</u> (NA20) Delphin Bay (NA30) Perenosa Bay (NA40) Seal Bay (NA50) Tonki Bay</p>	<p><u>Inner Marmot District</u> (IM20) Anton Larson Bay (IM30) Sharatin Bay (IM40) Kizhuyak Bay</p>
<p><u>South Afognak District</u> (SA10) Izhut Bay (SA20) Kitoi Bay (SA30) MacDonalds Lagoon (SA40) Danger Bay (SA50) Litnik (SA60) Inshore Marmot (Duck Bay)</p>	<p><u>Uyak District</u> (UY10) Offshore Uyak (UY20) Harvester Island (UY30) Inner Uyak Bay (UY31) Larsen Bay (UY32) Browns Lagoon (UY40) Zachar Bay (UY50) Spiridon Bay</p>						
<p><u>North Afognak District</u> (NA20) Delphin Bay (NA30) Perenosa Bay (NA40) Seal Bay (NA50) Tonki Bay</p>	<p><u>Inner Marmot District</u> (IM20) Anton Larson Bay (IM30) Sharatin Bay (IM40) Kizhuyak Bay</p>						
3	12:16 PM 4/17/96	12:21 PM 4/17/96	<u>Closure</u> at 12:21 PM 4/17 until further notice - Paramanof Bay Section (WA31)				
4	10:45 PM 4/17/96	11:00 PM 4/17/96	<u>Closure</u> at 11:00 PM 4/17 until further notice - Village Islands Section (UG30)				
5	10:20 AM 4/21/96	12:00 noon 4/20/96	<u>Closure</u> at 12:00 noon 4/20 until further notice - East Arm Uganik Section (UG33)				

-Continued-

Appendix A. (page 2 of 3)

E.O.#	Issued	Effective	Action Taken
6	8:00 PM 4/21/96	10:00 PM 4/21/96	<u>Closure</u> at 10:00 PM 4/21 until further notice - Barling Bay Section (EA24)
7	7:45 AM 4/26/96	8:30 AM 4/26/96	<u>Closure</u> at 8:30 AM 4/26 until further notice - Viekoda Bay Section (UG20)
8	4:00 PM 4/26/96	12:00 noon 4/26/96	<u>Closure</u> at 12:00 noon 4/26 until further notice - Malina Bay Section (WA20)
9	3:00 PM 4/30/96	12:00 noon 5/1/96	<u>Change in fishing periods</u> , beginning 12:00 noon 5/1 until further notice for <u>purse seiners</u> , 24 hours open followed by 24 hours closed
10	3:30 PM 5/2/96	12:00 noon 5/2/96	<u>Closure</u> at 12:00 noon 5/2 until further notice - Shearwater Bay Section (EA42)
11	6:30 AM 5/4/96	7:00 AM 5/4/96	<u>Closure</u> at 7:00 AM 5/4 until further notice - Sulua Bay Section (AL30)
		12:00 noon 5/4/96	<u>Closure</u> at 12:00 noon 5/4 until further notice - Foul Bay Section (WA32)
12	2:00 PM 5/4/96	12:00 noon 5/4/96	<u>Closure</u> at 12:00 noon 5/4 until further notice - South Arm Uganik Section (UG34)
13	10:15 PM 5/7/96	10:25 PM 5/7/96	<u>Closure</u> at 10:25 PM 5/7 until further notice - Outer Kiliuda Bay Section (EA43)
14	4:00 PM 5/8/96	12:00 noon 5/8/96	<u>Closure</u> at 12:00 noon 5/8 until further notice - Outer Ugak Bay Section (EA50)
15	9:40 PM 5/9/96	10:00 PM 5/9/96	<u>Closure</u> at 10:00 PM 5/9 until further notice - Three Saints Bay Section (EA21)
16	2:00 PM 5/10/96	12:00 noon 5/10/96	<u>Closure</u> at 12:00 noon 5/10 until further notice - West Uganik Passage Section (UG31) - Geese-Twoheaded Section (AL60)

-Continued-

Appendix A. (page 3 of 3)

E.O.#	Issued	Effective	Action Taken
17	1:00 PM 5/11/96	1:05 PM 5/11/96	<u>Closure</u> at 1:05 PM 5/11 until further notice - Inner Kiliuda Bay Section (EA44)
18	6:00 PM 5/13/96	8:30 PM 5/13/96	<u>Closure</u> at 8:30 PM 5/13 until further notice - Inner Katmai Section (MM10)
19	8:35 PM 5/13/96	10:00 PM 5/13/96	<u>Closure</u> at 10:00 PM 5/13 until further notice - Inner Ugak Bay Section (EA51)
20	5:00 PM 5/22/96	12:00 noon 5/22/96	<u>Closure</u> at 12:00 noon 5/22 until further notice - Portage Bay Section (AL31)
21	3:00 PM 6/4/96	12:00 noon 6/4/96	<u>Closure</u> at 12:00 noon 6/4 until further notice - Womans Bay Section (NE10)
22	8:00 AM 9/16/96	8:30 AM 9/16/96	<u>Closure</u> at 8:30 AM 9/16 for the remainder of the 1996/97 season - West Afognak Management Unit (Food/Bait Unit #1) - Uganik Management Unit (Food/Bait Unit #4) - North Mainland Management Unit (Food/Bait Unit #11) - Mid Mainland Management Unit (Food/Bait Unit #12) <u>NOTE:</u> The North Afognak Management Unit (F/B 2), South Afognak Management Unit (F/B 3), Uyak Management Unit (F/B 5), and the Inner Marmot Management Unit (F/B 10), will remain closed to food/bait herring for the entire 1996/97 season.
23	11:00 AM 10/30/96	12:00 noon 10/30/96	Closure at 12:00 noon 10/30 for the remainder of the 1996/97 season - Alitak District Management Unit (Food/Bait Unit # 7)
24	11:00 AM 11/3/96	12:00 noon 11/3/96	Closure at 12:00 noon 11/3 for the remainder of the 1996/97 season - Eastside Management Unit (Food/Bait Unit #8)

The Alaska Department of Fish and Game administers all programs and activities free from discrimination on the basis of sex, color, race, religion, national origin, age, marital status, pregnancy, parenthood, or disability. For information on alternative formats available for this and other department publications, contact the department ADA Coordinator at (voice) 907-465-4120, or (TDD) 907-465-3646. Any person who believes s/he has been discriminated against should write to: ADF&G, PO Box 25526, Juneau, AK 99802-5526; or O.E.O., U.S. Department of the Interior, Washington, DC 20240.