

Fishery Management Report No. 06-34

**Management Plan for the Lynn Canal (District 15)
Drift Gillnet Fishery, 2006**

by

Randall L. Bachman

June 2006

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the *Système International d'Unités* (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Measures (fisheries)	
centimeter	cm	Alaska Administrative Code	AAC	fork length	FL
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	mid-eye-to-fork	MEF
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	mid-eye-to-tail-fork	METF
hectare	ha	at	@	standard length	SL
kilogram	kg	compass directions:		total length	TL
kilometer	km	east	E	Mathematics, statistics	
liter	L	north	N	<i>all standard mathematical signs, symbols and abbreviations</i>	
meter	m	south	S	alternate hypothesis	H _A
milliliter	mL	west	W	base of natural logarithm	<i>e</i>
millimeter	mm	copyright	©	catch per unit effort	CPUE
Weights and measures (English)		corporate suffixes:		coefficient of variation	CV
cubic feet per second	ft ³ /s	Company	Co.	common test statistics	(F, t, χ^2 , etc.)
foot	ft	Corporation	Corp.	confidence interval	CI
gallon	gal	Incorporated	Inc.	correlation coefficient (multiple)	R
inch	in	Limited	Ltd.	correlation coefficient (simple)	r
mile	mi	District of Columbia	D.C.	covariance	cov
nautical mile	nmi	et alii (and others)	et al.	degree (angular)	°
ounce	oz	et cetera (and so forth)	etc.	degrees of freedom	df
pound	lb	exempli gratia		expected value	<i>E</i>
quart	qt	(for example)	e.g.	greater than	>
yard	yd	Federal Information Code	FIC	greater than or equal to	≥
Time and temperature		id est (that is)	i.e.	harvest per unit effort	HPUE
day	d	latitude or longitude	lat. or long.	less than	<
degrees Celsius	°C	monetary symbols		less than or equal to	≤
degrees Fahrenheit	°F	(U.S.)	\$, ¢	logarithm (natural)	ln
degrees kelvin	K	months (tables and figures): first three letters	Jan, ..., Dec	logarithm (base 10)	log
hour	h	registered trademark	®	logarithm (specify base)	log ₂ , etc.
minute	min	trademark	™	minute (angular)	'
second	s	United States (adjective)	U.S.	not significant	NS
Physics and chemistry		United States of America (noun)	USA	null hypothesis	H ₀
all atomic symbols		U.S.C.	United States Code	percent	%
alternating current	AC	U.S. state	use two-letter abbreviations (e.g., AK, WA)	probability	P
ampere	A			probability of a type I error (rejection of the null hypothesis when true)	α
calorie	cal			probability of a type II error (acceptance of the null hypothesis when false)	β
direct current	DC			second (angular)	"
hertz	Hz			standard deviation	SD
horsepower	hp			standard error	SE
hydrogen ion activity (negative log of)	pH			variance	
parts per million	ppm			population	Var
parts per thousand	ppt, ‰			sample	var
volts	V				
watts	W				

FISHERY MANAGEMENT REPORT NO. 06-34

**MANAGEMENT PLAN FOR THE LYNN CANAL (DISTRICT 15)
DRIFT GILLNET FISHERY, 2006**

by

Randall L. Bachman,

Alaska Department of Fish and Game, Division of Commercial Fisheries, Haines

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1565

June 2006

The Division of Sport Fish Fishery Management Reports series was established in 1989 for the publication of an overview of Division of Sport Fish management activities and goals in a specific geographic area. Since 2004, the Division of Commercial Fisheries has also used the Fishery Management Report series. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm>. This publication has undergone regional peer review.

*Randall L. Bachman,
Alaska Department of Fish and Game, Commercial Fisheries Division, Region 1
Mile 1 Haines Highway, P.O. Box 330, Haines, AK 99827-0330, USA*

This document should be cited as:

Bachman, R. L. 2006. Fishery management plan for the Lynn Canal (District 15) drift gillnet fishery, 2006. Alaska Department of Fish and Game, Fishery Management Report No. 06-34, Anchorage.

The Alaska Department of Fish and Game administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility, or if you desire further information please write to ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; U.S. Fish and Wildlife Service, 4040 N. Fairfax Drive, Suite 300 Webb, Arlington, VA 22203 or O.E.O., U.S. Department of the Interior, Washington DC 20240.

For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-6077, (TDD) 907-465-3646, or (FAX) 907-465-6078.

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	ii
LIST OF FIGURES.....	ii
LIST OF APPENDICES.....	iii
ABSTRACT.....	1
INTRODUCTION.....	1
FISHERY AREA.....	1
CONTRIBUTING STOCKS.....	1
Regulatory Decision Process.....	2
GENERAL GOAL.....	2
MANAGEMENT GOALS.....	2
2006 OUTLOOK.....	3
Chilkat River Drainage Sockeye Salmon.....	3
Chilkoot Sockeye Salmon.....	3
berners sockeye salmon.....	4
Summer Chum Salmon.....	4
Fall Chum Salmon.....	5
Coho Salmon.....	5
Chinook Salmon.....	5
Fishery Openings.....	5
Section 15-A.....	6
Section 15-B.....	6
Section 15-C.....	6
Other Comments.....	7
Stock Assessment Projects.....	7
REFERENCES CITED.....	9
TABLES AND FIGURES.....	11
APPENDIX A: LYNN CANAL AND CHILKAT RIVER KING SALMON FISHERY MANAGEMENT PLAN..	41
APPENDIX B: CALENDAR DATES FOR STATISTICAL WEEKS IN 2006.....	45
APPENDIX C: HISTORICAL AGE COMPOSITIONS OF SOCKEYE SALMON ESCAPEMENTS TO CHILKOOT LAKE, CHILKAT LAKE, CHILKAT RIVER MAINSTEM AND BERNERS BAY RIVERS.....	47
APPENDIX D: DATA COLLECTED FROM THE INSEASON INFORMATION SYSTEM TO DETERMINE FISHERY PERFORMANCE BY SPECIES.....	53
APPENDIX E: INCLUSIVE DATES OF OPERATION FOR CHILKOOT LAKE AND CHILKAT LAKE WEIRS AND CHILKAT RIVER FISH WHEELS.....	57

LIST OF TABLES

Table	Page
1. Historical catches of king, sockeye, coho, pink, and chum salmon in the District 15 (Lynn Canal) drift gillnet fishery, 1976–2005.....	12
2. Biological and sustainable escapement goals for Lynn Canal salmon stocks by species and location.	14
3. Yearly preseason smolt based forecast compared to the estimated total return of adult Chilkat Lake sockeye salmon, 1992, 1997–2006.....	15
4. Annual total return of Chilkat Lake sockeye salmon by week, 1976 to 2005.	16
5. Annual escapements of Chilkat Lake sockeye salmon by week, 1976–2005.....	18
6. Chilkat Lake sockeye salmon smolt age, weight, and length compositions (wild and enhanced components), 1989–1990 and 1994–2005.....	20
7. Percent age, average length and average weight composition of Chilkat Lake sockeye salmon smolt, 1989–1990 and 1994–2005.	21
8. Weekly and annual escapement of Chilkat River mainstem sockeye salmon, 1994–2005.	22
9. Annual total return of Chilkoot Lake sockeye salmon by week, 1978–2005.	23
10. Annual weir counts of Chilkoot Lake sockeye salmon by week, 1976–2005.	25
11. Selected data for Chilkoot Lake, annual autumn hydroacoustic pre-smolt estimates, total adult return, and average annual zooplankton densities for years 1987–2005.....	27
12. Annual harvests of Chilkat River mainstem and Berners Bay rivers, and other non-Chilkat or Chilkoot Lake, sockeye salmon by week, 1976– 2005.	28
13. Summary of releases of DIPAC chum salmon from Boat Harbor and Amalga Harbor, 1988–2005.	30
14. Summary of returns from DIPAC chum salmon enhancement projects in lower Lynn Canal, 1991–2004 with projections for 2005.....	31

LIST OF FIGURES

Figure	Page
1. Lynn Canal district, section and statistical area boundaries.	32
2. Upper Lynn Canal showing Chilkat and Chilkoot lakes. (Note: the statistical area 115-35 (formerly Burro Creek SHA) has been changed to represent the waters of Taiya Inlet north of Taiya Point.).....	33
3. Lynn Canal sockeye salmon weekly abundance by stock (Data from 1983-1992 average).....	34
4. Run timing (weekly proportion CPUE) of Chinook, coho, summer and fall chum, and pink salmon in the Lynn Canal drift gillnet fishery. Data for period 1992 to 2005.	34
5. Historical commercial catches of Chinook, sockeye, coho, pink, and chum (summer and fall) salmon in the District 15 (Lynn Canal) drift gillnet fishery, 1960 to 2005.....	35
6. Historical escapement and harvest of Chilkoot and Chilkat lake sockeye salmon, 1976 to 2005.	36
7. Yearly comparisons of Chilkoot Lake autumn hydroacoustic counts of juvenile sockeye salmon and average zooplankton densities, 1987–1991 and 1995–2005.	37
8. Peak aerial survey results for Sawmill Creek chum salmon, 1993-2005 compared with 1996-2005 average results.	37
9. Peak aerial survey results for Western Lynn Canal chum salmon streams combined, 1995–2005 compared to the 1996-2005 average.....	38
10. Total Chilkat River coho salmon fish wheel catch by year, 1994–2005 compared to the 1994-2005 average.	38
11. Average 1994–2005 run timing for Chilkat River sockeye salmon stocks at the Chilkat River fish wheels by fresh water age class.....	39
12. Map showing possible area in Lower Lynn Canal that may be opened for additional time during peak weeks of hatchery chum return (Statistical Weeks 28–31).....	40

LIST OF APPENDICES

Appendix	Page
A1. Lynn Canal and Chilkat River King Salmon Fishery Management Plan 5 AAC 33.384.....	42
B1. Calendar dates for statistical weeks in 2006.....	46
C1. Historical age composition of sockeye salmon escapements to Chilkat Lake, 1982–2005.....	48
C2. Historical age composition of sockeye salmon escapements to Chilkat River mainstem areas, 1984 to 2005.....	49
C3. Historical age composition of sockeye salmon escapements to Chilkoot Lake, 1982-2005.	50
C4. Historical age composition of sockeye salmon escapements to Berners Bay rivers, 1984 to 2005.....	51
D1. Data collected from the inseason information system to determine fishery performance by species.....	54

ABSTRACT

Generally, for 2006, returns of all salmon species to Lynn Canal is expected to be average to above average in run strength. Hatchery chum salmon returning to Amalga and Boat Harbor release sites are expected to be at least average in abundance. Sockeye salmon returns to major spawning areas in northern Lynn Canal are expected to be average. Pink salmon returns are expected to be well above average. Fall chum and coho returns are expected to be near average this year. This document describes plans for the management of the 2006 Lynn Canal (District 15) drift gillnet salmon fishery.

The purpose of this plan is to provide commercial fishers and processors with a general idea of how the fishery is managed, what options are available to the manager, the conditions that trigger major management actions and an idea of the expected run size of targeted salmon stocks. This plan will also serve as a reference that consolidates important historical harvest and escapement data and current fisheries information.

Key words: salmon, drift gillnet fishery, Lynn Canal (District 15), management plan.

INTRODUCTION

This document describes plans for the management of the 2006 Lynn Canal drift gillnet fishery. The purpose of this plan is to provide commercial fishers and others with a general idea of how the fishery is managed, what options are available to the manager, the conditions that trigger major management actions and an idea of the expected run size of targeted salmon stocks to the Lynn Canal district. This plan will also serve as a reference that consolidates important historical harvest and escapement data and current fisheries information.

The Lynn Canal drift gillnet fishery targets sockeye *Oncorhynchus nerka*, summer and fall chum *O. keta*, and coho *O. kisutch* salmon. Chinook *O. tshawytscha* and pink *O. gorbuscha* salmon are harvested as incidental catch. The sockeye run in Lynn Canal has historically been among the largest in Southeast Alaska. The coho run to the Chilkat River is among the largest in northern Southeast Alaska. Currently stocks of Lynn Canal sockeye, coho and fall chum salmon are meeting escapement goals and productive. During recent years, total returns of Chilkoot Lake sockeye salmon have improved and fishing effort has been directed on this stock. Returns of fall chum salmon have also improved and fishing effort has been directed on this stock. Fall chum salmon escapements based on fish wheel catch, escapement surveys and results from mark-recapture projects in 2002 through 2005, have indicated improved total returns of this stock.

FISHERY AREA

The Lynn Canal drift gillnet fishery occurs in the waters of District 15. The district is divided into three regulatory sections: 15-A (upper Lynn Canal), 15-B (Berners Bay), and 15-C (lower Lynn Canal). These regulatory sections are further divided into seven statistical areas (Figure 1).

CONTRIBUTING STOCKS

Stocks targeted by the gillnet fishery and the harvest timing are as follows:

1. Sockeye salmon contribute to the fishery from June through early September. The primary stocks originate in Chilkat and Chilkoot lakes (Figure 2). Sockeye salmon originating from Chilkat Lake, Chilkoot Lake and other locations are managed separately. Sockeye salmon from rivers within Berners Bay, the mainstem of the Chilkat River, and other small stocks are also harvested in the fishery. The return timing for the Chilkat and Chilkoot lake sockeye salmon stocks in the Lynn Canal drift gillnet fishery is presented in detail in Figure 3.

2. Douglas Island Pink and Chum (DIPAC) hatchery chum and wild pink salmon stocks are both harvested from late June through mid-July. The return timing for Chinook, summer chum, pink and coho salmon stocks to the Lynn Canal drift gillnet fishery is shown in Figure 4.
3. Fall chum and coho salmon are harvested from September through early-October. The primary fall chum salmon stocks originate in the Klehini and Chilkat rivers. The primary coho salmon stocks originate in the Chilkat River drainage and rivers within Berners Bay.

For 2006, the southeast Alaska drift gillnet fishery is limited to 10,057 “Treaty” Chinook salmon. This allocation does not include Alaskan hatchery produced fish. This amount is based on 2.9% of the Chinook all gear quota for Southeast Alaska. In District 15, the Lynn Canal and Chilkat River King Salmon Fishery Management Plan (5 AAC 33.384, shown in Appendix A1) will guide the management in Chilkat Inlet during the first three weeks of the season based on a preseason projected inriver forecast. This plan establishes management measures in subsistence, commercial, and sport fisheries, which harvest Chilkat River Chinook salmon. The return timing for coho, chum, pink and Chinook salmon stocks harvested by the Lynn Canal drift gillnet fishery is shown in Figure 4.

Historical catches for sockeye, coho, pink, chum, and Chinook salmon in the Lynn Canal drift gillnet fishery are shown in Table 1 and Figure 5.

REGULATORY DECISION PROCESS

The fishery will open at 12:01 p.m. on June 18, the third Sunday of June (statistical week 26, Appendix B1). Weekly fishing periods are established by emergency order and announced to the public by Alaska Department of Fish and Game (ADF&G) news releases that are generally issued on Thursday afternoons.

GENERAL GOAL

The overall management goal is to achieve desired spawning escapement levels while harvesting the available surplus for a long-term maximum sustainable yield of all Lynn Canal salmon stocks. Escapement to Chilkoot Lake is monitored by a weir located on the outlet of Chilkoot Lake. Escapements to Chilkat River and Chilkat Lake are monitored using fish wheels operated in the lower Chilkat River and a weir located near the entrance to Chilkat Lake. Other stocks in the general Lynn Canal area are monitored by aerial or foot surveys.

MANAGEMENT GOALS

Specific management goals for the 2006 Lynn Canal drift gillnet fishery are as follows (Table 2):

1. Obtain an escapement of between 50,000 and 90,000 (weir count units) sockeye salmon to Chilkoot Lake.
2. Obtain an escapement of between 80,000 and 200,000 (mark-recapture units) sockeye salmon to Chilkat Lake. The escapement will be monitored in season by the lower Chilkat River fish wheel project and the final escapement will be derived postseason from mark-recapture methods.
3. Obtain an escapement of between 1,750–3,500 three-ocean age and older Chinook salmon to the Chilkat River.

4. Obtain a peak foot escapement count between 4,000 and 9,200 coho salmon to Berners River.
5. Obtain a peak index stream count for Chilkat River drainage coho salmon that corresponds to 30,000–70,000 fish.
6. Provide for sufficient chum, coho, and pink salmon spawning escapements to the Chilkat, Chilkoot, and Berners rivers and other Lynn Canal systems, while harvesting those fish in excess of escapement needs.
7. Harvest all DIPAC hatchery-produced chum salmon available in the Boat Harbor Terminal Harvest Area while conserving wild stock summer chum salmon migrating to streams on the western shoreline of Lynn Canal.

2006 OUTLOOK

CHILKAT RIVER DRAINAGE SOCKEYE SALMON

The 2006 forecast return of Chilkat Lake sockeye salmon is approximately 175,000 fish (Table 3), the expected return is 90% of the 1976 to 2005 historical average of 210,000 fish (Table 4). The 2006 run size of Chilkat River mainstem sockeye salmon are expected to be above average during 2006. Based on less than favorable rearing conditions at Chilkat Lake but good parental brood year escapements, the total return of Chilkat Lake sockeye salmon should be just below average but stronger than the 2005 return.

Chilkat Lake escapement estimates of 131,000 and 132,000 during the 2000 and 2001 parent years were within the sustainable escapement goal range of 80,000–200,000 for the 2006 return (Table 5 and Figure 6). The dominant smolt years for the 2006 return (2003 and 2004), were estimated to be approximately 1.46 million fish for both years. These smolt abundance estimates are close the historical 1996–2005 average of 1.55 million (Table 6). The average weight and length of age-1.0 Chilkat Lake sockeye salmon smolt in 2003 and 2004 were below average (Table 7). Historically, 72% of the Chilkat Lake sockeye salmon escapements are 3-ocean age fish (31.9% are age-1.3 fish, 39.7% are age-2.3 fish and 0.3% are age-3.3 fish; Appendix C1).

Mark-recapture estimates of the Chilkat River mainstem sockeye salmon escapements in 2001, 2002, and 2003, (the dominant parent-years) were 21,900, 39,000, and 36,100 fish, respectively (Table 8). The estimates of abundance for the 2006 return were well below the historical 1994–2005 average of 34,700 fish for brood year 2001 but above average in years 2002 and 2003. The dominant age classes for this run includes age-0.2 (19.9%), age-0.3 (39.1%), and age-1.3 (29.8%) fish (Appendix C2). The proportion of age-0.2 fish of the 2005 escapement was above average indicating that the 2006 return of age-0.3 fish to the mainstem Chilkat River may be significantly larger than average. The Lower Chilkat River fish wheel project has been providing inseason stock assessment and postseason escapement estimates of Chilkat River mainstem sockeye salmon since 1994 (Bachman and McGregor 2001).

CHILKOOT SOCKEYE SALMON

Returns of Chilkoot Lake sockeye salmon in 2006 are expected to be near the long-term average of approximately 166,000 fish (Table 9). The Chilkoot Lake sockeye salmon weir count during the dominant parental brood year (2001) for the 2006 return was 76,300 fish, within the desired goal range (Table 10; Figure 6, top). The Chilkoot River weir is used to collect scale pattern

standards and monitor the escapement of this stock during the commercial fishing season (Kelley and Bachman 1999).

The 2002 fall hydroacoustic abundance estimate was above average, however, zooplankton abundance was below average during 2002 when the dominant brood year sockeye salmon juveniles would have been rearing in the lake (Table 11; Figure 7). Due to recent low zooplankton densities in Chilkoot Lake, ADF&G will implement management decisions in the commercial drift gillnet salmon fishery to achieve target escapement levels near the lower end of the escapement goal range for this stock.

The total return of 144,000 Chilkoot Lake sockeye salmon in 2001 was near the 1987–2005 average of 166,000 (Table 9). Age composition of the 2005 escapement was near average for most of the dominant age classes (Appendix C3). Given this information, the department is expecting an average return of Chilkoot Lake sockeye salmon. Management decisions will continue to be based on inseason data for the District 15 drift gillnet fishery.

BERNERS SOCKEYE SALMON

An average run of Berners Bay sockeye salmon is expected in 2006. Total escapement estimates are not available for Berners Bay sockeye salmon systems. Peak aerial escapements to Berners Bay streams were above average for all brood years. The average dominant age classes for Berners Bay streams are age-0.3 (15.4%), age-1.2 (13%), and age-1.3 (67.4%)(Appendix C4). The dominant age classes of the 2005 escapement were near the historical average indicating an average return in 2006. The 2002 and 2003 commercial harvests of Berners Bay and Chilkat River mainstem sockeye salmon were estimated at 10,500 and 12,600 fish respectively. These harvests were close to the historic 1976–2004 average harvest of 14,400 fish (Table 12).

SUMMER CHUM SALMON

Projections for the Boat Harbor Terminal Harvest Area in 2006 is for a return of approximately 192,000 fish. This forecast return is above the 2005 return and 1.4 times the 1991–2005 average (Table 14). The pre-season projection for the Amalga Harbor chum salmon return is approximately 1.33 million fish, 1.3 times the 1991–2005 average of 1.04 million fish (Table 14).

Based on parental-year escapement counts, the wild summer chum salmon return in 2006 should be average to above average in run strength but at a much lower scale than the hatchery summer chum salmon return.

The majority of the summer chum salmon harvest in lower Lynn Canal is comprised of hatchery fish from remote release sites at Boat Harbor and Amalga Harbor. Smaller numbers of wild chum salmon are produced from local area streams such as Sawmill Creek and other Berners Bay rivers on the eastern side of Lynn Canal. The Endicott, Beardslee, and St. James rivers on the western side of Lynn Canal are also important contributors to the wild summer chum harvest in the drift gillnet fishery.

Peak aerial escapement counts of summer chum salmon in Sawmill Creek in 2002, 2003, and 2004 were 720, 399, and 550 fish respectively. The peak aerial escapements are well below the 1995–2005 average for this index system for all brood years (Figure 8). Peak counts of chum salmon in western Lynn Canal streams for the 2006 return were generally above the previous 10-year average. All peak counts conducted during these brood years were near or exceeded the prior 10-year average (Figure 9).

FALL CHUM SALMON

The 2006 return of fall chum salmon is expected to be above average. The Chilkat River peak aerial survey counts were 3,200 and 61,800 fish for brood years 2001 and 2002. This is well below the peak aerial escapement average of 22,800 for the year 2001 but well above this average in 2002. Klehini River peak aerial counts were 1,500 and 1,600 fish respectively. The escapements during the dominant parental brood years were well above average.

The commercial harvest during the dominant parental brood years (2001 and 2002) was slightly below the 10-year average. Escapements of Chilkat River fall chum salmon since 1999 have improved. Management strategies designed to reduce harvests of these stocks have been successful. Fish wheel counts, mark-recapture estimates and aerial escapement surveys have indicated an increasing trend in escapement since 1999. This work has estimated escapements of 206,000 adult chum salmon in 2002, 166,000 in 2003, 310,000 in 2004 and 202,000 in 2005 (Bachman *In prep*).

COHO SALMON

The Lynn Canal coho salmon return is expected to be average during 2006. Coho salmon systems in the area include the Chilkat River, Berners River and Chilkoot River. Parent-year survey counts at the Chilkat River tributaries and Chilkoot River drainage were generally above the 10-year average. The 2002 mark-recapture estimate for Chilkat River drainage coho salmon was 209,300, the highest estimate on record. The 2002 and 2003 escapements to Berners Bay were well above the escapement goal range of 4,000 to 9,200 fish.

ADF&G Sport Fish Division has been conducting coho salmon smolt coded wire-tagging (CWT) studies on the Chilkat River to estimate smolt size, age structure, and production of coho salmon smolts since 1999 (Ericksen 2003). The 2005 trap catch per unit effort (CPUE) of coho smolt of 8.6 smolt/trap-day was near the 1999–2005 average of 7.7 fish/trap-day. The 2002 and 2003 Chilkat River fish wheel catches of 5,090 and 5,300 coho exceeded the 1994–2005 average for both years (Figure 10). The District 15 gillnet catches of 78,000 coho salmon in 2002 and 57,100 in 2003 (Table 1) were well above the previous 10-year average for both brood years. This data suggests an average coho salmon return for 2006.

CHINOOK SALMON

The 2006 preseason forecast for mature (\geq age 1.3) Chilkat River Chinook salmon is estimated to be near the upper range of the biological escapement goal range of 3,600 fish. Since the preseason forecast is projected to be above the escapement goal, the northern line in Chilkat Inlet will move northward to Glacier Point on the second week of the fishery and may be moved to the latitude of Cannery point by the third week of the season.

FISHERY OPENINGS

In 2006, ADF&G intends to manage the summer Lynn Canal drift gillnet fishery to obtain the mid-point of the escapement goal ranges for Chilkoot Lake and Chilkat Lake sockeye salmon. The department intends to manage the fishery to minimize harvest of wild stock summer chum salmon while harvesting expected large returns of hatchery chum salmon. The fall Lynn Canal drift gillnet fishery will be managed to conserve Klehini River (early-run) fall chum salmon while providing opportunity to harvest Chilkat River fall chum and coho salmon.

Section 15-A

Section 15-A will open for two days south of the latitude of Seduction Point beginning 12:01 p.m., Sunday June 18 (Statistical Week 25) with no mesh restriction. If the Chilkoot River weir count through June 15 is less than 2,500 sockeye salmon, the eastern side of Section 15-A will be closed. If the weir count is 2,500 sockeye salmon or greater on June 15, the eastern portion of 15-A may be opened. During the first three weeks of the season, Chilkat Inlet will be managed in accordance with the Chilkat River King Salmon Fishery Management Plan (Appendix A1). The first week of the season, Chilkat Inlet will be closed north of the latitude of Seduction Point. In week 27, Chilkat Inlet will be open south of the latitude of Glacier Point. In week 28, Chilkat Inlet will be open south of the latitude of Cannery Point. It is likely that the northern boundary line in Chilkat Inlet will remain at Cannery Point for the remainder of the summer season if escapements of Chilkat Lake sockeye salmon are projected to be within the desired goal range. If the Chilkoot Lake sockeye salmon return is strong, all of Section 15-A south of the latitude of Seduction point may be opened during the fourth week of the season for 2 or 3 days. Since ADF&G is forecasting an average return of sockeye salmon to the large sockeye producing systems, openings will be dictated by the results of various in season stock assessment programs operating on the Chilkat River and Chilkoot River drainages. Initial early returns to the Chilkat system are composed largely of Chilkat Lake returns. Chilkat mainstem returns overlap with Lake returns and peak in early to middle July followed by late run fish, which dominate August returns (Figure 11). Return timing is tied to freshwater age: mainstem sockeye salmon are predominantly age 0, Chilkat Lake early run fish are predominantly age 1, and Chilkat Lake late run fish are predominantly age 2.

Additional fishing opportunity in Chilkoot Inlet north of the latitude of Mud Bay Point for 2 or 3 days in weeks 31 through 37 may be possible if the Chilkoot Lake sockeye salmon return is as strong as expected. If the inseason information system indicates that the Chilkat Lake sockeye salmon return is not forecast to meet minimum escapement goals, limits in time and area of Section 15-A will be implemented until ADF&G can project sockeye escapement within desired goal ranges. Data from the Chilkat River fish wheel mark-recapture program will be used to judge run strength inseason and escapement levels postseason.

Fall fishery management in Section 15-A will occur from week 35 until the end of the season. As in recent years, the northern boundary line in Section 15-A will move southward in stages as the coho and fall chum stocks begin to migrate back to parental streams. Depending on effort levels, and coho and chum salmon run strength, fishing opportunity in Section 15-A will likely be managed similar to 2005.

Section 15-B

Based on the preseason forecast for coho salmon to Berners Bay, Section 15-B may be opened from week 38 to the end of the season south of the latitude of Cove Point (Figure 12) for 2 or 3 days each. Inseason information collected from coded wire tag recoveries and commercial harvest from various gear types will provide the data to manage fishing opportunity in Section 15-B.

Section 15-C

Section 15-C will open for two days beginning 12:01 p.m., Sunday, June 18 with no mesh restriction. If the Chilkoot River weir count is less than 2,500 sockeye salmon through June 15,

the eastern side of Section 15-C will be closed north of the latitude of Bridget Point. If such action is taken, the Boat Harbor area on the western side of Section 15-C would remain open.

Due to the expected average returns of Chilkat and Chilkoot Lake sockeye salmon, open fishing time in Section 15-C will be limited to 2 or 3 days (except for the Boat Harbor area). If inseason projections for the Chilkat and Chilkoot Lake sockeye salmon returns are below the escapement goal range, it is possible that time, area and gear restrictions will be enacted in Section 15-C during the summer season to boost escapement at desired levels.

If sockeye salmon escapement falls short of inseason escapement objectives, openings of the small area in eastern Section 15-C defined as: the waters of Section 15-C from the eastern shoreline of Lynn Canal at the latitude of Vanderbilt Reef Light to Vanderbilt Reef Light and east of a line from Vanderbilt Reef Light to Little Island Light, (Figure 12) may occur on the 3rd or 4th day during peak weeks (statistical weeks 27 through 31) of the hatchery chum salmon return. This strategy will be used to provide opportunity to harvest summer chum salmon while reducing the harvest of sockeye salmon in Section 15-C. The decision to use this strategy will be considered inseason based on Chilkat River fish wheel counts, Chilkoot Lake weir counts and results from site-specific sampling of the commercial fishery.

The Boat Harbor Terminal Harvest Area (THA) will be opened for extended periods beginning in week 28, (July 9). The Boat Harbor THA is shown in Figure 1 and is defined as: those waters within two nautical miles of the western shoreline of Lynn Canal south of the latitude of Danger Point at 58°41.73' N. latitude and north of a point 2.4 miles north of Point Whidbey at 58°37.05' N. latitude. The northern line of the Boat Harbor area will remain at the latitude of Danger Point through week 31 to protect wild Endicott River and other western Lynn Canal wild chum salmon stocks. This strategy has been in place since 2001. Escapements of wild chum salmon to the Endicott River have improved because of this action.

The area within the Boat Harbor area west of a line from the entrance to the Boat Harbor proper area will be opened continuously beginning the first week of the season. This strategy will be used to harvest expected large returns of hatchery chum salmon while fish are available with no risk to wild salmon.

Fall season management will begin in late August or early September in Section 15-C. Fall-season management will be based on coho and chum salmon overall run strength and fishing effort levels. Fishing effort will be directed at harvesting coho salmon in Section 15-C while limiting the harvest of early run fall chum salmon. Early run fall chum salmon are typically composed of fish returning to the Klehini River. Recent years escapement levels of chum salmon in the Klehini River have been low. Fishing time will be limited from 2 to 4 days each, beginning in statistical week 34.

Other Comments

To avoid gear conflicts, the District 15 drift gillnet fishery will not be open concurrently with the Juneau Golden North Salmon Derby. Consequently, during Statistical Week 33, the District 15 gillnet fishery will not open until Monday, August 7.

Stock Assessment Projects

The Haines Commercial Fisheries Division's salmon stock assessment projects include the Chilkoot River weir, Chilkat River fish wheels, Chilkat Lake weir and the marine fishery

performance project. Funding for the Chilkat River fish wheels and Chilkat Lake weir is through grants from the Southeast Sustainable Salmon Fund (SSSF). Funding for the Chilkoot River weir and marine fishery performance projects are supported through the general fund. SSSF funding to operate the Chilkat River fish wheel and Chilkat Lake weir projects are currently secured through 2006.

Information collection from the Chilkat River fish wheel, Chilkoot River weir and marine fishery performance projects form the inseason escapement and commercial catch information system used to make fishery decisions to manage the drift gillnet fishery in Lynn Canal.

Information from the lower Chilkat River fish wheel program is used to estimate the spawning escapements of Chilkat Lake and Chilkat River mainstem sockeye salmon through mark-recapture experiments. Sockeye salmon are marked at the fish wheels and ratios of marked to unmarked fish are determined from recaptured fish at the Chilkat Lake weir and on Chilkat River mainstem spawning areas. Since this program began in 1994, the average (1994–2005) escapement to Chilkat Lake and Chilkat River mainstem areas is estimated at 166,200 and 34,700, respectively, using mark-recapture methods.

A mark-recapture project utilizing the Chilkat River fish wheels began in 2002 to measure the abundance of fall chum salmon from June 8 through October 19. Over the past four years, this mark-recapture study has documented escapements of approximately 204,100 in 2002, 165,700 in 2003, 309,520 in 2004 and 202,000 in 2005. This information will be used to correlate fish wheel catch and aerial escapement surveys to estimates of abundance based on the 2002–2005 mark-recapture work.

ADF&G will conduct mark-recapture work on out-migrating Chilkat Lake sockeye salmon smolt during the spring of 2006 at the Chilkat Lake weir site. This information is critical to assess the rearing health of Chilkat Lake and to estimate the total emigration of Chilkat Lake sockeye salmon. This information forms the basis in forecasting future returns of Chilkat Lake sockeye salmon. ADF&G will operate the Chilkat Lake weir early in 2006 to sample and estimate the abundance of sockeye salmon smolt and estimate the returning adult Chilkat Lake sockeye salmon.

The Chilkoot Lake weir project has been in operation since 1976. The weir will operate beginning the first week in June to estimate the spawning escapement of this stock inseason.

As in previous years, the department's management crews as part of the marine fishery performance project will be on the fishing grounds during commercial fishing periods to sample sockeye and Chinook salmon and to monitor the fishery during each opening. The department requests that commercially caught sockeye and Chinook salmon be retained in separate fish holds or totes so department staff can collect scale and length data from targeted fish while monitoring the fishery. The sockeye salmon scale samples that are collected from the commercial gillnet fishery form the basis of our stock separation analysis. ADF&G vessels stand by on **channel 10 VHF** when on the fishing grounds.

Please report any commercial fisheries violations to the Bureau of Wildlife Enforcement at (907) 766-2533 (Haines), (907) 465-4000 (Juneau).

REFERENCES CITED

- Bachman R. L. *In prep.* Stock assessment studies of Chilkat River salmon stocks in 2003 and 2004. Alaska Department of Fish and Game, Division of Commercial Fisheries, Fishery Data Series Report, Anchorage.
- Bachman R. L. 2005. Stock assessment studies of Chilkat River salmon stocks in 2002. Alaska Department of Fish and Game, Division of Commercial Fisheries, Fishery Data Series Report 05-36, Anchorage.
- Bachman R. L. and A. J. McGregor. 2001. Stock assessment studies of Chilkat River salmon stocks in 1999. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 1J01-36, Juneau.
- Ericksen R. P. and S. A. Fleischman. *In prep.* Optimal production of coho salmon from the Chilkat River. Alaska Department of Fish and Game, Fishery Manuscript, Anchorage.
- Ericksen, R. P., and S. A. McPherson. 2004. Optimal production of Chinook salmon from the Chilkat River. Alaska Department of Fish and Game, Fishery Manuscript No. 04-01, Anchorage.
- Ericksen R. P. 2003. Smolt production and harvest of coho salmon from the Chilkat River, 2001–2002. Alaska Department of Fish and Game, Fishery Data Series, No. 03-28, Anchorage.
- Gieger, H. J., R. L. Bachman, S. C. Heintz, K. Jensen, T. A. Johnson, A. Piston, R. Riffe. 2005. Sockeye salmon stock status and escapement goals in Southeast Alaska [in] Der Hovanisian and H.J. Gieger, *editors*. Stock status and escapement goals for salmon in Southeast Alaska 2005. Alaska Department of Fish and Game, Special Publication No. 05-22, Anchorage.
- Kelley, M. S. and R. L. Bachman. 1999. Chilkoot River weir results. 1998. Alaska Department of Fish and Game, Commercial Fisheries Division, Regional Information Report 1J99-25, Juneau.
- McPherson, S. A. 1990. An inseason management system for sockeye salmon returns to Lynn Canal, Southeast Alaska. M. S. Thesis, University of Alaska Fairbanks.
- McPherson, S., D. Bernard, J. H. Clark, K. Pahlke, E. Jones, J. Der Hovanisian, J. Weller, and R. Ericksen. 2003. Stock status and escapement goals for Chinook salmon stocks in Southeast Alaska. Alaska Department of Fish and Game, Special Publication No. 03-01, Anchorage.
- Shaul, L., E. Jones and K. Crabtree. 2005. Coho salmon stock status and escapement goals in Southeast Alaska [in] Der Hovanisian and H.J. Gieger, *editors*. Stock status and escapement goals for salmon in Southeast Alaska 2005. Alaska Department of Fish and Game, Special Publication No. 05-22, Anchorage.

TABLES AND FIGURES

Table 1.—Historical catches of king, sockeye, coho, pink, and chum salmon in the District 15 (Lynn Canal) drift gillnet fishery, 1976–2005.

Year	King				Sockeye				Coho			
	15-A	15-B	15-C	Total	15-A	15-B	15-C	Total	15-A	15-B	15-C	Total
1976	2,044	0	24	2,068	122,735	0	3,887	126,622	49,167	0	22,817	71,984
1977	1,156	0	58	1,214	159,312	0	767	160,079	50,453	0	40,973	91,426
1978	457	0	79	536	105,491	0	2,989	108,480	26,084	0	27,081	53,165
1979	3,554	0	18	3,572	192,692	0	282	192,974	23,426	0	3,589	27,015
1980	434	0	6	440	53,096	0	891	53,987	26,120	0	2,778	28,898
1981	1,083	2	215	1,300	81,740	1,289	10,166	93,195	30,452	109	14,089	44,650
1982	5,878	1	66	5,945	268,290	160	5,432	273,882	47,719	78	24,573	72,370
1983	1,993	5	121	2,119	349,884	155	19,791	369,830	53,622	66	15,822	69,510
1984	5,822	13	264	6,099	320,277	3,759	10,546	334,582	43,637	58	24,520	68,215
1985	2,753	177	330	3,260	233,972	7,736	61,533	303,241	64,547	414	33,329	98,290
1986	2,141	41	590	2,772	248,264	1,100	40,541	289,905	48,046	4	34,071	82,121
1987	3,021	2	200	3,223	381,856	1,244	32,236	415,336	31,195	15	22,541	53,751
1988	1,136	27	94	1,257	327,330	17,469	7,000	351,799	50,984	410	30,142	81,536
1989	1,478	33	444	1,955	351,706	9,249	110,959	471,914	29,484	98	20,725	50,307
1990	364	16	290	670	248,878	3,612	104,928	357,418	36,260	48	26,764	63,072
1991	462	0	283	745	275,428	0	32,383	307,811	23,031	0	105,334	128,365
1992	225	0	385	610	230,229	0	55,806	286,035	30,021	0	78,732	108,753
1993	302	0	439	741	119,754	0	53,359	173,113	7,499	0	52,453	59,952
1994	253	4	723	980	111,061	80	60,588	171,729	55,925	13,805	71,034	140,764
1995	56	0	772	831	41,570	505	34,351	88,572	21,093	11,632	43,696	79,949
1996	106	0	491	642	65,031	0	41,354	149,961	16,525	0	29,885	52,658
1997	280	0	487	834	52,669	0	42,413	118,348	2,034	0	12,558	15,572
1998	375	0	304	679	114,467	0	20,470	134,937	7,003	0	19,115	26,118
1999	373	0	180	553	145,917	0	17,613	163,530	4,478	0	30,852	35,330
2000	140	0	157	297	76,732	0	32,648	109,380	7,652	0	27,984	35,636
2001	373	0	1,229	1,672	57,055	0	90,756	147,811	10,932	0	23,283	34,215
2002	64	0	518	582	41,677	6	40,326	82,014	23,823	12,574	41,544	77,941
2003	71	0	594	665	57,370	74	37,692	95,136	17,769	10,408	28,929	57,106
2004	365	0	440	805	73,944	130	77,171	151,245	13,323	5,822	32,815	51,960
2005	309	0	402	711	38,249	0	27,220	65,469	10,435	0	17,512	27,947
Ave.												
1996–05	246	0	480	744	72,311	21	42,766	121,783	11,397	2,880	26,448	41,448
1976–05	1,236	11	340	1,593	164,889	1,552	35,870	204,945	28,758	1,851	31,985	62,953

-continued-

Table 1–Page 2 of 2.

Year	Pink				Chum					
	15-A	15-B	15-C	Total	15-A	15-B	15-C	Total	Summer	Fall
1976	3,866	0	563	4,429	329,974	0	45,052	375,026	5,173	369,853
1977	130,644	0	216	130,860	152,923	0	48,711	201,634	5,577	196,057
1978	3,260	0	551	3,811	82,443	0	35,985	118,428	7,845	110,583
1979	28,752	0	11	28,763	225,713	0	17,119	242,832	7,006	235,826
1980	79,441	0	2,902	82,343	157,515	0	11,338	168,853	19,888	148,965
1981	112,471	4,107	20,692	137,270	90,619	508	26,248	117,375	13,215	104,160
1982	67,415	126	1,509	69,050	271,659	37	34,889	306,585	5,337	301,248
1983	134,319	452	22,775	157,546	311,510	1,116	28,519	341,145	19,303	321,842
1984	68,611	2,128	7,261	78,000	552,232	10,177	79,829	642,238	59,567	582,671
1985	169,644	6,079	63,357	239,080	582,649	12,377	103,784	698,810	77,806	621,004
1986	31,927	34	6,154	38,115	305,610	1,357	74,415	381,382	18,987	362,395
1987	124,066	430	41,255	165,751	295,663	548	96,727	392,938	26,698	366,240
1988	193,991	10,343	4,070	208,404	284,127	28,664	64,792	377,583	60,206	317,377
1989	61,365	291	48,798	110,454	90,735	3,508	29,388	123,631	28,813	94,818
1990	48,645	1,247	51,207	101,099	122,157	2,908	85,477	210,542	84,282	126,260
1991	3,815	0	1,657	5,472	100,121	0	110,068	210,189	100,627	109,562
1992	243,297	0	108,265	351,562	114,157	0	131,090	245,247	132,505	112,742
1993	680	0	10,656	11,336	62,190	0	244,376	306,566	229,284	77,282
1994	57,648	2	89,627	147,277	155,172	4,482	525,795	685,449	529,380	156,069
1995	883	0	14,641	5,799	62,206	1,332	494,792	568,368	493,279	75,089
1996	1,290	0	958	2,358	55,321	0	337,709	415,547	340,021	75,526
1997	13,601	0	36,864	32,962	28,410	0	425,122	461,614	431,699	29,915
1998	22,260	0	10,091	32,351	29,933	0	130,736	160,669	136,515	24,154
1999	36,989	0	25,748	62,737	46,947	0	303,947	350,894	290,325	60,569
2000	15,938	0	5,070	21,008	66,848	0	686,181	753,029	680,536	72,493
2001	26,709	0	41,009	67,718	66,024	0	377,501	443,525	358,987	84,538
2002	37,938	0	50,106	88,044	44,184	632	620,869	665,685	625,743	39,518
2003	33,936	0	19,685	53,621	35,487	2,114	356,287	393,888	348,820	45,672
2004	62,157	0	36,184	98,341	74,618	1,061	668,977	744,656	666,038	78,257
2005	89,755	0	120,078	209,833	63,048	0	263,847	326,895	240,055	86,840
Ave.										
1996–05	34,057	0	34,579	66,897	51,082	381	417,118	471,640	411,874	59,748
1976–05	63,510	841	28,065	91,513	162,007	2,361	215,319	381,041	201,451	179,584

Table 2.—Biological and sustainable escapement goals for Lynn Canal salmon stocks by species and location.

Species	Stock	Escapement Goal Type	Escapement Goal Range	Escapement Method
Sockeye ^a	Chilkoot Lake Total	Sustainable	50,000 to 90,000	Weir Count
Sockeye ^a	Chilkat Lake Total	Sustainable	80,000 to 200,000	Mark-Recapture Estimate
Coho ^b	Berners River	Biological	4,000 to 9,200	Peak Foot Count
Coho ^c	Chilkat River Combined	Biological	30,000 to 70,000	Sum of Peak Foot Index Counts
Chinook ^d	Chilkat River Combined	Biological	1,750 to 3,500	Mark-Recapture Estimate

^a Gieger et al. 2005

^b Shaul and Crabtree 2005

^c Ericksen and Feischman *In prep*

^d Ericksen and McPherson 2004

Table 3.—Yearly preseason smolt based forecast compared to the estimated total return of adult Chilkat Lake sockeye salmon, 1992, 1997–2006.

Year	Preseason Forecast	Estimated Total Return	Difference Forecast vs. TR	Percent Difference
1992	218,000	209,627	8,373	4%
1993 ^a	N/A	310,447	N/A	N/A
1994 ^a	N/A	275,752	N/A	N/A
1995 ^a	N/A	247,937	N/A	N/A
1996 ^a	N/A	359,232	N/A	N/A
1997	266,974	308,859	-41,885	-16%
1998	262,123	331,758	-69,635	-27%
1999	295,520	386,089	-90,569	-31%
2000	177,093	210,190	-33,097	-19%
2001	181,568	190,634	-9,066	-5%
2002	210,655	175,403	35,252	17%
2003	156,056	162,827	-6,771	-4%
2004	115,101	168,833	-53,732	-47%
2005	100,607	106,931	-6,324	-6%
2006	174,962	N/A	N/A	N/A
92, 97–2005 Avg.	198,370	225,115	-26,745	-13%

Note: Escapement for years 1994–2005 based on mark-recapture methods.

^a No forecast due to incomplete smolt data.

Table 4.—Annual total return of Chilkat Lake sockeye salmon by week, 1976 to 2005.

Stat	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
23														62		1
24	1		22	6										689	202	44
25	384	214	4,861	1,556	675	1,542	469		2,550	408	88			13,398	2,358	1,516
26	5,038	5,346	2,645	3,941	1,053	1,960	2,170	1,452	6,348	1,732	361	1,968	2,438	19,180	6,021	2,727
27	5,568	9,661	8,622	6,955	1,152	1,826	4,061	3,116	11,132	1,731	1,304	5,307	5,497	18,284	7,966	3,157
28	6,583	2,350	3,799	7,017	3,560	1,635	3,524	16,747	5,413	6,456	1,227	3,713	5,416	13,456	8,758	3,902
29	2,037	2,412	13,648	14,088	4,355	3,053	3,087	19,741	12,331	5,459	1,997	12,411	7,607	20,959	13,254	2,892
30	3,795	1,586	4,528	17,288	6,685	6,171	3,618	16,317	14,147	3,471	2,229	7,580	4,127	25,615	12,637	4,928
31	4,275	1,841	22,108	19,520	3,401	3,123	13,135	18,993	9,295	9,962	2,266	6,073	10,476	15,994	11,372	6,294
32	9,730	4,502	18,979	30,268	5,550	5,934	30,622	24,469	12,203	11,705	10,775	15,401	8,462	35,253	26,950	10,202
33	14,422	5,350	23,568	21,765	10,337	1,746	30,708	23,031	21,367	11,493	30,806	11,230	13,569	18,503	35,594	7,870
34	21,306	6,185	12,968	23,389	12,510	3,430	18,548	17,307	7,807	27,805	45,640	10,041	7,337	21,241	2,948	14,332
35	12,280	7,392	4,384	17,551	13,799	19,487	21,353	22,356	21,427	39,750	15,353	12,649	7,008	21,047	36,947	19,423
36	7,296	20,193	6,903	5,110	11,032	8,097	12,250	25,274	22,913	20,105	45,368	4,112	7,106	12,212	23,929	11,436
37	17,740	8,982	6,362	20,149	29,820	19,652	8,004	15,674	22,509	11,180	13,083	1,852	11,491	16,250	4,020	7,726
38–42	18,602	6,419	23,689	7,980	22,099	54,893	55,708	53,618	44,058	41,970	21,811	26,325	13,532	47,778	14,331	16,245
Yearly Total	129,057	82,433	157,086	196,583	126,028	132,549	207,257	258,095	213,500	193,227	192,308	118,662	104,066	299,921	207,287	112,695
Weekly Mean	8,604	5,888	10,472	13,106	9,002	9,468	14,804	19,853	15,250	13,802	13,736	9,128	8,005	18,745	13,819	7,043
Early Stock	37,411	27,912	79,212	100,639	26,431	25,244	60,686	100,835	73,419	40,924	20,247	52,453	44,023	162,890	89,518	35,663
Late Stock	91,646	54,521	77,874	95,944	99,597	107,305	146,571	157,260	140,081	152,303	172,061	66,209	60,043	137,031	117,769	77,032

-continued-

Table 4.—Page 2 of 2.

Stat															1976-05
Week	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	Average
23															32
24	10			57		476	592	258	53	105	683	144	87	157	211
25	53	2,358		4,516	987	5,003	4,308	1,127	3,861	2,046	6,595	1,875	1,237	904	2,496
26	3,452	2,886	6,261	7,021	5,954	6,569	14,634	10,292	18,395	8,353	8,177	2,920	3,810	2,776	5,529
27	6,280	6,120	10,224	10,474	11,981	15,157	29,804	20,235	21,008	10,802	10,290	3,940	9,299	3,836	8,826
28	5,310	9,787	15,900	14,557	34,411	8,338	31,533	46,640	18,335	14,305	10,457	3,244	16,586	5,469	10,948
29	12,216	9,417	16,880	11,359	30,905	8,469	36,090	35,884	16,039	23,472	12,363	4,519	16,485	5,863	12,643
30	10,681	10,541	14,476	20,856	31,177	15,775	31,506	39,289	15,403	26,331	14,267	5,321	14,643	6,020	13,034
31	7,825	9,317	17,780	19,782	34,786	23,640	32,112	28,573	16,273	19,647	16,056	6,598	13,620	5,896	13,668
32	15,267	13,613	21,151	22,516	58,568	31,728	41,282	36,690	21,336	27,652	23,266	17,393	20,781	7,876	20,671
33	25,623	26,504	46,225	20,818	39,784	17,349	36,814	33,663	18,377	16,358	16,560	26,531	20,962	6,468	21,113
34	15,416	33,161	29,319	44,587	37,510	20,648	31,761	35,772	18,609	11,091	14,304	24,585	18,984	7,395	19,865
35	36,915	38,108	35,280	28,202	31,720	26,064	31,529	35,087	11,137	10,136	13,927	25,053	15,236	7,373	21,266
36	22,689	29,950	32,105	17,769	20,683	22,670	19,649	19,154	12,845	10,546	15,655	23,323	15,124	11,933	17,248
37	8,234	5,901	26,137	23,374	18,089	29,896	27,720	15,695	6,044	3,396	7,806	11,063	3,356	11,904	13,770
38-42	39,656	112,784	4,012	2,048	2,676	77,078	46,224	42,015	12,474	6,395	4,996	10,510	7,968	23,066	28,699
Yearly Total	209,627	310,447	275,752	247,937	359,232	308,859	430,106	398,822	210,190	190,635	175,403	167,019	178,177	106,937	209,997
Weekly Mean	13,975	22,175	21,212	16,529	25,659	20,591	27,704	26,692	14,013	12,709	11,694	11,135	11,878	7,129	14,461
Early Stock	61,094	64,039	102,673	111,138	208,770	115,155	221,860	218,988	130,703	132,713	102,154	45,955	96,548	38,797	87,603
Late Stock	148,533	246,408	173,079	136,798	150,462	193,705	193,698	181,387	79,487	57,921	73,249	121,064	81,629	68,140	121,960

Table 5.—Annual escapements of Chilkat Lake sockeye salmon by week, 1976–2005.

Stat	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
23														62		1
24	1		22	6										689	202	44
25	0	214	476	44	72	3			302					5,802	639	305
26	433	305	1,302	698	887	0	31	368	1,441	7	4	88	59	10,690	3,615	901
27	944	572	8,622	6,930	1,152	5	532	1,248	5,436	98	2	1,777	2,015	7,845	1,660	1,600
28	2,437	773	2,751	2,081	3,560	141	605	11,144	623	1,317	602	2,197	496	2,295	4,353	1,971
29	1,140	207	11,816	8,576	4,355	549	461	15,284	3,280	1,141	139	5,601	9	8,126	9,566	503
30	2,055	542	1,310	4,068	4,575	1,071	2,515	8,935	6,011	334	20	2,542	722	15,810	2,380	2,812
31	2,816	711	1,814	1,413	2,100	1,002	1,743	10,750	929	812	24	1	1,969	3,161	1,449	2,234
32	310	1,184	40	2,056	2,100	266	3,496	6,865	141	2,029	1	123	1,965	4,340	1,925	3,724
33	2,740	725	1,078	5,895	2,100	729	509	4,254	2,971	157	3	1,776	200	11	380	1,821
34	9,810	968	1,634	7,288	5,666	1,450	4,073	5,589	1,417	1,555	138	1,875	566	3,207	2,948	4,295
35	4,283	1,269	1,246	11,212	6,910	767	5,151	1,433	14,899	4,434	736	6,193	280	7,582	7,167	10,732
36	6,799	18,711	5,670	3,639	10,351	4,967	1,575	5,475	18,015	3,271	1,006	1,618	469	8,379	9,647	5,380
37	17,483	8,664	6,106	19,464	29,613	18,652	6,091	10,526	18,512	3,372	5,364	27	7,973	15,019	259	2,260
38	9,655	144	7,747	12	10,739	1,113	20,378	21,097	21,106	12,639	6,943	259	2,254	34,155	664	3,264
39	5,584	5,821	9,469	2,353	7,015	6,134	25,516	9,455	17,510	17,688	3,796	18,033	2,747	2,713	4,465	1,873
40	0	234	6,334	1,413	3,374	32,516	7,467	9,398	2,252	5,258	3,762	6,165	4,551	2,936	3,552	1,091
41	3,001	0	91	2,125	778	10,222	78	7,305	424	2,009	831	0	655	3,053	4,456	1,427
42	238			1,316		4,502		5,081		1,603	576	318	663	4,600	904	6,651
Yearly Total	69,729	41,044	67,528	80,589	95,347	84,089	80,221	134,207	115,269	57,724	23,947	48,593	27,593	140,475	60,231	52,889
Weekly Mean	3,486	2,160	3,554	4,029	5,609	4,672	5,014	7,895	6,067	3,396	1,330	2,700	1,533	7,024	3,170	2,644
Early Stock	17,582	9,437	17,924	30,433	10,253	10,617	9,640	47,885	28,193	7,449	2,536	13,345	7,512	54,090	25,792	15,916
Late Stock	52,147	31,607	49,604	50,156	85,094	73,472	70,581	86,322	87,076	50,275	21,411	35,248	20,081	86,385	34,439	36,973

-continued-

Table 5.—Page 2 of 2.

Stat																1976–05
Week	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	Average
23	1															21
24	44	10			57		476	395	270	53	105	683	140	81	157	191
25	305	53	75		2,232		1,857	2,562	1,140	3,861	392	4,695	1,398	1,150	904	1,238
26	901	1,016	1,745	1,510	5,323	2,720	3,618	6,382	5,737	14,933	4,580	6,169	2,148	2,441	2,281	2,656
27	1,600	1,653	3,557	3,456	8,471	11,051	11,759	12,307	12,659	13,238	5,014	6,699	2,827	5,757	2,312	4,606
28	1,971	1,762	4,240	8,223	9,674	32,814	5,951	10,495	26,856	10,034	6,595	7,185	1,883	8,046	3,600	5,699
29	503	6,529	3,552	5,125	9,387	28,393	5,713	12,343	16,442	9,594	12,139	6,745	3,027	7,909	4,401	6,534
30	2,812	5,034	7,615	8,025	18,775	28,308	13,187	9,500	20,819	8,399	19,314	9,037	2,498	7,426	4,795	7,137
31	2,234	2,263	5,336	8,184	17,172	26,778	16,044	10,900	14,853	7,176	12,945	11,728	2,968	7,984	4,713	5,942
32	3,724	3,579	6,490	9,375	17,973	42,335	22,138	15,897	17,906	8,886	20,775	15,074	10,872	15,718	5,949	7,976
33	1,821	1,197	14,537	34,085	15,054	22,358	11,283	17,350	21,197	9,347	11,512	14,182	16,530	14,216	4,966	7,580
34	4,295	5,768	6,643	17,559	25,643	17,767	9,617	16,221	20,962	11,167	10,196	10,325	16,080	14,305	4,922	7,869
35	10,732	10,357	23,593	16,367	21,007	21,848	14,521	19,738	20,035	7,145	9,084	10,109	16,637	10,692	5,059	9,717
36	5,380	13,172	19,677	19,346	13,394	13,942	18,044	12,723	9,563	9,647	9,641	13,339	16,996	12,648	9,886	9,754
37	2,260	6,014	1,251	18,274	20,377	14,112	27,518	19,149	10,180	5,595	3,139	7,219	8,805	2,537	9,067	10,480
38	3,264	8,779	61,222	4,012		425	42,800	12,857	13,788	6,492	2,813	2,379	5,654	2,455	15,324	11,148
39	1,873	22,150	32,323				9,474	18,121	10,382	3,009	2,519	1,354	4,156	2,095	5,702	9,048
40	1,091	6,171	297				21,328	10,598	10,685	1,742	924	902		2,235		5,626
41	1,427	1,891	2,947				3,475	3,163	2,899	1,003		287		1,219		2,191
42	6,651	342	14,630					413						366		3,053
Yearly Total	52,889	97,740	209,730	153,540	184,541	262,852	238,803	211,114	236,374	131,322	131,687	128,111	112,619	119,280	84,039	114,004
Weekly Mean	2,644	5,144	10,487	10,236	13,182	17,523	11,940	11,111	13,132	7,296	7,746	7,468	7,039	6,923	5,252	6,497
Early Stock	15,916	23,096	47,147	43,897	89,065	172,400	80,744	80,781	116,682	76,175	81,859	68,015	27,761	56,512	29,112	42,509
Late Stock	36,973	74,644	162,583	109,643	95,476	90,451	158,059	130,333	119,692	55,147	49,828	60,096	84,858	62,768	54,927	71,495

Note: From 1994 through 2005, estimates of escapement are determined from mark-recapture methods; weekly escapement numbers are derived from fish wheel catch per unit effort (CPUE) and stock composition data.

Table 6.—Chilkat Lake sockeye salmon smolt age, weight, and length compositions (wild and enhanced components), 1989–1990 and 1994–2005.

Year	Total Outmigration ^a	Fry Stocked	Total Wild	Total Enhanced	% Enhanced	Enhanced Survival %	Wild age-1.0	Enhanced age-1.0	Wild age-2.0	Enhanced age-2.0	Wild age-3.0	Enhanced age-3.0
1989	2,000,000		2,000,000				1,520,000		480,000			
1990	2,600,000		2,600,000				702,000		1,898,000			
1994	2,367,891	4,400,000	2,367,891				1,207,624		1,160,267			
1995	1,897,413	2,393,558	1,210,977	686,436	36.0%	23.1%	403,217	686,436	801,223	n/a	6,537	
1996	2,869,160	2,691,311	2,269,741	599,419	21.0%	27.7%	939,393	269,365	1,325,183	330,054	5,165	
1997	1,515,859	2,806,858	1,039,634	476,225	31.0%	4.9%	113,201	98,786	918,711	377,439	7,722	
1998	1,386,118		1,115,700	270,418	19.5%	23.8%	666,224	220,892	340,569	33,683	108,907	15,843
1999	1,809,273		1,362,342	446,931	24.7%		620,377		716,718	446,931	25,247	
2000	1,629,883		1,629,883				115,214		1,509,020		5,649	
2001	1,398,802	2,698,874	1,398,802				657,269		694,397		47,136	
2002	434,411		432,608	1,803	0.4%		114,619	1,803	316,686		1,303	
2003	1,458,025		1,401,462	56,563	3.9%		840,998		549,390	56,563	11,075	
2004	1,457,990		1,457,990				831,210		624,685		2,096	
2005	1,585,693		1,585,693				1,379,553		206,140		n/a	
96–05 Ave. ^b	1,554,521	2,732,348	1,369,386	308,560	16.7%	18.8%	627,806	147,712	720,150	248,934	23,811	15,843

^a Total sockeye salmon smolt out migration estimated from mark-recapture techniques.

^b The 1995–2005 average smolt out migration used for the 2005 total out migration data point as the smolt migration was two weeks earlier than average.

Table 7.—Percent age, average length and average weight composition of Chilkat Lake sockeye salmon smolt, 1989–1990 and 1994–2005.

Year	Age %			AVG Length (mm)			AVG Weight (g)		
	age-1.0	age-2.0	age-3.0	age-1.0	age-2.0	age-3.0	age-1.0	age-2.0	age-3.0
1989	76.0%	24.0%		100.2	121.0		8.9	14.6	
1990	27.0%	73.0%		103.9	118.9		10.0	14.8	
1994	51.0%	49.0%		102.3	119.5		9.9	14.8	
1995	62.0%	37.0%	4.0%	92.5	115.4	147.4	7.1	13.2	27.2
1996	42.0%	58.0%	2.0%	86.3	107.2	185.0	5.7	10.3	56.0
1997	13.0%	86.0%	1.0%	95.2	101.2	154.5	7.0	8.8	34.4
1998	64.0%	27.0%	9.0%	92.7	109.4	138.3	7.3	11.2	22.7
1999	34.0%	64.0%	2.0%	88.1	107.6	155.8	5.3	9.5	37.7
2000	7.1%	92.6%	0.3%	93.8	104.8	120.4	7.1	9.4	14.3
2001	47.0%	49.6%	3.4%	92.5	113.4	131.5	6.8	11.8	19.0
2002	26.8%	72.9%	0.2%	85.5	92.7	175.0	5.2	6.3	38.7
2003	75.3%	24.1%	0.6%	88.9	111.4	136.9	5.9	11.4	21.1
2004	57.0%	42.8%	0.1%	87.2	93.8	115.0	5.6	6.8	12.5
2005	87.0%	13.0%	0.0%	93.0	116.0	n/a	6.9	14.1	n/a
89–90, 94–05 Ave.	47.8%	50.9%	2.1%	93.0	109.4	146.0	7.0	11.2	28.4

Table 8.—Weekly and annual escapement of Chilkat River mainstem sockeye salmon, 1994–2005.

Statistical													1994–05
Week	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	Average
23													
24		27		69	65		53	51	98	47	81		61
25		1,410		270	1,153	39	309	55	1,745	992	288	703	696
26	137	2,867	585	162	3,820	431	2,222	1,294	4,917	4,051	1,662	2,777	2,077
27	1,061	3,700	4,428	1,189	2,842	1,565	5,817	2,254	7,001	7,555	4,279	3,257	3,746
28	3,427	3,529	12,508	1,059	2,893	5,571	8,440	2,261	6,159	5,474	6,541	3,694	5,130
29	1,434	3,116	10,239	1,433	3,312	2,671	13,472	3,145	5,068	4,330	7,002	4,736	4,996
30	2,242	4,283	11,416	3,277	3,335	5,001	7,805	6,645	3,966	2,684	3,789	6,813	5,105
31	2,720	3,140	6,615	2,845	4,271	2,607	8,025	2,627	4,884	2,597	5,169	4,816	4,193
32	3,170	1,588	5,207	2,222	1,252	2,891	4,944	2,330	2,136	3,608	6,670	7,188	3,601
33	8,431	1,229	1,036	613	1,201	1,724	2,318	964	1,200	1,958	3,905	4,681	2,438
34	1,882	449	661	371	243	1,083	657	209	996	1,505	2,518	4,803	1,281
35	886	740	398	430	481	257	139	34	432	1,071	1,890	3,098	821
36	691		217	140		381	65	29	484	249	899	2,702	586
37	105		59	377	90			26			104	854	231
38				180		133					52	519	221
39												180	180
Total	26,186	26,078	53,369	14,637	24,958	24,354	54,266	21,924	39,086	36,121	44,849	50,824	34,721
Avg	2,182	2,173	4,447	976	1,920	1,873	4,174	1,566	3,007	2,779	2,990	3,388	2,623

Note: Estimates based on mark-recapture methods. Weekly estimates are calculated from stock proportions of sockeye salmon captured in the lower Chilkat River fish wheels.

Table 9.—Annual total return of Chilkoot Lake sockeye salmon by week, 1978–2005.

Statistical														
Week	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
23	844	3	0	0	0	0	333	8	25	11	0	571	328	1
24	1,957	8,738	0	25	252	467	3,349	6	101	176	95	4,266	2,060	471
25	3,796	4,802	1,312	3,394	14,437	2,764	13,273	630	414	198	1,082	26,973	5,062	8,300
26	1,007	2,188	1,479	4,255	13,272	10,175	14,204	6,975	647	21,421	6,097	15,106	14,736	7,186
27	6,677	2,832	1,824	2,309	6,972	6,636	12,092	3,372	2,992	23,211	28,807	13,475	9,912	5,030
28	2,404	12,032	2,241	3,346	7,306	7,186	18,878	6,926	4,685	8,025	20,534	28,206	9,938	9,064
29	4,984	1,097	5,894	16,810	10,680	7,929	28,068	2,856	4,025	51,328	29,550	45,814	13,233	13,531
30	9,173	8,064	10,184	24,110	28,921	41,318	61,590	8,196	7,809	54,190	25,275	37,381	46,320	30,131
31	6,348	29,879	10,225	20,685	56,819	61,978	56,888	37,270	13,506	51,040	54,337	65,123	31,621	53,137
32	6,128	56,146	27,834	12,128	41,839	45,339	46,017	33,668	29,464	95,943	46,488	48,270	36,726	57,302
33	3,449	21,877	28,288	17,209	29,943	44,734	26,207	32,265	33,637	47,338	49,678	35,796	32,794	44,373
34	2,207	10,019	14,261	11,611	21,130	25,253	27,087	55,628	49,703	49,361	25,032	10,998	13,553	41,469
35	359	5,801	11,971	5,567	10,965	31,197	13,338	17,265	27,309	19,521	28,384	8,312	18,492	24,411
36	136	1,379	1,205	4,751	2,272	26,034	7,454	12,367	15,178	6,455	13,495	4,337	12,215	11,310
37	123	634	518	915	1,865	7,794	3,258	2,817	7,277	1,466	4,577	1,245	4,568	6,968
38–42	126	319	122	49	892	3,008	173	1,102	1,682	496	1,811	890	630	1,995
Total	49,718	165,810	117,358	127,164	247,565	321,812	332,209	221,351	198,454	430,180	335,242	346,763	252,188	314,679

-continued-

Table 9.—Page 2 of 2.

Week	1978–05															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	Mean
23	1	31	65	309	185	0	873	0	1	0	89	102	15	41		142
24	471	4,744	249	2,687	295	129	2,317	117	59	174	265	2,005	342	233	417	1,286
25	8,300	8,775	2,592	1,117	1,747	1,862	13,611	327	143	413	3,928	3,427	709	1,640	1,270	4,571
26	7,186	9,426	13,123	8,630	1,507	5,389	8,785	824	859	2,637	9,025	4,118	1,629	996	3,583	6,760
27	5,030	21,317	11,730	8,852	1,332	3,574	5,899	969	2,181	2,800	9,965	4,105	4,090	3,285	2,984	7,472
28	9,064	10,118	12,017	7,004	2,164	5,987	4,825	909	1,270	4,696	8,109	7,457	5,789	8,910	4,635	8,024
29	13,531	15,498	9,274	3,760	1,069	3,060	4,762	1,241	1,326	6,157	20,119	9,984	7,982	13,370	5,051	12,088
30	30,131	16,556	8,745	3,351	1,754	12,819	7,866	1,864	2,184	7,546	21,434	13,098	12,623	19,284	8,581	18,942
31	53,137	22,984	10,374	7,558	1,944	12,518	8,212	2,067	3,048	8,566	31,855	11,090	28,349	20,488	5,699	25,843
32	57,302	25,652	9,309	3,951	561	9,726	3,775	2,091	2,633	7,696	24,341	9,722	24,257	23,907	7,749	26,381
33	44,373	30,146	7,798	4,431	795	5,213	2,786	1,469	1,329	7,272	8,378	4,048	11,720	21,505	7,261	20,062
34	41,469	15,382	7,946	4,518	633	3,180	4,405	1,293	1,378	3,033	2,523	5,341	2,888	6,466	11,869	15,292
35	24,411	16,460	4,436	3,640	502	3,363	2,997	818	3,058	3,201	2,204	3,448	4,086	9,716	10,019	10,387
36	11,310	7,161	3,030	2,182	562	1,226	1,588	280	2,517	3,297	1,267	3,850	1,630	8,071	7,707	5,820
37	6,968	2,703	2,010	721	76	1,056	468	237	1,225	602	271	839	657	4,000	2,847	2,205
38–42	1,995	837	553	119	29	499	0	34	331	139	12	4	13	224	826	604
Total	314,679	207,790	103,251	62,830	15,155	69,600	73,167	14,541	23,542	58,229	143,785	82,636	106,778	142,133	80,498	165,872

Table 10.—Annual weir counts of Chilkoot Lake sockeye salmon by week, 1976–2005.

Stat																
Week	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
23	124	14	844	3					333	8	25	11		571	328	1
24	623	9,572	1,957	8,738		25	252	467	3,349	6	101	176	95	4,266	2,060	471
25	241	35,751	1,368	2,730	391	1,108	12,220	2,764	11,100	104	163	198	1,082	21,300	2,778	5,599
26	3,579	11,150	274	469	1,157	2,177	9,440	8,860	7,444	4,681	224	16,583	1,506	2,466	12,190	3,083
27	735	3,361	6,677	407	1,824	559	2,623	4,062	4,406	783	857	6,879	22,846	1,009	1,893	2,097
28	397	6,970	1,311	309	2,241	606	1,981	3,304	9,993	463	3,650	3,365	5,872	913	1,980	2,528
29	1,752	1,844	2,526	95	5,894	7,346	5,095	4,090	6,738	810	2,328	7,000	4,389	2,122	0	5,436
30	4,091	1,854	7,650	2,871	9,239	15,951	17,574	21,548	11,917	3,601	5,467	8,134	2,554	2,942	4,989	21,990
31	28,061	9,016	3,465	22,765	8,294	9,006	20,806	12,747	9,610	19,778	11,438	8,998	5,416	3,614	1,853	17,870
32	13,587	9,561	5,157	31,000	20,860	9,963	13,358	4,507	8,020	9,832	21,563	9,944	5,824	4,313	1,995	7,317
33	11,827	6,059	2,316	16,091	21,333	15,631	8,287	3,614	5,522	12,501	12,276	5,899	5,683	2,157	4,255	8,229
34	5,205	1,019	1,469	5,140	12,968	10,659	4,938	2,720	11,185	7,013	11,839	16,978	10,851	2,793	13,553	4,115
35	346	372	155	3,880	10,669	5,028	2,655	3,016	3,435	4,432	6,348	6,018	6,650	3,067	13,734	5,077
36	49	403	56	933	1,077	4,519	1,518	4,366	4,474	2,817	5,416	3,918	4,544	1,840	9,147	3,988
37	118	103	106	427	479	794	1,404	2,604	2,891	1,546	5,071	738	2,646	876	2,128	1,879
38	410	2	83	8	45		822	1,070		480	762	217	759	232	365	416
39	142		12	70	36			502		145	409	112	381	216	5	294
40–42	10		28	10	5			102		26	87	17	176	203	71	248
Yearly Total	71,297	97,051	35,454	95,946	96,512	83,372	102,973	80,343	100,417	69,026	88,024	95,185	81,274	54,900	73,324	90,638
Early Stock Esc.	6,737	69,268	10,349	13,026	14,196	8,144	29,127	21,545	37,489	9,424	17,210	29,141	30,765	29,561	21,229	16,497
Late Stock Esc.	64,560	27,783	25,105	82,920	82,316	75,228	73,846	58,798	62,928	59,602	70,814	66,044	50,509	25,339	54,870	74,141

-continued-

Table 10.—Page 2 of 2.

Statistical	1976–05														
	Week	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
23	31	65	309	185		873		1		89	102	15	41		189
24	4,744	249	2687	295	129	2317	117	59	174	265	2,005	342	233	417	1,635
25	8,775	2,592	1,117	243	459	6,677	327	143	413	2,811	2,451	448	1,640	1,270	4,379
26	2,310	5,431	4,752	342	1,418	3,433	664	521	2,494	4,171	3,195	1,165		3,098	4,114
27	8,450	2,306	4,170	317	1,956	1,407	857	1,980	2,208	3,125	1,869	2,805	1,178	1,886	3,229
28	975	5,883	4,241	298	4,393	3,143	676	884	2,558	3,083	4,138	4,074	3,288	2,963	2,880
29	1,222	3,488	1,141	325	2,482	2,440	791	668	3,385	7,953	6,193	7,207	5,343	4,013	3,452
30	2,902	5,021	2,123	1,517	12,040	4,805	1,534	1,734	5,154	11,168	10,433	11,437	10,724	6,778	7,688
31	9,488	5,864	5,158	1,731	9,163	3,919	1,687	2,706	4,756	21,480	7,599	21,041	12,655	3,588	10,344
32	7,173	6,807	1,342	417	6,743	3,524	1,924	1,864	6,359	11,231	4,775	14,103	8,750	3,382	8,683
33	10,572	4,298	2,140	545	3,867	2,606	1,352	1,041	6,344	5,094	2,994	5,677	9,457	2,710	6,816
34	2,530	4,857	3,220	237	2,655	4,246	1,217	1,108	2,699	2,320	4,764	1,251	3,583	4,755	5,418
35	3,531	2,222	2,736	270	2,919	2,880	678	3,058	3,067	2,064	3,322	3,564	7,307	7,272	3,879
36	2,549	899	1,656	472	1,081	1,540	261	2,262	3,246	1,182	3,716	902	7,333	6,336	2,626
37	1,200	1,427	624	15	969	444	216	990	559	247	805	428	3,908	2,259	1,229
38	346	418			465		34	265	139			0	156	451	341
39	273														200
40–42															82
Yearly Total	67,071	51,827	37,416	7,209	50,739	44,254	12,335	19,284	43,555	76,283	58,361	74,459	75,596	51,178	66,177
Early Stock Esc.	25,285	16,526	17,276	1,680	8,355	17,850	2,641	3,588	7,847	13,544	16,733	8,834	6,339	9,634	17,328
Late Stock Esc.	41,786	35,301	20,140	5,529	42,384	26,404	9,694	15,696	35,708	62,739	41,628	65,610	69,216	41,544	48,939

Table 11—Selected data for Chilkoot Lake, annual autumn hydroacoustic pre-smolt estimates, total adult return, and average annual zooplankton densities for years 1987–2005.

Year	Survey Date	Estimated Juveniles	Total Return	Average zooplankton density (no./m²)	Average zoo biomass (mg/m²)
1987	30-Oct	1,344,951	430,180	172,295	207
1988	2-Oct	3,066,118	335,242	131,446	147.5
1989	16-Oct	874,794	346,763	46,872	135.5
1990	25-Oct	607,892	252,188	53,987	145.5
1991	22-Oct	475,404	314,679	9,751	25
1992	N/A	N/A	207,790	N/A	N/A
1993	N/A	N/A	103,251	N/A	N/A
1994	N/A	N/A	62,830	N/A	N/A
1995	6-Nov	260,797	15,155	26,579	84.7
1996	24-Oct	418,152	69,600	44,081	143.75
1997	22-Oct	755,060	73,167	15,063	46
1998	6-Oct	1,446,736	14,552	46,678	91.5
1999	14-Oct	351,096	23,542	14,329	46.25
2000	13-Oct	1,190,717	58,229	62,156	247
2001	17-Oct	696,000	143,785	88,791	275
2002	10-Oct	1,196,701	82,636	46,434	194
2003	10-Oct	1,384,754	106,778	46,788	155
2004	5-Nov	996,046	142,133	126,233	221
2005	9-Nov	247,243	80,498	3,265	7.7
Average		957,029	150,684	68,952	136

Table 12.—Annual harvests of Chilkat River mainstem and Berners Bay rivers, and other non-Chilkat or Chilkoot Lake, sockeye salmon by week, 1976– 2005.

Stat																
Week	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
25	60		548	504	381	143	44		355	134	16			3,214	1,823	2,213
26	694	2,653	1,759	1,328	56	101	210	49	514	1,688	599	734	968	3,381	1,783	6,782
27	963	1,330	207	0	725	145	145	255	491	5,173	1,233	6,958	6,611	2,440	6,998	4,097
28	1,194	332	386	494	158	150	155	294	383	6,691	4,365	983	4,889	1,742	2,221	2,470
29	375	848	316	501	73	181	175	105	309	273	738	872	5,100	2,030	1,054	3,451
30	735	116	577	1,414	0	116	172	268	561	522	897	263	1,057	1,725	4,601	1,012
31	204	0	486	1,942	76	154	549	1,204	706	746	597	330	1,316	2,922	4,669	1,729
32	227	0	0	0	75	67	128	740	536	448	903	350	442	1,956	4,251	1,138
33	151	0	269	165	8	0	329	663	244	377	948	111	348	366	3,088	224
34	132	98	74	492	3	14	0	256	73	68	825	121	101	494	0	151
35	76	0	29	195	3	0	0	78	130	48	206	22	100	233	297	635
36	8	0	6	35	0	0	0	42	48	0	87	0	122	98	216	0
37	0	0	1	14	0	0	0	1	0	10	0	7	23	19	40	38
38–42	23	0	0	32	0	0	1	0	0	0	0	0	33	5	3	24
Yearly Total	4,842	5,377	4,658	7,116	1,558	1,071	1,908	3,955	4,350	16,178	11,414	10,751	21,110	20,625	31,044	23,964
Weekly Mean	346	384	333	508	111	77	136	283	311	1,156	815	768	1,508	1,473	2,217	1,712

-continued-

Table 12.—Page 2 of 2.

Stat Week	1976–05														
	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	Mean
25				1,282	1,828	1,466				1,451	385	332			899
26	4,926	2,321	1,178	1,165	3,309	1,441	1,309	1,818	204	4,376	388	1,197	2,051	273	1,642
27	8,241	4,258	2,418	976	1,245	2,070	820	535	2,007	6,071	1,502	2,782	5,465	896	2,569
28	5,650	3,296	2,135	1,696	1,743	1,046	1,050	937	14,631	2,289	3,249	2,032	9,130	1,516	2,577
29	4,275	3,012	2,619	744	2,311	1,133	4,122	2,444	4,572	4,000	2,193	1,597	6,280	1,321	1,901
30	3,327	2,757	1,323	799	2,660	1,447	1,509	1,124	3,016	1,083	902	839	2,926	1,683	1,314
31	2,488	1,738	2,400	457	5,535	1,495	1,520	1,093	1,594	1,331	1,123	1,804	2,096	1,603	1,464
32	2,356	879	2,236	385	5,695	769	921	949	581	537	317	663	1,605	2,752	1,064
33	1,422	433	2,291	250	2,916	168	293	417	209	198	243	930	2,467	935	682
34	280	246	1,623	396	1,051	278	102	108	61	0	124	157	861	1,648	328
35	280	33	723	100	333	210	31	59	29	24	8	42	415	328	156
36	184	12	263	90	145	95	6	85	0	0	11	167	234	135	70
37	0	0	32	61	87	24	21	0	0	0	3	52	13	135	19
38-42	0	0	11	29	34	0	0	5	0	0	0	5	28	15	8
Yearly Total	33,429	18,985	19,252	8,430	28,893	11,642	11,704	9,575	26,903	21,361	10,446	12,599	33,571	13,240	14,332
Weekly Mean	2,388	1,460	1,481	602	2,064	832	900	737	2,069	1,526	746	900	2,582	1,018	1,048

Table 13.—Summary of releases of DIPAC chum salmon from Boat Harbor and Amalga Harbor, 1988–2005.

Brood Year	Release Year	Boat Harbor Releases	Boat Harbor Release Size (g)	Amalga Harbor Releases	Amalga Harbor Release Size (g)	Total Releases
1987	1988	5,170,000	N/A			5,170,000
1988	1989	8,508,356	0.77			8,508,356
1989	1990	8,300,782	1.31			8,300,782
1990	1991	9,337,000	0.88	34,744,923	0.87	44,081,923
1991	1992	6,709,659	0.62	35,918,054	1.08	42,627,713
1992	1993	9,545,177	0.75	36,147,451	1.23	45,692,628
1993	1994	6,464,450	0.86	34,817,531	1.38	41,281,981
1994	1995	8,931,491	1.06	34,472,077	1.49	43,403,568
1995	1996	8,536,780	0.7	34,979,646	1.22	43,516,426
1996	1997	7,759,020	1.4	34,535,728	1.33	42,294,748
1997	1998	7,217,000	1.45	49,155,073	1.52	56,372,073
1998	1999	9,262,694	1.32	7,655,324	3.44 (L/L ^a)	60,045,708
				43,127,690	1.53 (Reg ^b)	
1999	2000	9,010,000	1.61	8,722,507	4.04 (L/L)	62,228,963
				44,496,456	1.55 (Reg)	
2000	2001	14,883,720	1.17	7,604,465	4.07 (L/L)	60,911,856
				38,423,671	1.41 (Reg)	
2001	2002	11,263,498	0.69	17,452,832	0.72	28,716,331
2002	2003	5,400,000	2.62 (L/L)	17,400,000	4.02 (L/L)	47,100,000
		6,800,000	1.78 (Reg)	17,500,000	2.39 (Reg)	
2003	2004	5,960,363	3.24 (L/L)	12,006,165	4.20 (L/L)	50,618,272
		8,615,776	1.54 (Reg)	24,035,968	2.29 (Reg)	
2004	2005	6,100,000	(L/L)	11,500,000	(L/L)	63,000,000
		7,400,000	(Reg)	25,300,000	(Reg)	
2005	2006 ^c	14,000,000	(L/L, Reg)	12,000,000	(L/L)	61,000,000
				35,000,000	(Reg)	

Source: Douglas Island Pink and Chum Inc.

^a Late large release -Fry are held and fed for longer periods prior to release.

^b Regular release –Normal fry release timing

^c Planned releases.

Table 14.—Summary of returns from DIPAC chum salmon enhancement projects in lower Lynn Canal, 1991–2004 with projections for 2005.

Year	Total	% Estimated	Estimated	Boat Harbor			Amalga Harbor		
	Commercial Catch	Hatchery Contribution	Hatchery Contribution	Commercial Catch ^b	Cost Recovery	Total Return	Commercial Catch ^b	Cost Recovery	Total Return
1991	111,465	50.1%	55,818	55,818	0	55,818	0		
1992	162,231	52.9%	85,811	85,811	0	85,811	0		
1993	246,174	78.2%	192,446	192,446	0	192,446	0		
1994	568,850	81.4%	463,106	135,640	0	135,640	327,466	124,994	452,460
1995	499,167	91.2%	455,336	176,495	0	176,495	278,841	267,533	546,374
1996	340,021	78.2%	265,957	62,477	10,872	73,349	203,480	968,448	1,171,928
1997	431,699	87.8%	378,851	163,350	2,920	166,270	215,502	692,593	908,095
1998	136,515	83.4%	113,885	59,001	0	59,001	54,884	508,686	563,570
1999	290,325	85.5%	248,167	96,438	0	96,438	151,729	723,298	875,028
2000	680,536	88.6%	602,838	226,317	0	226,317	376,521	1,342,141	1,718,662
2001	358,987	85.1%	305,590	84,005	0	84,005	221,585	540,124	761,709
2002	630,486	94.5%	595,690	143,912	0	143,912	451,778	1,151,413	1,603,191
2003	348,820	96.8%	329,961	91,507	0	91,507	238,454	1,826,922	2,065,376
2004	688,471	91.7%	631,307	316,675	0	398,524	288,548	1,060,801	1,408,117
2005	240,055	92.1%	222,553	95,028	0	95,028	127,525	248,071	375,596
2006 ^a						192,000			1,331,000
91–05 Average	382,253	82.5%	329,821	132,328	919	138,704	195,754	787,919	1,037,509

Source: Douglas Island Pink and Chum Inc.

^a 2006 projected return.

^b Includes contribution to the Lynn Canal commercial drift gillnet fishery only.




Figure 1.—Lynn Canal district, section and statistical area boundaries.




Figure 2.—Upper Lynn Canal showing Chilkat and Chilkoot lakes. (Note: the statistical area 115-35 (formerly Burro Creek SHA) has been changed to represent the waters of Taiya Inlet north of Taiya Point.)




Figure 3.—Lynn Canal sockeye salmon weekly abundance by stock (Data from 1983–1992 average).




Figure 4.—Run timing (weekly proportion CPUE) of Chinook, coho, summer and fall chum, and pink salmon in the Lynn Canal drift gillnet fishery. Data for period 1992 to 2005.







Figure 5.—Historical commercial catches of Chinook, sockeye, coho, pink, and chum (summer and fall) salmon in the District 15 (Lynn Canal) drift gillnet fishery, 1960 to 2005.



Note: Escapements estimates in 1994–2005 in Chilkat Lake were based on mark-recapture estimates. Marine harvest of sockeye salmon for Chilkoot Lake in 1998 and 1999 was estimated to be 2,200 and 4,258 fish, respectively.

Figure 6.—Historical escapement and harvest of Chilkoot and Chilkat lake sockeye salmon, 1976 to 2005.



Source: D. Barto, ADF&G Commercial Fisheries Division, unpublished data.

Figure 7.—Yearly comparisons of Chilkoot Lake autumn hydroacoustic counts of juvenile sockeye salmon and average zooplankton densities, 1987–1991 and 1995–2005.




Figure 8.—Peak aerial survey results for Sawmill Creek chum salmon, 1993–2005 compared with 1996–2005 average results.




Figure 9.—Peak aerial survey results for Western Lynn Canal chum salmon streams combined, 1995–2005 compared to the 1996–2005 average.




Figure 10.—Total Chilkat River coho salmon fish wheel catch by year, 1994–2005 compared to the 1994–2005 average.




Figure 11.—Average 1994–2005 run timing for Chilkat River sockeye salmon stocks at the Chilkat River fish wheels by fresh water age class.




Figure 12.—Map showing possible area in Lower Lynn Canal that may be opened for additional time during peak weeks of hatchery chum return (Statistical Weeks 28–31).

**APPENDIX A: LYNN CANAL AND CHILKAT RIVER KING
SALMON FISHERY MANAGEMENT PLAN**

Appendix A1.—Lynn Canal and Chilkat River King Salmon Fishery Management Plan 5 AAC 33.384.

(a) The purpose of this management plan is to ensure biological spawning escapement requirements of king salmon to the Chilkat River. It is the intent of the Board of Fisheries (board) that the Chilkat River king salmon be harvested in the fisheries that have historically harvested them. The board, through this management plan, recognizes that the commercial drift gillnet fishery in Chilkat Inlet, and the subsistence fisheries in Chilkat Inlet and the Chilkat River are directed primarily toward sockeye salmon but catch king salmon incidentally. A secondary goal of this management plan is to provide a reasonable opportunity to harvest sockeye salmon in the Chilkat Inlet and Chilkat River subsistence fisheries while minimizing the incidental harvest of king salmon. This management plan provides ADF&G guidelines to preclude allocation conflicts between the various user groups of this resource. The department shall manage the Chilkat River king salmon stocks in a conservative manner consistent with sustained yield principles.

(b) ADF&G shall close the subsistence net fisheries in Chilkat Inlet north of a line extending from an ADF&G regulatory marker approximately one mile south of Anchorage Point to an ADF&G regulatory marker directly north of the Lenikof Cove boat ramp, through July 15. In the Chilkat River, excluding that portion of the river from Haines highway mile 19, continuing upstream to Well's Bridge, from approximately the third week of June through the fourth week of July.

(c) ADF&G shall manage the commercial and sport fisheries in Lynn Canal to achieve an inriver run goal of 1,850 to 3,600 king salmon in the Chilkat River upstream of the department fish wheels located approximately adjacent to mile 9 of the Haines highway. The inriver run goal provides for the following:

- (1) a biological escapement goal (BEG) of 1,750 to 3,500 large king salmon (three ocean age and older) to the Chilkat River; and
- (2) an incidental harvest of king salmon in the Chilkat River subsistence sockeye fishery.

(d) ADF&G will evaluate the inriver run of king salmon based on the following:

- (1) The preseason projected run of Chilkat River king salmon to Lynn Canal;
- (2) inseason fisheries performance; and
- (3) inriver stock assessment programs.

(e) ADF&G shall manage the commercial and drift gillnet and troll fisheries in Lynn Canal, and the sport king salmon fishery in Chilkat Inlet, as follows:

- (1) ADF&G shall close the commercial troll fishery in Chilkat Inlet north of a ADF&G regulatory marker immediately north of Seduction Point through July 14;
- (2) if the projected inriver run of king salmon to the Chilkat River is 1,850 fish (three ocean age and older) or less, ADF&G shall:

-continued-

(A) close the commercial drift gillnet fishery in Chilkat Inlet north of a ADF&G regulatory marker immediately north of Seduction Point through the first two weeks of the fishery; during the third and fourth week of the fishery, the Chilkat Inlet north of Glacier Point shall be closed; during the fifth week, the commercial drift gillnet fishery in Chilkat Inlet north of Cannery Point shall be closed; and

(B) close sport fishing for king salmon in Chilkat Inlet north of a ADF&G regulatory marker immediately north of Seduction Point through June 30; close king salmon fishing in Chilkat Inlet north of a line extending from an ADF&G regulatory marker one mile south of Anchorage Point to an ADF&G regulatory marker directly north of the Letnikof Cove boat ramp, through July 15; in the remainder of Chilkat Inlet north of Seduction Point, from July 1 – July 15, sport fisherman are allowed a bag and possession limit of one king salmon, 28 inches or greater in length;

(3) if the projected inriver run of king salmon to the Chilkat River is 1,850 to 3,600 fish ADF&G shall;

(A) close the commercial drift gillnet fishery in Chilkat Inlet north of a ADF&G regulatory marker immediately north of Seduction Point through the first two weeks of the fishery; during the third week of the fishery, close the area in Chilkat Inlet north of Glacier Point; during the fourth week, close the area in Chilkat Inlet north of Cannery Point; and

(B) close sport fishing for king salmon in Chilkat Inlet north of a line extending from an ADF&G regulatory marker approximately one mile south of Anchorage Point to an ADF&G regulatory marker directly north of the Lenikof Cove boat ramp from April 15 through July 15;

(4) if the projected inriver run of king salmon to the Chilkat River is greater than 3,600 fish ADF&G shall;

(A) close the commercial drift gillnet fishery in Chilkat Inlet north of a ADF&G regulatory marker immediately north of Seduction Point through the first week of the fishery; during the second week of the fishery, close the area in Chilkat Inlet north of Glacier Point; during the third week, close the area in Chilkat Inlet north of Cannery Point; and

(B) close sport fishing for king salmon in Chilkat Inlet north of a line extending from an ADF&G regulatory marker approximately one mile south of Anchorage Point to an ADF&G regulatory marker directly north of the Lenikof Cove boat ramp from April 15 through July 15; the commissioner may, through emergency order, increase the bag and possession limits of king salmon north of Seduction Point.

**APPENDIX B: CALENDAR DATES FOR STATISTICAL WEEKS IN
2006**

Appendix B1.—Calendar dates for statistical weeks in 2006.

Week	Beginning	Ending	Week	Beginning	Ending
1	1-Jan	7-Jan	28	9-Jul	15-Jul
2	8-Jan	14-Jan	29	16-Jul	22-Jul
3	15-Jan	21-Jan	30	23-Jul	29-Jul
4	22-Jan	28-Jan	31	30-Jul	5-Aug
5	29-Jan	4-Feb	32	6-Aug	12-Aug
6	5-Feb	11-Feb	33	13-Aug	19-Aug
7	12-Feb	18-Feb	34	20-Aug	26-Aug
8	19-Feb	25-Feb	35	27-Aug	2-Sep
9	26-Feb	4-Mar	36	3-Sep	9-Sep
10	5-Mar	11-Mar	37	10-Sep	16-Sep
11	12-Mar	18-Mar	38	17-Sep	23-Sep
12	19-Mar	25-Mar	39	24-Sep	30-Sep
13	26-Mar	1-Apr	40	1-Oct	7-Oct
14	2-Apr	8-Apr	41	8-Oct	14-Oct
15	9-Apr	15-Apr	42	15-Oct	21-Oct
16	16-Apr	22-Apr	43	22-Oct	28-Oct
17	23-Apr	29-Apr	44	29-Oct	4-Nov
18	30-Apr	6-May	45	5-Nov	11-Nov
19	7-May	13-May	46	12-Nov	18-Nov
20	14-May	20-May	47	19-Nov	25-Nov
21	21-May	27-May	48	26-Nov	2-Dec
22	28-May	3-Jun	49	3-Dec	9-Dec
23	4-Jun	10-Jun	50	10-Dec	16-Dec
24	11-Jun	17-Jun	51	17-Dec	23-Dec
25	18-Jun	24-Jun	52	24-Dec	30-Dec
26	25-Jun	1-Jul	53	31-Dec	31-Dec
27	2-Jul	8-Jul			

**APPENDIX C: HISTORICAL AGE COMPOSITIONS OF SOCKEYE
SALMON ESCAPEMENTS TO CHILKOOT LAKE, CHILKAT
LAKE, CHILKAT RIVER MAINSTEM AND BERNERS BAY
RIVERS.**

Appendix C1.—Historical age composition of sockeye salmon escapements to Chilkat Lake, 1982–2005.

Year	Sample	Percent by Age Class										
	Size	0.3	1.1	1.2	1.3	1.4	2.1	2.2	2.3	2.4	3.2	3.3
1982	1,630	0.1	0.4	2.3	12.9	0.0	2.6	45.3	34.8	0.0	1.3	0.1
1983	2,848	0.0	0.7	3.2	38.0	0.0	2.7	27.9	27.1	0.0	0.1	0.0
1984	2,728	0.0	0.1	1.5	22.8	0.0	1.5	53.6	20.2	0.0	0.2	0.0
1985	1,333	0.0	0.6	0.7	9.2	0.2	3.3	39.7	45.7	0.0	0.5	0.0
1986	940	0.0	0.0	1.7	1.6	0.0	0.5	20.6	73.1	0.0	1.9	0.5
1987	1,461	0.0	0.9	1.8	24.5	0.0	2.7	34.2	35.0	0.1	0.5	0.2
1988	1,918	0.1	0.0	0.8	47.3	0.2	0.0	7.9	43.4	0.1	0.1	0.1
1989	3,874	0.0	0.0	0.7	42.8	0.1	0.0	28.9	27.3	0.0	0.0	0.1
1990	2,635	0.1	0.0	1.8	14.0	0.5	0.0	24.8	58.0	0.1	0.8	0.0
1991	1,602	0.0	0.0	2.1	36.1	0.1	0.0	21.8	39.5	0.2	0.0	0.2
1992	2,505	0.1	0.0	1.1	40.8	0.2	0.0	16.9	40.7	0.0	0.0	0.1
1993	2,367	0.1	0.0	6.4	15.0	0.0	0.0	36.2	38.7	0.0	3.6	0.0
1994	2,187	0.0	0.0	2.6	58.6	0.1	0.0	11.4	26.6	0.0	0.1	0.6
1995	2,691	0.0	0.0	5.5	27.1	0.9	0.0	17.7	48.6	0.0	0.1	0.0
1996	308	0.0	0.0	10.4	67.5	0.0	0.0	8.8	13.3	0.0	0.0	0.0
1997	750	0.0	0.4	38.8	19.9	0.0	1.3	14.0	25.6	0.0	0.0	0.0
1998	1,198	0.0	0.1	4.9	69.4	0.0	0.3	19.0	6.0	0.2	0.0	0.0
1999	2,548	0.0	0.0	1.7	31.6	0.1	0.0	14.3	52.0	0.2	0.0	0.0
2000	2,316	0.0	0.0	2.4	5.1	0.3	0.0	7.8	81.4	0.1	2.8	0.0
2001	2,441	0.0	0.0	2.9	54.7	0.1	0.0	11.8	25.9	0.2	0.1	4.1
2002	2,504	0.0	0.0	2.5	26.5	0.4	0.1	20.1	50.3	0.0	0.0	0.0
2003	2,169	0.1	0.2	5.1	21.0	0.2	0.4	14.8	57.5	0.3	0.3	0.0
2004	3,004	0.1	0.1	3.5	49.7	0.0	0.2	17.2	28.4	0.2	0.1	0.4
2005	1,732	0.0	0.1	4.0	30.1	0.6	0.4	10.1	54.6	0.1	0.0	0.0
Average	2,070	0.0	0.2	4.5	31.9	0.2	0.7	21.9	39.7	0.1	0.5	0.3
SE		0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.0	0.0	0.0

Appendix C2.—Historical age composition of sockeye salmon escapements to Chilkat River mainstem areas, 1984 to 2005.

Year	Sample	Percent by Age Class										
	Size	0.1	0.2	0.3	0.4	1.1	1.2	1.3	1.4	2.2	2.3	2.4
1984	145	0.0	6.2	26.2	0.0	0.7	2.1	64.8	0.0	0.0	0.0	0.0
1985	136	0.0	14.7	42.6	0.0	0.0	0.0	39.7	0.7	1.5	0.7	0.0
1986	114	0.0	6.1	49.1	0.9	0.0	14.9	26.3	0.9	0.0	0.9	0.9
1987	51	0.0	9.8	9.8	0.0	0.0	3.9	74.5	0.0	0.0	2.0	0.0
1988	93	1.1	36.6	32.3	0.0	1.1	23.7	5.4	0.0	0.0	0.0	0.0
1989	195	0.0	4.6	27.2	0.0	0.0	3.6	63.1	0.0	0.5	1.0	0.0
1990	57	0.0	14.0	19.3	0.0	0.0	5.3	57.9	0.0	0.0	3.5	0.0
1991	310	0.6	21.0	52.3	0.0	1.0	9.7	15.5	0.0	0.0	0.0	0.0
1992	437	0.5	18.3	24.9	0.5	2.1	3.2	50.6	0.0	0.0	0.0	0.0
1993	67	0.0	1.5	85.1	0.0	0.0	0.0	13.4	0.2	0.0	0.0	0.0
1994	574	2.4	31.5	26.5	0.0	2.4	20.4	16.0	0.0	0.5	0.0	0.0
1995	474	0.0	29.7	36.1	0.2	0.0	12.4	21.5	0.0	0.0	0.0	0.0
1996	585	0.0	11.6	62.4	0.0	0.2	5.6	20.2	0.0	0.0	0.0	0.0
1997	437	0.2	18.3	62.9	0.0	0.0	3.2	15.3	0.0	0.0	0.0	0.0
1998	429	1.2	28.0	42.0	0.0	1.9	7.5	19.6	0.0	0.0	0.0	0.0
1999	334	5.1	47.3	14.4	0.0	4.2	20.1	8.4	0.0	0.6	0.0	0.0
2000	556	0.0	28.4	62.6	0.0	0.4	4.5	4.1	0.0	0.0	0.0	0.0
2001	438	0.0	3.4	49.8	0.0	0.0	6.8	40.0	0.0	0.0	0.0	0.0
2002	302	0.7	13.6	21.9	0.0	0.7	10.9	52.3	0.0	0.0	0.0	0.0
2003	547	0.0	47.9	21.4	0.0	2.6	13.3	14.4	0.0	0.0	0.2	0.0
2004	848	0.0	14.4	47.6	0.0	0.0	21.9	15.8	0.0	0.2	0.0	0.0
2005	672	0.0	31.0	43.0	0.0	0.0	8.2	16.8	0.0	0.0	0.0	0.0
Average	355	0.5	19.9	39.1	0.1	0.8	9.1	29.8	0.1	0.2	0.4	0.0
SE		0.0	0.2	0.2	0.0	0.0	0.1	0.3	0.0	0.0	0.0	0.0

Appendix C3.—Historical age composition of sockeye salmon escapements to Chilkoot Lake, 1982–2005.

Year	Sample Size	Percent By Age Class										
		0.3	1.1	1.2	1.3	1.4	2.1	2.2	2.3	2.4	3.2	3.3
1982	1,687	0.1	0.0	19.0	78.4	0.9	0.1	0.5	0.9	0.0	0.0	0.0
1983	1,790	0.1	0.1	12.0	60.4	0.2	0.1	1.4	25.8	0.0	0.0	0.0
1984	1,902	0.0	0.0	4.5	86.7	0.8	0.0	0.4	7.6	0.0	0.0	0.0
1985	1,623	0.0	0.1	12.2	66.4	2.4	0.0	2.6	15.9	0.3	0.1	0.0
1986	2,147	0.0	0.0	13.2	67.0	0.6	0.0	2.2	16.8	0.1	0.0	0.0
1987	2,207	0.0	0.0	8.4	69.2	0.2	0.0	2.2	19.8	0.1	0.0	0.0
1988	2,661	0.0	0.0	4.4	77.9	1.4	0.0	2.7	13.2	0.3	0.0	0.0
1989	2,586	0.0	0.0	4.5	54.9	1.2	0.0	5.0	33.5	0.4	0.0	0.5
1990	2,815	0.0	0.0	2.0	45.4	0.1	0.0	1.5	49.1	0.1	0.0	0.1
1991	2,297	0.0	0.0	12.5	55.9	0.4	0.0	4.9	25.9	0.3	0.0	0.1
1992	2,039	0.0	0.0	1.8	62.6	0.7	0.0	5.8	28.3	0.5	0.1	0.1
1993	2,075	0.0	0.0	2.6	35.6	0.3	0.0	1.8	59.0	0.4	0.0	0.2
1994	1,986	0.1	0.0	1.8	66.9	0.6	0.0	1.6	28.8	0.2	0.1	0.1
1995	606	0.0	0.0	44.1	30.7	0.8	0.0	3.8	20.0	0.7	0.0	0.0
1996	2,063	0.0	0.0	6.2	84.2	0.2	0.0	0.8	8.5	0.0	0.0	0.0
1997	2,111	0.0	0.0	2.2	90.1	0.1	0.0	0.4	7.1	0.0	0.0	0.0
1998	941	0.0	0.0	5.0	60.6	1.4	0.0	2.1	30.6	0.1	0.0	0.1
1999	2,033	0.0	0.0	28.8	46.5	0.2	0.0	8.1	16.3	0.2	0.0	0.0
2000	2,228	0.0	0.0	13.2	58.6	0.1	0.0	1.9	26.1	0.0	0.0	0.0
2001	2,345	0.3	0.0	4.8	89.8	0.0	0.0	0.2	4.9	0.0	0.0	0.0
2002	2,836	0.0	0.0	6.4	89.6	0.5	0.0	1.1	2.5	0.0	0.0	0.0
2003	2,611	0.0	0.0	41.3	45.0	0.4	0.0	4.2	9.1	0.0	0.0	0.0
2004	2,715	0.0	0.0	14.8	71.0	0.0	0.0	5.9	8.1	0.0	0.0	0.0
2005	2,730	0.0	0.0	19.9	67.5	0.0	0.0	3.9	8.6	0.0	0.0	0.0
Average	2,126	0.0	0.0	11.9	65.0	0.6	0.0	2.7	19.4	0.2	0.0	0.1
SE		0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0

Appendix C 4.—Historical age composition of sockeye salmon escapements to Berners Bay rivers, 1984 to 2005.

Year	Sample	Percent by Age Class										
	Size	0.1	0.2	0.3	0.4	1.1	1.2	1.3	1.4	2.1	2.2	2.3
1984	319	0.0	0.0	6.6	0.0	0.0	1.6	91.8	0.0	0.0	0.0	0.0
1985	84	3.6	10.7	4.8	1.2	4.8	11.9	61.9	0.0	0.0	1.2	0.0
1986	189	0.0	2.6	35.4	0.0	2.1	12.7	46.0	0.0	0.0	0.0	1.1
1987	133	0.0	0.0	32.3	0.0	0.0	4.5	62.4	0.0	0.0	0.0	8.0
1988	109	0.0	0.0	7.3	0.0	0.0	10.1	82.6	0.0	0.0	0.0	0.0
1989	285	0.0	1.8	2.1	0.0	1.1	6.7	85.3	0.0	0.4	0.7	2.1
1990	136	0.0	2.2	14.7	0.0	3.7	40.4	33.8	0.0	0.0	0.7	4.4
1991	409	0.7	3.2	27.6	0.0	1.5	15.6	50.1	0.2	0.0	0.2	0.7
1992	452	0.0	2.7	18.4	0.0	0.9	4.0	74.1	0.0	0.0	0.0	0.0
1993	451	0.0	1.3	30.8	0.0	0.9	14.4	51.7	0.0	0.0	0.2	0.7
1994	478	0.0	1.3	3.3	0.0	3.8	13.2	77.8	0.2	0.0	0.2	0.2
1995	333	0.0	2.7	13.5	0.0	0.3	35.7	45.3	1.2	0.0	0.6	0.6
1996	492	0.0	0.2	8.9	0.0	0.4	2.8	87.6	0.0	0.0	0.0	0.0
1997	218	0.5	0.9	38.5	0.0	0.9	11.0	48.2	0.0	0.0	0.0	0.0
1998	314	0.0	0.3	14.6	0.0	0.3	4.1	80.6	0.0	0.0	0.0	0.0
1999	297	0.3	0.3	6.7	0.0	1.0	8.1	82.2	0.0	0.0	0.0	1.3
2000	290	0.0	4.5	19.3	0.0	0.3	12.1	63.4	0.0	0.0	0.0	0.3
2001	339	0.0	1.2	17.7	0.3	3.2	4.7	72.9	0.0	0.0	0.0	0.0
2002	235	0.4	1.3	5.1	0.0	0.9	24.3	68.1	0.0	0.0	0.0	0.0
2003	278	0.0	1.1	11.2	0.0	1.1	17.1	72.7	0.0	0.0	0.0	7.6
2004	187	0.0	0.5	5.3	0.0	1.1	17.1	75.9	0.0	0.0	0.0	0.0
2005	119	0.8	5.0	13.4	0.0	2.5	15.9	61.6	0.8	0.0	0.0	0.0
Average	287	0.3	1.8	15.4	0.1	1.3	13.0	67.4	0.1	0.0	0.2	1.3
SE		0.0	0.0	0.1	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.0

**APPENDIX D: DATA COLLECTED FROM THE INSEASON
INFORMATION SYSTEM TO DETERMINE FISHERY
PERFORMANCE BY SPECIES**

Appendix D1.—Data collected from the inseason information system to determine fishery performance by species.

Sockeye Salmon

- a. Inseason abundance forecasts: Forecasts will be obtained by comparing current-year total return information (catch plus escapement), and expanding those results by historical run timing percentages for each stock.
- b. Escapement tracking: Daily escapements are tracked at the Chilkoot River weir. The weir provides timely data for inseason assessment as fish pass that weir within one week of fishery. Chilkat Lake/River sockeye salmon escapements will be monitored using two fish wheels in the lower Chilkat River. Fish wheel catch alone is not a definitive index of abundance but current year data will be compared to historic data. The Chilkat weir will be operated to provide a site for mark-recovery, biological sampling, and a timely preliminary estimate of escapement into Chilkat Lake.
- c. Inseason catch figures: Inseason catch figures are from the ADF&G fish ticket system. In the first 24 hours of an opening interview data from the fleet is used to estimate catches. After that time a sub sample of deliveries is expanded to total effort to estimate weekly catch.
- d. Stock contributions: Inseason catch stock contributions are estimated each week from random scale samples. Estimates are made for three groups: Chilkoot Lake, Chilkat Lake, and a combination of Berners Bay and Chilkat mainstem. Postseason stock contributions are made to add to the historic database from which models are derived. Escapements are sampled for scales to determine age structure of spawners in order to combine with catch data for spawner-recruit and preseason databases.
- e. Fishery monitoring: Site-specific fishery performance data and scale sampling are used to monitor migration paths and identify areas of overlap between stocks. Information is also provided on fish buildups in specific areas.

Fall and Summer Chum and Pink Salmon

- a. Inseason catch figures: Inseason catch data are obtained from the ADF&G fish ticket system. In the first 24 hours of an opening interview data from the fleet are used to estimate total harvests. After that time, a sub sample of deliveries is expanded to total effort to estimate catches.
- b. Aerial surveys and fish wheel catch: Escapement rates and distribution are monitored by aerial survey inseason when feasible and throughout the peak spawning period. Current fish wheel catches of salmon are compared to historical levels to estimate escapement rates.
- c. Fishery monitoring: Collect catch data and other fishery performance information such as effort level, fishing conditions, influence of northerly winds on rate of entry into Chilkat River, and observations of fish buildups.
- d. Fish wheel index: Lower Chilkat River fish wheel catch rates will be compared to the historical database to be used as an indicator of abundance.

-continued-

Coho Salmon

- a. Inseason catch figures: Inseason catch figures from the ADF&G fish ticket system.

- b. Aerial and foot surveys: Peak spawner counts are not obtained until postseason. However, in some index systems, Berners River and Chilkoot Lake, early season surveys provide an indication of escapement rates when water levels and conditions allow.

- c. Fishery monitoring: Availability of coho salmon is judged by comparing current CPUE and catch to the historical average and by the relative abundance of coho salmon in specific areas.

- d. Fish wheel index: Lower Chilkat River fish wheel catch rates will be compared to the historical database to be used as an indicator of abundance.

- e. Berners River coded-wire tag monitoring and inseason projections: Coded-wire tag recoveries for Berners River and Chilkat River coho salmon are monitored inseason by commercial fisheries staff. Data collected from this program will be used to project the total return of Berners and Chilkat River coho salmon. This system is used as an index for other Lynn Canal coho salmon stocks.

**APPENDIX E: INCLUSIVE DATES OF OPERATION FOR
CHILKOOT LAKE AND CHILKAT LAKE WEIRS AND CHILKAT
RIVER FISH WHEELS.**

Appendix E1.—Inclusive dates of operation for Chilkoot and Chilkat weirs and Chilkat River fish wheels, 1976 to 2005.

Year	Chilkoot Lake Weir Operation	Chilkat Lake Weir Operation	Chilkat River Fish Wheel Operation
1976	5/30–11/2	6/3–10/21	
1977	5/28–9/11	6/3–9/27	8/21–10/21
1978	6/6–11/7	6/05–11/05	8/14–11/9
1979	6/9–11/5	6/9–11/11	
1980	6/15–10/5	6/15–10/08	
1981	6/10–10/12	6/11–10/22	
1982	6/3–9/16	6/24–10/06	10/5–10/26
1983	6/4–11/13	6/22–11/12	8/9–10/3
1984	6/3–9/14	6/9–10/07	
1985	6/5–10/21	6/23–10/22	
1986	6/6–10/29	6/16–11/14	
1987	6/4–11/2	6/19–11/20	
1988	6/9–11/12	6/18–11/14	
1989	6/4–10/30	6/5–10/28	
1990	6/3–10/30	6/6–11/13	8/14–10/25
1991	6/7–10/8	7/10–10/24	5/8–7/20
1992	6/2–9/26	6/8–10/15	
1993	6/3–9/30	6/13–10/14	
1994	6/4–9/24	5/20–10/5	6/18–9/11
1995	6/5–9/11	6/8–10/9	6/16–9/16
1996	6/6–9/11	Weir not operated	6/22–9/16
1997	6/4–9/9	Weir not operated	6/11–10/09
1998	6/4–9/13	6/9–10/13	6/9–10/13
1999	6/4–9/13	6/30–10/28	6/7–10/08
2000	6/3–9/12	6/16–10/18	6/9–10/07
2001	6/7–9/12	6/19–10/13	6/6–10/07
2002	6/8–9/11	6/23–10/18	6/7–10/19
2003	6/6–9/9	6/27–10/10	6/6–10/21
2004	6/3–9/12	7/6–10/13	6/7–10/19
2005	6/6–6/12	6/28–10/12	6/7–10/11