

**Fishery Data Series No. 11-01**

---

---

**Stock Assessment of Arctic Grayling in the Delta  
River, 2008**

by

**Andrew D. Gryska**

---

---

January 2011

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



## Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the *Système International d'Unités* (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

<b>Weights and measures (metric)</b>		<b>General</b>		<b>Mathematics, statistics</b>	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	$H_A$
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	$e$
hectare	ha	at	@	catch per unit effort	CPUE
kilogram	kg	compass directions:		coefficient of variation	CV
kilometer	km	east	E	common test statistics	(F, t, $\chi^2$ , etc.)
liter	L	north	N	confidence interval	CI
meter	m	south	S	correlation coefficient	
milliliter	mL	west	W	(multiple)	R
millimeter	mm	copyright	©	correlation coefficient	
		corporate suffixes:		(simple)	r
<b>Weights and measures (English)</b>		Company	Co.	covariance	cov
cubic feet per second	ft <sup>3</sup> /s	Corporation	Corp.	degree (angular)	°
foot	ft	Incorporated	Inc.	degrees of freedom	df
gallon	gal	Limited	Ltd.	expected value	$E$
inch	in	District of Columbia	D.C.	greater than	>
mile	mi	et alii (and others)	et al.	greater than or equal to	≥
nautical mile	nmi	et cetera (and so forth)	etc.	harvest per unit effort	HPUE
ounce	oz	exempli gratia		less than	<
pound	lb	(for example)	e.g.	less than or equal to	≤
quart	qt	Federal Information Code	FIC	logarithm (natural)	ln
yard	yd	id est (that is)	i.e.	logarithm (base 10)	log
		latitude or longitude	lat. or long.	logarithm (specify base)	log <sub>2</sub> , etc.
<b>Time and temperature</b>		monetary symbols		minute (angular)	'
day	d	(U.S.)	\$, ¢	not significant	NS
degrees Celsius	°C	months (tables and figures): first three letters	Jan, ..., Dec	null hypothesis	$H_0$
degrees Fahrenheit	°F	registered trademark	®	percent	%
degrees kelvin	K	trademark	™	probability	P
hour	h	United States (adjective)	U.S.	probability of a type I error	
minute	min	United States of America (noun)	USA	(rejection of the null hypothesis when true)	$\alpha$
second	s	U.S.C.	United States Code	probability of a type II error	
		U.S. state	use two-letter abbreviations (e.g., AK, WA)	(acceptance of the null hypothesis when false)	$\beta$
<b>Physics and chemistry</b>				second (angular)	"
all atomic symbols				standard deviation	SD
alternating current	AC			standard error	SE
ampere	A			variance	
calorie	cal			population	Var
direct current	DC			sample	var
hertz	Hz				
horsepower	hp				
hydrogen ion activity	pH				
(negative log of)					
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

***FISHERY DATA SERIES NO. 11-01***

**STOCK ASSESSMENT OF ARCTIC GRAYLING IN THE DELTA RIVER,  
2008**

by  
Andrew D. Gyska  
Division of Sport Fish, Fairbanks

Alaska Department of Fish and Game  
Division of Sport Fish, Research and Technical Services  
333 Raspberry Road, Anchorage, Alaska, 99518-1565

January 2011

This investigation was partially financed by the Federal Aid in Sport Fish Restoration Act (16 U.S.C. 777-777K) under Project F-10-24, Job No. R-3-2(e) and in-kind funding through the U.S. Bureau of Land Management in association with Agreement No. BLM L09AC15726.

ADF&G Fishery Data Series was established in 1987 for the publication of Division of Sport Fish technically oriented results for a single project or group of closely related projects, and in 2004 became a joint divisional series with the Division of Commercial Fisheries. Fishery Data Series reports are intended for fishery and other technical professionals and are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm> This publication has undergone editorial and peer review.

*Andrew D. Gryska,  
Alaska Department of Fish and Game, Division of Sport Fish,  
1300 College Road, Fairbanks, AK 99701-1599 USA*

*This document should be cited as:*

*Gryska, A. D. 2011. Stock assessment Arctic grayling in the Delta River, 2008. Alaska Department of Fish and Game, Fishery Data Series No. 11-01, Anchorage.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375.

# TABLE OF CONTENTS

	<b>Page</b>
LIST OF TABLES.....	ii
LIST OF FIGURES .....	ii
LIST OF APPENDICES .....	ii
ABSTRACT .....	1
INTRODUCTION.....	1
OBJECTIVES.....	5
METHODS.....	5
EXPERIMENTAL AND SAMPLING DESIGN .....	5
Data Collection.....	8
Data Analysis.....	8
Abundance Estimate .....	8
Length Composition .....	9
RESULTS.....	9
Movement.....	9
Abundance Estimate.....	9
Length Composition .....	10
DISCUSSION.....	15
ACKNOWLEDGEMENTS.....	16
REFERENCES CITED .....	17
APPENDIX A EQUATIONS AND STATISTICAL METHODOLOGY .....	19
APPENDIX B DATA FILE LISTING.....	27

## LIST OF TABLES

Table	Page
1. Estimated sport fishing effort (angler days) for all species of fish, and estimates of sport fishing catch and harvest of Arctic grayling in the Delta River, Alaska.....	3
2. Results of diagnostics used to detect and correct for size-selective sampling (Appendix A2) for estimating abundance and length compositions of Arctic grayling in the Delta River, July 2008. ....	11
3. Results of consistency tests for the Petersen estimator (Appendix A3) for estimating abundance of Arctic grayling in the Delta River, July 2008.....	12
4. Number of Arctic grayling 240–269 mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.....	12
5. Number of Arctic grayling 270–299 mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.....	13
6. Number of Arctic grayling $\geq 300$ mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.....	13
7. Number of Arctic grayling $\geq 330$ mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.....	14
8. Number of representative fish sampled ( $n$ ), estimated, estimated abundance ( $\hat{N}_k$ ), and estimated proportion of abundance ( $\hat{p}[\hat{N}_k]$ ), by length category for the population of Arctic grayling ( $\geq 240$ mm FL) in the Delta River, July 2008.....	14
9. Number of representative fish sampled ( $n$ ), estimated abundance ( $\hat{N}_k$ ), and estimated proportion of abundance ( $\hat{p}[\hat{N}_k]$ ), by length category for the population of Arctic grayling ( $\geq 270$ mm FL) in the Delta River, July 2008.....	15

## LIST OF FIGURES

Figure	Page
1. Location of the Delta River drainage. ....	2
2. Delta River study area and adjacent mining claims of Pure Nickel Inc. (demarcated approximately by shaded area).....	4
3. Delta River study area with demarcation of each 1.05-km angling section. For consistency tests data from angling sections were pooled in the following manner: I (1–4), II (5–9), and III (10–16). ....	7
4. Proportions of recaptured Arctic grayling ( $n = 87$ ) that moved (km) upstream (positive values) or downstream (negative values) in the Delta River study area, July 2008. ....	10

## LIST OF APPENDICES

Appendix	Page
A1. Equations for calculating estimates of abundance and its variance using the Bailey-modified Petersen estimator.....	20
A2. Detection of size and/or sex selective sampling during a two-sample mark recapture experiment and its effects on estimation of population size and population composition. ....	21
A3. Tests of consistency for the Petersen estimator (from Seber 1982, page 438). ....	23
A4. Equations for estimating length composition and their variances for the population.....	24
B1. Data files for population estimate of Arctic grayling captured in the Delta River, 2008. ....	28

## ABSTRACT

The Alaska Department of Fish and Game and the U.S. Bureau of Land Management cooperated in a stock assessment of the Arctic grayling *Thymallus arcticus* population inhabiting a 17-km portion of the Delta River federally designated as Wild and Scenic and located in the central Alaska Range. The study was conducted during July 2008 using a mark-recapture experiment to estimate abundance and length composition of the population.

Abundance was estimated for Arctic grayling  $\geq 240$  mm FL ( $\hat{N} = 44,212$ ; SE = 9,108),  $\geq 270$  mm FL ( $\hat{N} = 23,152$ ; SE = 3,189), and  $\geq 330$  mm FL ( $\hat{N} = 5,864$ ; SE = 818). Of Arctic grayling  $\geq 240$  mm FL, 48% were 240 – 269 mm FL and 24% were 270 – 299 mm FL, and among Arctic grayling  $\geq 270$  mm FL, 55% were 270 – 299 mm FL. The density of Arctic grayling  $\geq 240$  and  $\geq 270$  mm FL in the Delta River was the greatest ever observed among published density estimates for Alaskan fluvial Arctic grayling.

Key words: Arctic grayling, *Thymallus arcticus*, abundance, length composition, hook-and-line, mark-recapture, Delta River, Alaska.

## INTRODUCTION

The Delta River begins at the outlet of Lower Tangle Lake in the Alaska Range about 100 km south-southwest of Delta Junction, Alaska. A large portion of the river and drainage have been designated a wild and scenic river by the US Bureau of Land Management (BLM); specifically, the Tangle Lakes portion of the drainage is classified as scenic, the Delta River from Lower Tangle Lake to mile 212 Richardson Highway is classified as wild, and the remaining portion through Black Rapids is classified as recreational (Figure 1). Initially, the river is a clear stream rarely prone to high muddy water due to the buffering of the large Tangle Lakes system. The river traverses 3.1 km from Lower Tangle Lake before it passes over a series of waterfalls, which are neither navigable nor allow fish to pass upstream. From the waterfalls, the river travels 17 km before the glacial Eureka Creek tributary intercepts the Delta River and it is the first of several glacial tributaries to render the Delta River glacially turbid for the remainder of its course to the Tanana River.

Each year, numerous recreational floaters and anglers travel through the Tangle Lakes and the Upper Delta River, almost all of whom take-out at Mile 212. For anglers, Arctic grayling *Thymallus arcticus* and lake trout *Salvelinus namaycush* (primarily in the Tangle Lakes) comprise most of the catch, though burbot *Lota lota* and round whitefish *Prosopium cylindraceum* are also caught to a lesser degree. Other species in the drainage include longnose sucker *Catostomus catostomus*, Dolly Varden *Salvelinus malma*, slimy sculpin *Cottus cognatus*, and at the Delta River mouth a spawning population of chum salmon *Oncorhynchus keta* (Peckham 1976). Despite the number of floaters, the level of harvest is moderate (Table 1), and accordingly the regulations for Arctic grayling in the Delta River drainage are as liberal as allowed in the Arctic grayling management plan (Swanton and Wuttig *In prep*). The regulations are:

1. a daily bag and possession limit of five Arctic grayling;
2. any size Arctic grayling may be retained; and,
3. no seasonal spawning period closure.




Figure 1.—Location of the Delta River drainage.

Table 1.—Estimated sport fishing effort (angler days) for all species of fish, and estimates of sport fishing catch and harvest of Arctic grayling in the Delta River, Alaska.

Year <sup>a</sup>	All species	Arctic grayling	
	Angler days	Harvest	Catch
1996	654	291	3,215
1997	553	770	4,112
1998	410	160	5,562
1999	1,023	640	5,924
2000	551	243	2,813
2002	663	121	5,600
2005	847	259	2,593
2006	882	304	3,317
		Average	
	698	349	4,142

<sup>a</sup> Data from Howe et al. 2001a-d; Jennings et al., 2006, 2009 a-b; Walker et al. 2003.

In addition to being a recreationally significant area, the Delta and Eureka drainages hold large deposits of economically important minerals and gold has been intermittently mined by small operations. The scope of the mining operations may increase dramatically with the development of the Pure Nickel Inc. MAN Alaska Project that has located significant amounts of platinum group elements, nickel, gold, cobalt and copper in an area ~280-mi<sup>2</sup> (Figure 2). Actual mining is projected to begin within as little as five years and this potential development increased the need for more comprehensive understanding of resident fishes that may be affected by the mining area.

This need prompted a cooperative project between the BLM and ADF&G, Division of Sport Fish, for which two mutual informational needs were identified: improved understanding of the Arctic grayling population within the Delta River between the falls and Eureka Creek and more comprehensive information on fish distributions within the drainage. Information on this Arctic grayling population was very limited and based on a few small-scaled studies during which fish were sampled (Peckham 1974, 1975; Baker 1989; Holmes et al. 1990). Distribution information of fish species was collected by Carlton (1976) and Peckham (1976), but large areas of the drainage to the west of the river were not sampled.

The cooperative project had three goals to be completed over a 2-year period: 1) estimate population size of Arctic grayling between the falls and Eureka Creek; 2) determine seasonal distribution and habitat use of those Arctic grayling using radiotelemetry; and 3) conduct species inventory of streams and lakes within an established Upper Delta River study area via helicopter transport. This report focuses on the first goal related to estimating Arctic grayling population size in the Delta River.




Figure 2.—Delta River study area and adjacent mining claims of Pure Nickel Inc. (demarcated approximately by shaded area).

## OBJECTIVES

The research objectives for 2008 were to:

1. estimate the abundance of Arctic grayling  $\geq 270$  and  $\geq 330$  mm FL in the Delta River study area (17 km) during July 2008, such that the estimates were within 25% of the actual abundance 95% of the time; and,
2. estimate the length composition (in 10-mm intervals) of the Arctic grayling  $\geq 270$  mm FL in the Delta River study area (17 km) during July 2008, such that the estimates were within five percentage points of the true value 95% of the time.

The study area encompassed a 17-km reach of the Delta River from the falls to the mouth of glacial Eureka Creek (Figure 2) and it wholly contained the grayling fishery in the Delta River downstream of the falls. The size limits identified in the objectives, 270 and 330 mm FL, are commonly used standards in Arctic grayling stock assessments or management objectives within ADF&G Region III. The 270-mm length limit is related to the minimum harvest length often used in regulations. The 330-mm length limit is the length at which Arctic grayling begin to be considered large by anglers and the abundance of fish of this size is often used to manage or evaluate Interior Alaska fisheries. Because the length at which Arctic grayling recruit to the gear can range between 200 and 270 mm (Gryska 2004a, b; Wuttig 2004), all fish  $\geq 200$  mm FL were tagged in the event abundance at a lower length limit could be estimated.

## METHODS

### EXPERIMENTAL AND SAMPLING DESIGN

During July 2008, the study was designed to estimate abundance and length composition of Arctic grayling within the 17-km study area of the Delta River (Figure 2) using two-event Petersen mark-recapture techniques for a closed population (Seber 1982). The study was designed to satisfy the following assumptions:

1. the population was closed (Arctic grayling did not enter the population, via growth or immigration, or leave the population, via death or emigration, during the experiment);
2. all Arctic grayling had a similar probability of capture in the first event or in the second event, or marked and unmarked Arctic grayling mixed completely between events;
3. marking of Arctic grayling did not affect the probability of capture in the second event;
4. marked Arctic grayling were identifiable during the second event; and,
5. all marked Arctic grayling were reported when recovered in the second event.

The estimator used was a modification of the general form of the Petersen estimator:

$$\hat{N} = \frac{n_1 n_2}{m_2}, \quad (1)$$

where:

$n_1$  = the number of Arctic grayling marked and released during the first event;

$n_2$  = the number of Arctic grayling examined for marks during the second event; and,

$m_2$  = the number of marked Arctic grayling recaptured during the second event.

The sampling design and data collected allowed the validity of the five assumptions to be ensured or tested. The specific form of the estimator was determined from the experimental design and the results of diagnostic tests performed to evaluate if the assumptions were met (Appendices A1, A2, and A3).

The study area was 17 km in length and divided into 16 approximately 1.05-km sections (Figure 3). This division served to guide sampling and provided a minimum geographic scale at which to conduct diagnostic tests. The first event occurred during July 8–11, and the second during July 15–18. During each event, three two-person crews sampled in a downstream progression, covering two to five sections a day (variation due to logistical constraints, hydrographic conditions, and densities of Arctic grayling). Daily, each crew was assigned to a section or more to capture fish using hook-and-line gear. The sampling schedule resulted in a 3-day hiatus between events or a 6-day hiatus for each specific reach of river sampled, which was deemed long enough to ensure that capture probability for each fish did not vary between events as a result of being captured and marked (DeCicco 1997). The distribution and allocation of sampling effort were planned to ensure adequate sample sizes were attained, no segments of the population were isolated from the experiment, and the study area was sampled uniformly.

The selection of the sampling area and timing of the experiment ensured that the movement of fish did not violate the assumption of closure due to combined emigration and immigration. No emigration or immigration across the lower boundary of the study area was expected because it was at the mouth of the glacial Eureka Creek, after which the Delta River becomes very turbid and is not preferred habitat for summer feeding. The upper boundary was a waterfall preventing upstream migration and believed to not have much downstream migration based on a previous study (Peckham 1976). During the summer feeding period, Arctic grayling were expected to have only localized movements (e.g., < 1.6 rkm), as has been observed during previous experiments (Ridder and Gryska 2000; Gryska 2001). The duration of the hiatus was designed to: 1) eliminate potential bias due to growth recruitment and mortality; 2) allow for localized mixing of marked and unmarked fish to eliminate isolated pockets of fish; and, 3) allow for marked fish to recover from the effects of handling between events.

During each event, all sampling areas were accessed using canoes or a power boat (one crew during second event). The upper boundaries of individual sampling sections were reached by floating in a canoe, after which each crew waded through its assigned angling section, excepting the lowermost sections (14–16) which were primarily fished from a boat because the channel is deeper and more incised. All waters were angled, and in an effort to subject fish to more similar capture probabilities, the work day was adjusted such that areas of high fish densities were fished for longer periods than low density areas. Areas of higher fish densities were identified by visually observing aggregations of Arctic grayling and by evaluating catch rates and preferred habitat (e.g. heads of pools).




Figure 3.–Delta River study area with demarcation of each 1.05-km angling section. For consistency tests data from angling sections were pooled in the following manner: I (1–4), II (5–9), and III (10–16).

The terminal gear consisted of a combination of flies (dry and wet) and rubber-bodied jigs. The degree to which each gear was used was left to each angler's discretion. Typically, jigs were used most often. At each angling location, captured Arctic grayling were temporarily held 1–10 minutes in a five-gallon bucket until data were collected. Afterwards they were generally released at or within 25 m of their capture locations and in no cases were fish displaced by more than 100 m or from a distinct feature such as a pool.

Sample size objectives for estimating abundance were established using methods in Robson and Regier (1964) and for length composition using the criteria developed by Thompson (1987) for multinomial proportions.

## **DATA COLLECTION**

After angling a portion of a section (25–50 m) all temporarily held fish were measured for length (mm FL), and carefully examined for marks. In the first event, all fish  $\geq 200$  mm FL were tagged with an individually numbered Floy<sup>TM</sup> FD-94 internal anchor tag (brown color, white print, numbered between 7,001 and 8,961) placed at the insertion of the dorsal fin so that the tag locks between the posterior interneural rays and received an upper caudal finclip to identify tag loss. To eliminate duplicate sampling in the second event, all fish received a lower caudal finclip. All fish in both events were carefully inspected for attendant Floy<sup>TM</sup> tags and fin clips and had their capture/release locations recorded using a GPS (latitude and longitude coordinate as decimal degrees, NAD27 Alaska datum). Fish captured in the first event that exhibited signs of injury, excessive stress, or imminent death were not marked and censored from the experiment.

## **DATA ANALYSIS**

### **Abundance Estimate**

When capturing fish in a river using angling equipment it is inherently difficult to approximate the taking of a simple random sample (i.e., a random sample without replacement). Therefore, samples from the Delta River were taken systematically in the sense of progressively moving downstream and sampling proportionally to the abundance of fish present (discussed above with respect to Assumption 2). Under these circumstances the Bailey-modified Petersen estimator (Appendix A1; Bailey 1951, 1952) is preferred over the Chapman-modified Petersen estimator (Chapman 1951) for estimating abundance.

Relative to Assumption 1, closure was not tested directly but inferred from examination of the movement of recaptured Arctic grayling within the study area. The data were examined for evidence of movement away from or towards the boundaries of the study area to provide evidence of significant immigration and emigration.

Violations of Assumption 2 relative to size were evaluated using Kolmogorov-Smirnov (K-S) two-sample tests with a significance level of  $\alpha = 0.05$ . There were four possible outcomes of these tests relative to evaluating size selective sampling (either one of the two samples, both, or neither of the samples had size selectivity) and two possible actions for abundance estimation (length stratify or not). The tests and possible actions for data analysis are outlined in Appendix A2. If stratification by size was required, capture probability by location were examined for each length stratum.

The tests for consistency of the Petersen estimator (Seber 1982; Appendix A3) were used to determine if, for each identified length stratum, stratification by location was required due to

spatiotemporal effects and to determine the appropriate abundance estimator: the pooled Bailey-modified Petersen estimator, the completely stratified Bailey-modified Petersen estimator, or a partially stratified estimator (Darroch 1961). Documentation of release location by section for each fish permitted the examination of multiple geographic stratification schemes for purposes of assumption testing, and testing was performed at the scale of a cluster defined by grouping of adjacent angling sections (sections 1–4, 5–9, and 10–16), which also corresponded to different hydrological characteristics in the river. Sections 1–4 have a relatively narrow and swift channels (e.g. <15 m); sections 5–9 are relatively moderate in width (e.g. 15–25 m) with well defined pool-riffle sequences that meander in a wider valley; and the lower most sections are generally wide (e.g. 25 to 50 m) with few discernible riffles and elongated, deep pools. This grouping strategy also provided a sufficient number of recaptures for diagnostic testing to ensure negligible statistical bias in  $\hat{N}$  (Seber 1982) and accommodated localized movements (i.e. within a 1-km radius) of Arctic grayling.

### **Length Composition**

Length composition of the population was estimated in 10-mm length categories using the procedures outlined in Appendix A4.

## **RESULTS**

### **MOVEMENT**

Because fish were released relatively close to their capture location (within about 25 m), movement was defined as a fish that was recaptured  $\geq 0.5$  km from its release location. Using this definition of movement, 69 of the 87 (79%) recaptured Arctic grayling had not moved (Figure 4), and 7 (8%) other recaptured Arctic grayling had moved between 0.5 and 1.5 km. Only 2 (2%) fish moved  $> 5$  km, having moved 6 and 8 km downstream. The virtual lack of any large or directional movements indicated that no meaningful immigration or emigration occurred during the experiment.

### **ABUNDANCE ESTIMATE**

In the 17-km study area, 3,597 Arctic grayling  $\geq 136$  mm FL were captured ( $n_1 = 1,767$ ,  $n_2 = 1,830$ ,  $m_2 = 87$ ), but the smallest recapture was 242 mm FL and abundance was estimated for fish  $\geq 240$  mm FL. Two hooking mortalities occurred during the second event. K-S tests (Appendix A2) results indicated Case IV for Arctic grayling  $\geq 240$  and  $\geq 270$  mm FL (Table 2). Although a single stratum of 240–299 mm FL was found to be Case I due to a marginally acceptable p-value (0.061) for one test and a borderline p-value (0.047) for the other, the stratum was assigned a Case IV (Appendix A2 has a more extensive discussion for this reasoning) because the D-value (0.35) was large, there were few recaptures ( $m_2 = 15$ ), and the capture probability for Arctic grayling 240–269 mm FL was half of that for those 270–299 mm FL (Tables 4 and 5). Abundance estimates were stratified by length in the following categories: 240–269 mm FL, 270–299 mm FL,  $\geq 300$  mm FL, and  $\geq 330$  mm FL. These strata were Case I, which indicated that there was no size selective sampling during both capture events and the data from both events were pooled for composition estimates of each stratum (Table 2).

Among all size groups, at least one consistency test (Appendix A3) failed to be rejected (Tables 3–7). Therefore, there was no need to geographically stratify and the Bailey-modified Petersen estimator was used to calculate abundance estimates for each stratum, which were pooled for total abundance.

Estimated abundances of Arctic grayling were:

- a. 240–269 mm FL was 21,060 (SE = 8,531);
- b. 270–299 mm FL was 10,465 (SE = 2,856);
- c.  $\geq 240$  mm FL was 44,212 (SE = 9,108);
- d.  $\geq 270$  mm FL was 23,152 (SE = 3,189);
- e.  $\geq 300$  mm FL was 12,686 (SE = 1,419); and,
- f.  $\geq 330$  mm FL was 5,864 (SE = 818).

## LENGTH COMPOSITION

A substantial proportion of the estimated population of Arctic grayling  $\geq 240$  mm FL was 240 to 269 mm FL (48%; Table 8), although that estimate of abundance was most imprecise. The more precise estimates of abundance of Arctic grayling  $\geq 270$  mm FL indicated that abundance of fish 270–299 mm FL comprised 45% of the estimated population,  $\geq 300$  mm FL comprised 55% of the estimated population, and  $\geq 330$  mm FL comprised 30% of the estimated population (Table 9).




Figure 4.—Proportions of recaptured Arctic grayling (n = 87) that moved (km) upstream (positive values) or downstream (negative values) in the Delta River study area, July 2008.

Table 2.—Results of diagnostics used to detect and correct for size-selective sampling (Appendix A2) for estimating abundance and length compositions of Arctic grayling in the Delta River, July 2008.

Stratum	Comparison and Test Statistic			Result
	M vs. R	C vs. R	M vs. C	
≥240 mm FL	D = 0.263 P-value < 0.001 Reject H <sub>0</sub>	D = 0.269 P-value < 0.001 Reject H <sub>0</sub>	D = 0.031 P-value = 0.382 Fail to Reject H <sub>0</sub>	Case IV, stratify
240–299 mm FL	D = 0.342 P-value = 0.061 Reject H <sub>0</sub>	D = 0.354 P-value = 0.047 Reject H <sub>0</sub>	D = 0.047 P-value = 0.426 Fail to reject H <sub>0</sub>	Case IV, stratify
240–269 mm FL	D = 0.194 P-value = 0.998 Fail to reject H <sub>0</sub>	D = 0.202 P-value = 0.997 Fail to reject H <sub>0</sub>	D = 0.052 P-value = 0.737 Fail to reject H <sub>0</sub>	Case I, do not stratify, use lengths and ages from both events for composition analysis
270–299 mm FL	D = 0.270 P-value = 0.405 Fail to reject H <sub>0</sub>	D = 0.328 P-value = 0.192 Fail to reject H <sub>0</sub>	D = 0.071 P-value = 0.303 Fail to reject H <sub>0</sub>	Case I, do not stratify, use lengths and ages from both events for composition analysis
≥300 mm FL	D = 0.139 P-value = 0.124 Fail to reject H <sub>0</sub>	D = 0.093 P-value = 0.566 Fail to reject H <sub>0</sub>	D = 0.060 P-value = 0.061 Fail to reject H <sub>0</sub>	Case I, do not stratify, use lengths and ages from both events for composition analysis
≥330 mm FL	D = 0.125 P-value = 0.491 Fail to reject H <sub>0</sub>	D = 0.073 P-value = 0.972 Fail to reject H <sub>0</sub>	D = 0.063 P-value = 0.242 Fail to reject H <sub>0</sub>	Case I, do not stratify, use lengths and ages from both events for composition analysis

Table 3.—Results of consistency tests for the Petersen estimator (Appendix A3) for estimating abundance of Arctic grayling in the Delta River, July 2008.

Stratum	Consistency Test		
	I	II	III
	Complete Mixing	Equal probability of Capture, 1 <sup>st</sup> Event	Equal Probability of Capture, 2 <sup>nd</sup> Event
240–269 mm FL	$\chi^2 = 5.61$ P-value = 0.23	$\chi^2 = 1.69$ P-value = 0.43	$\chi^2 = 0.81$ P-value = 0.67
270–299 mm FL	$\chi^2 = 15.52$ P-value = 0.02	$\chi^2 = 1.46$ P-value = 0.48	$\chi^2 = 1.25$ P-value = 0.53
≥300 mm FL	$\chi^2 = 92.52$ P-value <0.01	$\chi^2 = 3.56$ P-value = 0.17	$\chi^2 = 5.04$ P-value = 0.08
≥330 mm FL	$\chi^2 = 66.77$ P-value <0.01	$\chi^2 = 2.79$ P-value = 0.25	$\chi^2 = 2.22$ P-value = 0.33

Table 4.—Number of Arctic grayling 240–269 mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.

		Cluster where recaptured			$m_2$	$n_1$	$m_2/n_1^c$
		I	II	III			
Cluster <sup>a</sup> where marked	I	2	0	0	2	102	0.02
	II	0	1	0	1	146	0.01
	III	0	1	0	1	77	0.01
	$m_2$	2	2	0			
	$n_2$	87	165	71			
	$(m_2/n_2)^b$	0.02	0.01	0.00			

<sup>a</sup> Cluster refers to a grouping of adjacent sections: Cluster I = Sections 1-4; Cluster II = Sections 5-9, and Cluster III = Sections 10-16.

<sup>b</sup> Probability of capture during first event.

<sup>c</sup> Probability of capture during second event.

Table 5.–Number of Arctic grayling 270-299 mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.

		Cluster where recaptured			$m_2$	$n_1$	$m_2/n_1^c$
		I	II	III			
Cluster <sup>a</sup> where marked	I	4	1	0	5	110	0.05
	II	0	5	0	5	156	0.03
	II	0	0	1	1	68	0.01
	$m_2$	4	6	1			
	$n_2$	93	200	82			
	$(m_2/n_2)^b$	0.04	0.03	0.01			

<sup>a</sup> Cluster refers to a grouping of adjacent sections: Cluster I = Sections 1-4; Cluster II = Sections 5-9, and Cluster III = Sections 10-16.

<sup>b</sup> Probability of capture during first event.

<sup>c</sup> Probability of capture during second event.

Table 6.–Number of Arctic grayling  $\geq 300$  mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.

		Cluster where recaptured			$m_2$	$n_1$	$m_2/n_1^c$
		I	II	III			
Cluster <sup>a</sup> where marked	I	27	1	1	29	317	0.09
	II	0	36	3	39	496	0.08
	II	0	0	4	4	132	0.03
	$m_2$	27	37	8			
	$n_2$	321	467	191			
	$(m_2/n_2)^b$	0.08	0.08	0.04			

<sup>a</sup> Cluster refers to a grouping of adjacent sections: Cluster I = Sections 1-4; Cluster II = Sections 5-9, and Cluster III = Sections 10-16.

<sup>b</sup> Probability of capture during first event.

<sup>c</sup> Probability of capture during second event.

Table 7.—Number of Arctic grayling  $\geq 330$  mm FL marked ( $n_1$ ), examined ( $n_2$ ), and recaptured ( $m_2$ ) by cluster in the Delta River study area, July 2008.

		Cluster where recaptured			$m_2$	$n_1$	$m_2/n_1^c$
		I	II	III			
Cluster <sup>a</sup> where marked	I	19	0	0	19	167	0.11
	II	0	21	2	23	269	0.09
	II	0	0	3	3	58	0.05
	$m_2$	19	21	5			
	$n_2$	181	261	103			
	$(m_2/n_2)^b$	0.10	0.08	0.05			

<sup>a</sup> Cluster refers to a grouping of adjacent sections: Cluster I = Sections 1-4; Cluster II = Sections 5-9, and Cluster III = Sections 10-16.

<sup>b</sup> Probability of capture during first event.

<sup>c</sup> Probability of capture during second event.

Table 8.—Number of representative fish sampled ( $n$ ), estimated, estimated abundance ( $\hat{N}_k$ ), and estimated proportion of abundance ( $\hat{p}[\hat{N}_k]$ ), by length category for the population of Arctic grayling ( $\geq 240$  mm FL) in the Delta River, July 2008.

Length (mm FL)	$n$	$\hat{N}_k$	$\hat{SE}[\hat{N}_k]$	$\hat{p}[\hat{N}_k]$	$SE \hat{p}[\hat{N}_k]$
240–249	179	5,818	2,381	0.13	0.003
250–259	244	7,930	3,233	0.18	0.005
260–269	225	7,313	2,984	0.17	0.005
270–279	210	3,100	863	0.07	<0.001
280–289	228	3,365	935	0.08	<0.001
290–299	271	4,000	1,107	0.09	0.001
300–309	303	1,999	247	0.05	<0.001
310–319	293	1,933	240	0.04	<0.001
320–329	289	1,907	237	0.04	<0.001
330–339	289	1,907	237	0.04	<0.001
340–349	253	1,669	210	0.04	<0.001
350–359	205	1,352	175	0.03	<0.001
360–369	141	930	128	0.02	<0.001
370–379	91	600	91	0.01	<0.001
380–389	34	224	45	0.01	<0.001
390–399	21	139	34	<0.01	<0.001
400–409	4	26	13	<0.01	<0.001

Table 9.–Number of representative fish sampled (n), estimated abundance ( $\hat{N}_k$ ), and estimated proportion of abundance ( $\hat{p}[\hat{N}_k]$ ), by length category for the population of Arctic grayling ( $\geq 270$  mm FL) in the Delta River, July 2008.

Length (mm FL)	n	$\hat{N}_k$	$\hat{SE}[\hat{N}_k]$	$\hat{p}[\hat{N}_k]$	SE $\hat{p}[\hat{N}_k]$
270–279	210	3,100	863	0.13	<0.001
280–289	228	3,365	935	0.15	0.001
290–299	271	4,000	1,107	0.17	0.001
300–309	303	1,999	247	0.09	<0.001
310–319	293	1,933	240	0.08	<0.001
320–329	289	1,907	237	0.08	<0.001
330–339	289	1,907	237	0.08	<0.001
340–349	253	1,669	210	0.07	<0.001
350–359	205	1,352	175	0.06	<0.001
360–369	141	930	128	0.04	<0.001
370–379	91	600	91	0.03	<0.001
380–389	34	224	45	0.01	<0.001
390–399	21	139	34	0.01	<0.001
400–409	4	26	13	<0.01	<0.001

## DISCUSSION

The study goal was to estimate the abundance and length composition of the population in the study area, and both objectives were met. Estimates of abundance were calculated for Arctic grayling  $\geq 270$  and  $\geq 330$  mm FL, as well as for  $\geq 240$  mm FL. Estimates of abundance were expected to be within 25% of the actual abundance 95% of the time, but both objective estimates were slightly less precise having a precision of 27%. The abundance estimate for  $\geq 300$  mm FL stratum did have a precision of 22%. The estimate for 240–269 mm FL stratum was very imprecise (79%) lowering the precision of the abundance estimate for  $\geq 240$  mm stratum to 40%. Overall, the estimates were satisfactorily precise. However, given the density of Arctic grayling present in the study area, substantially more effort would be needed to have produced more precise estimates, particularly for fish of smaller sizes (i.e.  $\leq 270$  mm FL).

Because this was the first abundance estimate of Arctic grayling in the Delta River, there are no previous estimates with which to compare. However, comparing to the numerous other Arctic grayling abundance estimates throughout the state during the previous 50 years of statehood, it is clear that this population is uniquely abundant as only two other estimates had similarly high densities. Roguski and Tack (1970) estimated 36,985 Arctic grayling  $\geq 145$  mm FL in the riverine section between Long Tangle and Lower Tangle lakes, but the veracity of the estimate is questionable because the presentation in the report lacks essential information to validate the estimate. Additionally, it is unclear if a viable estimate is possible because Ridder (1992), after four years of study, concluded that estimating abundance in the large Tangle Lakes system was extremely difficult due to movements of Arctic grayling as well as the logistical difficulties of executing appropriate sampling effort.

The only defensible estimate with similar densities was for the Delta Clearwater River (DCR) during 2006 (Wuttig and Gryska 2010). For that estimate, a 22-km study area contained 14,799 (SE = 2,204) Arctic grayling  $\geq 270$  mm FL and 11,311 (SE = 1,513) Arctic grayling  $\geq 330$  mm FL. Both the Delta River and DCR are short (terminating at a glacial river or creek), clear, cool, exhibit stable flows and are uniquely productive. The spring-fed DCR is nutrient rich supporting the highest production levels of benthic algae and invertebrate drift documented in Interior Alaska (LaPerriere 1994). The ecology of the Delta River has not been studied, but it is suspected that the long series of lakes upstream buffers turbid flows and supplements production downstream, for example, via transport of nutrients, organic matter, and invertebrates. The hydrological and ecological conditions of both rivers appear to be conducive to supporting high densities of Arctic grayling, which makes sense given that Arctic grayling are physiologically adapted to cold water and are visual predators, primarily consuming invertebrates (Northcote 1995).

For management purposes, there is no concern for the Arctic grayling population because the estimated abundance was exceptionally large relative to angler effort and harvest.

## **ACKNOWLEDGEMENTS**

The author thanks Brian Collyard, Rick Queen, Tim Viavant, David Lorrington, John Burr, Kyle Schuman, and James Riedman for their assistance in sampling and logistical support. Thanks to Don Roach, Matt Evenson, and Klaus Wuttig for their supervisory support, Rachael Kvapil for the editing and formatting of this report for publication, and Anton Antonovich for his biometric review. This project and report were made possible by partial funding provided by the U.S. Fish and Wildlife Service through the Federal Aid in Sport Fish Restoration Act (16 U.S.C. 777-777K) under project F-10-24, Job No. R-3-2(e) and in-kind funding through the U.S. Bureau of Land Management in association with Agreement No. BLM L09AC15726.

## REFERENCES CITED

- Bailey, N. T. J. 1951. On estimating the size of mobile populations from capture-recapture data. *Biometrika* 38:293-306.
- Bailey, N. T. J. 1952. Improvements in the interpretation of recapture data. *Journal of Animal Ecology* 21:120-127.
- Baker, T. T. 1989. Stock assessment of Arctic grayling in the Tangle Lakes and River system. Alaska Department of Fish and Game, Division of Sport Fish, Fishery Data Series No. 92, Anchorage, Alaska. 54 pp.
- Carlton, M. 1976. Delta River – Tangle Lakes fisheries inventory: 1976. eds. R. S. Kavanagh and J. W. Bosworth. Western Interstate Commission for Higher Learning. Boulder, CO. 39pp.
- Chapman, D. G. 1951. Some properties for the hypergeometric distribution with applications to zoological censuses. *Univ. Calif. Public. Stat.* 1, 131-160.
- Cochran, W. G. 1977. *Sampling Techniques*, 3<sup>rd</sup> edition. John Wiley. New York.
- Conover, W. J. 1980. *Practical nonparametric statistics*, 2nd edition. John Wiley. New York.
- Darroch, J. N. 1961. The two-sample capture-recapture census when tagging and sampling are stratified. *Biometrika* 48:241-260.
- DeCicco, A. L. 1997. Assessment of Arctic grayling in selected streams of the Seward Peninsula, 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-15, Anchorage.
- Goodman, L. A. 1960. On the exact variance of products. *Journal of the American Statistical Association* 55:708-713.
- Gryska, A. D. 2001. Abundance and length and age composition of Arctic grayling in the Delta Clearwater River, 2000. Alaska Department of Fish and Game, Fishery Data Series No. 01-27, Anchorage.
- Gryska, A. D. 2004a. Abundance and length and age composition of Arctic grayling in the Richardson Clearwater River, 2001. Alaska Department of Fish and Game, Fishery Data Series No. 04-03, Anchorage.
- Gryska, A. D. 2004b. Abundance and length and age composition of Arctic grayling in the Snake River, 2001. Alaska Department of Fish and Game, Fishery Data Series No. 04-15, Anchorage.
- Holmes, R., D. N. McBride, T. Viavant, J. B. Reynolds. 1990. Electrofishing induced mortality and injury to rainbow trout, Arctic grayling, humpback whitefish, least cisco, and northern pike. Alaska Department of Fish and Game, Division of Sport Fish. Fishery Manuscript No. 90-3, Anchorage, Alaska.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001a. Participation, catch, and harvest in Alaska sport fisheries during 1999. Alaska Department of Fish and Game, Fishery Data Series No. 01-8, Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001b. Revised Edition: Harvest, catch, and participation in Alaska sport fisheries during 1996. Alaska Department of Fish and Game, Fishery Data Series No. 97-29 (revised), Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001c. Revised Edition: Harvest, catch, and participation in Alaska sport fisheries during 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-25 (revised), Anchorage.
- Howe, A. L., R. J. Walker, C. Olness, K. Sundet, and A. E. Bingham. 2001d. Revised Edition: Participation, catch, and harvest in Alaska sport fisheries during 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-41 (revised), Anchorage.
- Jennings, G. B., K. Sundet, A. E. Bingham, and H. K. Sigurdsson. 2006. Participation, catch, and harvest in Alaska sport fisheries during 2002. Alaska Department of Fish and Game, Fishery Data Series No. 06-34, Anchorage.
- Jennings, G. B., K. Sundet, and A. E. Bingham. 2009a. Estimates of participation, catch, and harvest in Alaska sport fisheries during 2005. Alaska Department of Fish and Game, Fishery Data Series No. 09-47, Anchorage.

## REFERENCES CITED (Continued)

- Jennings, G. B., K. Sundet, and A. E. Bingham. 2009b. Participation, catch, and harvest in Alaska sport fisheries during 2006. Alaska Department of Fish and Game, Fishery Data Series No. 09-54, Anchorage.
- Laperriere, J. D. 1994. Benthic ecology of a spring-fed river of interior Alaska. *Freshwater Biology* 32 (2): 349–357.
- Northcote, T. G. 1995. Comparative biology and management of Arctic and European grayling (Salmonidae, Thymallus). *Reviews in Fish Biology and Fisheries* 5:141-194.
- Peckham, R. D. 1974. Evaluation of interior Alaska waters and Sportfish with emphasis on stocked lakes. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Performance Report, 1973-1974, Project F-9-6, 15(G-III-E):1-34.
- Peckham, R. D. 1975. Evaluation of interior Alaska waters and Sportfish with emphasis on stocked lakes. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Performance Report, 1974-1975, Project F-9-7, 16(G-III-E):52-77.
- Peckham, R. D. 1976. Inventory and cataloging of the sport fish and sport fish waters in the interior Alaska, Delta River drainage. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Performance Report, 1975-1976, Project F-9-8, 17(G-I-M):69-116.
- Ridder, W. P. 1992. Abundance, composition, and exploitation of selected Arctic grayling spawning stocks in the Tangle Lakes system, 1991. Alaska Department of Fish and Game, Fishery Data Series No. 92-6, Anchorage.
- Ridder, W. P. and A. D. Gryska. 2000. Abundance and composition of Arctic grayling in the Delta Clearwater River, 1999. Alaska Department of Fish and Game, Fishery Data Series No. 00-38, Anchorage.
- Robson, D. S., and H. A. Regier. 1964. Sample size in Petersen mark-recapture experiments. *Transactions of the American Fisheries Society* 93:215-226.
- Roguski, E. A. and S. L. Tack. 1970. Investigation of the Tanana River and Tangle Lakes fisheries: migratory and population study. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual Report of Progress, 1969-1970, Project F-9-2, 11(16B).
- Seber, G. A. F. 1982. The estimation of animal abundance and related parameters, second edition. Charles Griffin and Co., Ltd. London, U.K.
- Swanton, C. O., and K. G. Wuttig. *In prep.* A management plan and policies for Arctic grayling fisheries within the Arctic-Yukon-Kuskokwim region. Alaska Department of Fish and Game, Sport Fish Division, 1300 College Road, Fairbanks, AK.
- Thompson, S. K. 1987. Sample size for estimating multinomial proportions. *American Statistician* 41(1):122-132.
- Walker, R. J., C. Olnes, K. Sundet, A. L. Howe, and A. E. Bingham. 2003. Participation, catch, and harvest in Alaska sport fisheries during 2000. Alaska Department of Fish and Game, Fishery Data Series No. 03-05, Anchorage.
- Wuttig, K. G. 2004. Stock assessment of Arctic grayling in the headwaters of the Chatanika and Chena rivers during 2002. Alaska Department of Fish and Game, Fishery Data Series No. 04-22, Anchorage.
- Wuttig, K., and A. D. Gryska. 2010. Abundance and length composition of Arctic grayling in the Delta Clearwater River, 2006. Alaska Department of Fish and Game, Fishery Data Series No. 10-84, Anchorage.

**APPENDIX A**  
**EQUATIONS AND STATISTICAL METHODOLOGY**

Appendix A1.—Equations for calculating estimates of abundance and its variance using the Bailey-modified Petersen estimator.

---

The Bailey-modified Petersen estimator (Bailey 1951 and 1952) was used because the sampling design called for a systematic downstream progression, fishing each pool and run and attempting to subject all fish to the same probability of capture while sampling with replacement. The Bailey modification to the Petersen estimator may be used even when the assumption of a random sample for the second sample is false when a systematic sample is provided:

- 1) there is uniform mixing of marked and unmarked fish; and,
- 2) all fish, whether marked or unmarked, have the same probability of capture (Seber 1982).

The abundance of Arctic grayling was estimated as:

$$\hat{N} = \frac{n_1(n_2 + 1)}{m_2 + 1}, \quad (\text{A1-1})$$

where:

$n_1$  = the number of Arctic grayling marked and released alive during the first event;

$n_2$  = the number of Arctic grayling examined for marks during the second event; and,

$m_2$  = the number of Arctic grayling marked in the first event that were recaptured during the second event;  
and

The variance was estimated as (Seber 1982):

$$\hat{V}[\hat{N}] = \frac{n_1^2(n_2 + 1)(n_2 - m_2)}{(m_2 + 1)^2(m_2 + 2)}. \quad (\text{A1-2})$$

Appendix A2.—Detection of size and/or sex selective sampling during a two-sample mark recapture experiment and its effects on estimation of population size and population composition.

Size selective sampling: The Kolmogorov-Smirnov two sample test (Conover 1980) is used to detect significant evidence that size selective sampling occurred during the first and/or second sampling events. The second sampling event is evaluated by comparing the length frequency distribution of all fish marked during the first event (M) with that of marked fish recaptured during the second event (R) by using the null test hypothesis of no difference. The first sampling event is evaluated by comparing the length frequency distribution of all fish inspected for marks during the second event (C) with that of R. A third test that compares M and C is then conducted and used to evaluate the results of the first two tests when sample sizes are small. Guidelines for small sample sizes are <30 for R and <100 for M or C.

Sex selective sampling: Contingency table analysis (Chi<sup>2</sup>-test) is generally used to detect significant evidence that sex selective sampling occurred during the first and/or second sampling events. The counts of observed males to females are compared between M&R, C&R, and M&C using the null hypothesis that the probability that a sampled fish is male or female is independent of sample. If the proportions by gender are estimated for a sample (usually C), rather than observed for all fish in the sample, contingency table analysis is not appropriate and the proportions of females (or males) are then compared between samples using a two sample test (e.g. Student's t-test).

<b>M vs. R</b>	<b>C vs. R</b>	<b>M vs. C</b>
----------------	----------------	----------------

*Case I:*

Fail to reject H <sub>0</sub>	Fail to reject H <sub>0</sub>	Fail to reject H <sub>0</sub>
-------------------------------	-------------------------------	-------------------------------

There is no size/sex selectivity detected during either sampling event.

*Case II:*

Reject H <sub>0</sub>	Fail to reject H <sub>0</sub>	Reject H <sub>0</sub>
-----------------------	-------------------------------	-----------------------

There is no size/sex selectivity detected during the first event but there is during the second event sampling.

*Case III:*

Fail to reject H <sub>0</sub>	Reject H <sub>0</sub>	Reject H <sub>0</sub>
-------------------------------	-----------------------	-----------------------

There is no size/sex selectivity detected during the second event but there is during the first event sampling.

*Case IV:*

Reject H <sub>0</sub>	Reject H <sub>0</sub>	Either result possible
-----------------------	-----------------------	------------------------

There is size/sex selectivity detected during both the first and second sampling events.

*Evaluation Required:*

Fail to reject H <sub>0</sub>	Fail to reject H <sub>0</sub>	Reject H <sub>0</sub>
-------------------------------	-------------------------------	-----------------------

Sample sizes and powers of tests must be considered:

A. If sample sizes for M vs. R and C vs. R tests are not small and sample sizes for M vs. C test are very large, the M vs. C test is likely detecting small differences which have little potential to result in bias during estimation. *Case I* is appropriate.

B. If a) sample sizes for M vs. R are small, b) the M vs. R p-value is not large (~0.20 or less), and c) the C vs. R sample sizes are not small and/or the C vs. R p-value is fairly large (~0.30 or more), the rejection of the null in the M vs. C test was likely the result of size/sex selectivity during the second event which the M vs. R test was not powerful enough to detect. *Case I* may be considered but *Case II* is the recommended, conservative interpretation.

-continued-

C. If a) sample sizes for C vs. R are small, b) the C vs. R p-value is not large (~0.20 or less), and c) the M vs. R sample sizes are not small and/or the M vs. R p-value is fairly large (~0.30 or more), the rejection of the null in the M vs. C test was likely the result of size/sex selectivity during the first event which the C vs. R test was not powerful enough to detect. *Case I* may be considered but *Case III* is the recommended, conservative interpretation.

D. If a) sample sizes for C vs. R and M vs. R are both small, and b) both the C vs. R and M vs. R p-values are not large (~0.20 or less), the rejection of the null in the M vs. C test may be the result of size/sex selectivity during both events which the C vs. R and M vs. R tests were not powerful enough to detect. *Cases I, II, or III* may be considered but *Case IV* is the recommended, conservative interpretation.

*Case I.* Abundance is calculated using a Petersen-type model from the entire data set without stratification. Composition parameters may be estimated after pooling length, sex, and age data from both sampling events.

*Case II.* Abundance is calculated using a Petersen-type model from the entire data set without stratification. Composition parameters may be estimated using length, sex, and age data from the first sampling event without stratification. If composition is estimated from second event data or after pooling both sampling events, data must first be stratified to eliminate variability in capture probability (detected by the M vs. R test) within strata. Composition parameters are estimated within strata, and abundance for each stratum needs to be estimated using a Petersen-type formula. Overall composition parameters are estimated by combining stratum estimates weighted by estimated stratum abundance according to the formulae below.

*Case III.* Abundance is calculated using a Petersen-type model from the entire data set without stratification. Composition parameters may be estimated using length, sex, and age data from the second sampling event without stratification. If composition is estimated from first event data or after pooling both sampling events, data must first be stratified to eliminate variability in capture probability (detected by the C vs. R test) within strata. Composition parameters are estimated within strata, and abundance for each stratum needs to be estimated using a Petersen-type formula. Overall composition parameters are estimated by combining stratum estimates weighted by estimated stratum abundance according to the formulae below.

*Case IV.* Data must be stratified to eliminate variability in capture probability within strata for at least one or both sampling events. Abundance is calculated using a Petersen-type model for each stratum, and estimates are summed across strata to estimate overall abundance. Composition parameters may be estimated within the strata as determined above, but only using data from sampling events where stratification has eliminated variability in capture probabilities within strata. If data from both sampling events are to be used, further stratification may be necessary to meet the condition of capture homogeneity within strata for both events. Overall composition parameters are estimated by combining stratum estimates weighted by estimated stratum abundance.

If stratification by sex or length is necessary prior to estimating composition parameters, then an overall composition parameters ( $p_k$ ) is estimated by combining within stratum composition estimates using:

$$\hat{p}_k = \sum_{i=1}^j \frac{\hat{N}_i}{\hat{N}_\Sigma} \hat{p}_{ik}; \text{ and,} \quad (\text{A2-1})$$

$$\hat{V}[\hat{p}_k] \approx \frac{1}{\hat{N}_\Sigma^2} \sum_{i=1}^j \left( \hat{N}_i^2 \hat{V}[\hat{p}_{ik}] + (\hat{p}_{ik} - \hat{p}_k)^2 \hat{V}[\hat{N}_i] \right). \quad (\text{A2-2})$$

where:

- $j$  = the number of sex/size strata;
- $\hat{p}_{ik}$  = the estimated proportion of fish that were age or size  $k$  among fish in stratum  $i$ ;
- $\hat{N}_i$  = the estimated abundance in stratum  $i$ ; and,
- $\hat{N}_\Sigma$  = sum of the  $\hat{N}_i$  across strata.

---

## Tests of consistency for Petersen Estimator

Of the following conditions, at least one must be fulfilled to meet assumptions of a Petersen estimator:

1. Marked fish mix completely with unmarked fish between events;
2. Every fish has an equal probability of being captured and marked during event 1; or,
3. Every fish has an equal probability of being captured and examined during event 2.

To evaluate these three assumptions, the chi-square statistic will be used to examine the following contingency tables as recommended by Seber (1982). At least one null hypothesis needs to be accepted for assumptions of the Petersen model (Bailey 1951, 1952; Chapman 1951) to be valid. If all three tests are rejected, a geographically stratified estimator (Darroch 1961) should be used to estimate abundance.

### I.-Test for complete mixing<sup>a</sup>

Section Where Marked	Section Where Recaptured				Not Recaptured (n <sub>1</sub> -m <sub>2</sub> )
	A	B	...	F	
A					
B					
...					
F					

### II.-Test for equal probability of capture during the first event<sup>b</sup>

	Section Where Examined			
	A	B	...	F
Marked (m <sub>2</sub> )				
Unmarked (n <sub>2</sub> -m <sub>2</sub> )				

### III.-Test for equal probability of capture during the second event<sup>c</sup>

	Section Where Marked			
	A	B	...	F
Recaptured (m <sub>2</sub> )				
Not Recaptured (n <sub>1</sub> -m <sub>2</sub> )				

<sup>a</sup> This tests the hypothesis that movement probabilities ( $\theta$ ) from section  $i$  ( $i = 1, 2, \dots, s$ ) to section  $j$  ( $j = 1, 2, \dots, t$ ) are the same among sections:  $H_0: \theta_{ij} = \theta_j$ .

<sup>b</sup> This tests the hypothesis of homogeneity on the columns of the 2-by-t contingency table with respect to the marked to unmarked ratio among sections:  $H_0: \sum_i a_i \theta_{ij} = k U_j$ , where  $k$  = total marks released/total unmarked in the population,  $U_j$  = total unmarked fish in stratum  $j$  at the time of sampling, and  $a_i$  = number of marked fish released in stratum  $i$ .

<sup>c</sup> This tests the hypothesis of homogeneity on the columns of this 2-by-s contingency table with respect to recapture probabilities among sections:  $H_0: \sum_j \theta_{ij} p_j = d$ , where  $p_j$  is the probability of capturing a fish in section  $j$  during the second event, and  $d$  is a constant.

Appendix A4.–Equations for estimating length composition and their variances for the population.

For Case I-III scenarios (Appendix A2), the proportions of Arctic grayling within each age or length class  $k$  were estimated:

$$\hat{p}_k = \frac{n_k}{n} \quad (\text{A4-1})$$

where:

$n_k$  = the number of Arctic grayling sampled within age or length class  $k$  and,

$n$  = the total number of Arctic grayling sampled.

When calculating  $n$  and  $n_k$  the diagnostic test results were used to determine which fish were included (Appendix A2). For Case I, fish from both capture events are used.

The variance of each proportion was estimated as (from Cochran 1977):

$$\hat{V}[\hat{p}_k] = \frac{\hat{p}_k(1-\hat{p}_k)}{n-1}. \quad (\text{A4-2})$$

The abundance of Arctic grayling in each length or age category,  $k$ , in the population was then estimated:

$$\hat{N}_k = \hat{p}_k \hat{N}, \quad (\text{A4-3})$$

where:

$\hat{N}$  = the estimated overall abundance (Appendix A1).

The variance for  $\hat{N}_k$  was then estimated using the formulation for the exact variance of the product of two independent random variables (Goodman 1960):

$$\hat{V}[\hat{N}_k] = \hat{V}[\hat{p}_k] \hat{N}^2 + \hat{V}[\hat{N}] \hat{p}_k^2 - \hat{V}[\hat{p}_k] \hat{V}[\hat{N}]. \quad (\text{A4-4})$$

For the Case IV scenario (Appendix A2), that requiring stratification by size or sex, the proportions of Arctic grayling within each age or length class  $k$  were estimated by first calculating:

$$\hat{p}_{jk} = \frac{n_{jk}}{n_j} \quad (\text{A4-5})$$

where:

$n_j$  = the number sampled from size stratum  $j$  in the mark-recapture experiment;

$n_{jk}$  = the number sampled from size stratum  $j$  that are in length or age category  $k$ ; and,

$\hat{p}_{jk}$  = the estimated proportion of length or age category  $k$  fish in size stratum  $j$ .

When calculating  $n_j$  and  $n_{jk}$  the within stratum diagnostic test results were used to determine which fish were included in the analysis following the rules for  $n$  and  $n_k$  provided above.

The variance calculation for  $\hat{p}_{jk}$  is equation 2 substituting  $\hat{p}_{jk}$  for  $\hat{p}_k$  and  $n_j$  for  $n$ .

The estimated abundance of fish in length or age category  $k$  in the population is then:

-continued-

$$\hat{N}_k = \sum_{j=1}^s \hat{p}_{jk} \hat{N}_j \quad (\text{A4-6})$$

where:

$\hat{N}_j$  = the estimated abundance in size stratum  $j$ ; and,

$s$  = the number of size strata.

The variance for  $\hat{N}_k$  will be estimated using the formulation for the exact variance of the product of two independent random variables (Goodman 1960):

$$\hat{V}[\hat{N}_k] = \sum_{j=1}^s \left( \hat{V}[\hat{p}_{jk}] \hat{N}_j^2 + \hat{V}[\hat{N}_j] \hat{p}_{jk}^2 - \hat{V}[\hat{p}_{jk}] \hat{V}[\hat{N}_j] \right). \quad (\text{A4-7})$$

The estimated proportion of the population in length or age category  $k$  ( $\hat{p}_k$ ) is then:

$$\hat{p}_k = \hat{N}_k / \hat{N} \quad (\text{A4-8})$$

where:  $\hat{N} = \sum_{j=1}^s \hat{N}_j$ .

Variance of the estimated proportion can be approximated with the delta method (Seber 1982):

$$\hat{V}[\hat{p}_k] \approx \sum_{j=1}^s \left\{ \left( \frac{\hat{N}_j}{\hat{N}} \right)^2 \hat{V}[\hat{p}_{jk}] \right\} + \frac{\sum_{j=1}^s \left\{ \hat{V}[\hat{N}_j] (\hat{p}_{jk} - \hat{p}_k)^2 \right\}}{\hat{N}^2}. \quad (\text{A4-9})$$



**APPENDIX B**  
**DATA FILE LISTING**

Appendix B1.–Data files for population estimate of Arctic grayling captured in the Delta River, 2008.

---

File Name <sup>a</sup>
Delta River Arctic grayling population estimate data files for archive-2008.xls

---

<sup>a</sup> Data files are archived at and are available from the Alaska Department of Fish and Game, Sport Fish Division, 1300 College Road, Fairbanks, Alaska 99701-1599.