

Fishery Management Report No. 05-31

**2005 Yukon Area Subsistence, Personal Use, and
Commercial Salmon Fisheries Outlook and
Management Strategies**

by

Fred J. Bue

and

Tracy L. Lingnau

May 2005

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Measures (fisheries)	
centimeter	cm	Alaska Administrative		fork length	FL
deciliter	dL	Code	AAC	mid-eye-to-fork	MEF
gram	g	all commonly accepted		mid-eye-to-tail-fork	METF
hectare	ha	abbreviations	e.g., Mr., Mrs., AM, PM, etc.	standard length	SL
kilogram	kg			total length	TL
kilometer	km	all commonly accepted			
liter	L	professional titles	e.g., Dr., Ph.D., R.N., etc.		
meter	m	at	@	Mathematics, statistics	
milliliter	mL	compass directions:		<i>all standard mathematical</i>	
millimeter	mm	east	E	<i>signs, symbols and</i>	
		north	N	<i>abbreviations</i>	
		south	S	alternate hypothesis	H _A
		west	W	base of natural logarithm	<i>e</i>
		copyright	©	catch per unit effort	CPUE
		corporate suffixes:		coefficient of variation	CV
		Company	Co.	common test statistics	(F, t, χ^2 , etc.)
		Corporation	Corp.	confidence interval	CI
		Incorporated	Inc.	correlation coefficient	
		Limited	Ltd.	(multiple)	R
		District of Columbia	D.C.	correlation coefficient	
		et alii (and others)	et al.	(simple)	r
		et cetera (and so forth)	etc.	covariance	cov
		exempli gratia		degree (angular)	°
		(for example)	e.g.	degrees of freedom	df
		Federal Information		expected value	<i>E</i>
		Code	FIC	greater than	>
		id est (that is)	i.e.	greater than or equal to	≥
		latitude or longitude	lat. or long.	harvest per unit effort	HPUE
		monetary symbols		less than	<
		(U.S.)	\$, ¢	less than or equal to	≤
		months (tables and		logarithm (natural)	ln
		figures): first three		logarithm (base 10)	log
		letters	Jan, ..., Dec	logarithm (specify base)	log ₂ , etc.
		registered trademark	®	minute (angular)	'
		trademark	™	not significant	NS
		United States		null hypothesis	H ₀
		(adjective)	U.S.	percent	%
		United States of		probability	P
		America (noun)	USA	probability of a type I error	
		U.S.C.	United States	(rejection of the null	
			Code	hypothesis when true)	α
			use two-letter	probability of a type II error	
			abbreviations	(acceptance of the null	
			(e.g., AK, WA)	hypothesis when false)	β
				second (angular)	"
				standard deviation	SD
				standard error	SE
				variance	
				population	Var
				sample	var

Weights and measures (English)

cubic feet per second	ft ³ /s
foot	ft
gallon	gal
inch	in
mile	mi
nautical mile	nmi
ounce	oz
pound	lb
quart	qt
yard	yd

Time and temperature

day	d
degrees Celsius	°C
degrees Fahrenheit	°F
degrees kelvin	K
hour	h
minute	min
second	s

Physics and chemistry

all atomic symbols	
alternating current	AC
ampere	A
calorie	cal
direct current	DC
hertz	Hz
horsepower	hp
hydrogen ion activity	pH
(negative log of)	
parts per million	ppm
parts per thousand	ppt, ‰
volts	V
watts	W

FISHERY MANAGEMENT REPORT NO. 05-31

**2005 YUKON AREA SUBSISTENCE, PERSONAL USE, AND
COMMERCIAL SALMON FISHERIES OUTLOOK AND MANAGEMENT
STRATEGIES**

by

Fred J. Bue,
Division of Commercial Fisheries, Fairbanks
and
Tracy L. Lingnau
Division of Commercial Fisheries, Anchorage

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1599

May 2005

The Division of Sport Fish Fishery Management Reports series was established in 1989 for the publication of an overview of Division of Sport Fish management activities and goals in a specific geographic area. Since 2004, the Division of Commercial Fisheries has also used the Fishery Management Report series. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm>. This publication has undergone regional peer review.

*Fred J. Bue,
Alaska Department of Fish and Game, Division of Commercial Fisheries,
1300 College Road, Fairbanks, AK 99701-1599, USA*

and

*Tracy L. Lingnau
Alaska Department of Fish and Game, Division of Commercial Fisheries,
333 Raspberry Road, Anchorage, AK 99518-1599, USA*

This document should be cited as:

Bue, F. J., and T. L. Lingnau. 2005. 2005 Yukon Area subsistence, personal use, and commercial salmon fisheries outlook and management strategies. Alaska Department of Fish and Game, Fishery Management Report No. 05-31, Anchorage.

The Alaska Department of Fish and Game administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility, or if you desire further information please write to ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; U.S. Fish and Wildlife Service, 4040 N. Fairfax Drive, Suite 300 Webb, Arlington, VA 22203 or O.E.O., U.S. Department of the Interior, Washington DC 20240.

For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-6077, (TDD) 907-465-3646, or (FAX) 907-465-6078.

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	ii
LIST OF FIGURES.....	ii
PREFACE.....	iii
ABSTRACT.....	1
1.0 INTRODUCTION.....	1
2.0 OUTLOOK FOR 2005.....	1
2.1 Chinook Salmon.....	1
2.2 Summer Chum Salmon.....	2
2.3 Fall Chum Salmon.....	2
2.4 Coho Salmon.....	4
3.0 MANAGEMENT STRATEGY FOR 2005.....	4
3.1 Alaska Board Of Fisheries Actions.....	6
3.2 Subsistence Fishery.....	6
3.2.1 Districts 1, 2, and 3.....	8
3.2.2 District 4.....	8
3.2.3 District 5.....	8
3.2.4 District 6.....	9
3.3 Personal Use Fishery.....	9
3.4 Commercial Fishery and Reporting Requirements.....	10
3.5 Chinook and Summer Chum Salmon Commercial Season.....	10
3.5.1 Districts 1, 2, and 3.....	11
3.5.2 District 4.....	12
3.5.3 Anvik River Management Area.....	12
3.5.4 District 5.....	12
3.5.5 District 6.....	12
3.6 Fall Chum and Coho Salmon Commercial Season.....	13
3.6.1 Districts 1, 2, and 3.....	14
3.6.2 District 4.....	14
3.6.3 Subdistricts 5-B, 5-C, and 5-D.....	14
3.6.4 Subdistrict 5-A and District 6.....	14
3.7 Guideline Harvest Ranges.....	15
4.0 U.S./CANADA YUKON RIVER SALMON PANEL AGREEMENT.....	15
REFERENCES CITED.....	17
FIGURES.....	19

LIST OF TABLES

Table		Page
1.	Proportions of expected fall chum salmon runs.	3
2.	Approximate age composition of fall chum salmon expected in 2005.....	3
3.	Escapement goals for Chinook salmon.	5
4.	Escapement goals for summer chum salmon.	5
5.	Escapement goals for fall chum salmon.	6
6.	Yukon Area subsistence salmon fishing schedule.....	7
7.	Summary of the Summer Chum Salmon Management Plan.	11
8.	Summary of the Fall Chum Salmon Management Plan.	13
9.	Guideline harvest ranges and midpoints for commercial harvest of Yukon River Chinook, summer, and fall chum salmon.	16

LIST OF FIGURES

Figure		Page
1.	Yukon Area communities and fishing districts.	20
2.	Yukon River Chinook salmon subsistence and commercial harvests compared to the recent 10-year average, and the lower ANS range.	21
3.	Yukon River summer chum salmon subsistence and commercial harvests compared to the recent 10-year average, and the lower ANS range.	22
4.	Subsistence and commercial harvest of fall chum salmon, 1961 to present.....	23
5.	Subsistence and commercial harvest of coho salmon, 1961 to present.	24
6.	Yukon River fall chum salmon management plan overview and comparison with historical run sizes.....	25

PREFACE

The following important information is for fishers participating in subsistence, personal use and commercial fisheries in the Yukon Area during the 2005 season. Fishers may contact ADF&G, Division of Commercial Fisheries staff at the office locations listed below.

Emmonak Seasonal Field Office

(June through August)

Alaska Department of Fish and Game

Division of Commercial Fisheries

P.O. Box 127

Emmonak, Alaska 99581

Telephone: (907) 949-1320

Fax: (907) 949-1830

Recorded Information: (907) 949-1731

Fairbanks Office

Alaska Department of Fish and Game

Division of Commercial Fisheries

1300 College Road

Fairbanks, Alaska 99701

Telephone: (907) 459-7274

Fax: (907) 452-1668

Anchorage Office

Alaska Department of Fish and Game

Division of Commercial Fisheries

333 Raspberry Road

Anchorage, Alaska 99518

Telephone: (907) 267-2109

Fax: (907) 267-2442

For a recording of the current fishing schedule call toll free 1-866-479-7387 or in the Fairbanks area call (907) 459-7387.

For Subsistence and Personal Use Permit Reporting call (907) 459-7388.

Notice to Fishermen

Waters subject to ANILCA Title VIII (including waters in which the United States has identified a reserved water right) or “federal subsistence jurisdiction”. Subject to federal restrictions and closures, waters subject to ANILCA Title VIII are open to fishing under state regulations. If you are subsistence fishing in waters under federal subsistence jurisdiction, you must comply with federal subsistence regulations and some state permit conditions or state regulations may be pre-empted.

To familiarize yourself with the federal regulations you should consult the *Subsistence Management Regulations for the Harvest of Fish and Shellfish on Federal Public Lands and Waters in Alaska* for details. Copies may be obtained at federal offices. Calling the federal agencies is also recommended as inseason closures or temporary regulatory changes can occur at anytime and may not be reflected in the annual regulatory publication.

For more information, or a copy of federal regulations, please contact:

U. S. Fish and Wildlife Service, Office of Subsistence Management:1-800-801-5108

Or any of the following agencies:

National Park Service 907-257-2649

U. S. Fish and Wildlife Service1-800-478-1456

Bureau of Land Management..... 907-271-5960

National Forests: U.S. Forest Service 907-586-8806

ABSTRACT

This management plan provides an overview of the expected salmon outlooks, management issues, and harvest strategies for Yukon River summer and fall salmon fisheries in 2005. Subsistence, personal use, and commercial fisheries occur throughout the Yukon Area. The Yukon Area includes all waters of the Yukon River drainage in Alaska and all coastal waters of Alaska from Point Romanof south to Naskonat Peninsula. These outlooks and management strategies are for salmon runs expected in 2005. Management strategies may change inseason based upon assessment of salmon runs.

Key words: Yukon, Chinook, summer chum, fall chum, coho, outlook, strategy, management plan

1.0 INTRODUCTION

This document provides the 2005 outlook for Yukon Area salmon runs, as well as management strategies for the subsistence, personal use, and commercial salmon fisheries. Subsistence fishing in portions of the Yukon Area is under dual management authority of the Alaska Department of Fish & Game (ADF&G) and the U.S. Fish and Wildlife Service (USFWS). Fishermen are reminded that they should consult both the State of Alaska fishing regulations and the Federal Subsistence Management Regulations for Federal Public Lands before fishing in the Yukon Area.

The Yukon Area includes all waters of Alaska within the Yukon River drainage and coastal waters from Point Romanof, northeast of Kotlik, to the Naskonat Peninsula. For management purposes the Yukon Area is divided into 7 districts and 10 subdistricts (Figure 1). Commercial fishing may be allowed along the entire 1,224 miles of the Yukon River in Alaska and along the lower 225 miles of the Tanana River. The Coastal District includes the majority of coastal marine waters within the Yukon Area and is only open to subsistence fishing. The Lower Yukon Area (Districts 1, 2, and 3) includes coastal waters of the Yukon River delta and that portion of the Yukon River drainage downstream of Old Paradise Village (river mile 301). The Upper Yukon Area (Districts 4, 5, and 6) is the Alaskan portion of the Yukon River drainage upstream of Old Paradise Village.

Chinook *Oncorhynchus tshawytscha*, chum *O. keta*, and coho *O. kisutch* salmon are harvested in Yukon River commercial, subsistence and personal use fisheries. In recent years Chinook salmon have become the primary commercial species. Yukon River chum salmon consists of an earlier and typically more abundant summer chum salmon run and a later fall chum salmon run. No directed commercial fishing has occurred for pink *O. gorbuscha* salmon within the Yukon River. Aboriginal, commercial, domestic, and sport salmon fisheries also occur in Canada, which are managed by the Canadian Department of Fisheries and Oceans (DFO).

2.0 OUTLOOK FOR 2005

2.1 CHINOOK SALMON

Yukon River Chinook salmon return primarily as age-5 and age-6 fish¹, although age-4 and age-7 fish also contribute to the run. Spawning ground escapements in 1999, the brood year producing 6-year-old fish returning in 2005, were above the upper end of the escapement goals in the Chena and Salcha Rivers but below the escapement objective in Canada. The 5-year-old

¹ Salmon ages contained in this document represent the combined freshwater and saltwater age.

component in 2004 was below average and the 4-year-old component in 2004 was above average. The previous 2 years (2003 and 2004) runs have been near average indicating good production from the poor runs of 1998–2000.

Assuming a below normal return of 6-year-old fish, and a stronger return of 5-year-old fish, the 2005 run is expected to be below average and not as strong as the 2003 and 2004 runs. Given the uncertainties associated with recent declines in productivity, it is anticipated the run will provide for escapements, support a normal subsistence harvest, and a below average commercial harvest. Fishery management will be based upon inseason assessments of the run. Similar to 2004, if inseason indicators of run strength suggest sufficient abundance exists to have a commercial fishery, the commercial harvest in Alaska could range from 20,000 to 60,000 Chinook salmon. This commercial harvest range is below the 10-year (1995–2004) average of approximately 63,400 Chinook salmon (Figure 2).

2.2 SUMMER CHUM SALMON

Summer chum salmon runs in 2005 are dependent on escapements from 2001 (age-4 fish) and 2000 (age-5 fish). The 2000 and 2001 summer chum salmon runs were the poorest on record and none of the escapement goals were met either year. Since 2001, the summer chum salmon runs have exhibited steady improvements with harvestable surpluses in the last 3 years (2002–2004). Weak Yukon River summer chum salmon runs from 1998 through 2001 are attributed to reduced productivity and is not the result of low levels of parent year escapements. In 2004, a record number of 3-year-old summer chum salmon were observed throughout the Arctic-Yukon-Kuskokwim (AYK) Region. The BASIS (Bering-Aleutian Salmon International Survey) study and Bering Sea trawl bycatch have both observed significant increases in chum salmon. Although not all of these salmon are bound for Western Alaska, it is an indicator that the health of the Bering Sea has improved considerably and chum salmon production may have increased significantly.

If ocean conditions are more conducive to survival, it is anticipated the run will be below average to average and provide for escapements, support a normal subsistence harvest, and possibly a below average to average commercial harvest. If inseason indicators of abundance suggest a harvestable surplus exists for a commercial fishery, the commercial harvest in Alaska could range from 100,000 to 500,000 summer chum salmon depending on market conditions.

2.3 FALL CHUM SALMON

Yukon River fall chum salmon estimated escapements for the period 1974 through 2000 have ranged from approximately 180,000 (1982) to 1,500,000 (1975), based upon expansion of escapement assessments for selected stocks to approximate overall abundance. Escapements in these years resulted in subsequent returns that ranged in size from approximately 312,000 (1996 production) to 1,400,000 (1975 production) fish, using the same approach to approximate overall escapement. Corresponding return-per-spawner rates range from 0.3 to 3.2, averaging 1.8 for all years combined (1974–1998).

Western Alaska experienced dramatic declines in salmon runs from 1997 through 2002 with a record low in 2000. Weak salmon runs during this time period have been attributed to reduced productivity in the marine environment and is not the result of low levels of parent year escapements. Beginning in 1999, the fall chum salmon run projections have been presented as a range that includes the normal point projection as the high end in response to previous run

failures. The low end was determined by reducing the normal point projection by the average ratio of observed to predicted returns from 1998 through the current year. Production has since improved based on the 2003 and 2004 runs. Therefore the low end for 2005 was determined based on a 4-year (one chum salmon life cycle) average ratio of observed to predicted returns from 2001 to 2004. The proportions of the expected runs are shown in Table 1.

Table 1.—Proportions of expected fall chum salmon runs.

Year	Expected Run Size	Estimated Run Size	Proportion of Expected Run
1998	880,000	329,000	0.37
1999	1,197,000	424,000	0.35
2000	1,137,000	241,000	0.21
2001	962,000	383,000	0.40
2002	646,000	414,000	0.65
2003	647,000	750,000	1.16
2004	672,000	543,000	0.81
All Year Average 98–04			0.57
4-Year Average 01–04			0.75

Yukon River fall chum salmon return primarily as age-4 or age-5 fish, although age-3 and age-6 fish also contribute to the run. The 2005 run will be comprised of the parent years 1999 to 2002. Estimates of return per spawner based on brood year return were used to estimate production for 1999 and 2000 and an auto-regressive Ricker spawner-recruit model was used to predict the returns from 2001 and 2002. The point estimate utilizes the 1974 to 1983 even/odd maturity schedules to represent years of higher production. Using the 4-year average proportion of 0.75 results in tightening up the lower end of the range and together result in a 2005 run size projection in the range of 584,000 to 776,000 fall chum salmon, with the following approximate age composition (Table 2).

Table 2.—Approximate age composition of fall chum salmon expected in 2005.

Brood Year	Escapement	Estimated Production (R/S)^a	Estimated Production	Contribution based on age	2005 Return
1999	292,185	2.96	864,868	0.4%	2,720
2000	212,376	1.85	392,896	11.3%	87,940
2001	337,559	2.40	810,142	80.9%	628,230
2002	392,920	2.11	829,061	7.4%	57,518
Total expected run (unadjusted)					776,409
Total expressed as a range using 2001 to 2004 forecasted vs. observed return (75%):					584,000 to 776,000

^a R/S = return per spawner.

The escapements from the 1999 and 2000 parent years that will contribute to the 2005 run were extremely poor and below the minimum drainage-wide escapement goal of 300,000 fall chum salmon. The major contributor to the 2005 fall chum salmon run is anticipated to be age-4 fish returning from the 2001 parent year. Both 2001 and 2002 escapements were within the drainage-wide escapement goal range but in the lower third. This will be the first year of returns from the 2002 brood year. Stocks within the Tanana River drainage may have been affected by a magnitude 7.9 earthquake which occurred on November 3, 2002. The epicenter was located

within the Alaska Range on the Denali fault line and could have affected fall chum and coho salmon eggs incubating in the gravels from the Toklat River in the Kantishna River drainage to the upper Tanana River mainstem including the Delta River area.

In 2004, there was an exceptional return of approximately 120,000 age-3 fish from the 2001 brood year (1974–1998 brood year return average is 29,000 age-3 fish). This component was the second highest on record and may represent further evidence of improved conditions in the marine environment. The return of age-4 fish from odd-numbered brood years during the time period 1974 to 1998 averaged 721,000 chum salmon, and ranged from a low of 245,000 for brood year 1997 to a high of 1,225,000 for brood year 1975. Based on high productive years 1974 to 1983 the return of odd-numbered brood years averaged 1,256,000 chum salmon.

The projection for the 2005 fall chum salmon run is based on evident improvements in production observed in 2003 and 2004 runs. The age-4 component is typically expected to dominate the return more than normal during odd-numbered years. The data also supports continued above average returns of age-3 fish. If age-3, age-4, and age-5 fish trend upwards in 2005, a strong fall chum salmon run size may surpass the upper end of the projected range. The 2005 projected run size range should support normal subsistence fishing activities and should provide some opportunity for commercial fishing where markets exist. The run will be monitored inseason to determine the run strength and the amount of harvest that could be provided based on the levels stipulated in 5 AAC 01.249 *Alaska Yukon River Drainage Fall Chum Salmon Management Plan*.

2.4 COHO SALMON

Coho salmon primarily return as age-4 fish and overlap in run timing with fall chum salmon. The major contributor to the 2005 coho salmon run will be from the 2001 parent year. Based on Pilot Station sonar operations from 1995, and 1997 through 2004, the 2001 return was slightly below average and early in run timing. The evaluations of escapement in the Andreafsky, Kantishna, and upper portions of the Tanana River drainage were above average in abundance in 2001. The monitored spawning areas in the Nenana River drainage were the only areas below average, which could have been a factor of survey conditions. In 2001 the Delta Clearwater River (DCR), the major producer of coho salmon in the upper Tanana River drainage, had its third highest escapement in 33 years. Based on the DCR escapements, coho salmon abundance has improved throughout this time period with a significant increase within the last decade. Assuming average production, the 2005 coho salmon run is anticipated to be average to above average based on excellent escapements in 2001.

The *Yukon River Coho Salmon Management Plan* (5 AAC 05.369) allows a directed commercial coho salmon fishery under certain conditions. Directed coho salmon fishing is dependent on the assessed abundance for both coho and fall chum salmon. Based on the projected 2005 fall chum salmon run expectations and the anticipated strong coho salmon outlook, the combined 2005 Yukon River fall runs should be adequate to support a commercial coho salmon harvest of 10,000 to 75,000 fish.

3.0 MANAGEMENT STRATEGY FOR 2005

ADF&G manages Yukon Area salmon according to policies and regulations established by the Alaska Board of Fisheries (BOF). Management of the Yukon Area commercial salmon fishery is complex due the mixed stock nature of the fishery, increased efficiency of the commercial fleet,

allocation issues, and the complication of State/Federal dual management regimes for the subsistence fishery in about half the drainage. The Yukon River Drainage Subsistence Salmon Fishery Management Protocol provides guidelines for coordinated management with federal agencies concerning subsistence fisheries in waters subject to Federal reserved water rights within the Yukon River drainage. However, state and federal managers may not agree on specific management actions, which results in differing regulations for waters subject to applicable federal management.

The *Policy for Statewide Salmon Escapement Goals* (Escapement Goal Policy: 5 AAC 39.223) and the *Policy for the Management of Sustainable Salmon Fisheries* (Sustainable Salmon Policy: 5 AAC 39.222) define various levels of escapement in a manner consistent with sustained yield. Escapement objectives that were previously estimated in the absence of a stock specific catch estimate and used as an index, or as an escapement estimate, are now defined as a Sustainable Escapement Goal (SEG). Tables 3, 4, and 5 list Biological Escapement Goals (BEGs), SEGs, and Optimal Escapement Goals (OEGs) that will be used for inseason management and postseason assessment. The Canadian Chinook salmon objective and fall chum salmon escapement objectives are based on limited scientific information and is not a SEG or a BEG. These objectives are negotiated by the U.S./Canada Panel annually as stipulated in the treaty agreement.

Table 3.—Escapement goals for Chinook salmon.

Stream	Goal	Type of Goal
East Fork Andreafsky River Aerial Survey	960–1,900	SEG
West Fork Andreafsky River Aerial Survey	640–1,600	SEG
Anvik River Aerial Survey	1,100–1,700	SEG
Nulato River Aerial Survey	940–1,900	SEG
Gisasa River Aerial Survey	420–1,100	SEG
Chena River Tower	2,800–5,700	BEG
Salcha River Tower	3,300–6,500	BEG
Canada Mainstem Tagging Rebuilding Goal	28,000	*

* The US-Canada Panel agreed to a spawning escapement objective of 28,000 in Canada for 2005.

Table 4.—Escapement goals for summer chum salmon.

Stream	Goal	Type of Goal
East Fork Andreafsky River Weir	65,000–135,000	BEG
Anvik River Sonar	400,000–800,000	BEG
Drainage-wide Escapement (above Pilot Station)*	800,000–1,600,000	OEG

* Inriver run goal: This is a specific management objective for salmon stocks that are subject to harvest upstream of the point where escapement is estimated.

Table 5.—Escapement goals for fall chum salmon.

Stream	Goal	Type of Goal
Drainage-wide Escapement	300,000–600,000	BEG
Tanana River drainage	61,000–136,000	BEG
Upper Tanana River	46,000–103,000	BEG
Delta River	6,000–13,000	BEG
Toklat River	15,000–33,000	BEG
Upper Yukon Tributaries	152,000–312,000	BEG
Chandalar River	74,000–152,000	BEG
Sheenjek River	50,000–104,000	BEG
Fishing Branch	24,000	*
Canadian Yukon River Mainstem	>65,000	**

*Interim Goal agreed to by the Yukon River Panel for the 2005 season.

**Rebuilding goal agreed to by the Yukon River Panel for the 2005 season.

3.1 ALASKA BOARD OF FISHERIES ACTIONS

In response to the guidelines established in the Sustainable Salmon Policy, the BOF determined the Yukon River Chinook and fall chum salmon stocks to be yield concerns during the September, 2000 work session. This determination was based on the inability, despite the use of specific management measures, to maintain expected yields, or harvestable surpluses, above the stock's escapement needs since 1998 and the anticipated low harvest level in 2001. In addition, the BOF classified the Yukon River summer chum and Toklat and Fishing Branch River fall chum salmon stocks as management concerns. The determination of the management concerns was based on the chronic inability to meet existing escapement goals for the summer chum stock since 1998 and for the Toklat and Fishing Branch Rivers fall chum salmon stocks since 1997.

During the January 2004 BOF meeting, the BOF addressed stocks of concern action plan options and regulatory proposals. Chinook salmon will continue as a stock of concern at the yield level and summer chum at the management level. The drainage-wide fall chum salmon stock also remains unchanged as a yield concern. In 2001, the Toklat River fall chum salmon stock was declared a management concern because it did not meet the established escapement goal at the time. Since then, the BEG analysis has been finalized and the BOF found that the Toklat River stock does not meet the criteria of a management concern. Also, since the Fishing Branch River fall chum salmon stock is under the purview of the US/Canada Panel, the BOF removed the stock as a management concern. However, both the Toklat and Fishing Branch River fall chum salmon stocks are included in the drainage-wide Yukon River fall chum salmon stock classification as a yield concern.

3.2 SUBSISTENCE FISHERY

Subsistence fishing has the highest priority among all uses of the resource in the State of Alaska and occurs throughout most of the Yukon River Area. When the salmon stocks are abundant and commercial fishing will occur, it is necessary to place some restrictions on the subsistence fishery in order to enforce commercial fishing regulations. For example, subsistence salmon fishing is closed in most areas 24 hours prior to the commercial salmon fishing season to discourage the illegal sale of subsistence caught salmon or salmon roe. However, substantially

more fishing time is allowed throughout the fishing season for subsistence than for commercial purposes.

The subsistence salmon fishery will be based on a schedule and implemented chronologically by ADF&G consistent with migratory timing as the run progresses upstream. The subsistence salmon fishing schedule is based on current or past fishing schedules and should provide reasonable opportunity for subsistence during years of normal to below average runs. The objectives of the schedule are to 1) reduce harvest early in the run when there is a higher level of uncertainty, 2) spread the harvest throughout the run to reduce harvest impacts on any particular component of the run and 3) spread subsistence fishing opportunity among all users during years of low salmon runs. Table 6 shows the subsistence fishing schedule and it is listed in regulation 5AAC 01.210 & 5AAC 05.360. Depending on run strength, the schedule is subject to change.

Table 6.–Yukon Area subsistence salmon fishing schedule.

Note: this schedule is subject to change depending on run strength.

Area	Regulatory subsistence salmon fishing periods	Schedule to begin	Days of the week
Coastal District	7 days/week	by regulation	M/T/W/TH/F/SA/SU – 24 hours
District 1	Two 36-hour periods/week	May 30, 2005	Mon. 8 pm to Wed. 8 am /Thu. 8 pm to Sat. 8 am
District 2	Two 36-hour periods/week	June 1, 2005	Wed. 8 pm to Fri. 8 am / Sun. 8 pm to Tue. 8 am
District 3	Two 36-hour periods/week	June 3, 2005	Fri. 8 am to Sat. 8 pm / Tue. 8 am to Wed. 8 pm
District 4	Two 48-hour periods/week	June 12, 2005	Sun. 6 pm to Tue. 6 pm / Wed. 6 pm to Fri. 6 pm
Koyukuk River	7 days/week	By Regulation	M/T/W/TH/F/SA/SU – 24 hours
Subdistricts 5-A, B, C	Two 48-hour periods/week	June 21, 2005	Tue. 6 pm to Thu. 6 pm /Fri. 6 pm to Sun. 6 pm
Subdistrict 5-D	7 days/week	By Regulation	M/T/W/TH/F/SA/SU – 24 hours
District 6	Two 42-hour periods/week	By Regulation	Mon. 6 pm to Wed. Noon /Fri. 6 pm to Sun. Noon
Old Minto Area	5 days/week	By Regulation	Friday 6pm to Wednesday 6pm

Once it has been determined there is a harvestable surplus of salmon in excess of subsistence uses, the subsistence fishing schedule can be reverted to the schedule specified in 5AAC 01.210, (c–h) FISHING SEASONS AND PERIODS, unless modified by emergency order.

During closed subsistence salmon fishing periods, subsistence fishing for whitefish, suckers, and other non-salmon species will be allowed throughout the drainage. Gillnets with greater than 4 inch mesh must be removed from the water and fish wheels may not be operated during closed subsistence salmon fishing periods in an effort to avoid salmon species. In addition, gillnets used to take species other than salmon during subsistence salmon closures are limited to 60 feet in length. This opportunity to target non-salmon species, while protecting the salmon stocks of concern, may be discontinued if found ineffective at reducing salmon harvest. Subsistence fishing permits are required for the Tanana River drainage upstream of the Wood River, portions of District 5 in the upper Yukon River drainage near the Haul Road Bridge, and from above the village of Fort Yukon to the U.S./Canada border.

The summer and fall chum salmon management plans adopted by the BOF provide guidelines for managing subsistence salmon fisheries based on inseason run projections. If subsistence harvest reductions are necessary, efforts will be made to spread the burden of conservation throughout the drainage. Potential harvest reduction measures include gear restrictions, reductions in fishing time, or extended periods of closed fishing. Conservation of salmon may require fish wheels to be equipped with a live box or live chute.

ADF&G requires fishermen to keep track of their subsistence salmon harvests on their permit in permit areas. Subsistence fishers in permit areas are reminded that they must have their permit in possession while fishing. In non-permit areas, ADF&G conducts a postseason harvest survey and encourages fishermen to use catch calendars to keep track of their harvest. Non-permitted fishermen who did not receive a subsistence salmon calendar by mail may obtain one by contacting ADF&G in Emmonak or Fairbanks. ADF&G has prepaid postage for the calendar in an effort to encourage fishermen to use and return catch calendars. Additionally, a \$200 lottery will be conducted following the season for which all households that have returned properly filled out calendars will be entered.

3.2.1 Districts 1, 2, and 3

The subsistence salmon fishing schedule in Districts 1, 2, and 3 will begin with two, 36-hour periods per week. During the Chinook and summer chum salmon commercial fishing season, subsistence salmon fishing will be closed 18 hours before, during, and 12 hours following a commercial salmon fishing period. During the fall season, subsistence salmon fishing will be closed 12 hours before, during, and 12 hours following each District 1, 2, or 3 commercial salmon fishing period. During the commercial season, dorsal fins are required to be removed from subsistence caught Chinook salmon.

3.2.2 District 4

The subsistence salmon fishing schedule in District 4 is two, 48-hour periods per week. Regulations separate subsistence fishing periods from commercial fishing periods in Subdistrict 4-A. During the commercial salmon fishing season, subsistence salmon fishing will be closed 12 hours before, during, and 12 hours following a Subdistrict 4-A commercial salmon fishing period. If the commercial salmon fishing season is opened in Subdistricts 4-B and 4-C, managers will attempt to coincide allowable commercial salmon fishing periods with the traditional subsistence salmon fishing schedule. When ADF&G announces a commercial fishing closure that will last longer than 5 days in duration during the commercial salmon season in District 4, subsistence salmon fishing may occur 5 days per week, unless modified by emergency order.

From November 1 through June 31, waters open for subsistence fishing in the Koyukuk River drainage are expanded to include the Middle Fork of the Koyukuk River upstream of its confluence with the North Fork, and the South Fork of the Koyukuk River upstream from the mouth of the Jim River. A household subsistence fishing permit is required as a condition of this increased fishing opportunity. Only gillnet gear is allowed and the mesh size may not exceed 3½ inches. This was done in an effort to protect salmon species.

3.2.3 District 5

The Subdistrict 5-A, 5-B, and 5-C subsistence fishing schedule is two, 48-hour fishing periods per week. In Subdistrict 5-D, subsistence salmon fishermen may harvest salmon 7 days per week throughout the season unless restricted by emergency order. Attempts will be made to coincide

the subsistence salmon fishing schedule with commercial periods. Additionally, “subsistence only” salmon fishing periods may also be scheduled. When ADF&G announces a commercial fishing closure that will last longer than 5 days in duration during the commercial salmon season in Subdistricts 5-A, 5-B, and 5-C, subsistence salmon fishing may occur 5 days per week, unless modified by emergency order.

Regulations require subsistence fishermen to obtain subsistence fishing permits in portions of District 5. Permits are required on the Yukon River from the western tip of Garnet Island to the Dall River including the village of Rampart and the Haul Road bridge area. Permits are also required for portions of the Yukon River near the communities of Circle and Eagle. Subsistence fishermen must obtain a permit prior to subsistence fishing which can be done by contacting ADF&G’s office in Fairbanks. Permits can be issued in person, by mail, and more recently by email. All permit holders are required to report harvest information on their permits and return their permits to ADF&G at the end of the fishing season.

3.2.4 District 6

Within the majority of Subdistricts 6-A and 6-B, the subsistence salmon fishing schedule is two, 42-hour periods per week from 6:00 p.m. Monday until 12 noon Wednesday and from 6:00 p.m. Friday until 12 noon Sunday. An exception is within the Old Minto Area, where subsistence salmon fishing is allowed 5 days a week from 6:00 p.m. Friday until 6:00 p.m. Wednesday.

Regulations require subsistence salmon permits in District 6, the Tanana River drainage, except for Subdistrict 6-C, which is managed under personal use regulations (see Section 3.3). Subsistence salmon fishermen can obtain a permit by contacting ADF&G’s office in Fairbanks. Subsistence permit holders in that portion of Subdistrict 6-B, from a point 3 miles upstream of the mouth of Totchaket Slough to the upper boundary of Subdistrict 6-B, are required to report the number of salmon taken to ADF&G each week. Permit holders can report their weekly catch on a message recording at (907) 459-7388. All Tanana River subsistence permit holders are required to record their harvest information on their permit and return expired permits to ADF&G’s office in Fairbanks at the end of the fishing season.

3.3 PERSONAL USE FISHERY

Subdistrict 6-C falls entirely within the Fairbanks Non-subsistence Area and is managed under personal use regulations. Personal use salmon fishing permits are required in Subdistrict 6-C and can be obtained from ADF&G’s office in Fairbanks. Personal use fishermen must possess a valid State of Alaska resident sport fishing license and report their harvests to ADF&G each week. Only one permit per household is allowed annually. The annual possession limit per permit holder is 10 king salmon and 75 chum salmon for periods through August 15, and 75 chum and coho salmon combined for periods after August 15. Subdistrict 6-C fishery harvest limits are 750 Chinook, 5,000 summer chum, and 5,200 fall chum and coho salmon combined. If a harvest limit is reached inseason, the Subdistrict 6-C personal use fishery will be closed.

The personal use fishing schedule is two, 42-hour periods/week by regulation and fishing is from 6:00 p.m. Monday until noon Wednesday and 6:00 p.m. Friday until noon Sunday. Whitefish and suckers may also be taken with dip nets under personal use fishing regulations and a separate personal use whitefish/sucker permit is required.

3.4 COMMERCIAL FISHERY AND REPORTING REQUIREMENTS

All processors, buyers, catcher/processors, and catcher/sellers of salmon are required to register with ADF&G before operating in the Yukon Area. Processors, buyers, and catcher/sellers in Districts 1, 2, and 3 must register with ADF&G's office in Emmonak. Processors, buyers, and catcher/sellers in Districts 4, 5, and 6 must register with ADF&G's office in Fairbanks. Registered salmon buyers are required to provide a verbal report of their salmon purchases within 18 hours following the closure of a commercial fishing period. Buyers may verbally report harvest information in the Upper Yukon Area after office hours by calling a 24-hour message recording at (907) 459-7388. Buyers are also required to mail or deliver fish tickets to ADF&G within 24 hours following closure of a commercial fishing period in the Lower Yukon Area. In the Upper Yukon Area, buyers are required to mail or deliver fish tickets to ADF&G within 36 hours following the closure of a commercial fishing period. If there is incomplete reporting, ADF&G may delay additional commercial fishing periods until the needed harvest reports are received. In addition, it is very important for buyers to accurately report on each fish ticket the statistical area where salmon were harvested.

All salmon caught by CFEC permit holders during commercial periods in which salmon roe was sold shall be reported as numbers of fish on fish tickets in numbers of males and females. Buyers are requested to ensure this information is reported on fish tickets. Regulations also require commercial fishermen in Subdistrict 6-C to report, on each fish ticket, the number of salmon harvested but not sold during commercial fishing periods.

3.5 CHINOOK AND SUMMER CHUM SALMON COMMERCIAL SEASON

Inseason Chinook salmon run assessment will be based on lower river test fisheries, subsistence catch reports, age and sex composition, and preliminary passage and escapement monitoring information. As in years past, ADF&G will participate in Yukon River Drainage Fisheries Association (YRDFA) teleconferences inseason to gather information from the public, disseminate project information, and to discuss run status and management actions. The YRDFA teleconferences have provided an excellent venue for not only distributing information, but also to provide tentative management actions for YRDFA and public participants to comment on. More recently, management decisions have been made with recommendations from these teleconferences.

There is a Chinook Salmon and a Summer Chum Salmon Management Plan, which guides ADF&G's management actions. Table 7 summarizes the summer chum salmon management plan. If the abundance of summer chum salmon in 2005 is similar to that in 2004, a directed summer chum salmon fishery will be allowed provided indications that escapements will be met and markets are available to sell the catch.

Table 7.–Summary of the Summer Chum Salmon Management Plan.

<i>Summer Chum Salmon Management Plan Overview</i>					
Projected Run Size ¹	RECOMMENDED MANAGEMENT ACTION				Targeted Drainage-wide Escapement
	Commercial	Personal Use	Sport	Subsistence	
600,000 or less	Closure	Closure	Closure	Closure ²	≤600,000
600,001 to 700,000	Closure	Closure	Closure	Possible Restrictions ²	
700,001 to 1,000,000	Restrictions ²	Restrictions ²	Restrictions ²	Normal Fishing Schedules	
Greater than 1,000,000	Open ³	Open	Open	Normal Fishing Schedules	≥800,000

¹ PROJECTED RUN SIZE: Mainstem river sonar passage estimate plus the estimated harvests below the sonar site and the Andreafsky River escapement.

² The fishery may be opened or less restrictive in areas that indicator(s) suggest the escapement goal(s) in that area will be achieved.

³ DRAINAGE-WIDE COMMERCIAL FISHERIES: The harvestable surplus will be distributed by district or subdistrict in proportion to the guidelines harvest levels established in 5AAC 05.362 (f) and (g) and 5 AAC 05.365 if buying capacity allows.

3.5.1 Districts 1, 2, and 3

If the Chinook salmon run in 2005 is similar to the run in 2004, a directed Chinook salmon commercial fishery will be possible. Typically, the first commercial opening occurs after the first quarter point of the run. This management strategy provides for passage of a portion of the early run segment through the lower river districts before commercial fishing starts. Lower Yukon River set net test fishing catch per unit effort (CPUE) and Pilot Station sonar data will be used for relative timing and abundance information. The opening of the fishing season is normally announced 48 hours in advance to provide fishermen and buyers adequate time to prepare.

Directed Chinook salmon commercial fishing periods with unrestricted mesh size gillnets are anticipated to be 6 hours in duration, but may be as short as 4 hours. The fishery has become more actively managed by fishing the front and back side of the pulses, allowing the peak to move through the lower Yukon, especially early in the commercial fishing season. Actively managing the commercial fishery also allows fishing periods to be scheduled further apart, if necessary, to spread the commercial harvest across the run. This strategy reduces harvest impacts on any particular portion of below average runs. Districts 1 and 2 have a combined guideline harvest range and it may not be possible to allow an equal amount of fishing time for each district.

Although it is anticipated a summer chum salmon directed commercial fishery may be possible, it is unlikely there will be commercial openings in Districts 1-3 because of poor market conditions. If a market exists, 6 inches or smaller mesh size gillnets will be utilized to direct the harvest toward summer chum salmon.

Commercial fishing periods may be established in District 3 based on input from buyers and fishermen. Regulations allow a permit holder registered in District 3 to transfer to District 1 or 2 following a 72-hour waiting period. Only one district transfer per fisher is allowed in the Lower Yukon Area prior to July 15.

3.5.2 District 4

The first District 4 commercial fishing period usually occurs between June 18 and June 25. A BOF regulation passed in January 2004 allows fish wheel permit holders to use set gillnets during commercial fishing periods established by emergency order. This allows commercial fishermen, who normally target chum salmon, to participate and direct their efforts towards Chinook salmon. Management for summer chum salmon will be based on the Yukon River Summer Chum Salmon Management Plan as outlined on page 11 in this document.

If commercial fishing is allowed, it is anticipated Subdistricts 4-B and 4-C would be placed on a schedule of one or two 24-hour periods per week beginning at 6:00 p.m. Sunday and/or 6:00 p.m. Wednesday. Subdistricts 4-B and 4-C may open earlier than Subdistrict 4-A to allow harvest of earlier migrating Chinook salmon. Historically, there is a much lower harvest of summer chum salmon in Subdistricts 4-B and 4-C than in Subdistrict 4-A when directing harvest at Chinook salmon. If subsistence salmon fishing opportunities in District 4 are not sufficient to meet subsistence needs due to the commercial fishing schedule, additional subsistence-only fishing time will be allowed.

3.5.3 Anvik River Management Area

The Anvik River may be opened to summer chum salmon commercial fishing if the escapement exceeds 500,000 fish. Fishing periods in the Anvik River will be based upon size of the surplus available for commercial harvest and the availability of a commercial market. The intent is to allow a harvest of Anvik River summer chum salmon that is in excess of the spawning escapement goal and to decrease harvest pressure on non-Anvik River summer chum salmon stocks. Permit holders are reminded that all Chinook salmon caught during Anvik River commercial fishing periods must be released alive.

3.5.4 District 5

Assessment of run abundance and timing from downstream districts, along with subsistence catch reports, will be used to determine a season opening. By regulation, no commercial fishing will be allowed in Subdistrict 5-A during the Chinook and summer chum salmon fishing season.

If commercial fishing is allowed in Subdistricts 5-B and 5-C, it is anticipated that fishing periods would be 12-hours in duration. For Subdistrict 5-D, 24-hour commercial fishing periods are typical. This allows ADF&G to monitor and maintain the harvest within guideline harvest ranges. In years with average returns and run timing, the first commercial fishing period usually occurs between June 25 and July 5 in Subdistricts 5-B and 5-C, and between July 1 and July 10 in Subdistrict 5-D.

3.5.5 District 6

District 6 is managed inseason using salmon run strength and timing indicators in the Tanana River drainage including test fish wheel catches near the village of Nenana and escapement information on Chinook and summer chum salmon collected by tower counting projects on the Chena and Salcha Rivers. The District 6 commercial fishing season typically opens in early to mid-July. ADF&G can exceed the upper end of the guideline harvest ranges in years when it has been determined that escapement goals and subsistence needs will be met.

Directed summer chum salmon commercial fishing periods would likely occur later in July and into August and will depend on inseason run assessment. The length and duration of commercial fishing periods will depend on run strength and buyer capacity.

3.6 FALL CHUM AND COHO SALMON COMMERCIAL SEASON

The 2005 fall chum salmon outlook is for an average to below average run and coho salmon are expected to be average to above average. A commercial fishery for both chum and coho salmon is anticipated. In managing the 2005 Yukon River fall chum salmon fishery, ADF&G will follow guidelines provided by the Board in 5 AAC 01.249 *Yukon River Drainage Fall Chum Salmon Management Plan*, which was amended by the BOF in January 2004. The plan stipulates that directed fall chum salmon commercial fisheries may only be allowed on the projected surplus of the run above 600,000 fall chum salmon for the entire Yukon River drainage. There is an exception to this plan where commercial fishing may be allowed in portions of the drainage where escapement and subsistence needs are projected to be exceeded and when the drainage-wide run is projected to be greater than 500,000. Since the 2005 preseason projection is for a run size ranging from 584,000 to 776,000 fall chum salmon, it is anticipated that the run will likely provide sufficient abundance for escapement and subsistence needs with surplus available for commercial harvest. Table 8 summarizes the fall chum salmon management plan.

Table 8.—Summary of the Fall Chum Salmon Management Plan.

<i>Fall Chum Salmon Management Plan Overview</i>					
Projected Run Size ¹	RECOMMENDED MANAGEMENT ACTION				Targeted Drainage-wide Escapement
	Commercial	Personal Use	Sport	Subsistence	
300,000 or less	Closure	Closure	Closure	Closure ²	
300,000 to 500,000	Closure	Closure ²	Closure ²	Possible Restrictions ^{2,3}	
500,000 to 600,000	Restrictions ²	Open	Open	Pre-2001 Fishing Schedules	300,000 to 600,000
Greater than 600,000	Open ⁴	Open	Open	Pre-2001 Fishing Schedules	

¹ PROJECTED RUN SIZES use the best available data (including preseason projections, mainstem river sonar passage estimates, test fisheries indices, subsistence and commercial fishing reports, and passage estimates from escapement monitoring projects)

² The fishery may be opened or less restrictive in areas that indicator(s) suggest the escapement goal(s) in that area will be achieved.

³ Subsistence fishing will be managed to achieve a minimum drainage-wide escapement goal of 300,000.

⁴ DRAINAGE-WIDE COMMERCIAL FISHERIES may be open and the harvestable surplus above 600,000 will be distributed by district or subdistrict (in proportion to the guidelines harvest levels established in 5AAC 05.365 and 5 AAC 05.367.

The 2005 coho salmon fishery will be managed consistent with regulation 5 AAC 05.369 *Yukon River Coho Salmon Management Plan*. The plan allows a directed coho salmon commercial fishery under certain conditions due to overlapping run timing with fall chum salmon. Therefore, the commercial harvest of coho salmon will likely be dependent upon the abundance of fall chum salmon and accompanying management strategies used to harvest fall chum salmon. In 2005 it is anticipated that some commercial fishing for both fall chum and coho salmon will occur simultaneously.

ADF&G will monitor the fall salmon runs inseason by using the lower Yukon River drift gillnet test fishery near Emmonak, the Mountain Village drift gillnet test fishery, Pilot Station sonar

passage estimates, subsistence catch reports, and, if available, commercial catch statistics. This information, in combination with the preseason expectation and the performance of the summer chum salmon run, will be the basis for initial management decisions for Districts 1, 2, and 3 commercial fisheries. The fall season outlook is for a commercial harvest of fall chum salmon of 20,000 to 150,000 and a potential coho salmon harvest of 10,000 to 75,000. Due to conservation concerns for fall chum salmon and the irregular entry pattern, it is likely the decision to prosecute a commercial fishery will not be made until near the midpoint in the fall chum salmon run around late July or early August for the lower river to build confidence in run assessment.

3.6.1 Districts 1, 2, and 3

The majority of fall chum salmon enter the Yukon River from mid-July through early September in erratic pulses usually lasting 2 to 3 days. Typically, four or five large pulses occur each season. These pulses are often associated with onshore wind events and/or high tides. This entry pattern makes it difficult to accurately assess run strength inseason, particularly early in the season. Because of this unpredictable entry pattern and recent conservation measures, the decision to allow a commercial fishery would not be made until near the midpoint of the run in Districts 1, 2, and 3.

Regulations require District 1 commercial fishermen to register for the coastal *Set Net Only Area* prior to opening of the fall commercial season. Registration “sign-in” sheets will be available at Lower Yukon Area village post offices and at ADF&G’s field office in Emmonak. There are provisions that allow fishermen to transfer into and out of the *Set Net Only Area*. After initial registration for the *Set Net Only Area*, a permit holder may not commercially fish for salmon in the remainder of District 1, or in another district, until 72 hours after re-registration with ADF&G. After the first fall season commercial fishing period, a permit holder not registered for the *Set Net Only Area* may transfer to the *Set Net Only Area* after re-registration with ADF&G. The re-registration and 72-hour waiting period begins at the time the notification is received and noted by ADF&G.

3.6.2 District 4

In January 2001, the BOF took regulatory action to include Subdistrict 4-A with the Subdistricts 4-B and 4-C guideline harvest range of 5,000 to 40,000 fall chum salmon. In years with average run timing and a commercially harvestable surplus, the first District 4 fall season commercial fishing period normally occurs in early to mid-August. In the event a directed coho salmon commercial fishery is allowed, a commercial fishing period in Subdistrict 4-A may only occur on or after August 20 and would close by September 15. No more than 32 hours of commercial fishing time may be allowed per week. It is anticipated that there will be a surplus available for commercial harvest, however weak market conditions may limit commercial harvests in the district for 2005.

3.6.3 Subdistricts 5-B, 5-C, and 5-D

In years with average run timing and a commercially harvestable surplus, the first fall season commercial fishing period in Subdistricts 5-B and 5-C normally occurs in mid-August with Subdistrict 5-D starting later in August or early September. Although some fall chum salmon may be available, there is only a small chance of a commercial fall salmon fishery in these subdistricts during the 2005 season due to weak market conditions.

3.6.4 Subdistrict 5-A and District 6

Management of Subdistrict 5-A and District 6 is outlined in regulation 5 AAC 05.367 *Tanana River Salmon Management Plan*. This management plan directs ADF&G to manage Subdistrict

5-A and District 6 based on the stock status and timing of salmon bound for, and into the Tanana River. It is believed the majority of fall chum and coho salmon harvested in Subdistrict 5-A are bound for the Tanana River, located in District 6.

ADF&G will initially manage the fall season in Subdistrict 5-A and District 6 based on the run strength and timing of the overall Yukon River fall chum salmon run. As the run progresses into the Tanana River, the Tanana and Kantishna River mark-recapture tagging projects will be used to assess the run size in the Tanana River portion of the drainage along with other harvest and test fish wheel information.

The Tanana River management plan adopted by the BOF allows Subdistrict 5-A commercial activities only during the fall season. In most years, the Subdistrict 5-A commercial fishery would be managed for a guideline harvest range of 0 to 4,000 pounds of fall chum salmon roe. No waste of carcasses will be permitted. In adopting this regulation, the Board recognized that carcasses produced by this commercial roe fishery should be incorporated into the subsistence take in the village of Tanana.

Depending on the inseason Tanana River fall chum salmon run strength and timing indicators, ADF&G does have the authority to manage Subdistrict 5-A and District 6 for a different harvest level within the guideline harvest range or to exceed the guideline harvest range. The first fall season commercial salmon fishing period normally occurs in early to mid-September.

3.7 GUIDELINE HARVEST RANGES

Table 9 shows a summary of the guideline harvest ranges for all districts in the Yukon Area for the commercial harvest of Chinook, summer, and fall chum salmon.

4.0 U.S./CANADA YUKON RIVER SALMON PANEL AGREEMENT

Negotiations were initiated in 1985 between the U.S. and Canada regarding a Yukon River salmon treaty. In December 2002, the United States and Canada signed an agreement that set salmon harvest share target ranges based on a postseason assessment of run strength for Chinook and fall chum salmon into the Canadian mainstem of the Yukon River. The Alaskan and Canadian fisheries will be managed consistent with stock rebuilding and conservation objectives that have been jointly developed.

For the 2005 season, the U.S./Canada panel has recommended to forego the Canadian Yukon River Mainstem Chinook salmon escapement goal of 33,000–44,000 and have agreed to a rebuilding spawning escapement goal of 28,000 Chinook salmon (Joint Technical Committee of the Yukon River US/Canada Panel *in prep*). The panel agreed to a fall chum salmon Canadian Yukon River Mainstem spawning escapement rebuilding objective of 65,000 chum salmon, and an interim spawning escapement goal of 24,000 chum salmon into the Fishing Branch River. Management plans were laid out to rebuild the Canadian Mainstem Chinook salmon stocks over two life cycles and rebuild the Canadian Mainstem fall chum salmon stocks over three lifecycles which is considered a 12-year period. The Fishing Branch River escapement target of 24,000 represents a small step towards rebuilding the stock. The intent of these lowered goals is to minimize hardships in the subsistence and aboriginal fisheries while continuing efforts towards rebuilding the stocks to higher levels.

Table 9.—Guideline harvest ranges and midpoints for commercial harvest of Yukon River Chinook, summer, and fall chum salmon.

Chinook Salmon						
District or Subdistrict	Guideline Harvest Range ^a					
	Lower		Midpoint		Upper	
	Numbers	Percent	Numbers	Percent	Numbers	Percent
1 and 2	0 to 60,000	89.1	90,000	91.6	120,000	92.9
3	0 to 1,800	2.7	2,000	2.0	2,200	1.7
4	0 to 2,250	3.3	2,550	2.6	2,850	2.2
5-B & 5-C	0 to 2,400	3.6	2,600	2.6	2,800	2.2
5-D	0 to 300	0.4	400	0.4	500	0.4
6	0 to 600	0.9	700	0.7	800	0.6
Total	67,350	100.0	98,250	100.0	129,150	100.0

Summer Chum Salmon						
District or Subdistrict	Guideline Harvest Range ^b					
	Lower		Midpoint		Upper	
	Numbers	Percent	Numbers	Percent	Numbers	Percent
1 and 2	0 to 251,000	62.8	503,000	62.9	755,000	62.9
3	0 to 6,000	1.5	12,500	1.6	19,000	1.6
4-A ^c	0 to 113,000	28.3	225,500	28.2	338,000	28.2
4-B & 4-C	0 to 16,000	4.0	31,500	3.9	47,000	3.9
5-B, -C, -D	0 to 1,000	0.3	2,000	0.3	3,000	0.3
6	0 to 13,000	3.3	25,500	3.2	38,000	3.2
Total	400,000	100.0	800,000	100.0	1,200,000	100.0

Anvik River Management Area roe cap of 100,000 pounds. ^d

Fall Chum Salmon						
District or Subdistrict	Guideline Harvest Range ^e					
	Lower		Midpoint		Upper	
	Numbers	Percent	Numbers	Percent	Numbers	Percent
1, 2, and 3	60,000	82.5	140,000	71.2	220,000	68.6
4	5,000	6.9	22,500	11.4	40,000	12.5
5-B, -C, -D	4,000	5.5	20,000	10.2	36,000	11.2
6	1,000	1.4	2,500	1.3	4,000	1.2
6	2,750	3.8	11,625	5.9	20,500	6.4
Total	72,750	100.0	196,625	100.0	320,500	100.0

Subdistrict 5A range of 0 to 4,000 pounds of roe. ^f

^a The Chinook salmon guideline harvest ranges have been in effect since 1981.

^b Summer chum salmon guideline harvest ranges were established in February 1990 based on the average harvest shares from 1975–1989.

^c Or the equivalent roe poundage of 61,000 to 183,000 pounds or some combination of fish and pounds of roe.

^d The current Anvik River Management Area roe cap was established in March 1996.

^e The current fall chum salmon guideline harvest ranges were established in 1990.

^f Subdistrict 5-A was removed from the guideline harvest ranges for Chinook and summer chum and a separate guideline harvest range of 0–4,000 pounds of fall chum salmon roe was established in November 1998.

REFERENCES CITED

Joint Technical Committee of the Yukon River US/Canada Panel. *In Prep.* Yukon River salmon 2004 season summary and 2005 season outlook. Alaska Department of Fish and Game, Division of Commercial Fisheries, Joint Technical Committee Report, Anchorage.

FIGURES








Figure 1.—Yukon Area communities and fishing districts.





Note: ANS = Amount necessary for subsistence. 2004 subsistence data preliminary.

Figure 2.—Yukon River Chinook salmon subsistence and commercial harvests compared to the recent 10-year average, and the lower ANS range.





Note: ANS = Amount necessary for subsistence. 2004 subsistence data preliminary.

Figure 3.—Yukon River summer chum salmon subsistence and commercial harvests compared to the recent 10-year average, and the lower ANS range.



Note: 2004 subsistence data preliminary.

Figure 4.—Subsistence and commercial harvest of fall chum salmon, 1961 to present.



Note: 2004 subsistence data preliminary.

Figure 5.—Subsistence and commercial harvest of coho salmon, 1961 to present.




Figure 6.–Yukon River fall chum salmon management plan overview and comparison with historical run sizes.