

Fishery Management Report No. 92-2

**Annual Management Report for Sport Fisheries in
the Arctic-Yukon-Kuskokwim Region, 1990**

by

William D. Arvey

June 1992

Alaska Department of Fish and Game

Division of Sport Fish



FISHERY MANAGEMENT REPORT NO. 92-2

ANNUAL MANAGEMENT REPORT FOR
SPORT FISHERIES IN THE
ARCTIC-YUKON-KUSKOKWIM REGION, 1990¹

By

William D. Arvey

Alaska Department of Fish and Game
Division of Sport Fish
Anchorage, Alaska

June 1992

¹ Some of the data included in this report were collected under various jobs of project F-10-6 of the Federal Aid in Fish Restoration Act (16 U.S.C. 777-777K).

The Fishery Management Reports series was established in 1989 for the publication of an overview of Division of Sport Fish management activities or management goals in a specific geographic area. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Distribution is to state and local publication distribution centers, libraries and individuals and, on request, to other libraries, agencies, and individuals. This publication has undergone editorial and peer review.

The Alaska Department of Fish and Game receives federal funding. All of its public programs and activities are operated free from discrimination on the basis of race, religion, sex, color, national origin, age, or handicap. Any person who believes he or she has been discriminated against by this agency should write to:

OEO
U.S. Department of the Interior
Washington, D.C. 20240

TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES.....	iv
LIST OF FIGURES.....	vi
LIST OF APPENDICES.....	vii
ABSTRACT.....	1
PREFACE.....	2
INTRODUCTION.....	3
TANANA AREA DESCRIPTION.....	3
Geographic and Geologic Setting.....	3
Lake and Stream Resources.....	10
Climate.....	10
Species of Importance to the Sport Fishery.....	11
Tanana Area Sport Fisheries.....	11
Chinook Salmon.....	14
Coho Salmon.....	18
Arctic Grayling.....	18
Northern Pike.....	20
Lake Trout.....	21
Burbot.....	25
Whitefish.....	26
Sheefish.....	27
Rainbow Trout.....	27
ARCTIC, YUKON, AND KUSKOKWIM AREA DESCRIPTION.....	28
Geographic and Geologic Setting.....	28
Lake and Stream Resources.....	29
Yukon River Sub-area.....	29
Kuskokwim River Sub-area.....	47
Seward Peninsula/Norton Sound Sub-area.....	48
Northwest Alaska Sub-area.....	49
North Slope Brooks Range Sub-area.....	51
AYK Area Climate.....	56

TABLE OF CONTENTS (Continued)

	<u>Page</u>
Species of Importance to the Sport Fishery.....	56
AYK Area Sport Fisheries.....	56
Chinook Salmon.....	56
Coho Salmon.....	57
Pink Salmon.....	57
Arctic Grayling.....	62
Northern Pike.....	63
Lake Trout.....	64
Dolly Varden/Arctic Char.....	67
Burbot.....	68
Whitefish.....	68
Sheefish.....	69
Rainbow Trout.....	70
STAFF ORGANIZATION.....	70
Regional Staff Responsibilities.....	70
Synopsis of Published Reports	72
MANAGEMENT ACTIVITIES.....	78
1990 In-Season Regulatory Measures.....	78
Emergency Orders and Regulations.....	78
News Releases, Announcements, and Articles.....	79
Preseason and Postseason Regulatory Activities.....	81
Advisory Committees.....	81
Alaska Board of Fisheries.....	81
AYK Sport Fishing Regulations.....	83
AYK SPORT FISHERIES ENHANCEMENT.....	83
Interior Alaska Lake Stocking Program.....	83
1990 Stocking Results.....	84
Anadromous Fish Releases by State Hatcheries.....	88

TABLE OF CONTENTS (Continued)

	<u>Page</u>
LAND USE, HABITAT AND WATER QUALITY.....	88
Habitat-Related Fisheries Issues.....	88
Placer Mining.....	90
Gravel Mining.....	91
Industrial Metals Development.....	91
Oil and Gas Development.....	91
Timberland Development.....	92
Highway Development.....	92
Commercial Utilization of Fisheries Resources.....	92
Wilderness Lodges and Guiding.....	92
Commercial Fisheries.....	93
Land Withdrawals, Status, and Planning.....	94
ANILCA.....	94
National Parks, Monuments, and Preserves.....	94
National Wildlife Refuges.....	96
Wild and Scenic Rivers.....	99
Wilderness Land Designations.....	99
Natural Factors Affecting Sport Fisheries.....	99
Fires.....	99
Snow Pack Assessment.....	100
Stream Discharge Assessment.....	100
Mean Air Temperature and Precipitation.....	101
ACKNOWLEDGEMENTS.....	101
LITERATURE CITED.....	104
APPENDIX A.....	114
APPENDIX B.....	117
APPENDIX C.....	122
APPENDIX D.....	130

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Comparison of 1990 Alaska Sport Fish Survey and Tanana River drainage creel survey harvest estimates.....	12
2. Number of sport anglers, fishing trips, angler days and total freshwater fish harvested in the Tanana River drainage, the AYK Area and the entire state of Alaska, 1990.....	13
3. AYK Region creel surveys, 1990.....	15
4. Tanana Area sport fish harvest and effort by fisheries and species, 1990.....	16
5. Arctic-Yukon-Kuskokwim Region sport fish harvest by species, 1978-1990.....	22
6. Yukon River sub-area sport fish harvest and effort by fisheries and species, 1990.....	58
7. Kuskokwim River sub-area sport fish harvest and effort by fisheries and species, 1990.....	59
8. Seward Peninsula-Norton Sound sub-area sport fish harvest and effort by fisheries and species, 1990.....	61
9. Northwest Alaska sub-area sport fish harvest and effort by fisheries and species, 1990.....	65
10. North Slope Brooks Range sub-area sport fish harvest and effort by fisheries and species, 1990.....	66
11. Number and size of rainbow trout stocked in AYK Region waters in 1990.....	85
12. Number and size of Arctic grayling stocked in AYK Region waters in 1990.....	86
13. Number and size of Arctic char, lake trout, sheefish, sockeye salmon, and coho salmon stocked in AYK Region waters in 1990.....	87

LIST OF TABLES (Continued)

<u>Table</u>	<u>Page</u>
14. Gaging station records of mean and maximum discharge for 1990, and mean and maximum discharge for the period of record, for 11 rivers in AYK.....	89

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. The Arctic-Yukon-Kuskokwim Region. Dashed lines indicate boundaries between harvest reporting areas U - Z.....	4
2. The Tanana River drainage.....	5
3. Waters and highways of the middle Tanana River valley.....	6
4. Tanana River waters near Fairbanks.....	7
5. Map of the Tangle Lakes system.....	8
6. Tributaries of the upper Tanana River.....	9
7. Middle Yukon River and Porcupine River drainages.....	30
8. Major highways in interior Alaska.....	31
9. Lower Yukon River and eastern Norton Sound drainages.....	32
10. Middle Yukon River and Koyukuk River drainages.....	33
11. Upper Koyukuk River and North Slope Haul Road.....	34
12. Porcupine River drainage.....	35
13. Chandalar River system.....	36
14. Waters of the lower Kuskokwim River valley.....	40
15. Waters of the upper Kuskokwim River valley.....	41
16. Waters of the Seward Peninsula.....	42
17. Kotzebue Sound and surrounding area.....	43
18. Waters of the Noatak River.....	44
19. Waters of the upper Kobuk River.....	45
20. Federal land designations within the A-Y-K region.....	46
21. Waters of the Arctic Slope.....	52
22. Principal tributaries of the Colville River.....	53

LIST OF FIGURES (Continued)

<u>Figure</u>	<u>Page</u>
23. Principal tributaries of the Sagavanirktok River.....	55
24. Organizational chart for A-Y-K Sport Fish staff, January, 1990.....	71
25. Monthly mean air temperature for the 1990 water year compared to monthly mean air temperature, 1958-1990, in four AYK locations.....	102
26. Mean monthly precipitation for water year 1990 compared to mean monthly precipitation, 1958-1990, in four AYK locations.....	103

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
A. List of Wild and Scenic Rivers in the Arctic-Yukon-Kuskokwim Region.....	115
B. Emergency orders issued in 1990 for the AYK Region.....	117
C. AYK Region sport fishing regulations summary for 1990.....	123
D. Partial listing of wilderness lodges, guiding and outfitting operations in the AYK Region.....	131

ABSTRACT

Recreational fisheries taking place in northern, northwestern, western, and interior Alaska, an area referred to as the Arctic-Yukon-Kuskokwim Region, are described and reviewed for the calendar year 1990. Species important to the sport fisheries are identified, species distribution is discussed, as is stock status and sport fishing harvest. Regulatory actions affecting the regional sport fishery in 1990 are described along with other management, research, and enhancement activities. Climatic factors of importance to area fisheries are summarized, and federal land status within the region is described.

KEY WORDS: Arctic, Yukon, Kuskokwim, Tanana River, sport fishery, fishery management, recreation, harvest, effort, abundance, regulations.

PREFACE

This is the fifth Annual Management Report for sport fisheries in the Arctic-Yukon-Kuskokwim Region (AYK Region). Statistics pertinent to recreational fisheries in the AYK Region are presented, Alaska Department of Fish and Game (ADF&G) research and reporting activities are described, stock status information is introduced, regulatory changes in 1990 are documented, and other natural and man-influenced factors that may affect fish survival and production during the reporting period are briefly summarized. In addition, long term trends in abundance and/or exploitation are described. Arvey et al. (1990, 1991), Arvey and Parker (1991), and Arvey (1991) present similar information for the years 1986, 1987, 1988, and 1989. The reader is advised to consult other cited regional fisheries data reports for specific information. A brief summary of reports completed by regional staff during the reporting year, along with complete citations is included here under the section entitled "Synopses of Published Reports".

INTRODUCTION

The AYK Region encompasses the majority of the landmass of the state of Alaska (Figure 1). Included within the region are some 1,061,000 km², the state's largest river systems (Yukon, Kuskokwim, Colville, and Noatak), thousands of lakes, and thousands of miles of coastline and streams. Regional coastline boundaries extend from Cape Newenham in the southwest, around all of western, northwestern and northern Alaska to the Canadian border on the Arctic Ocean (Figure 1). The region as a whole is very sparsely populated, with the exception of population centers located in the Tanana River valley. Fairbanks (population about 31,000) is the largest of these communities. The Fairbanks North Star Borough Census Area contains about 78,000 people. Other population centers in the region include the Yukon-Koyukuk Census Area with 8,500 people, Nome Census Area with 8,300 people, Southeast Fairbanks Census Area with 6,000 people, Northwest Arctic Borough with 6,100 people, Wade Hampton Census Area with 5,800 people, and the North Slope Borough with 6,000 people (Alaska Department of Labor 1991).

The State of Alaska sport fishery program divides the AYK Region into two separate fishery management areas; the Tanana Area, and the AYK Area. The Tanana River drainage is a separate management area because it contains population centers that have greater impacts upon fishery resources. Intensive, stock specific studies have been required in the Tanana Area to provide needed biological information for fishery management because of higher exploitation rates in this area. Beginning in 1990, the Sport Fish Division assigned separate management responsibility to the upper Tanana River Valley and to the Norton Sound/Kotzebue portion of the AYK Area.

TANANA AREA DESCRIPTION

Prior to 1990, harvest reports for the Tanana Area included all southern drainages of the Yukon River from its confluence with the Tanana River near Tanana, east to the Canadian border. They also included the Alaskan portion of the Fortymile and Sixtymile River drainages, the entire Tanana River watershed, and the Alaskan portion of the White River drainage. Beginning in 1990, only the Tanana River watershed is considered to be part of the Tanana Area for the purpose of harvest reporting. All other portions of the upper Yukon River are now included in the AYK Area.

Geographic and Geologic Setting

The Tanana River basin (Figures 2-6) is an area of approximately 113,900 km² (11.4 million ha). The main river is a large glacial stream formed at the confluence of the Chisana and Nebesna rivers near Tok. After flowing downstream in a general northwesterly direction for some 917 km, it joins the Yukon River at Tanana. It is the second largest tributary of the Yukon River; the Porcupine River is slightly larger. The Tanana River receives most of its water volume and sediment loads from streams draining the glaciers of the Alaska Range and the Wrangell Mountains. All major tributaries flowing into the north side of the Tanana River originate in the Tanana Hills uplands and are clear in both winter and summer except for relatively short periods after




Figure 1. The Arctic-Yukon-Kuskokwim Region. Dashed lines indicate boundaries between harvest reporting areas U - Z.




Figure 2. The Tanana River drainage.




Figure 3. Waters and highways of the middle Tanana River valley.




Figure 4. Tanana River waters near Fairbanks.




Figure 5. Map of the Tangle Lakes system.




Figure 6. Tributaries of the upper Tanana River.

heavy rains and during the spring period of snowpack melt. They include the Goodpaster, Salcha, Chena, Chatanika, and Tolovana rivers. Rivers flowing from the Alaska Range and the Wrangell Mountains and entering the south side of the Tanana River are of glacial origin. They include the Chisana, Nabesna, Tok, Delta, Nenana, and Kantishna rivers (Figure 2).

Lake and Stream Resources

Large alluvial aquifers associated with porous floodplain gravels influence fish production by storing water and providing more stable winter stream flows in the upper Tanana River and some of its tributaries from Delta Junction upstream and in the Toklat River, tributary to the Kantishna River. All the large aquifers are located on the south side of the Tanana River and are associated with sub-surface water flows from the north slope of the Alaska Range. The Delta Clearwater and Richardson Clearwater rivers (Figures 2 and 3) are the most important sport fishing streams that originate from these aquifers. The few on-stream lakes present (lakes that drain into a stream tributary to the Tanana River or into the Tanana River itself) in the drainage are of insufficient volume to sustain flow in the Tanana River during winter or through dry summers. Headwater glaciers provide water storage that maintains stream flows in dry summers (Selkregg 1976).

Lake development in the Tanana basin is not as extensive as in many other parts of Alaska. Some 20 lakes within the drainage exceed 26 km² in surface area (Selkregg 1976). Most of the water bodies within the system do not contain sufficient volume to influence Tanana River flows, but many are important for sport fishing because of wild or stocked species present in the lakes. Primary lakes for sport fishing within the Tanana River drainage are Harding, Birch, Chena, Quartz, Volkmar, George, Fielding, and Tangle lakes (Figures 3 and 4). Volkmar and George lakes do not have roadside access. Chena Lakes (Figure 2) in the lower Chena River basin are man-made, resulting from gravel removal to erect flood control structures during the 1970's. Many of the waters listed are shown in Figures 3 and 4, and the lakes of the upper Delta River are shown in Figure 5.

Lakes are generally ice covered by late October and breakup can occur in late June or early July. By late winter, ice thickness typically ranges from 81 to 102 cm on interior Alaska lakes.

Lakes were formed in various ways. Some (e.g. Harding, Healy, and George lakes) were created by silt from the Tanana River damming streams draining the surrounding hills. The melting of buried ice masses can sometimes result in lake formation in the sub-glacial soil. Lakes also form when permafrost melts due to vegetative disturbance. Vegetation insulates permafrost soils, and melting can occur when it is removed (Selkregg 1976).

Climate

Climate in the area is one of harsh contrasts, with spring-like conditions occurring as early as mid-April and snowfall, with sub-freezing temperatures, occurring as late as June. Summers are three months in duration and are characterized by long daylight hours and temperatures that occasionally exceed

32°C. The fall season may extend through early October, with snowfall and decreasing temperatures. Winter lasts from mid-November until mid-March, and during this time temperatures may fall to -57°C. Annual precipitation averages around 28 cm, the majority falling between June and September (USDA 1990).

Species of Importance to the Sport Fishery

There are 17 fish species known in the Tanana area of which 10 are important sport species. They include: chinook salmon *Oncorhynchus tshawytscha*, coho salmon *Oncorhynchus kisutch*, Arctic grayling *Thymallus arcticus*, burbot *Lota lota*, lake trout *Salvelinus namaycush*, inconnu (sheefish) *Stenodus leucichthys*, least cisco *Coregonus sardinella*, humpback whitefish *Coregonus pidschian*, and northern pike *Esox lucius*. Rainbow trout *Oncorhynchus mykiss* are not native to the drainage, but have been stocked in several locations.

Tanana Area Sport Fisheries

The statewide harvest survey (SHS), compiled annually since 1977 by Mills (1979-1991) serves as the basic reference of effort and harvest in AYK Region fisheries. Many fisheries in the Tanana Area are not monitored by ADF&G creel surveys, and several are monitored only to obtain CPUE (catch-per-unit-effort) and biological catch composition data, but not seasonal harvest totals. Because so few fisheries can be monitored by creel survey, the statewide survey is the only source of harvest information. Comparison of available harvest estimates from creel surveys with those from the SHS, (Table 1) indicates that similar harvest estimates generally resulted, with the exception of Piledriver Slough rainbow trout and Arctic grayling, where the estimates disagree substantially.

Most harvest and effort data presented in this report are based upon the Alaska statewide sport fish harvest report (Mills 1991) in which the harvest for 30 separate locations within the Tanana drainage was estimated. SHS estimates are based upon responses to postal surveys sent to a random sample of resident and nonresident Sport Fish license holders. Approximately 4% of all sport fishing license holders were surveyed annually until 1989, when the sample size was approximately doubled and 8.7% of all anglers received questionnaires. A total of 30,261 questionnaires was mailed in 1990, followed by two reminders to non-respondents. Responses were obtained from more than 57% of the individuals that received the first mailing.

Catch, as opposed to harvest was estimated and reported separately for the first time in 1990 for all areas statewide (Mills, 1991).

Results of the 1990 SHS for the Tanana River drainage fisheries were based on responses received from several hundred people (Mills 1991). In 1990, the proportion of the total number of angler-days spent in the Tanana Area was 75% and 10% of the total number of all angler-days spent in the AYK Region and the entire state of Alaska, respectively (Table 2). Of the 1,185,603 fish harvested in freshwater (includes anadromous salmonids) in the state, 128,680 (11%) were taken from the Tanana River drainage, and 42,735 (4%) were taken from the AYK Area (Table 2).

Table 1. Comparison of 1990 Alaska Sport Fish Survey and Tanana River drainage creel survey harvest estimates.

Fishery	Creel survey	Harvest survey ^a	
	Harvest	Harvest	95% Confidence Interval
Piledriver Slough			
rainbow trout	2,366	8,052	ND
Arctic grayling	317	2,380	ND
Delta Clearwater River			
Arctic grayling	1,096	1,772	ND
Chatanika River			
whitefish	6,501	5,216	2,245 - 9,250
Salcha River			
chinook salmon	200	291	168 - 414
Chena River			
chinook salmon	24	64 ^b	28 - 105

^a Statewide sport fish harvest survey (Mills 1990).

^b Estimate is for both large and small anadromous chinook salmon.

Table 2. Number of sport anglers, fishing trips, angler days and total freshwater^a fish harvested in the Tanana River drainage, AYK and the entire state of Alaska, 1990^b.

	Tanana Drainage	AYK ^d	Alaska
Number Sport Anglers ^c	38,031	9,768	424,873
Number Fishing Trips ^c	131,831	31,721	1,874,753
Number Angler Days ^c	184,887	60,742	2,453,284
Total Fish Harvested	128,680	42,735	1,185,603

^a Includes anadromous salmonids.

^b Mills 1991.

^c Includes both marine and freshwater; harvest is freshwater only.

^d Exclusive of Tanana River drainage.

An average of approximately 72% of the sport fish harvested in the AYK Region is taken from the Tanana Area (Mills 1977-1991). Effort in angler-days in the Tanana Area has increased from 100,000 in 1977 to about 185,000 in 1990. Since 1988 the harvest of stocked fish has exceeded that of wild stocks in the Tanana Area. A brief description of the sport fisheries in the Tanana Area follows.

Chinook Salmon:

Sport fishing for sea-run chinook salmon in the Tanana River drainage is largely limited to the lower sections of the Salcha and Chena rivers and to the upper Chatanika River, since these are essentially the only road-accessible stocks of sufficient size to support recreational harvests. Annual sport harvests in the Salcha River since 1980 have ranged between zero (1988) and 904 (1980) chinook salmon, while harvests in the Chena and Chatanika rivers are generally smaller. Harvest estimates by the SHS, of chinook salmon in both the Chena and Salcha rivers, were similar in 1990 to those obtained by the creel survey study (Table 1). The sport harvest estimated by creel survey for the lower Chena River was lower (24 chinook salmon, Table 3) than the harvest estimated by the SHS (64 chinook salmon, Mills 1991). The creel survey estimated a sport harvest of 200 chinook salmon in the Salcha River in 1990 (Table 3), compared to an estimated 291 by the SHS (Mills 1991; Table 4).

Separate harvest estimates for chinook salmon less than 41 cm (16 in) have been made in the SHS since 1988. These estimates include only small anadromous chinook salmon. Harvest estimates of small chinook salmon taken from stocked lakes are included with landlocked coho salmon. Mills (1991) estimates that 56 small anadromous chinook salmon were harvested in the Tanana River drainage in 1990, the majority from the Salcha River, where 264 large chinook salmon were estimated in the sport harvest.

The Salcha River spawning escapement in 1990 was estimated¹ to be within the range desired by ADF&G for reproductive purposes (ADF&G 1991). Escapements estimated by aerial survey that range from 1,500 to 3,500 adult spawning fish are considered desirable, and the peak aerial escapement estimate of 3,446 chinook salmon in 1990 is near the upper limit of the range. Since 1980, escapements have ranged from as few as 1,031 to as many as 6,757 chinook salmon under good survey conditions in the Salcha River. The average escapement from 1980 to 1990 (11 data points), is 2,696 chinook salmon. Chena River chinook salmon escapement in 1990 was estimated to be 1,436 fish. Estimates of escapement from aerial surveys conducted under good conditions have ranged from 1,280 to 2,553 fish during the 1980 to 1990 period, with an average (nine data points) of 1,971 fish per year. Escapement objectives (estimated by aerial survey) for the Chena River have been set at 1,000 to 1,700 chinook salmon, and the 1990 escapement estimate fell within that range.

In addition to aerial survey estimates of escapement, mark-recapture experiments to estimate spawner abundance have been conducted since 1986 in the Chena River and 1987 in the Salcha River by the Sport Fish Division.

¹ Estimated by fixed wing aerial survey

Table 3. AYK Region creel surveys, 1990

Water Body/Fishery	Species ^a	Date(s)	Effort (hours)	HPUE ^b	Harvest
Chatanika River	LC	14 Sept-10 Oct	4,894 ^b	ND	5,396
Chatanika River	HW	14 Sept-10 Oct	4,894 ^b	ND	957
Chatanika River	RW	14 Sept-10 Oct	4,894 ^b	ND	148
Chena River	KS	07 Jul- 29 Jul	1,699	0.15	24
Delta Clearwater River	GR	01 Jun- 12 Aug	2,703	ND	1,096
Piledriver Slough	RT	1 May- 12 Aug	8,484	ND	2,366
Piledriver Slough	GR	1 May- 31 Aug	8,494	ND	317
Salcha River	KS	07 Jul- 29 Jul	5,796	0.98	200
Harding Lake	NP	18 May- 17 Jun	815 ^c	ND	15

^a GR: Arctic grayling; HW: Humpback whitefish; KS: chinook salmon; LC: least cisco; RW: round whitefish; RT: rainbow trout; NP: northern pike.

^b Harvest per unit effort data for all whitefish species.

^c Incomplete season data.

Table 4. Tanana Area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	KI ^d	KS	SS	LL	CS	LT	DV AC	RT	GR	WF
Upper Chena River (river miles 77 to 88)	1,131	1,283	1,889	0	0	0	0	0	0	0	0	0	0
Remainder of the Upper Chena River ^e	2,566	3,747	5,060	0	0	0	0	0	0	0	0	945	17
Lower Chena River ^f	5,840	11,310	18,957	5	59	0	0	0	0	0	0	3,494	68
Badger Slough ^g	1,485	2,937	3,455	0	0	0	0	0	0	0	0	68	51
Piledriver Slough	6,313	23,024	27,705	0	5	0	0	0	0	0	8,052	2,380	51
Nenana River Drainage	541	658	724	0	0	261	0	0	51	17	0	490	34
Chatanika River	5,469	7,883	11,801	14	23	0	0	14	0	0	0	4,237	5,216
Salcha River	4,828	8,086	9,783	27	264	0	0	27	0	0	0	1,992	68
Delta Clearwater River	2,498	3,663	4,853	0	0	1,375	0	0	0	0	0	1,772	0
Goodpaster River	861	1,097	2,083	0	0	0	0	0	0	0	0	760	186
Brushkana Creek	523	354	725	0	0	0	0	0	0	0	0	574	0
Lower Tanana River	439	524	866	0	0	0	0	0	0	0	0	118	0
Middle Tanana River	1,047	1,756	2,433	5	23	12	0	0	0	17	0	68	0
Upper Tanana River	557	1,182	1,294	0	9	0	0	0	0	0	0	68	0
Shaw Creek	591	1,300	1,462	0	0	0	0	0	0	0	0	203	34
Richardson Clearwater River	338	371	518	0	0	0	0	0	0	0	0	523	0
Delta River (below Tangle Lakes)	422	321	958	0	0	0	0	0	51	17	0	304	0
Other Streams	2,926	6,555	7,653	0	0	0	0	0	68	153	0	3,734	811
Birch Lake	7,528	10,196	15,541	0	0	0	3,308	0	0	0	15,901	0	0
Quartz Lake	9,824	12,829	19,746	0	0	0	7,309	0	0	0	20,847	0	0
Fielding Lake	1,047	608	1,255	0	0	0	0	0	186	0	0	1,097	0
Minto Flats	709	608	932	0	0	0	0	0	0	0	0	0	17
Tangle Lakes	2,836	2,414	5,228	0	0	0	0	0	236	0	0	2,853	135
Chena Lake (Lake only)	5,115	8,980	12,875	0	0	0	2,313	0	0	0	8,558	0	0
Harding Lake	2,650	3,410	3,895	0	0	0	0	0	51	304	354	17	0
Dune Lake	625	422	815	0	0	0	422	0	0	0	591	304	169
East Twin Lake	523	861	1,035	0	0	0	0	0	0	0	0	0	0
George Lake	658	760	1,540	0	0	0	0	0	0	0	0	17	0
Koole Lake	304	540	686	0	0	71	0	0	0	0	574	0	0
Other Lakes	6,724	14,152	19,120	5	0	0	591	14	253	322	9,266	2,396	34
TOTAL	38,031^h	131,831	184,887	56	383	1,719	13,943	55	896	830	64,143	28,414	6,891

-continued-

Table 4. (Page 2 of 2).

	SF	NP	BB	OTHER
Upper Chena River (river miles 77 to 88)	0	0	0	0
Remainder of the Upper Chena River ^e	0	68	0	0
Lower Chena River ^f	0	68	304	0
Badger Slough ^g	0	68	0	0
Piledriver Slough	34	68	456	0
Nenana River Drainage	0	17	68	0
Chatanika River	0	388	17	0
Salcha River	0	135	203	0
Delta Clearwater River	0	17	0	0
Goodpaster River	0	17	0	0
Brushkana Creek	0	0	0	0
Lower Tanana River	17	405	236	0
Middle Tanana River	0	17	912	0
Upper Tanana River	0	0	641	0
Shaw Creek	0	0	354	0
Richardson Clearwater River	0	0	0	0
Delta River (below Tangle Lakes)	0	0	0	0
Other Streams	0	507	388	34
Birch Lake	0	0	0	0
Quartz Lake	0	0	0	0
Fielding Lake	0	0	0	0
Minto Flats	17	1,182	0	0
Tangle Lakes	0	0	17	0
Chena Lake (Lake only)	0	0	0	0
Harding Lake	0	591	17	0
Dune Lake	0	0	0	0
East Twin Lake	0	760	0	0
George Lake	0	945	34	0
Koole Lake	0	0	0	0
Other Lakes	101	2,095	152	0
TOTAL	169	7,348	3,799	34

^a Tanana River Drainage (Area U): The entire Tanana River watershed, excluding all portions of the Yukon and Kuskokwim rivers watersheds.

^b KS: chinook salmon; SS: coho salmon; LL: landlocked coho or chinook salmon; CS: chum salmon; LT: lake trout; DV/AC: Dolly Varden or Arctic char; RT: rainbow trout; GR: Arctic grayling; WF: whitefish of various species; SF: Inconnu (sheefish); NP: northern pike; and, BB: burbot.

^c From Mills 1991.

^d Chinook salmon less than 410 mm (16 inches).

^e The Chena River and tributaries accessed from the Chena Hot Springs Road beyond Milepost 25 on the road.

^f The Chena River and tributaries from the mouth upstream to Milepost 25 of the Chena Hot Springs Road.

^g All parts of Badger Slough (sometimes called Chena Slough).

^h Angler totals may not equal sum of sites due to some anglers fishing at more than one site.

Results of the abundance estimates in 1990 indicated that 10,728 adult chinook salmon returned to the Salcha River, compared to the peak aerial survey count of 3,446 fish (Burkholder 1991b). It was estimated that 5,603 adult chinook salmon returned to the Chena River in 1990, compared to the peak aerial survey count of 1,436 fish (Evenson 1991a).

Coho Salmon:

Anadromous coho salmon are taken in the Delta Clearwater River and from various creeks in the Nenana River drainage. Coho salmon spawn in other clear spring-fed tributaries of the Tanana River where little sport effort occurs. The Kantishna River and Toklat River tributaries such as Moose, Barton, and Geiger creeks, and the Sushana River (Figure 2), where artesian waters are found, support coho salmon populations. The largest sport harvest in the Tanana Area occurred in 1988 when an estimated 2,237 anadromous coho salmon were taken. The sport harvest in 1990 is estimated to have been 1,719 fish (Table 4). The Delta Clearwater River supports both the largest coho spawning population and the largest recreational harvest in the Tanana Area, where an estimated 1,375 fish were taken in 1990 (Table 4). The fishery in the Delta Clearwater River for coho salmon typically takes place from mid-September until the end of October.

Delta Clearwater River escapements have generally increased since 1985 when 5,358 coho salmon were estimated in the spawning population. Comparable escapement sizes for 1988, 1989 and 1990 were 21,600, 11,000, and 8,325 fish respectively (Whitmore et al. 1990; ADF&G 1991). Escapement since 1977 has ranged from 3,946 fish in 1980 to 22,300 fish in 1987. Commercial fishing restrictions in the lower Yukon River salmon fisheries, combined with low natural mortality have apparently favored large escapements of this species to spawning grounds in drainages of the Yukon and Tanana rivers.

Stocking of landlocked coho salmon in Quartz, Birch, Chena and several smaller lakes has been popular with anglers in the Tanana River valley. An estimated 13,943 such fish were harvested from stocked lakes in 1990, with 7,309 taken from Quartz Lake alone (Table 4).

Arctic Grayling:

Arctic grayling are widely distributed in streams and lakes of the Tanana Area. Trout-like feeding characteristics and their susceptibility to lures and flies make the Arctic grayling a favorite of anglers. Essentially all major clear water tributaries to the Tanana River support Arctic grayling populations. The largest Arctic grayling fisheries in the Tanana drainage occur in the Chatanika, Chena, Salcha, Goodpaster, Delta Clearwater and Richardson Clearwater rivers; Badger and Piledriver sloughs; and Fielding and Tangle lakes. The Tangle lakes in the headwaters of the Delta River support the largest lacustrine fishery for Arctic grayling in Alaska.

Historically, the Chena River supported the largest Arctic grayling fishery in the state, due to an abundant stock, proximity of the Chena River to Fairbanks, and the extensive road access available. The average annual harvest of Arctic grayling in the Chena River exceeded 21,000 fish during the

eight years from 1977 to 1984. The harvest declined from a peak of 42,000 in 1980 to a low of 3,090 in 1987. Clark (1987) attributed declines in abundance and fishing success in the Chena River since 1984 to both sport fishing over-harvest and to reduced recruitment because of unfavorable environmental conditions (primarily high river discharge during the natal year). The estimated 1989 sport harvest in the Chena River and its tributaries was 13,737 Arctic grayling. The 1990 harvest was estimated at 4,507 fish, (Table 4) 33% of the 1989 harvest estimate, comparable to the 1988 harvest estimate of 5,328. Modest increases in survival and recruitment estimates over recent previous years for Chena River Arctic grayling were reported by Clark (1990) for 1988-1989. Clark (1991) also reported substantial increases in survival and recruitment in 1989-1990, exceeding the long term average. Abundance of fish 3 years old and older in the lower 152 km of the Chena River (estimated in July) was estimated to be 29,130 fish, compared to 16,236 fish estimated in 1989.

In 1990, the sport harvest of Arctic grayling in the Chena River was regulated by a minimum length limit of 30.5 cm (12 in) and a bag and possession limit of two fish, among other restrictions. The reduction in bag limit from five to two fish daily is probably responsible for the major reduction in estimated harvest, in spite of the fact that population estimates had increased from 1989 to 1990. Generally, the estimated harvest from the Chena River has declined drastically since 1984, in spite of the fact that effort levels remain high.

Concern for the conservation of several Tanana River drainage Arctic grayling stocks were identified from stock status studies conducted during the latter half of the 1980's decade. Shaw Creek, the Delta Clearwater River, Richardson Clearwater River and Chena River Arctic grayling stocks were considered in danger of over-exploitation. Since early assessments, stocks in all but the Chena River appear to have stabilized at healthy levels.

In 1975, because of increased fishing effort associated with the influx of people for the construction of the trans-Alaska pipeline, the daily bag limit in the Tanana River drainage was decreased from 10 to five Arctic grayling. The reduction in bag limit in 1975 to five fish daily and 10 in possession did not prevent the decline of important Arctic grayling stocks nor did it increase or stabilize fishing success. Further restrictions were enacted in 1987, including a decrease in the possession limit to five Arctic grayling daily, corresponding to the daily bag limit, and institution of a minimum length limit of 30.5 cm in the Chena River. Conservation measures were also enacted for the Chena River and other stocks in the Tanana River drainage. Further restrictions were imposed by the BOF for 1990, including a bag limit reduction to two fish daily and in possession, and use of un-baited, single-hook, artificial lures upstream of the Chena River flood control dam.

Harvest estimates of Arctic grayling in the Tanana Area for 1990 were the lowest recorded since the statewide harvest survey was initiated in 1977, with a total of 28,414 fish, 10,000 fewer fish than were estimated in 1987, the previous low harvest year. The largest harvests from single systems in the area were taken from the Chatanika and Chena rivers in 1990.

In spite of (or perhaps, because of) large historic annual harvests in the Tanana Area, there have been no trophy Arctic grayling (larger than 1.4 kg, 3.0 lb) registered since the inception of the trophy program in the mid-1960's, while 125 individuals have been recorded from rest of the state. Growth rates of individual Arctic grayling in the Tanana River drainage are considered to be typical for Alaska, however, growth rates of Arctic grayling in Bristol Bay, and the Seward Peninsula, where the majority of the trophy fish have been taken, are exceptionally high. It is also possible that Tanana River drainage populations have been maintained at smaller individual size from steady fishing and natural mortality, so that even though growth rates are normal, most individuals are removed before reaching minimum trophy size.

Northern Pike:

Northern pike are harvested by anglers using hook and line gear in summer and winter as well as with spears during the winter. The majority of the Tanana Area harvest comes from lakes that have relatively good access. Important fishing areas are found in Minto Flats, northwest of Fairbanks (Figure 2) and in Harding, George, Healy and East Twin lakes (Figures 3 and 4). Through-the-ice fisheries during the two months just prior to spring break-up, when northern pike have concentrated for spawning, account for a significant portion of the annual fishing mortality.

In 1990, the estimated sport harvest of northern pike in Harding Lake, Minto Flats (including Chatanika River), George Lake, and East Twin Lake was 591, 1,570, 945, and 760 fish respectively, accounting for about 53% of the harvest of 7,348 northern pike in the Tanana River drainage (Table 4). Harvest estimates for the Tanana Area declined 35% from more than 11,000 fish in 1989 to 7,348 in 1990. The harvest estimate for 1990 is the lowest since inception of the SHS in 1977.

Additional lakes and streams utilized by sportsmen in pursuit of this species include West Twin and Wien lakes in the Kantishna River drainage, Fish Lake near the Tanana-Yukon confluence, Volkmar and Healy lakes near Delta, Wellesley, Dog, Jatahmund, Island, and Deadman lakes, Moose and Scotty creeks in the vicinity of Northway (Figures 2 and 6), and other tributary streams of the Tanana River including the Chena River.

Although the 1990 harvest was low, generally harvest and effort have increased for northern pike in the past ten years. Angler surveys indicate that this species is the second most sought-after indigenous sport fish species in interior Alaska (Holmes and Pearse 1987). The total sport harvest of northern pike within the Tanana River drainage has remained relatively stable over the past ten years, from 8,000 to 12,000 fish annually although the distribution of the harvest has varied among various fishing sites. Because of increased effort, the catch per angler day has decreased, and some stocks have been over-fished. The estimated harvest of northern pike in the Tanana River drainage in 1990 (7,348 fish) constituted about 46% of the estimated statewide harvest (15,985) of this species (Mills 1991).

Recent studies of northern pike populations within the Tanana River drainage indicated that exploitation rates are higher than sustainable in some

populations. Even in populations where exploitation rates are not judged to be excessive (less than 20% per year) as in Volkmar Lake, the number of large fish has declined under only moderate harvest pressure. The Minto Flats population has been of special concern due to high additive estimated rates of exploitation in summer and winter subsistence and sport fisheries. Concern for increased harvest in the Minto Flats (Tolovana River drainage) resulted in a winter closure of the sport fishery beginning in the winter of 1987. New regulations, including a reduction in the daily bag limit from 10 to five fish and a winter sport fishery closure, have helped reduce sport harvests of northern pike in the Minto Flats since 1987. Harvest estimates were 1,492 northern pike in 1988, 1,684 in 1989, and 1,182 in 1990. Historical harvests from this wetlands complex were much higher, with estimates exceeding 4,600 and 4,900 northern pike in 1985 and 1986. A total of 58 (33%) of the statewide registered trophy northern pike (minimum weight of 6.8 kg, 15 lbs) was taken from the Tanana River drainage. The Chatanika River, Tolovana River and Minto Flats (considered parts of the Minto Flats complex) account for 23 (38%) of the Tanana River drainage trophy northern pike records, with 13 (21%) from East Twin Lake and eight (13%) from Volkmar Lake. The number of large pike taken from each area may be more reflective of relative fishing effort than of size and growth characteristics of the respective stock.

Lake Trout:

Lake trout occur in many lakes and some streams of the Delta River and upper Tanana River drainages (Burr 1987). They most frequently inhabit deep, oligotrophic mountain lakes and are rarely found at lower elevations of the Tanana River drainage. Lakes of the Delta River drainage (Figure 5) including Fielding, Landmark Gap, Glacier, Sevenmile, and the Tangle lakes contain lake trout. Transplanted lake trout occur in Harding Lake (Figure 4) near Fairbanks and although the small numbers do not support a large fishery, some large individuals have been taken. On average, 65% of the AYK Region lake trout harvest is from lakes in the Tanana River drainage. The regional lake trout harvest increased at an annual average rate of 27% from 1978 to 1985. An apparent major decline in abundance occurred and was first observed in 1986 in waters of the Tanana River drainage. Research in both southcentral and interior Alaska indicates that most of the road accessible stocks have been overharvested in recent years.

Lake trout are a long lived, slow growing and late maturing species, and the impact of even modest fishing pressure can be significant. Lake trout 25 years of age and older are not uncommon and individuals estimated to be older than 50 years are recorded for Alaska (Burr 1987). Trophy lake trout weighing 8.7 kg (20 lbs) or more are typically 20 or more years old (Burr 1987). Lake trout inhabiting high elevation lakes in the Alaska Range migrate into shallow rocky shoals to spawn in late fall. Lake trout spawn for the first time at ages ranging from 5 to 12 years of age, depending apparently on growth conditions. Alternate year spawning may be more normal than spawning in consecutive years in interior and northern Alaska.

Estimated lake trout harvest in the Tanana River drainage peaked at about 3,100 fish in 1982. Harvest declined to 713 and 652 in 1986 and 1987, after the bag limit was reduced from 12 to two lake trout per day (Table 5). After

Table 5. Arctic-Yukon-Kuskokwim Region sport fish harvests by species, 1978-1990^a.

Species	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<u>Tanana Area:</u>													
Chinook Salmon	163	515	941	763	984	1,048	338	1,356	781	502	853	963	439
Sea-Run Coho Salmon	139	25	67	45	52	147	831	796	1,374	1,231	2,237	1,596	1,719
Landlocked Coho/ Chinook Salmon	22,412	36,073	25,733	57,294	43,374	34,255	29,245	41,042	24,061	26,566	32,342	18,614	13,943
Sockeye Salmon	0	0	0	0	0	0	0	0	0	0	0	0	0
Pink Salmon	0	0	0	0	0	0	0	0	0	0	0	0	0
Chum Salmon	158	219	483	595	698	649	585	1,255	693	620	491	1,134	55
Rainbow Trout	6,406	5,186	19,584	24,571	26,186	20,664	34,022	33,432	31,270	31,824	78,345	74,675	64,143
Lake Trout	603	946	1,264	1,721	3,104	2,937	2,104	2,984	713	652	2,221	1,932	896
Dolly Varden/ Arctic Char	524	364	524	572	482	293	350	1,230	200	36	909	913	830
Arctic Grayling	83,275	70,243	80,150	75,288	81,753	92,363	83,626	63,560	45,981	38,480	52,659	54,823	28,414
Northern Pike	7,838	7,975	9,452	9,941	9,822	10,225	9,607	12,090	11,934	9,471	11,986	11,330	7,348
Whitefish	6,573	5,159	5,987	4,873	8,643	8,311	11,658	20,230	26,810	26,435	11,775	16,935	6,891
Burbot	1,383	1,979	2,700	4,122	3,887	5,040	5,556	4,795	5,142	3,855	3,733	4,357	3,799
Sheefish	234	279	96	93	127	157	338	420	72	235	982	643	169
Other Fish	81	79	0	108	10	21	39	0	171	0	0	130	34
Total	129,789	129,042	146,981	179,986	179,122	176,110	178,299	183,190	149,202	139,907	198,533	188,045	128,680

-continued-

Table 5. (Page 2 of 3).

Species	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<u>AYK Area:</u>													
Chinook Salmon	932	644	939	1,112	1,530	2,649	1,481	1,331	2,079	1,691	2,965	2,450	1,366
Sea-Run Coho Salmon	1,308	2,958	3,469	2,087	5,909	5,785	11,205	2,396	7,319	7,210	9,713	8,655	4,819
Landlocked Coho/Chinook Salmon	0	0	0	0	0	0	0	0	0	58	0	0	0
Sockeye Salmon	85	126	112	117	430	261	650	169	439	1,364	1,528	456	818
Pink Salmon	8,328	2,918	7,844	3,118	14,214	5,286	8,712	1,206	3,404	1,322	3,859	3,765	7,994
Chum Salmon	1,834	1,482	2,290	3,045	5,083	4,049	2,689	1,781	3,643	2,148	3,201	4,748	2,091
Rainbow Trout	362	401	835	982	796	1,783	1,455	659	504	592	1,599	757	435
Lake Trout	497	655	1,025	1,100	2,023	1,157	1,520	2,370	2,537	461	509	1,955	847
Dolly Varden/Arctic Char	4,014	8,144	8,273	8,176	13,647	20,324	12,882	13,430	10,173	12,333	11,238	13,359	7,431
Arctic Grayling	11,289	19,229	20,396	20,892	27,043	30,800	15,516	17,666	19,744	19,476	16,302	17,215	9,092
Northern Pike	3,915	4,004	6,190	5,184	7,435	8,609	4,610	3,613	7,062	4,751	7,838	5,853	4,982
Whitefish	909	855	1,705	576	3,708	4,746	234	630	4,960	724	1,855	1,997	810
Burbot	506	118	663	684	1,896	555	377	420	469	162	145	537	1,684
Sheefish	1,057	1,263	2,315	2,146	3,154	3,166	3,609	2,100	3,649	2,362	2,239	1,663	581
Smelt	0	0	0	0	0	0	0	8,750	464	7,080	2,476	2,424	1,709
Halibut	0	0	0	0	0	0	0	62	0	36	0	0	144
Other Fish	1,212	2,218	3,513	3,124	8,551	8,806	1,844	1,336	1,178	0	371	241	580
Total	36,248	45,015	59,569	52,343	95,419	97,976	66,784	57,919	67,624	61,770	65,838	66,075	45,383

-continued-

Table 5. (Page 3 of 3).

Species	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<u>AYK Region (both management areas):</u>													
Chinook Salmon	1,095	1,159	1,880	1,875	2,514	3,697	1,819	2,687	2,860	2,193	3,818	3,413	1,805
Sea-Run Coho Salmon	1,447	2,983	3,536	2,132	5,961	5,932	12,036	3,192	8,693	8,441	11,950	10,251	6,610
Landlocked Coho/Chinook Salmon	22,412	36,073	25,733	57,294	43,374	34,255	29,245	41,042	24,061	26,624	32,342	18,614	13,943
Sockeye Salmon	85	126	112	117	430	261	650	169	439	1,364	1,528	456	818
Pink Salmon	8,328	2,918	7,844	3,118	14,214	5,286	8,712	1,206	3,404	1,322	3,859	3,765	7,994
Chum Salmon	1,992	1,701	2,773	3,640	5,781	4,698	3,274	3,036	4,336	2,768	3,692	5,882	2,146
Rainbow Trout	6,768	5,587	20,419	25,553	26,982	22,447	35,477	34,091	31,774	32,416	79,944	75,432	64,618
Lake Trout	1,100	1,601	2,289	2,821	5,127	4,094	3,624	5,354	3,250	1,113	2,730	3,605	1,743
Dolly Varden/Arctic Char	4,538	8,508	8,797	8,748	14,129	20,617	13,232	14,660	10,373	12,369	12,147	14,272	8,261
Arctic Grayling	94,564	89,472	100,546	96,180	108,796	123,163	99,142	81,226	65,725	57,956	68,961	71,637	37,506
Northern Pike	11,753	11,979	15,642	15,125	17,257	18,834	14,217	15,703	18,996	14,222	19,824	17,183	12,330
Whitefish	7,482	6,014	7,692	5,449	12,351	13,057	11,892	20,860	31,770	27,159	13,630	18,932	7,701
Burbot	1,889	2,097	3,363	4,806	5,783	5,595	5,933	5,215	5,611	4,017	3,878	4,894	5,483
Sheefish	1,291	1,542	2,411	2,239	3,281	3,323	3,947	2,520	3,721	2,597	3,221	2,306	750
Smelt	0	0	0	0	0	0	0	8,750	464	7,080	2,476	2,424	1,709
Halibut	0	0	0	0	0	0	0	62	0	36	0	0	144
Other Fish	1,293	2,297	3,513	3,232	8,561	8,827	1,883	1,336	1,349	0	371	371	614
Total	166,037	174,057	206,550	232,329	274,541	274,086	245,083	241,109	216,826	201,677	264,371	253,437	174,175

^a From Mills 1991

two years of reduced harvest, Mills (1989) estimated that the 1988 harvest of lake trout had increased to 2,221 fish (Table 5). However, further investigation of responses to the SHS showed that harvests of lake trout were reported from lakes known to contain only stocked rainbow trout and/or Arctic char, particularly from those lakes on Eielson Air Force and Fort Greely Army bases. Lake trout harvest estimates for the Tanana Area in 1989 total 1,932 fish. After eliminating some responses, a more accurate estimate of 1989 native lake trout in the Tanana River drainage is 1,498 fish. Lake trout harvest estimates for the Tanana Area in 1990 total 896 fish of which 253 are reported from un-specified lake locations (Table 4). Responses to the SHS have not been examined to determine whether any harvest reports included in the un-specified category can be omitted from the Tanana Area total because of similar reporting errors. A substantial harvest reduction from previous years is evident.

Five trophy lake trout are recorded from the Tanana Area, three taken in Harding Lake, one each in Fielding and Upper Tangle lakes.

Burbot:

Participation in burbot fisheries has increased in recent years. The majority of the AYK Region harvest has occurred in waters of the Tanana Area. Local residents using baited setlines or hand-held fishing gear are the primary participants. Most fishing in the Tanana River near Fairbanks occurs during the winter months while in the upper Tanana River drainage, a major portion of the annual harvest occurs in spring and summer. Burbot fishing occurs in streams, such as the Tanana, Chena and Tolovana rivers, and in lakes. In past years, the most heavily fished lakes were Fielding, Harding, and Tangle lakes. Since 1987, bag limits in these lakes were reduced to two fish daily, and use of set lines was eliminated. The Tanana River supports one of the largest burbot fisheries in the state, and only the burbot fisheries in the Glennallen area are of comparable effort and harvest levels. Although exploitation rates of burbot in the Tanana River are not considered excessive, studies suggest low stock abundance in most of the lakes examined. Population density of burbot in lakes declined dramatically in the early 1980's due to unsustainable rates of sport fishing exploitation. Studies in Fielding Lake indicate that significant recovery of the population has taken place, probably due to minimum fishing pressure (Lafferty et al. 1991). Burbot stocks in the Tanana River are exploited most heavily near population centers such as Fairbanks, Delta Junction, and near Northway. Burbot movements within the Tanana River tend to minimize effects of concentrated local fishing effort, and stocks in the Tanana River appear to be lightly exploited (Evenson 1990a, 1991b).

To prevent further declines in burbot populations in lakes of the Tanana drainage, the Department implemented emergency regulations in 1987 that prohibited the use of set lines from 15 May to 15 October, and reduced the bag and possession limit in all Tanana drainage lakes to five fish. Also, a ban on the use of set lines throughout the entire year was enacted for Harding, Fielding, T, and Tangle lakes along with a further reduction in the bag and possession to two burbot daily in these waters. The estimated harvest of burbot in the Tanana River drainage by sport anglers in 1990 was 3,799 fish.

The majority (2,093 fish) of the harvest was taken in the Tanana River and the lower Chena River (Table 4).

Of the 178 trophy burbot registered in Alaska, (minimum size 3.6 kg, 8 lbs) 90 (51%) were taken in the Tanana Area, and the majority of those were taken near Fairbanks in the Tanana (55, 31%) and Chena (20, 11%) rivers.

Whitefish:

Most of the statewide recreational whitefish harvest occurs in the AYK Region. The Tanana Area sport harvest of whitefish is almost entirely from the Chatanika River, tributary to the Tolovana River, itself a tributary of the Tanana River, where an active spear fishery occurs in the fall. Hook and line techniques are also used here and in other places to capture whitefish, in which small baited hooks are drifted along the stream bottom.

The estimated (SHS) 1990 harvest of Tanana Area whitefish was 6,891 fish, with more than 5,000 taken from the Chatanika River (Table 4). This represents a substantial harvest reduction (59% from 1989) and is the result of reduced stock abundance and an early closure of the fishery on October 11 by emergency order. Creel survey (Hallberg and Bingham 1991) estimates indicated a harvest of 6,501 whitefish of all species in the Chatanika River spear fishery in 1990, of which 5,396 were least cisco and 957 were humpback whitefish, and 148 were round whitefish *Prosopium cylindraceum* (Table 1, Table 3). Estimates of harvest obtained from the creel survey are thus in substantial agreement with those from the SHS.

The Chatanika River supports spawning populations of humpback whitefish, least cisco, and round whitefish. During late summer and fall, these fish migrate upstream from Minto Flats to spawn. By freeze-up in approximately mid-October, adult whitefish have departed for wintering areas that are as yet unidentified, and which may be located downstream of the Chatanika River. The importance of the Chatanika River as habitat for whitefish other than during spawning and the egg-fry development stages is not fully understood.

Harvest levels in the Tanana Area increased steadily during the 1981 to 1989 period, ranging from 5,449 fish in 1981 to 26,810 fish in 1986. From 1977 until 1989, harvest of whitefish from the Chatanika River increased at an average annual rate of 34%, the fastest growing recreational fishery in the Tanana River drainage (Hallberg and Holmes 1987). Approximately 4,894 angler-hours of effort were expended to spear whitefish in 1990, a decrease of about 18% from the 1989 level. Estimated maximum exploitation rate of humpback whitefish in 1990 was 14%, versus 37% for least cisco. Abundance estimates upon which the calculation of exploitation rates are based were restricted to a section of the Chatanika River near the location of the fishery, and consequently, the abundance estimate cannot be considered germane to the entire river and its stocks of whitefish (Timmons 1990). Since 1987 a daily bag limit of 15 whitefish for the waters of the Tanana River drainage has been in effect. At the time of their implementation, it was hoped that the new regulations would not only reduce harvest but also limit exploitation rates to no more than 20%, a level thought to be sustainable for these species.

Sheefish:

Spawning stocks of sheefish in the Tanana River drainage have been documented only in the upper Chatanika River (Alt 1987). Tagging studies from 1967 to 1971 indicated that sheefish that spawned in the Chatanika River also spent the summer feeding in Minto Flats (Alt 1987). Recaptures in the lower Chena River and at Nenana, of fish that were tagged in the Chatanika River, showed that sheefish disperse widely in the areas adjacent to spawning. Sheefish are widely distributed in the Tanana River drainage during the open water season, from the Tanana River mouth to more than 300 km upstream of Fairbanks. They have also been found at the mouths of the Bearpaw and Toklat rivers in the Kantishna River drainage. Typically sheefish are taken in the lower reaches of clear water tributaries such as the Chena, Chatanika, Tolovana, and Tatalina rivers as well as others. Total sport harvest of sheefish in the Tanana River drainage in 1990 was estimated to be 169 fish (Table 4).

Attempts to stock lakes with sheefish in the Tanana River drainage to create new sport fisheries have met with little success. Harding Lake was stocked nearly annually from 1982 through 1989 (with the exception of 1984). No returns to the sport fishery were documented and stocking for the purpose of fisheries enhancement will discontinue in future years. While growth and survival of stocked sheefish in lakes has been generally disappointing, good growth and survival of stocked sheefish was reported from Four Mile Lake (Figure 3) along the Taylor Highway from stocking events in 1968 and 1969 (Alt 1981). The appearance of new age classes led Peckham and Doxey (1983) to speculate that natural reproduction had occurred.

Rainbow Trout:

Rainbow trout are not indigenous to the Yukon River drainage but have been introduced in several locations, including about 75 Tanana Area lakes since the 1950's. There is no evidence that successful natural reproduction has taken place in any of the stocked locations.

Piledriver Slough has been stocked with rainbow trout since 1987. The slough was formerly connected to the Tanana River and is located about 30 km south of Fairbanks (Figure 2). Water in the slough became clear when the Army Corps of Engineers blocked Tanana River water from entering the upper end at several locations in 1976. The slough was blocked in conjunction with the Army Corps of Engineers Chena Flood Control Project to prevent spillage of high water discharge from the Tanana River into the floodway channel during construction. The temporary dikes are still in place, although they have not been maintained. Piledriver Slough, fed by groundwater from the Tanana River valley, re-established itself as a clear-water tributary to Moose Creek which discharges directly into the Tanana River. Arctic grayling, whitefish and long-nose suckers *Catostomus catostomus* were found inhabiting Piledriver Slough within a year after its upper end was blocked. The objective of stocking was to create a stream rainbow trout fishery in Alaska's interior, thus providing more diversity in angling opportunity for interior anglers. This was the first documented time rainbow trout had been released into flowing waters in interior Alaska since statehood.

The harvest of rainbow trout in Piledriver Slough by sport anglers in 1990 was estimated to be 8,052 fish, while approximately 64,000 rainbow trout were taken in the Tanana Area including Piledriver Slough (Table 4). Angler effort (number of days fishing) on Piledriver Slough in 1990 exceeded 27,000 days, the largest amount of effort for any single water body in the Tanana Area (Table 4).

Substantial harvests of rainbow trout occur in Quartz, Birch, and Chena lakes. The largest harvest occurred in Quartz Lake, where an estimated 20,847 rainbow trout were taken (Table 4). The Tanana Area harvest of rainbow trout in 1988 of more than 78,000 fish represented an historical high value; the 1990 harvest estimate of 64,000 fish was about 18% smaller (Table 4). The steady rise in harvest of this species reflects the successful expansion of the stocking program in the Tanana River valley.

Trophy rainbow trout (minimum size 6.8 kg, 15 lbs) have not been recorded from interior Alaska streams or lakes, and most of the registered trophy fish are native anadromous rainbow trout (steelhead) taken in coastal streams. Nevertheless, good growth rates and size have been achieved in some enhanced lake situations. The largest rainbow trout recorded in the Tanana Area was taken in 1980 from Quartz Lake at a size of 4.5 kg (9.8 lbs). Rainbow trout exceeding 2.3 kg (5 lbs) are commonly taken from area lakes.

ARCTIC, YUKON, AND KUSKOKWIM AREA DESCRIPTION

Excluding the Tanana River drainage which comprises a little over 10% of the land area of the AYK Region, the AYK Area consists of some 870,000 km² (58% of the entire land area of Alaska) of extremely varied topography, climate, and zoogeography. Land ownership and jurisdictions fragment this huge area into a complex mosaic. The federal government is the major land manager through its jurisdiction over lands in National Parks and Preserves, National Wildlife Refuges, Wild and Scenic Rivers, as well as other classifications of federal lands. Native corporations, State of Alaska and private lands comprise the rest, approximately 40% of the landmass of the state. The State of Alaska, by virtue of the Statehood Act retains authority to manage fisheries and wildlife on all lands and waters of the state (approximately 42,300 ha, compared to 88,100 ha under federal jurisdiction). For purposes of reporting and organizing statistics in the SHS, the AYK Area is subdivided into five sub-areas; Yukon (Y), Kuskokwim (V), Seward Peninsula-Norton Sound (W), Northwest Alaska (X), and North Slope of the Brooks Range (Z) (Figure 1).

Geographic and Geologic Setting

Dominant features of the huge landmass that lies north of the Alaska Range divide include the Alaska Range itself which provides water for streams in the Kuskokwim drainage and to the Tanana River and its tributaries. The Brooks Range and its drainages supplies water to the Noatak, Kobuk, Colville, Koyukuk, and Porcupine rivers as well as to many other streams that drain directly into the Yukon River or the Arctic Ocean and the Chukchi Sea.

Lake and Stream Resources

Sport fishing waters and opportunities are extremely varied as could be expected in an area so large and diverse. In the following section the primary fishing waters and species of interest will be briefly characterized in each of the five sub-areas within the AYK Area. It is recognized that not all streams, lakes, or fish stocks of importance receive attention in this cursory treatment.

Yukon River Sub-area:

The Yukon is the largest river in Alaska and its drainage constitutes the fifth largest in North America. The river originates in the basin and range domain of the southern Yukon Territories and northern British Columbia, and flows over 3,700 km northwest to its mouth on the Bering Sea coast. Additional Canadian flows to the upper Yukon River watershed are added from glacial streams such as the White River which originates in the Wrangell and St. Elias Mountain ranges. Approximately one-third of the Yukon River watershed is in Canada. The total drainage area of the Yukon River is approximately 855,000 km², including the area in Canada. Approximately three-fourths of the land area of the AYK Region is in the Yukon River drainage. The entire mainstem of the Yukon River up to the confluence of the White River (Figure 8) in Canada is turbid from glacial silt entrained in the waters draining the Alaska, St. Elias, and Wrangell Mountain ranges.

The Yukon sub-area (statewide harvest Area Y; Figure 1) includes drainages of the Yukon River from the south side of the Brooks Range to the Bering Sea, from Naskonat Peninsula north to Pastol Bay; from the Canadian border west to the Bering Sea. This sub-area does not include any portion of the Tanana or Kuskokwim river watersheds. Map figures that include the Yukon River and major drainages include Figures 7 - 13. It should be noted that prior to 1990 the Lower Yukon and Kuskokwim rivers were combined into one sub-area for reporting purposes henceforth separate harvest reporting will be utilized for the two rivers.

The primary flowing waters of the sub-area are the mainstems of the Yukon River and its tributaries. Clear water streams with sport fishing potential that are part of the Yukon River drainage are extremely numerous and extend to third and fourth order tributaries. Although the main stem of the river flows for approximately 3,200 km, (with the upper third in Canada) this report includes only Alaskan waters. Beginning near the Yukon River mouth, (Figure 9) the east and west forks of the Andreafsky River are both high quality sport fishing streams and have been designated as Wild and Scenic Rivers (Wild and Scenic Rivers Act 1968; ANILCA 1980). All the Pacific salmon species, with the exception of sockeye salmon *Oncorhynchus nerka*, occur in the rivers as do Arctic grayling, Dolly Varden *Salvelinus malma*, and northern pike (in sloughs and lakes off the rivers). Each fork of the Andreafsky River is in itself a major stream and they drain extensive remote areas of the Nulato Hills between the Yukon River Delta and Norton Sound.

The lower Yukon River provides a migratory corridor for all the species of resident, anadromous, and semi-anadromous fishes of the drainage. In




Figure 7. Middle Yukon River and Porcupine River drainages.




Figure 8. Major highways in interior Alaska.




Figure 9. Lower Yukon River and eastern Norton Sound drainages.




Figure 10. Middle Yukon River and Koyukuk River drainages.




Figure 11. Upper Koyukuk River and North Slope Haul Road.




Figure 12. Porcupine River drainage.




Figure 13. Chandalar River system.

addition, many species, such as sheefish, northern pike, several whitefish species, burbot and longnose suckers utilize the mainstem lower river for rearing and feeding, particularly in winter months. For some species such as burbot, the mainstem provides year-round habitat.

The Innoko River and its tributaries drain a large area of flat wetlands and foothills of the Kuskokwim Mountains. The confluence of the Innoko River with the Yukon River is near the village of Holy Cross. The Innoko River system contains numerous northern pike and whitefish as well as other species. A small sockeye salmon stock may spawn in the system, in addition to chum *Oncorhynchus keta*, chinook and coho salmon, but there is no evidence that the Innoko River is important for salmon production when compared to other known productive streams in the Yukon River drainage.

The Anvik River, which enters the Yukon River near the village of Anvik about 515 km upstream from the mouth, is a highly productive stream. The river courses eastward from its drainage area in the Nulato Hills for about 130 km and although it is primarily a rapid runoff stream, artesian upwelling helps stabilize winter flows and water temperatures. Besides supporting the largest chum salmon spawning stock in the Yukon River drainage, with over a million individuals spawning in some years (Whitmore et al. 1987), the stream supports large numbers of chinook and coho salmon, Arctic grayling, Dolly Varden, and some northern pike. Fishing quality is excellent, but few anglers use the stream during the summer season because of its remoteness and difficult access.

The Kaltag and Rodo river mouths, and Bishop Creek mouth support sheefish and Arctic grayling fisheries during summer and early fall months. Sport fishing for northern pike is common in sloughs and lakes near the main stems of the Yukon and Koyukuk rivers as well as in the extensive Kaiyuh Flats southeast of Galena.

The Nowitna River, whose confluence with the Yukon River is upstream from Ruby, is a major clear tributary which enters the Yukon from the south and drains the north slope of the Kuskokwim Mountains. It was designated as a Wild and Scenic River in 1980 (Alaska National Interest Lands Conservation Act, P.L. 96-487), and supports a significant amount of recreational fishing. Most sport fishing is done by Fairbanks residents using personal riverboats or aircraft to reach the river. Good angling for sheefish, northern pike and Arctic grayling can be found in the system, which consists of several branches. Most of the main stem and major tributaries are included in the Nowitna National Wildlife Refuge (USFWS 1987b). Sheefish spawn in the Sulukna River tributary (Alt 1987).

Few lakes of sufficient area or depth to influence winter flow volume or temperature are present in the Alaska Yukon River drainage. The majority of the lakes in the lower Yukon drainage developed as a consequence of the thawing of saturated permafrost soils and as a result, these lakes are mostly shallow and not supportive of primary sport species such as lake trout. There are thousands of such lakes in the deltas and floodplains of the drainage. Many provide summer feeding and rearing for various whitefish species, as well as for northern pike and occasionally, sheefish. Fish utilizing shallow thaw

lakes for summer feeding generally move into primary tributaries and main stems of the major rivers prior to freeze-up.

A major portion of the Yukon River drainage along the south slope of the Brooks Range is contained within the boundaries of the Gates of the Arctic National Park and Preserve. Most of the streams in the sub-area drain to the south from the Brooks Range into the Porcupine Koyukuk, and Yukon rivers (Figures 10, 11, 12, and 13). Significant flowing waters include the Alatna River, and other Koyukuk River tributaries such as the Gisasa, Kateel, Dulbi, Huslia, Indian, Kanuti, Hogatza, Dakli, Henshaw, John, Wild, North Fork, Tinayguk, South Fork, Middle Fork, and Jim rivers. To the east are the Dall, Hodzana and Hadweenzic rivers, the Chandalar River with several tributaries and forks, the Christian River, and the lower Porcupine River with tributaries such as the Sheenjek, Coleen, Black, and Little Black rivers. The Dalton Highway (North Slope Haul Road) bisects the sub-area in a north-south direction (Figure 11), and provides access for recreational fishermen to several streams of the area, including the Ray River, the Middle Fork and South Fork of the Koyukuk River, as well as Prospect Creek and Jim River of the upper Koyukuk River system.

The Nulato River enters the Yukon River near Nulato, about 775 km from the mouth of the Yukon River. Smaller and more difficult to navigate than the Anvik River, the stream nevertheless has sport fishing potential for salmon, Arctic grayling, Dolly Varden and northern pike. The stream receives some seasonal sport fishing use at the present time from anglers stationed at a U.S. Air Force station in Galena.

The mouth of the Melozitna River supports fisheries for sheefish and Arctic grayling during summer and early fall. Geothermal hot springs occur on one of the creeks of the Melozitna River, and a permanent lodge there caters to hunters and fishermen. The Melozitna River is also utilized frequently by local fishermen for Arctic grayling, particularly in the lower 16 km below rapids which effectively isolate the upper reaches of this stream.

The Koyukuk River, one of the largest first order tributaries of the Yukon, enters the Yukon River downstream from Galena, about 820 km from the mouth (Figures 10 and 11). The main stem of the Koyukuk River is turbid in its lower reaches from tannic stain, bank erosion and leaching. Lower Koyukuk River tributaries such as the Gisasa, Kateel, Dulbi, and Indian rivers are not well known outside of the local area but seasonally provide good sport fishing opportunities. Sheefish are taken at the mouths of several streams including the Kateel and Dulbi rivers and where John Junior Slough meets the Koyukuk River about 32 km upstream from the mouth. Arctic grayling are common in clear tributary streams and local residents of nearby villages as well as military personnel stationed at the Galena Air Station fish for them. Sport fishing for northern pike takes place commonly in sloughs and lakes near the main stems of the Yukon and Koyukuk rivers as well as in the extensive Kaiyuh Flats southeast of Galena. Since sheefish spawn in the main stem of the Koyukuk River near Hughes, there are both immature and adult mature prespawning individuals present in the lower Koyukuk River throughout the summer prior to the September spawning period (Alt 1987).

Other Yukon River tributaries below the Porcupine River confluence that support sport fishing include the Tozitna River, Ray River, Dall River, Hodzana River, Hadweenzic River, Chandalar River (Figures 7 and 13), and Christian River (Figure 12). Upper Koyukuk River tributaries that cross the Dalton Highway (North Slope Haul Road) are illustrated in Figure 11.

The Yukon Flats is an extensive wilderness wetland between Circle and Stevens Village below the confluence of the Porcupine River. Thousands of shallow thaw lakes have developed throughout the flats. Many of the lakes support fish populations, at least seasonally, especially those with occasional connections to the sloughs and streams in the area. Northern pike and whitefish species are most common to this area, but sheefish and Arctic grayling also occur in some waterways (USFWS 1985). Salmon production is very limited in the Yukon Flats proper. The area between Beaver Village and Fort Yukon on the Yukon River main stem is known to support sheefish spawning (Alt 1987). This sub-area contains approximately half of the Yukon Flats National Wildlife Refuge, (Figures 13 and 20) and there are literally thousands of lakes present of various sizes and origins. The lakes of the area are categorized roughly (USFWS 1985) into: (1) foothill lakes (formed from streams, with sufficient depth for fish habitation); (2) tundra lakes (which are shallow and often freeze to the bottom); and (3) lowland lakes of three types: oxbow lakes with river connections and deep enough to support fish, mud lakes (shallow, and suitable only for fish rearing), and lakes created from beaver activity. There is currently little documentation available on resident fishes that utilize these vast wetlands. The U.S. Fish and Wildlife Service is conducting inventories and lake surveys in the waters of the Yukon Flats Refuge to provide information on this subject.

The Porcupine River (Figure 12) is the largest Yukon River tributary, draining an immense area of the eastern Brooks Range through the Sheenjek and Coleen rivers, the British Mountains through the Old Crow River, the Richardson Mountains in Canada through the Bell, Eagle, and Rock rivers, and the northern Ogilvie Mountains in north-central Yukon Territories through the East Porcupine Fork and its tributaries. The Black River which drains the southeastern slopes of the Ogilvie Mountains is one of its major Alaskan tributaries. The Little Black River drains a lowland area south of the Black River parallel to the main stem of the Yukon River.

Other major tributaries between Fort Yukon and the Canadian border are all above Circle City, and include the Charley, Seventymile, and Fortymile rivers on the south side of the Yukon River and the Nation, Kandik, and Tatonduk rivers entering the north side of the Yukon River (Figure 8). Parts of Birch and Beaver creeks as well as parts of the Charley and Fortymile rivers are designated as National Wild and Scenic Rivers (Appendix A). The major species for sport fishing on the rivers upstream of Fort Yukon are Arctic grayling in the upper stream reaches and northern pike in the lower, slower sections.

Several large mountain lakes are present. They include Iniakuk, Wild, Big, Twin, Chandalar, Ackerman, and Old John lakes (Figures 11, 12 and 13). All are believed to contain lake trout populations as well as Arctic grayling and other species of whitefish and cisco in most cases.




Figure 14. Waters of the lower Kuskokwim River valley.




Figure 15. Waters of the upper Kuskokwim River valley.




Figure 16. Waters of the Seward Peninsula.




Figure 17. Kotzebue Sound and surrounding area.




Figure 18. Waters of the Noatak River.




Figure 19. Waters of the upper Kobuk River.




Figure 20. Federal land designations within the A-Y-K region.

Kuskokwim River Sub-area:

The Kuskokwim River sub-area (Statewide harvest area V, Figure 1) includes the Kuskokwim River watershed and all waters flowing into Kuskokwim Bay; adjacent salt water from Cape Newenham north to the Naskonat Peninsula (north of Nelson Island) and Nunivak Island².

The Holitna River is the most productive stream for sport fishing in the Kuskokwim River drainage (Figure 14) above the Aniak River confluence, because of the diversity and abundance of its resident and anadromous species. Approximately six fishing guides provide services on the river to about 75 clients per year (Rue et al. 1987). No permanent lodge or tourist structures exist on the river. The Holitna River supports populations of Dolly Varden, Arctic grayling, northern pike, burbot, sheefish, various whitefish species and all five pacific salmon species. Rainbow trout do not occur in the Kuskokwim River drainage upstream of the Aniak River. Dolly Varden, coho salmon, and chinook salmon are the primary sport fish species in the Holitna River, although feeding sheefish are present in the summer as far upstream as the Hoholitna River (Alt 1987) and are sought by some anglers. The Stony, Swift, Gagaryah, Tatlawiksuk, Cheeneetnuk, and Hoholitna rivers represent some of the other important middle Kuskokwim River tributaries. All originate in the Alaska Range and its foothills (Figure 14). Fishery resources of these streams are incompletely documented, and because of remoteness and limited access, they are thought to receive only light recreational use from sport anglers.

Upstream of McGrath, (Figure 15) in the upper Kuskokwim drainage, there are many tributaries that originate in the Alaska Range, such as the Big River, the Middle, South, Windy, Big Salmon, Slow, and East forks of the Kuskokwim River, as well as the Tonsona and Little Tonsona rivers and Highpower Creek near Telida. The North Fork, Nixon Fork, and Takotna rivers originate in the Kuskokwim Mountains west of the Kuskokwim River. Chinook, coho, and chum salmon spawn in streams of the upper Kuskokwim drainage, as do sheefish. Sheefish spawning has been documented in Big River and in Highpower Creek (Alt 1987). Although most of the primary sport fish species occur in the middle and upper Kuskokwim River drainage, (with the exception of rainbow trout), sport fishing effort is extremely light on most streams and fish stocks. Most fishing exploitation in the middle and upper part of the drainage occurs in local subsistence fisheries that mainly target salmon and whitefish.

Lake development in the Kuskokwim River drainage above the Aniak River is sparse, and there are few documented lakes with high potential for recreational fisheries. The fisheries resources in two lakes (Telequana and Two; Figure 14) in the upper Stony River were surveyed by Russell (1980), and Whitefish Lake in the upper Hoholitna River was surveyed in 1977 by Baxter (1977). Lake trout, Arctic grayling, northern pike, and various whitefish species were present in all lakes surveyed. Dolly Varden were noted in Two

² The Sport Fish Division assigns management responsibility for Kuskokwim Bay and Kuskokwim River waters upstream to Aniak to its Southcentral Region headquartered in Anchorage. Responsibility for these areas is assigned to Sport Fisheries staff stationed in Dillingham.

and Telequana lakes but not in Whitefish Lake. Recreational angling occurs in Telequana and Two lakes, both by guided and unguided fishermen. Little information is available regarding sport fishing opportunities and species available in other lakes of the drainage. Most of the lakes in the upper drainage are shallow tundra lakes, unsuitable for supporting year-round resident fish populations. Big Lake near Sleetmute, and Tundra and Trout lakes near Lime Village on the Stony River support year-round fish populations because of their size.

Seward Peninsula/Norton Sound Sub-area:

The Seward Peninsula-Norton Sound sub-area (statewide harvest Area W; Figure 1) includes all waters north of the Yukon River drainage and south of the Selawik River-Kotzebue Sound area and west of the Yukon-Koyukuk River drainages. This area includes Pastol Bay and all salt water north and west of it in Norton Sound as well as salt water adjacent to the Seward Peninsula, including Spafarief Bay in Kotzebue Sound and the southern half of Eschscholtz Bay (ADF&G 1984).

Primary sport fishing streams in eastern Norton Sound (Figure 9) include several that drain the Nulato Hills which separate Norton Sound from the Yukon and Koyukuk River valleys. They include the Unalakleet, Shaktoolik, Inglutalik, and Ungalik rivers. The Unalakleet River is the largest and most heavily utilized of these, and it supports a sport fishery during summer months. A permanent lodge is established on the lower Unalakleet River, and guide service is available from it and other sources. The river and its tributaries support populations of Arctic grayling and Dolly Varden as well as chinook, coho, and pink salmon *Oncorhynchus gorbuscha*. Other area streams also support those species, but are not as intensively fished, primarily because of the limited access and facilities available to non-local fishermen. The Koyuk River main stem carries an abundance of entrained material, including tannic stain, reducing water clarity. The stream terminates in Norton Bay at the extreme eastern corner of Norton Sound. It offers little potential for sport fishing except for northern pike and Arctic grayling in some clear water tributaries.

Several high quality sport fishing streams are located along the southern half of the Seward Peninsula from Koyuk to Teller, (Figure 16) including the Tubutulik, Kwiniuk, Fish, Niukluk, Bonanza, Eldorado, Nome, Snake, Sinuk, Pilgrim, Agiapuk, and Kuzitrin rivers. Road access from Nome exists to many of these streams. Arctic grayling, Dolly Varden, and coho salmon occur in these streams, and chinook salmon, pink salmon, chum salmon, burbot or northern pike are found in many others. Small, perhaps remnant, sockeye salmon stocks are also present in the Pilgrim and Sinuk rivers. Trophy Arctic grayling, larger than 1.4 kg (3 lbs) are present in many streams on the Seward Peninsula, including the Sinuk, Nome, American, Tubutulik, Fish, Pilgrim and Kuzitrin rivers as well as others. Many of the largest Arctic grayling recorded as trophies for Alaska have been taken from streams on the Seward Peninsula. Of the 125 largest fish registered from 1967 to 1990 in the ADF&G trophy fish program, 28 were taken in waters of the Seward Peninsula. Nineteen of the 28 registered trophy Arctic grayling from the Seward Peninsula were taken from the Sinuk River.

Most of the streams draining the northern half of the Seward Peninsula have low sport fishing potential due to relatively small flow volumes, difficult access, and poorer quality of water and fisheries habitat.

Most of the lakes on the Seward Peninsula were created either by thaw action in river floodplains or by glaciers in the mountains of the central and western Seward Peninsula. The largest inland body of water on the Seward Peninsula is Imuruk Lake (Figure 17) in the north-central portion of the peninsula. It is approximately 32 km² in area, and was probably formed when volcanic lava originating in the nearby area cut off drainage streams causing water to back up into a local depression. The lake presently drains northward through the Inmachuk River. Salmon spawn at the outlet in the fall and the lake supports whitefish and Dolly Varden.

Other lake waters with recreational fishing potential are smaller glacial lakes in the Imuruk Basin watershed and in the Kigluaik Mountains east of Nome. Some contain populations of lake resident Arctic char *Salvelinus alpinus*, (Kretsinger 1987) while other lakes and streams in this area contain anadromous Dolly Varden. Salmon Lake, located about 150 km northeast of Nome in the headwaters of the Pilgrim River, contains Dolly Varden, Arctic grayling, round whitefish and a remnant stock of sockeye salmon. Since the lake can be reached by road from the town of Nome, it receives use for sport fishing, and during the first half of the century was an important recreation and fishing area for gold miners in the area. Subsistence fishing for salmon in Salmon Lake has been prohibited for many years because the stock was practically eliminated by early fisheries. Salmon sport fishing in the lake and it's tributaries is presently prohibited.

Northwest Alaska Sub-area:

The Northwest Alaska Area (statewide harvest Area X; Figure 1) includes all waters and drainages of the Kotzebue area, including drainages of the Selawik, Kobuk, Noatak, Wulik, and Kivalina rivers. The area also includes all salt water in the northern half of Eschscholtz Bay, including the Chamisso Island area and the northern half of Kotzebue Sound to and including Point Hope (ADF&G 1984). The eastward limit of the sub-area extends to the Alatna River.

The most important streams of the Northwest Alaska sub-area (Figures 17, 18 and 19) are the Noatak and Kobuk rivers, both of which drain large areas of the southern slope of the western Brooks Range. Each has a drainage area of approximately 31,000 km² and stream length of from 560 km (Kobuk) to 640 km (Noatak; U.S. Army Corps of Engineers 1967). The third largest drainage is that of the Selawik River, with an approximate area of 11,700 km². The Noatak River is slightly turbid at most times during the summer months from entrained glacial silt carried from mountain glaciers in the Brooks Range, while waters of the Kobuk and Selawik rivers are more clear. Abundant groundwater resources are found in both the Noatak and Kobuk rivers as water-bearing gravel aquifers on the lower main stem of the Noatak River and in tributaries of the Kobuk River. These aquifers tend to stabilize flows and water temperature fluctuations and provide water storage within the systems.

The Noatak River is designated as a National Wild and Scenic River, and most of the drainage is included in the Noatak National Preserve (Figure 18). The extreme upper headwaters of both the Noatak and Kobuk rivers are included in the Gates of the Arctic National Park. A part of the lower Kobuk Valley between Kiana and Ambler is included in the Kobuk National Park, and the Salmon River tributary, as well as the upper main stem of the Kobuk River are National Wild and Scenic Rivers as is the Selawik River. Much of the Selawik River valley is part of the Selawik National Preserve.

These three large river systems contain abundant fisheries resources. The Noatak River produces a large run of late chum salmon that are the primary species for the Kotzebue-based commercial fishery. Many thousands of anadromous Dolly Varden overwinter and spawn in the river. Whitefish and northern pike are resident in the Noatak River. Alt (1987) reports that sheefish use the river for feeding but do not spawn there.

Both the Selawik and Kobuk rivers support spawning populations of sheefish in their upper main stems. The brackish delta systems which have formed at the river mouths serve as overwinter feeding areas for juvenile as well as adult sheefish. Trophy sheefish are taken from these waters especially in the upper Kobuk River during the fall when large mature spawners congregate near spawning areas in the main stem. The Alaska state record sheefish was taken in 1986 from the upper Kobuk River (mouth of the Pah River) and weighed 24 kg (53 lbs). Abundant whitefish utilize the rivers and delta areas, including Selawik Lake and Hotham Inlet (Kobuk Lake). Dolly Varden occur in some tributaries to the Kobuk River, as do northern pike in sloughs and connecting lakes to the lower river. Lake trout inhabit deeper lakes of the upper Kobuk River watershed. The Wulik and Kivalina rivers, which empty into the Chukchi Sea near the village of Kivalina, are well known as trophy streams for Dolly Varden.

Sport fishing effort in Northwest Alaska is relatively light compared to most other areas in the state. Heaviest use occurs on the Noatak, Kobuk, and Wulik rivers. Many visitors to Gates of the Arctic National Park and Kobuk Valley National Park participate in float trips on the Kobuk River from Walker Lake to Kobuk village (Alt 1984; ADF&G 1986; NPS 1984, 1985a). A lodge on Walker Lake promotes lake trout and Arctic char fishing. A small amount of shore fishing with hook and line for sheefish takes place near Kotzebue in the summer.

Guided and unguided anglers and river floaters use the Noatak River as do Kotzebue area residents who boat to different parts of the river to fish. The most popular fishing area is the Kelly River, but other tributaries such as the Nimiuktuk and Kugururok rivers are also used occasionally primarily for Dolly Varden fishing (Alt 1978).

Raft, canoe, and kayak trips are increasingly popular recreational uses on the Noatak River. Many parties put in at Matcharak Lake (Figure 18) and portage to the river, or land wheel aircraft on gravel bars farther upstream. Arctic grayling, Dolly Varden and lake trout are available in the upper Noatak River, but downstream from the Nimiuktuk River, chum salmon and northern pike also occur. Lake trout occur in Matcharak, Feniak, and Desperation lakes and in

other lakes in the middle and upper Noatak drainage. Most lakes in the area are accessible only by floatplane.

The lower floodplains of the Kobuk and Selawik rivers, especially in the vicinity of the Kobuk River delta, and the lower Noatak River (upstream of the lower canyon of the Noatak) contain hundreds of shallow thaw lakes of various sizes. Fisheries resources of the waters in this area are poorly inventoried, but large populations of whitefishes, northern pike, and sheefish are known to be seasonally present. The mountains in the upper Kobuk River drainage (Figure 19) contain several relatively large, deep lakes which were formed by glacial action. Lake trout, Arctic grayling, Arctic char and perhaps one or two whitefish species inhabit these lakes. They include Walker Lake, Nutuvukti Lake, Minakokosa Lake, Lake Selby and Narvak Lake.

Lakes of the upper Noatak River (Figure 18) were surveyed by Alt (1978), with a brief inventory of 13 lakes in the upper drainage. Fish were present in all lakes surveyed, and round whitefish lake trout and Arctic grayling were the most common species. Least cisco, northern pike, Dolly Varden, slimy sculpin *Cottus cognatus*, salmon (chum and sockeye), and ninespine stickleback *Pungitius pungitius* were also found.

North Slope Brooks Range Sub-area:

The north slope of the Brooks Range sub-area (statewide harvest Area Z; Figure 1) includes all waters north of the Brooks Range divide flowing into the Beaufort and Chukchi Seas from Point Hope on the west to the Canadian border on the east including adjacent saltwater areas.

The northern-most part of Alaska is characterized by its broad Arctic coastal plain, which abuts the Arctic Ocean and Beaufort Sea, and by the foothills and mountains which form the Brooks Range (Figure 21). The central and eastern Brooks Range consist of rugged, glaciated, east-trending ridges with summits rising to elevations of 4,350 to 5,000 m. The Delong Mountains on the western flank of the Brooks Range consist of glaciated ridges, 1,865 to 2,500 m in elevation, which drain northward into the Chukchi Sea. Only a few small lakes and no glaciers exist in the Delong Mountains although they were glaciated during the ice age. From the central and eastern Brooks Range, the mountain rivers flow northward to the Beaufort Sea. As Selkregg (1976) points out, although several large rock basin lakes lie at the mouths of glaciated valleys on both sides of the range, there are surprisingly few lakes for a glaciated area. Although most of the streams that cross the foothills flow northward from their sources in the range, the region's largest stream, the Colville River flows eastward for more than 320 km before turning north onto the coastal plain (Figures 21 and 22). The drainage area of the Colville River is about 62,000 km², a little more than half of the area drained by the Tanana River. Most streams east of the Colville River are braided and cross broad gravel flats that are often blocked in winter by auffs (fields of ice that form continuously downstream from spring water sources) that cause local flooding (Selkregg 1976). The upper valleys of major rivers flowing from the Brooks Range often contain morainal lakes.




Figure 21. Waters of the Arctic Slope.




Figure 22. Principal tributaries of the Colville River.

The coastal plain is an area of low relief and poor drainage due to underlying permafrost and a shallow active layer, factors that lead to moisture entrapment near the surface. Rivers that cross the plain originate in the hills or mountains to the south. In the west, more than half of the plain is covered by oriented thaw lakes aligned to the north-northwest on their long axes. Ice-wedge polygons are found throughout the coastal plain section.

Major flowing waters of the coastal plain from west to east, include the Kukpowruk, Utukok, Kuk, Meade, and Itpikpuk rivers (Figure 21). The Colville River has several major tributaries, including the Killik, Chandler, Anaktuvuk, and Itkillik rivers (Figure 22). Streams east of the Colville River include the Kuparuk, Sagavanirktok (Figure 23), Canning, Hulahula, and Kongakut rivers (Figure 21).

The North Slope is accessible by air travel or by driving the Dalton Highway north from Fairbanks. The highway was originally built in 1974 to support construction of the Trans-Alaska oil pipeline. Sport fishing was closed for 8 km (5 miles) on either side of the pipeline beginning in 1978 to prevent rapid fisheries depletion by construction workers along the Trans-Alaska Pipeline. The closure was rescinded in 1980 by the Alaska Board of Fisheries when the Haul Road Corridor was opened for fishing for all species except sheefish and salmon (Bendock 1980). Since June 1981, the highway south of Disaster Creek (near Dietrich, Figure 11) has been open to travel by the general public (Bendock 1982). The Alaska Board of Fisheries opened the Haul Road Corridor for sheefish fishing in 1987, but salmon fishing remains closed.

The Dalton Highway crosses tributaries of the Sagavanirktok, Toolik, and Kuparuk rivers and parallels the Sagavanirktok River for about 160 km (100 miles) south of Prudhoe Bay, providing access to sportfishing opportunities for Arctic grayling, lake trout, Dolly Varden and Arctic char (Figure 11). There are numerous small lakes between Ribdon River (Figure 23) (tributary to the Sagavanirktok River) and Galbraith Lake (Figure 11) that provide good fishing opportunities for these species .

Light sportfishing effort also takes place on the Colville, Kongakut, Canning, Ivishak, Echooka, Killik, and Anaktuvuk rivers, as well as the Hula Hula River (ADF&G 1986). Some fishing may also be done by parties floating the Nigu and Etivluk rivers (Bendock 1983). A significant proportion of the sportfishing in this sub-area is by persons engaged in hunting activities as their main pursuit.

Lakes such as Toolik Lake in the upper Kuparuk River and Galbraith Lake in the upper Sagavanirktok River are two of the most frequently fished lakes north of Atigun Pass (Bendock and Burr 1984). Other popular lakes along the Haul Road include Island, Campsite, and Tea lakes. Lakes outside the Dalton Highway Corridor that receive sportfishing effort include Elusive, Shainin, Itkillik, Cascade, Kurupa, and Chandler lakes (Figures 11, 13, and 22; Furniss 1974; NPS 1985a).




Figure 23. Principal tributaries of the Sagavanirktok River.

AYK Area Climate

Because of geographic and topographic diversity and size, annual climatic parameters vary considerably throughout the area. Except for the immediate coastal areas, a continental type of climate prevails over much of the AYK Area, with warm summers, cold winters and little precipitation. Annual precipitation is usually sparse except for the areas under coastal influence. In lower latitudes, the amount of precipitation generally received in the AYK Area would result in arid conditions similar to those found in many deserts of the world. Because of cooler temperatures in Alaska, and the fact that a great deal of moisture is trapped in permafrost soils, desertification has not occurred, and, in fact, much of the area gives the appearance of having received abundant moisture. Typical summer weather lasts from mid-June until late August or early September, with rainy weather typical during August and September. Snowfall is usually light even in the coastal areas of the AYK Area, as compared to other areas of the state.

Species of Importance to the Sport Fishery

In addition to the species listed for the Tanana Area, Dolly Varden and Arctic char are important to sport fishing in many waters of the AYK Area. Wild stocks of rainbow trout occur as far upstream in the Kuskokwim River drainage as the Aniak River and its tributaries. Rainbow trout do not occur naturally north of the Kuskokwim River, although they have been stocked in two lakes of the AYK Area. Pink salmon are an important sport fish species in the Norton Sound and Seward Peninsula sub-area where there is sport fishing effort in both freshwater and marine waters. Additional species of whitefish that are of importance to fisheries in the AYK Area include the broad whitefish, *Coregonus nasus*, Arctic cisco, *Coregonus autumnalis*, and Bering cisco, *Coregonus laurettae*. All other species listed under the Tanana Area are also found in the AYK Area.

AYK Area Sport Fisheries

A brief description of AYK Area sport fisheries follows.

Chinook Salmon:

Chinook salmon spawn throughout the Kuskokwim and Yukon River drainages and in streams of eastern Norton Sound and the southern Seward Peninsula. Chinook salmon occur in streams north of the Seward Peninsula, but no stocks identified are sufficiently abundant to support commercial or sport fisheries. Concentrated sport fishing occurs in only a few streams in the AYK Area, and the majority of the sport harvest is taken by local residents. More intensive sport fishing occurs in Norton Sound on the Unalakleet River and on the Seward Peninsula in parts of the Fish River system where commercial guiding and lodging facilities have been developed to promote sport fishing for salmon. Guided fishing for chinook salmon also takes place on the Holitna River in the Kuskokwim drainage. The Salmon River (Kuskokwim River tributary) near Nikolai and McGrath also supports a sport fishery on a chinook salmon spawning stock by local residents (Stokes 1985). An undocumented amount of angling for chinook salmon takes place on both the Andreafsky and Anvik rivers. Few

chinook salmon are harvested in the AYK and Tanana Areas compared to other management areas in Alaska (Arvey and Mills 1991; Mills 1991). Since 1980 the AYK Area chinook salmon sport harvest has ranged from about 1,000 to 3,000 fish, (combined large and small sea-run chinook salmon) with the majority taken from streams in the Yukon and Kuskokwim sub-areas (Mills 1991)³. An estimated total of 706 large sea-run chinook salmon were harvested in 1990 in the AYK Area. In addition, an estimated 660 small (less than 71 cm in length) sea-run chinook salmon were harvested in the AYK Area (Tables 6 - 10). When totalled, the estimated 1990 harvest of both small and large chinook salmon in the AYK Area is 1,366 fish.

Coho Salmon:

Coho salmon are distributed widely south of the Brooks Range in the AYK Area, however, they are more abundant in the Kuskokwim River drainage and in drainages to the south than in drainages north of the Kuskokwim River. Returns of coho salmon to the Kuskokwim River may be the largest to a single river in Alaska. Approximately 660,000 coho salmon were harvested in the 1986 Kuskokwim River commercial fishery, historically the largest commercial harvest for this system (Francisco et al. 1987). Western Alaska coho salmon are thought to spawn primarily in spring-fed portions of streams. The upper Kuskokwim River and its tributaries that drain the northern slopes of the Alaska Range are extensively underlain with alluvial gravels as a result of outwash from the Alaska Range. The resulting gravel aquifers provide high quality spring water for spawning and rearing of coho salmon in the Kuskokwim drainage, and fall chum salmon in the Yukon and Tanana River drainages. SHS results in 1990 indicate that the majority of the coho salmon recreational harvest in the AYK Area (4,891 fish) was from streams in the Seward Peninsula-Norton Sound sub-area (3,305 fish), followed by harvests in the Kuskokwim River sub-area (1,346 fish) (Tables 7 and 8). The sport harvest of coho salmon from the Seward Peninsula-Norton Sound sub-area in 1990 was composed of 1,826 fish which were taken from the Unalakleet River (Table 8).

Coho salmon are locally abundant north of the Yukon River in Norton Sound, where coastal and stream fisheries occur at least as far north as Teller. Several streams of eastern Norton Sound (Figures 9 and 16) including the Unalakleet, Shaktoolik, Fish, and Niukluk rivers support spawning stocks of coho salmon, as do many of the streams in the Nome area, Port Clarence, and Safety Sound. Active sport fisheries occur in these areas for coho salmon. The Nome and Unalakleet rivers accounted for the majority of the harvest. Coho salmon occur north of Port Clarence, but as with other freshwater rearing salmon species, their abundance decreases markedly at higher latitudes.

Pink Salmon:

This species rarely undertakes extensive freshwater migrations (more than 160 km) and as a consequence, it is not abundant upstream of the lower main stems of the major river systems of the AYK Area. For example, pink salmon seldom ascend the Yukon River beyond the Anvik River (513 km), or the

³ Reported harvest figures include fish taken from lower Kuskokwim River and Kuskokwim Bay streams.

Table 6. Yukon River sub-area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	KI ^d	KS	SS	CS	LT	DV AC	GR	WF	SF	NP	BB	OTHER
Jim River	305	372	973	0	0	0	0	0	17	660	152	0	0	0	0
Other Haul Road Streams	898	1,067	1,776	0	0	0	0	0	68	1,438	51	17	187	0	0
Yukon River	644	916	2,159	105	0	0	76	0	34	135	0	51	372	85	0
Other Streams	3,443	5,286	7,589	0	0	228	341	0	187	2,440	119	255	1,694	271	0
Haul Road Lakes	68	609	432	0	0	0	0	0	0	186	0	0	0	0	0
Other Lakes	694	1,237	2,610	0	0	0	0	220	85	102	0	0	221	153	0
TOTAL	4,162^e	9,487	15,539	105	0	228	417	220	391	4,961	322	323	2,474	509	0

^a Yukon River Drainage (Area Y): All Yukon River drainages (including the Alaska portion of the White River drainage but excluding the Tanana River drainage), from the south side of the Brooks Range to the Bering Sea; and from the Canadian border to the Bering Sea; and all drainages of the Koyukuk and Alatna Rivers.

^b KS: chinook salmon; SS: coho salmon; LL: landlocked coho or chinook salmon; RS: sockeye salmon; PS: pink salmon; CS: chum salmon; LT: lake trout; DV/AC: Dolly Varden or Arctic char; RT: rainbow trout; GR: Arctic grayling; WF:various whitefish; SF: sheefish; NP: northern pike; BB: burbot; and, SM: smelt.

^c From Mills (1990).

^d King salmon less than 28 inches.

^e Angler totals may not equal sum of sites due to some anglers fishing at more than one site.

Table 7. Kuskokwim River sub-area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	KI ^d	KS	SS	LL	RS	PS	CS	LT	DV AC	RT	GR	WF	SF	NP
SALTWATER:																	
Boat	123	246	226	0	0	12	0	24	0	0	0	18	0	0	0	0	0
Shoreline	71	194	119	0	0	0	0	12	0	14	0	0	0	0	0	0	0
SALTWATER TOTAL	176^e	440	345	0	0	12	0	36	0	14	0	18	0	0	0	0	0
FRESHWATER:																	
Kanektok River	932	1,337	4,525	299	204	644	0	462	145	202	0	1,020	281	123	0	0	0
Aniak River	493	1,214	1,964	95	190	182	0	49	29	159	18	598	35	422	0	0	18
Other Streams	1,568	1,973	8,515	55	54	520	0	73	173	374	0	161	159	442	88	107	213
Lakes	230	299	509	0	0	0	0	0	0	0	54	0	0	353	0	0	0
FRESHWATER TOTAL	2,639^e	4,823	15,513	449	448	1,346	0	584	347	735	72	1,779	475	1,340	88	107	231
GRAND TOTAL	2,691^e	5,263	15,858	449	448	1,358	0	620	347	749	72	1,797	475	1,340	88	107	231

-continued-

Table 7. (Page 2 of 2).

	BB	SM	HA	OTHER
SALTWATER:				
Boat	0	0	144	0
Shoreline	0	0	0	0
SALTWATER TOTAL	0	0	144	0
FRESHWATER:				
Kanektok River	0	211	0	0
Aniak River	0	0	0	0
Other Streams	1,125	0	0	0
Lakes	0	0	0	0
FRESHWATER TOTAL	1,125	211	0	0
GRAND TOTAL	1,125	211	144	0

- ^a Kuskokwim River Drainage (Area V): The Kuskokwim River drainage and all waters flowing into Kuskokwim Bay; adjacent saltwater from Cape Newenham north to the Naskonat Peninsula (north of Nelson Island). Does not include the Yukon, Tanana or Koyukuk River drainages.
- ^b KS: chinook salmon; SS: coho salmon; LL: landlocked coho or chinook salmon; RS: sockeye salmon; PS: pink salmon; CS: chum salmon; LT: lake trout; DV/AC: Dolly Varden or Arctic char; RT: rainbow trout; GR: Arctic grayling; WF: various whitefish; SF: sheefish; NP: northern pike; BB: burbot; SM: smelt; and, HA: halibut.
- ^c From Mills (1991).
- ^d King salmon less than 711 mm (28 inches).
- ^e Angler totals may not equal sum of sites due to some anglers fishing at more than one site.

Table 8. Seward Peninsula-Norton Sound sub-area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	KL ^d	KS	SS	RS	PS	CS	DV AC	GR	WF	NP	BB	SM	OTHER
SALTWATER:																
Boat	348	381	1,322	10	10	81	0	69	41	17	0	0	0	0	0	99
Shoreline	348	978	1,644	10	0	221	105	597	68	166	0	0	0	0	166	415
SALTWATER TOTAL	647^e	1,359	2,966	20	10	302	105	666	109	183	0	0	0	0	166	514
FRESHWATER:																
Nome River	1,559	3,979	6,609	29	0	384	0	2,637	95	1,078	33	0	0	0	0	0
Pilgrim River	779	1,144	1,627	0	19	81	93	208	41	166	415	0	1,194	33	0	0
Unalakleet River	465	1,625	3,957	57	219	1,826	0	1,180	298	614	99	0	0	0	332	0
Fish-Niukluk River System	580	829	2,059	0	0	267	0	638	216	348	415	0	17	0	0	0
Other Streams	1,460	3,966	4,468	0	10	445	0	2,318	166	1,227	416	299	746	0	497	0
Lakes	84	84	113	0	0	0	0	0	0	332	0	0	0	0	0	66
FRESHWATER TOTAL	2,686^e	11,627	18,833	86	248	3,003	93	6,981	816	3,765	1,378	299	1,957	33	829	66
GRAND TOTAL	3,233^e	12,986	21,799	106	258	3,305	198	7,647	925	3,948	1,378	299	1,957	33	995	580

^a Seward Peninsula-Norton Sound (Area W): All drainage area north of the Yukon River drainage, including all saltwater north and west of Pastol Bay in Norton Sound; and, south of the Selawik River drainage. Does not include the Selawik River.

^b KS: chinook salmon; SS: coho salmon; RS: sockeye salmon; PS: pink salmon; CS: chum salmon; DV/AC: Dolly Varden or Arctic char; GR: Arctic grayling; WF: various whitefish; NP: northern pike; BB: burbot; and, SM: smelt.

^c From Mills 1991.

^d King salmon less than 711 mm (28 inches).

^e Angler totals may not equal sum of sites due to some anglers fishing at more than one site.

Kuskokwim River beyond the Holitna River (540 km). Estimated 1990 harvest of AYK Area pink salmon was 7,994 fish, of which all but 347 were taken in the Seward Peninsula/Norton Sound sub-area (Table 5).

Pink salmon are a target species in Norton Sound sport fisheries where harvest estimates have ranged from 1,100 (1985) to more than 13,000 (1982). The recent (1981-1990) ten year average sport harvest is 4,948 fish. Even-year harvests for the five years of this period averaged 7,159 fish, while harvests during odd years averaged 2,738 fish. Pink salmon utilize numerous streams in Norton Sound for spawning, and in some years large returns are documented. Extremely large returns have been recorded for the Unalakleet River, the Kwiniuk River and Tubutulik River (Lean et al. 1986). For example more than 6 million pink salmon are estimated to have spawned in the Unalakleet River in 1984 (Lean 1985). Returns of this species are extremely variable even in more southerly latitudes. The common two-year cycle of pink salmon return abundance is not as pronounced in Norton Sound and in more northerly areas, although the magnitude of return variation is large because of climatic variation at higher latitudes. Snow cover, depth, and mean monthly temperatures during the winter months all affect ice thickness and the amount of frozen versus unfrozen groundwater in local streams of the area. Salmon egg and fry survival rates are directly affected by the amount of freezing in the stream gravels where incubation takes place.

Non-rearing species such as pink and chum salmon may have an advantage in northern streams where freshwater productivity is much lower than in lower latitudes. Sport and subsistence fisheries in the Nome area for pink salmon in local streams such as the Nome, Snake and Sinuk rivers are active in late July and early August.

Arctic Grayling:

Arctic grayling inhabit waters throughout the AYK Area where they are perhaps the most popular and ubiquitous sport fish species. Most of the freshwater drainages that have been surveyed along the Arctic coast contain Arctic grayling (USFWS 1982). It is the principal species inhabiting foothill lakes and streams on the Seward Peninsula. Arctic grayling occur in lakes and streams emptying into the Chukchi Sea between Kotzebue and Barrow (ADF&G 1978, 1986). Arctic grayling also migrate through, or inhabit during parts of the year, main stems and tributaries of the largest rivers such as the Yukon, Kuskokwim, Porcupine, Koyukuk, Kuskokwim, Noatak, Kobuk and Colville rivers.

Arctic grayling typically spawn in smaller, clear headwater streams with gravel bottoms and low stream gradients, usually during May and early June. After spawning, the adults disperse throughout the streams for summer feeding. Juveniles and sub-adults frequently feed during summer months in far upstream reaches that become dewatered in winter. They overwinter in lower stretches of tributaries where water and oxygen concentrations are adequate, as well as in lakes and spring-fed portions of streams. Summary descriptions of distribution, life history and abundance of Arctic grayling in the AYK Region are provided in Alaska Habitat Management Guides for the Interior, Western and Arctic Regions (ADF&G 1986).

SHS results indicate that since 1977, the Arctic grayling harvest in the AYK Area has ranged from about 9,000 fish in 1990 to more than 30,000 in 1983. Since 1983, estimated harvests have declined, and have ranged from 15,500 in 1984 to 17,215 in 1989, with 9,092 being taken in 1990 (Table 5). The Arctic grayling harvest is usually taken fairly evenly from the five sub-areas, with from about 1,000 to 9,000 Arctic grayling taken in each sub-area (Tables 6 - 10). The largest harvest estimate in 1990 (4,961 fish) was for the Yukon River sub-area (Table 6). The 1990 harvest distribution in each of the AYK sub-areas was less even than usual.

Northern Pike:

Sloughs, interconnected lakes, and the lower sections of large rivers throughout most of the AYK Area are inhabited by northern pike. Lowland areas of the Yukon and Kuskokwim rivers are particularly noted for large northern pike. Northern pike are abundant in all parts of the AYK Area containing appropriate habitat except on the north slope of the Brooks Range, where their distribution is limited. Bendock and Burr (1985) reported the presence of northern pike on the Arctic coastal plain west of the Colville River, in rivers and lakes draining into Admiralty and Smith Bays (Figure 21) and in middle reaches of the Killik River, tributary to the Colville River.

During summer, northern pike are generally distributed near shore in shallow water that contains aquatic vegetation and a mud bottom. Northern pike have some tolerance for salinity and they are taken frequently in brackish waters of the Yukon-Kuskokwim Delta. They are not known to feed or travel extensively in marine or coastal waters outside the major rivers. During winter, northern pike congregate in deep, well-oxygenated waters found in the lower reaches of tributaries or other areas of sufficient water flow (Hallberg 1984).

Most of the recreational harvest of northern pike is taken with hook and line. Spearing, bow and arrow, and hand jigging techniques are also legal means and account for a small proportion of the total harvest. Northern pike sport fishing occurs in the Kuskokwim River drainage from McGrath to downstream of Bethel, including the Takotna, Nixon Fork, Holitna, and Johnson rivers. Most recreational fishing for northern pike along the Yukon River takes place upstream from Galena. Popular areas include the Yukon Flats near Fort Yukon, Koyukuk River, Beaver Creek, Birch Creek, Dall River, Hess Creek, Tozitna River, Melozitna River, and Nowitna River. The Pilgrim, Kuzitrin, and Fish rivers in the Nome area are popular as well.

Much of the recreational and some of the subsistence harvest in the AYK Area is taken during winter months through the ice with hook and line gear. Sport fishing for northern pike has gained in popularity since the early 1960's. Northern pike are eagerly sought by fishermen in areas that have good boat access. They are often fished in the fall in combination with hunting activities.

The estimated sport harvest of northern pike in the AYK Area has ranged from about 2,000 fish in 1977 to more than 8,600 fish in 1983 (Mills 1979-1991). The estimated harvest in 1990 was 4,982 fish (Table 5). Generally, the

largest harvests have been taken in the Yukon and Kuskokwim river sub-areas. Estimated 1990 harvests from these sub-areas was 2,474 and 231 northern pike respectively. In 1990, an estimated harvest of 1,957 northern pike was taken in the Seward Peninsula/Norton Sound sub-area. (Tables 6 - 10).

Little is known concerning the status of northern pike stocks in the AYK Area, but because of remoteness and restricted access, fishing effort is light except on those stocks near towns and villages where angling and subsistence gill netting effort may be more intense. Northern pike populations close to the Yukon River Haul Road Bridge have experienced more angling pressure because the recent opening of the road has allowed easy boat access to people living in the Fairbanks area. Northern pike population studies conducted in the Tanana River drainage suggest that abundance and stock composition parameters such as age and size composition respond negatively and rapidly when annual harvest exploitation rates exceed 16%.

Lake Trout:

Historically, approximately 35% of the total AYK Region harvest of lake trout has occurred in the AYK Area. In 1986, the percentage taken outside the Tanana River drainage was much larger (88%) due to declines in catch and perhaps abundance within the Tanana River drainage. Harvest in 1990 for the entire region totalled 1,743 fish⁴, of which 847 fish (49%) were taken from the AYK Area (Table 5). Harvests for the AYK Area since 1977 have ranged from about 500 fish in 1977 to about 2,500 in 1986 (Table 5).

ADF&G studies indicate that lake trout resources have been overharvested in many of the more accessible waters of south-central and interior Alaska. Specific life history features (slow growth, delayed maturity, and non-consecutive spawning) combined with the short growing season at higher latitudes and altitudes increases the vulnerability of the species to overharvest.

Burr (1987) described the distribution of lake trout in Alaska. Lake trout are most frequently associated with deep, oligotrophic lakes in the mountains and are rarely found at lower elevations of the Yukon or Kuskokwim basins (Redick 1967; Morrow 1980). In northwest Alaska, lake trout occur in lakes and streams of the Brooks Range in the Noatak and Kobuk River drainages. Lake trout are found in most drainages that flow into the Yukon River from the Brooks Range. Lake trout distribution is primarily restricted to lakes at higher elevations in these drainages. Lake trout are widely distributed on the north slope of the Brooks Range. They occur most frequently in mountain and foothill lakes, but they also occur in streams of the Colville, Sagavanirktok, and Canning River drainages. Lake trout generally do not occur in the lowland lakes of the Arctic coastal plain, but they occur commonly in central coastal plain lakes between the Ikpikpuk and Colville rivers.

ADF&G has conducted little research on this species in areas outside of the Tanana Valley and near Glennallen. Lake trout research is being conducted by federal agencies, such as USFWS, NPS (National Park Service), and BLM (Bureau

⁴ See remarks on harvest estimate accuracy for the Tanana Area, page 25.

Table 9. Northwest Alaska sub-area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	KL ^d	KS	SS	RS	PS	CS	LT	DV AC	GR	WF	SF	NP	BB
SALTWATER:																
Shoreline	34	101	100	0	0	0	0	0	0	0	0	0	0	0	0	0
SALTWATER TOTAL	34	101	100	0	0	0	0	0	0	0	0	0	0	0	0	0
FRESHWATER:																
Kobuk River	420	286	1,306	0	0	0	0	0	0	17	17	67	0	151	151	0
Noatak River	504	621	1,306	0	0	0	0	0	0	168	218	269	84	0	17	0
Other Streams	269	338	719	0	0	0	0	0	0	0	302	286	0	0	17	0
Lakes	135	118	351	0	0	0	0	0	0	202	269	0	0	0	135	0
FRESHWATER TOTAL	991^e	1,363	3,682	0	0	0	0	0	0	387	806	622	84	151	320	0
GRAND TOTAL	1,008^e	1,464	3,782	0	0	0	0	0	0	387	806	622	84	151	320	0

^a Northwest Alaska (Area X): Kotzebue area including drainages of Selawik, Kobuk, Noatak, Wulik, and Kivalina Rivers. All saltwater in the northern half of Kotzebue Sound to and including Point Hope.

^b KS: chinook salmon; SS: coho salmon; RS: sockeye salmon; PS: pink salmon; CS: chum salmon; LT: lake trout; DV/AC: Dolly Varden or Arctic char; GR: Arctic grayling; WF: various whitefish; SF: sheefish; NP: northern pike; BB: burbot; and, SM: smelt.

^c From Mills 1990.

^d King salmon less than 711 mm (28 inches).

Table 10. North Slope Brooks Range sub-area^a sport fish harvest and effort by fisheries and species^b, 1990^c.

	Anglers	Trips	Days Fished	PS	CS	LT	DV AC	GR	WF	SF	NP	BB	SM
SALTWATER:													
Boat	34	17	34	0	0	0	0	0	0	0	0	0	0
Shoreline	152	303	285	0	0	0	102	0	0	0	0	0	503
SALTWATER TOTAL	185^d	320	319	0	0	0	102	0	0	0	0	0	503
FRESHWATER:													
Sagavanirktok River	218	621	493	0	0	0	101	151	0	0	0	17	0
Other Haul Road Streams	252	353	399	0	0	17	0	118	0	0	0	0	0
Other Streams	437	740	1,474	0	0	0	168	438	17	0	0	0	0
Haul Road Lakes	269	369	852	0	0	101	0	67	0	0	0	0	0
Other Lakes	151	118	227	0	0	50	118	17	0	0	0	0	0
FRESHWATER TOTAL	1,057^d	2,201	3,445	0	0	168	387	791	17	0	0	17	0
GRAND TOTAL	1,158^d	2,521	3,764	0	0	168	489	791	17	0	0	17	503

^a North Slope Brooks Range (Area Z): All Alaskan waters, including drainages, north of the Brooks Range and flowing into the Beaufort and Chukchi Seas to the north and east of Point Hope. Does not include Point Hope.

^b PS: pink salmon; CS: chum salmon; LT: lake trout; DV/AC: Dolly Varden or Arctic char; GR: Arctic grayling; WF: various whitefish; SF: sheefish; NP: northern pike; BB: burbot; and, SM: smelt.

^c From Mills 1991.

^d Angler totals may not equal sum of sites due to some anglers fishing at more than one site.

of Land Management) in National Wildlife Refuges, National Parks, National Preserves and other unclassified lands.

Dolly Varden/Arctic Char:

The majority of the harvest of Arctic char or Dolly Varden (collectively referred to as char in this section) in the region occurs in the AYK Area since only the dwarf stream resident form is found in the upper Yukon and Tanana River systems. Char occur in most of the waters of western and Arctic Alaska, either in the anadromous, river resident, lake resident, or stream dwarf forms. Char are a target species for subsistence and sport fisheries in waters of the Arctic, Kotzebue vicinity, Seward Peninsula, and Norton Sound.

Two species are recognized within the AYK Area based upon meristic characters, (gill raker and pyloric caecae counts), life history features and the occurrence of anadromy (Behnke 1980). Dolly Varden are the dominant species from Bristol Bay north to the Arctic coast, and occur in either the anadromous, stream resident or stream dwarf form, while Arctic char occur only as residents in foothill lakes of the North Slope (Chandler Lake Campsite Lakes, etc; Figures 11 and 22), the Kobuk River drainage (Walker and Selby lakes; Figure 18), the Seward Peninsula (lakes of the Kigluaik Mountains; Figure 16), and the Kuskokwim Mountains (Aniak, Kisaralik, Kagati, and Goodnews lakes; Figure 14). The majority of char in the AYK Area are Dolly Varden of either the stream resident or anadromous type. They occur throughout the area but are most abundant in tributaries of the lower Yukon and Kuskokwim rivers, Norton Sound, Northwest Alaska, and along the north slope of the Brooks Range and the Arctic coastal plain. Morrow (1980) distinguishes two distinct forms of Dolly Varden in Alaska based upon gill raker and vertebral counts. The southern form usually occurs only south of the Alaska Range, however examination of specimens collected in the upper reaches of the Copper, Tanana, Nenana, and Susitna Rivers indicates that non-anadromous southern and northern forms occur on both sides of the Alaska Range passes. Morrow (1980) argues that headwater transfer, which may still be occurring, is responsible for the mixing of southern and northern forms in these areas.

Sport harvests of char in the AYK Area between 1977 and 1990 have ranged from approximately 4,000 fish to more than 20,000 fish (Table 5). This represents from less than 5% to about 20% of the statewide total harvest (Mills 1979-1991). The estimated harvest in 1990 was 7,431 fish from the AYK Area, with the majority of the harvest in the Seward Peninsula-Norton Sound sub-area (Tables 6 - 10). Typically, the largest percentage of the harvest is taken in the Seward Peninsula-Norton Sound sub-area. High quality sport fishing for char is available in northwest Alaska, particularly in the Wulik, Kivalina, and Noatak rivers north of Kotzebue, when trophy Dolly Varden move into the streams for either overwintering or spawning in the fall. The Noatak River in northwest Alaska supports populations of spawning char in its tributaries. Important spawning tributaries include the Kelly River, Kogururok River, and the Nimiuktuk River (DeCicco 1985). Incidental commercial harvest as well as directed subsistence harvests account for the highest proportion of the annual fishing mortality in northwest Alaska in most years (Bernard and DeCicco 1987).

During the period from 1967 to 1990, 23 trophy char (26% of the Alaska total) were registered from the AYK Area, including the state record char, (7.9 kg; 17 lbs 8 oz) taken from the Wulik River in 1968. All except one of the trophy char taken in the AYK Area came from northwest Alaska.

Burbot:

Burbot are distributed throughout the AYK Area in all major rivers and many of the lakes and minor waterways. It is an important fisheries resource for subsistence economies of rural Alaska and for sport fisheries as well. Burbot are members of the cod family, *Gadidae*, and spawn in midwinter under the ice of rivers and lakes. Sport fishing interest and intensity has increased for the species in recent years, particularly near settlements where burbot fishing provides an outdoor wintertime activity for many people. Reported annual sport harvests of burbot in the AYK Area since 1978 have ranged from just over 100 fish to approximately 2,000 fish (Table 5). The majority of the harvest and fishing effort occurs in the winter with lines set through the ice, although hand-held lines with rod and reel are also used in summer and winter months. The majority of the burbot sport harvest in the AYK Region takes place in the Tanana River drainage. The estimated 1990 AYK Area harvest of burbot was 1,684 fish (Table 5).

Only three of the 178 Alaska trophy burbot registered from 1967 to 1990, were taken in the AYK Area, a fact that is no doubt reflective of lack of sport fishing effort in much of the area.

Whitefish:

Although members of the whitefish family, *Coregonidae*, are seldom considered to have substantial recreational value, as a group they constitute an extremely important fisheries resource in the AYK Area. They are taken year around by subsistence fisheries and are utilized for human consumption as well as for dog food. In addition, various whitefish species provide a forage base for many of the predatory fishes that support important sport fisheries such as northern pike, burbot, and sheefish. The most important whitefish species in northern, western and interior Alaska include the humpback whitefish, broad whitefish, round whitefish, Arctic cisco, least cisco, and Bering cisco.

Recreational fisheries throughout the area account for a very small proportion of the total harvest of all species of whitefish. An estimated harvest of 810 whitefish was taken by sport anglers in 1990 in the AYK Area (Table 5). The magnitude of the whitefish subsistence harvest is not well documented except in a few specific instances, but is believed to be orders of magnitude greater than the sport harvest in the AYK Area. Where salmon are not available, and during the winter months, whitefish are the major source of fish for subsistence use. Such a situation prevails over the entire North Slope, and in many of the remote villages of interior Alaska located beyond the upstream limits of the salmon runs. Limited commercial fisheries for whitefish exist in the AYK Area.

Sheefish:

Sheefish are large, predatory whitefish found throughout western, interior, and northwestern Alaska. They do not occur in streams of the North Slope nor in Norton Sound north of the Koyuk River. Alt (1987) identified nine stocks of sheefish, with anadromous-estuarine stocks occurring in the Kuskokwim, lower Yukon, Koyuk, Kobuk-Selawik rivers, and resident non-anadromous stocks in Yukon River tributaries of the Nowitna, Tanana River (Minto Flats), Porcupine, and Salmon Fork of the Black River, as well as the main stem of the upper Yukon River. The distribution of this species in Alaska is limited to the AYK Region.

Sheefish are harvested by subsistence, commercial, and recreational users with subsistence harvests exceeding all others. The major commercial fishery (Kotzebue Sound) is limited by a harvest quota of 11,350 kg or approximately 3,300 fish annually (Lean 1985). Reported sales of sheefish from this commercial fishery have only once (in 1977-78) reached the allowed quota (Lean et al. 1986). During the period from 1978-1990, estimates of the recreational harvest in the AYK Area ranged from about 600 to about 3,600 fish (Table 5). The harvest estimate for the AYK Area in 1990 was 581 sheefish, the lowest since the survey was initiated in 1977. Kobuk and Selawik River sheefish are the most abundant and heavily fished stocks. Harvest occurs on a year-round basis from subsistence, commercial and sport fishermen, from spawning grounds in the upper Kobuk River to the coastal inlets, Hotham Inlet and Selawik Lake near Kotzebue and Selawik villages. The data base for sheefish stocks of the Kobuk and Selawik rivers is not adequate to allow precise estimates of sustainable yields for the stocks. The size of the spawning stocks has been poorly documented as has the harvest and biological composition of the harvest. Because of life history features that favor late maturation, slow growth, nonconsecutive spawning, and the existing harvest pressure on major stocks, there is concern that the Kobuk and Selawik river stocks may be experiencing harvests beyond sustainable levels. The question cannot be addressed until further biological information becomes available.

Sheefish generally overwinter in lower reaches of rivers and in estuarine waters, migrate upstream in summer to feeding grounds, and migrate further upstream to spawning grounds in the late summer and fall. Migrations of over 1,650 km have been documented. Sheefish spawn in late September and early October at water temperatures of 0 - 5°C. Aerial survey estimates of the number of spawning sheefish on the Kobuk and Selawik rivers are available for seven years between 1966 and 1979. The highest aerial survey count for the Kobuk River was 8,166 spawning sheefish in 1971 (Alt 1987). A total of 1,243 spawners was observed in the Selawik River in 1968, one of two years when surveys were made on this stream. It is not certain that spawning grounds have been identified for all major sheefish stocks. Availability of spawning habitat with desired current (2 m per sec), water depth (2 m), and bottom substrate of differentially-sized gravels may be the most critical factor limiting sheefish distribution and abundance (Alt 1987).

Rainbow trout:

In only two known instances have rainbow trout been introduced into waters of the AYK Area. Approximately 3,000 fingerling were released into a man-made, landlocked reservoir known as Webster Reservoir in Prudhoe Bay in 1977. Fate of the transplanted fish was not assessed in subsequent years. Since the reservoir is used as an industrial water source in the winter, water levels are often drawn down severely, increasing the likelihood of winter kill from hypoxia or freeze-out. It is assumed that the transplant failed and that no surviving fish remain.

In a second instance, rainbow trout were released into Roy's Pond near Central in 1987. The pond is a 15.38 ha (38 ac) waterbody with an outlet stream that does not flow except at extreme high water, into Crooked Creek, tributary to Birch Creek in the Yukon River drainage. The 1987 release consisted of 10,000 fingerling rainbow trout. The success of the stocking in producing catchable rainbow trout in the pond in subsequent years has not been assessed. During an ADF&G fisheries survey of Mammoth Creek, tributary to Crooked Creek, in 1990, a rainbow trout adult of about 235 mm in fork length was captured (Townsend 1991). There are anecdotal reports that other rainbow trout were captured in Crooked Creek also. There is therefore strong evidence that the rainbow trout planted in Roy's Pond have escaped into the Birch Creek drainage of the Yukon River.

In addition to authorized releases of rainbow trout, there may have been accidental or purposeful, but unauthorized stockings taking place. An anecdotal report from a trapper (Cheney pers. comm. ⁵) in the Farewell area, near McGrath, indicates that rainbow trout may inhabit portions of the South Fork of the Kuskokwim River near Farewell. The lower Jones River and the South Fork above the confluence with the Jones River are reported to contain the species. These fish may represent escaped fish from an unauthorized stocking into Farewell Lake.

STAFF ORGANIZATION

Regional Staff Responsibilities

Organization of the region staff is outlined in Figure 24. All activities were directed by the Regional Supervisor (J. Clark) who delegated appropriate tasks to the Administrative Assistant (E. Nielsen), the Research Supervisor (P. Merritt) and the following Fishery Biologist III's: W. Arvey, J. Hallberg, and M. Kramer. Lake stocking activities were the responsibility of M. Doxey. Area management responsibility and emergency order authority was vested with four positions, W. Arvey (AYK Area), J. Hallberg (Lower Tanana Area), F. Parker (Upper Tanana Area), and F. DeCicco (Northwest Alaska Area). Each area manager conducted fisheries research projects in the respective areas. The AYK Area biologist was responsible for a study of Dolly Varden near the North Slope Haul Road; the Lower Tanana River Area biologist

⁵ Cheney, Lee. 1991. Personal Communication. ADF&G, F.R.E.D. Division, Big Lake Hatchery, P.O. Box 520509, Big Lake AK 99652.




Figure 24. Organizational chart for A-Y-K Sport Fish staff, January 1990.

conducted creel survey studies in the Fairbanks area; the Upper Tanana Area biologist conducted studies of lake burbot, and northern pike; and the Northwest Area biologist conducted studies on Arctic grayling and Dolly Varden. Special research was conducted by W. Ridder on stock assessment of Arctic grayling in the Salcha, Chatanika and Goodpaster rivers, and in the Tangle lakes, all in the upper Tanana Valley. Arctic grayling research in the Salcha, Chena, and Chatanika rivers near Fairbanks, and in Fielding Lake near Delta was conducted by R. Clark. Char research was conducted in northwest Alaska by A. DeCicco, who also directed Arctic grayling studies on the Seward Peninsula. Lake trout research in the Tanana River drainage and Copper River was conducted by J. Burr. Burbot research in rivers of the upper Tanana Valley was conducted by M. Evenson. F. Parker conducted a study of burbot in lakes of interior Alaska. Northern pike studies were conducted by G. Pearse and A. Burkholder in the Tanana River drainage. Evaluation of the fish stocking program was conducted by C. Skaugstad, M. Doxey, and J. Clark. Adult chinook salmon abundance in the Salcha and Chena rivers was estimated by A. Burkholder and M. Evenson. Fisheries information and education as well as the fisheries access program were the duties of M. Kramer.

Synopses of Published Reports

The intent of the following section is to provide a brief overview of AYK Region fisheries studies completed during the reporting year, however some of the reports are from work completed in other years. Since the majority of AYK Region activities are documented by technical reports following each field season, the report itself is cited, followed by a brief synopsis.

Arvey, W. D., M. J. Kramer, J. E. Hallberg, J. F. Parker, and A. L. DeCicco. 1991. Annual Management Report for Arctic-Yukon-Kuskokwim Region, 1987. Alaska Department of Fish and Game. Fishery Management Report No. 91-1. 139 pp.

Important species to the sport fisheries in this region are identified, zoogeographic distribution of species is discussed, as is stock status and harvest data from the 1987 calendar year. Regulatory actions affecting the regional sport fishery in 1987 are described along with other management, research, stocking and enhancement activities. Climatic factors of importance to area fisheries are summarized and federal land status within the region is described.

Arvey, W. D., and J. F. Parker. 1991. Annual Management Report for Arctic-Yukon-Kuskokwim Region, 1988. Alaska Department of Fish and Game. Fishery Management Report No. 91-2. 117 pp.

Important species to the sport fisheries in this region are identified, zoogeographic distribution of species is discussed, as is stock status and harvest data for calendar year 1988. Regulatory actions affecting the regional sport fishery in 1988 are described along with other management, research, stocking and enhancement activities. Climatic factors of importance to area fisheries are summarized and federal land status within the region is described.

Arvey, W.D. 1991. Stock status of anadromous Dolly Varden in Waters of Alaska's North Slope. Alaska Department of Fish and Game. Fishery Data Series No. 91-3. 20 pp.

Estimates of overwintering abundance in key streams of the North Slope are presented, along with estimates of the sport fishery harvest, the subsistence harvest, and mortality from research activities on the North Slope. Sport fishery harvest are estimated to range from about 1,000 to about 3,000 Dolly Varden annually.

Burkholder, A. 1991a. Abundance and composition of northern pike, Harding Lake, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-9. 33 pp.

Abundance of fish greater than 449 mm fork length was estimated to be 1,283 fish. Density was estimated to be 1.28 fish per hectare. Age six northern pike comprised 33% of the sample while seven year old fish comprised 31%. The oldest northern pike was 10 years of age. Study results suggested that the existing northern pike population could not sustain the levels of recreational harvest estimated in recent years.

Burkholder, A. 1991b. Stock composition of northern pike captured in Minto Flats during 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-23. 19 pp.

A total of 3,752 northern pike were captured in the spring, and 264 were captured in the fall. Insufficient numbers of fish captured in the fall sampling event precluded the completion of an abundance estimate in 1990. Age two (two years old) fish were most abundant in the spring sampling while fish of age six were most abundant in the fall sampling event.

Burkholder, A. 1991c. Abundance, egg production, and age-sex-length composition of the chinook salmon escapement in the Salcha River, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-5. 32 pp.

The estimate of abundance was 10,728 fish. Age composition estimates indicated that 49% of the spawners were six years old, having spent four years in the ocean, and 25% were five years old, having spent three years at sea. Aerial surveys of spawner abundance indicated 3,744 fish present during the highest count, about 35% of the mark-recapture point estimate. Egg production from the escapement was estimated at 52 million eggs (standard error = 2.7 million).

Burr, J. M. 1991. Lake trout population studies in interior Alaska, 1990, including abundance estimates of lake trout in Glacier, Sevenmile, and Paxson lakes during 1989. Alaska Department of Fish and Game. Fishery Data Series No. 91-7. 50 pp.

Size composition was widely variable between lakes. Most large lake trout (greater than 715 mm fork length) were found in Paxson Lake. A few large fish

were captured in Fielding Lake, but most fish from the Tanana River drainage were less than 500 mm. Ages ranged from 0 to 33 years.

Abundance for 1989 was estimated using mark-recapture experiments in Sevenmile, Glacier, and Paxson lakes. Abundance of fish greater than 275 mm fork length was 1,665 fish (50 fish per hectare) in Sevenmile Lake and 3,142 fish (18 per hectare) greater than 250 mm in Glacier Lake. Estimated abundance of spawning lake trout in Paxson Lake was 5,066 fish (three fish per hectare).

Clark J. H. 1991. Abundance, length, and age composition of sockeye salmon, Arctic char, and least cisco in pelagic waters of Harding Lake, Alaska, 1989 and 1990.. Alaska Department of Fish and Game. Fishery Data Series No. 91-4. 25 pp.

Applying the area-swept method to 1989 tow net catches, the presence of 17,278 sockeye salmon and 28,984 least cisco was estimated in the pelagic zone of Harding Lake. Survival rate of sockeye salmon stocked as fry in June 1989 through the first four months of residence was estimated to be 3.3%. Survival rate of fingerling sockeye salmon from the fall of 1988 until fall of 1989 (12 months) was estimated to be 2.7%. In 1990, it was estimated that zero sockeye salmon, 61,891 least cisco and 998 stocked Arctic char resided in the pelagic zone of Harding Lake. The sockeye salmon fry stocked in 1988, 1989, and 1990 had evidently died before fall 1990.

Clark J. H., Viavant T.R., Skaugstad C. and McKinley T.R. 1991. Growth, survival, and costs of rearing game fish in floating net-pens at Harding Lake, Alaska, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-2. 25 pp.

Arctic grayling and sockeye salmon stocked as sac fry grew fastest (18 % and 15% weekly gain in length) but also had the lowest survival (less than 10%). Arctic grayling and rainbow trout stocked in net-pens as fingerlings grew 11% and 10% in length per week, with survival of 99%. Rainbow trout stocked at an average length of 215 mm grew an average of three percent weekly with no mortality. Arctic char fingerlings grew an average of five percent weekly with greater than 99% survival. Lake trout fingerlings grew three percent weekly with survival at 98%. The average cost of producing 1 kilogram of fish in the net-pen facility was \$15.

Clark R. A. 1991. Stock assessment of Arctic grayling in Fielding Lake. Alaska Department of Fish and Game. Fishery Data Series No. 91-11. 62 pp.

Abundance estimates (fish greater than 199 mm fork length) ranged from 4,512 fish in 1987 to 8,837 fish in 1988. Recruitment ranged from 895 fish to 5,043 fish during the five year period. Onset of sexual maturity of Fielding Lake Arctic grayling was at four years (fifth summer) and 265 mm fork length. Available data suggest a relatively stable Arctic grayling population in Fielding Lake, although causative mechanisms of variability in recruitment are unknown at the present.

Clark R. A. 1991. Stock status of Chena River Arctic grayling during 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-35. 58 pp.

Estimated abundance of Arctic grayling in the lower 152 km of the Chena River was 31,815 fish greater than 149 mm fork length. Annual recruitment between 1989 and 1990 was 16,881 Arctic grayling and survival during this period was 75.4%. Although recruitment and survival rates appear to be increasing, increases in recreational fishing effort and harvest have caused a decline in the abundance of mature (age five and older) fish. Annual harvests of mature Arctic grayling must be reduced to prevent further declines in abundance.

Clark R. A. Fleming, D. M. and W. P. Ridder. 1991. Stock assessment of Arctic grayling in the Salcha, Chatanika, and Goodpaster rivers. Alaska Department of Fish and Game. Fishery Data Series No. 91-15. 85 pp.

Population abundance in a 36.8 km section of the Salcha River was 5,792 fish greater than 149 mm fork length. Population abundance in the lower 50 km of the Goodpaster River was 7,113 fish greater than 149 mm fork length. Age 3 fish were most abundant in the sections sampled in all three rivers. All historical data on age, size, and sex compositions, harvest and effort, and population abundance from 1952 to 1990 are presented.

DeCicco A. L. 1991. Seward Peninsula Arctic grayling study, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-24. 45 pp.

Estimated abundance of Arctic grayling greater than 149 mm in fork length in a 16 km section of the Niukluk River was 3,739 fish or 234 fish per km. Fork length of fish captured in the Niukluk River averaged 344 mm. Abundance of fish greater than 149 mm fork length in a 24 km section of the Fish River was 2,909 fish or 121 fish per km. Fork length of Fish River Arctic grayling averaged 369 mm. Estimated abundance in a 12 km section of the Pilgrim River of fish greater than 149 mm fork length was 1,764 fish or 147 fish per km. Fork length of Pilgrim River fish averaged 390 mm. The estimated abundance of Arctic grayling greater than 249 mm in fork length in a 40 km section of the Sinuk River during 1989 was 1,120 fish or 28 fish per km. Fork length of fish from the Sinuk River averaged 443 mm. Mean fork length-at-age was greatest in the Sinuk River and least in the Fish and Niukluk rivers.

Evenson M. J. 1990b. Age and length at sexual maturity of burbot in the Tanana River, Alaska. Alaska Department of Fish and Game. Fishery Manuscript No. 90-2. 10 pp.

A total of 351 burbot were collected between November and February. Ages ranged from 4 to 18 years and total length ranged from 368 to 1,076 mm. Onset of sexual maturity was first noted at age 4 and 452 mm for males and age 6 and 498 mm for females. Burbot in nonspawning condition were found in almost all age and length classes sampled, supporting the hypothesis that adult burbot may not spawn each year.

Evenson M. J. 1991a. Abundance, egg production, and age-sex-size composition of the chinook salmon escapement in the Chena River, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-6. 35 pp.

The estimated abundance was 5,603 chinook salmon, 2,633 of which were females and 2,970 males. The highest count of chinook salmon during aerial spawner counts was 1,436, about 26% of the mark-recapture point estimate. Estimated egg production for the 1990 escapement was about 25 million eggs.

Evenson M. J. 1991b. Abundance and size composition of burbot in rivers of interior Alaska during 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-33. 44 pp.

Abundance of burbot 400 mm and larger in a 40 km section of the Tanana River near Northway was estimated at 4,803 fish, and 1,966 fish for a 24 km section of the lower Chena River. Estimates of mean total length were 552 mm for the Chena River section, 617 mm for the Northway section, and 553 mm for a 32 km section of the Tanana River near Fairbanks. Catch-per-unit of effort data are presented.

Fleming D. F. and W. P. Ridder 1991. Stock assessment of Arctic grayling in Mineral Lake Outlet, 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-22. 25 pp.

Age and size composition estimates from 1988 and 1990 indicated slight reductions in older or larger Arctic grayling. Maturity was achieved in 50% of sampled fish at 236 mm fork length, or age four.

Hallberg, J. E. and A. E. Bingham. 1991. Creel surveys conducted in Interior Alaska during 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-56. 98 pp.

Catch, effort, harvest, and age and length composition were estimated by creel surveys conducted on Piledriver Slough rainbow trout and Arctic grayling, Harding Lake northern pike, Delta Clearwater River Arctic grayling, lower Chena River chinook salmon, Salcha River chinook salmon and Chatanika River whitefish fisheries.

Lafferty R., Parker, J. F., and D. R. Bernard. 1991. Stock assessment and biological characteristics of burbot in lakes of interior Alaska during 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-57. 72 pp.

Abundance and/or indices of abundance were estimated for populations of burbot in 16 lakes in interior Alaska. Abundance of fully recruited burbot estimated with mark-recapture experiments was greatest in Susitna Lake (4,659 fish) and lowest in T Lake (87 fish). Three independent methods of burbot stock assessment indicate that burbot stocks in lake Louise and Paxson Lake are stabilizing. The Fielding Lake population has recovered significantly since 1987 due to increased recruitment. The population in Sevenmile Lake has not recovered, probably due to high elevation and extreme environmental conditions.

Ridder W. P. 1991. Summary of recaptures of Arctic grayling tagged in the middle Tanana River drainage, 1977 through 1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-34. 97 pp.

Approximately 3,800 recoveries of tagged Arctic grayling were used to summarize the seasonal movements. Recoveries came from releases of about 19,000 fish tagged in 13 locations, encompassing silted rapid run-off and bogged streams to spring-fed systems for feeding. Notable migratory patterns include: from Caribou Creek to the Richardson Clearwater River; and, from the Goodpaster and Volkmar rivers to the Delta Clearwater River. High specificity to feeding areas, especially the Delta and Richardson Clearwater rivers was found.

Skaugstad C. and B. McCracken 1991. Fecundity of chinook salmon, Tanana River, Alaska. Alaska Department of Fish and Game. Fishery Data Series No. 91-8. 23 pp.

Fecundity was estimated for 49 chinook salmon collected from the Tanana River in 1989. Mean fecundities for chinook salmon that spent three, four, or five years in the ocean were 8,500, 9,100, and 11,900 eggs. Fecundity increased with age, however, mean fecundities for fish spending three or four years in the ocean were not significantly different.

Timmons L. S. 1991. Opinions and regulatory preferences of holders of sport fishing licenses resident in the Arctic-Yukon-Kuskokwim area of Alaska in 1988. Alaska Department of Fish and Game. Fishery Data Series No. 91-1. 75 pp.

Motivations, regulatory preferences, and profiles of holders of sport fish licenses were examined using a postal questionnaire. Over half the respondents rated overall enjoyment as excellent. Most respondents were motivated by either obtaining food or non-success reasons, and none of the respondents listed catching a trophy fish as a primary motive for sport fishing. Minimum length limits and time/area closures were approved most often as a means to improve fishing and as emergency regulations. Complete closure was the least preferred regulation.

Timmons L. S. 1991. Evaluation of the rainbow trout stocking program for Piledriver Slough. Alaska Department of Fish and Game. Fishery Data Series No. 91-25. 62 pp.

The number of rainbow trout stocked annually during the 1987-1990 period ranged from 20,000 to 79,481 fish. Fish were stocked at several locations as both fingerlings and subcatchables in size. Evaluation of stocking was by using statewide harvest surveys, creel surveys, and field sampling. After initiation of stocking, Piledriver Slough quickly became one of the most popular sport fishing locations in interior Alaska in terms of effort and, a large number of rainbow trout were harvested. Growth and survival of stocked fish were generally low and only those stocked as catchables contributed significantly to the fishery. In 1990, four percent of the rainbow trout

sampled were gravid females, and none were found that were conclusively age 0 rainbow trout resulting from natural reproduction.

Timmons L. S. and R. A. Clark 1991. Stock status of Piledriver Slough Arctic grayling. Alaska Department of Fish and Game. Fishery Data Series No. 91-37. 44 pp.

French Creek, Moose Creek and Garrison Slough, which are interconnected with Piledriver Slough, were also sampled in this study. Abundance in 1990 was estimated to be 16,435 Arctic grayling over 149 mm fork length. Age composition was estimated.

Viavant T. and J. H. Clark 1991a. Distribution and relative abundance of stocked species in Harding Lake, 1987-1990. Alaska Department of Fish and Game. Fishery Data Series No. 91-20. 31 pp.

Catch-per-net-night of fishing effort was highest for northern pike in the shallow and deep littoral zones, highest for burbot and least cisco in the pelagic zone, and highest for Arctic char and lake trout in the various benthic zones. Of all the introductions made into Harding Lake during recent years, Arctic char stocked as juveniles or adults have been the most successful. Fish stocked smaller than 10 g have not been detected in appreciable numbers because survival of these fish has been negligible.

Viavant T. and J. H. Clark 1991b. Experimental hook and line fishing at Harding Lake, Alaska, 1989-1991. Alaska Department of Fish and Game. Fishery Data Series No. 91-38. 47 pp.

Catch-pr-unit-effort was low for most locations and gear types (generally less than one fish-per-angler-hour). Baited lures were more effective in winter than bait or unbaited lures alone. Green lures were more effective than other colors. Salmon eggs were more effective as bait than shrimp or corn. Bait was more effective in summer sampling than spinners, and worked best in water deeper than 10 m, and when fished near bottom. Catches were low or zero in water greater than 10 m in depth in winter.

MANAGEMENT ACTIVITIES

1990 In-Season Regulatory Measures

ADF&G has limited authority to rapidly enact changes in regulations without formal hearings and passage by the Alaska Board of Fisheries. Such authority is normally reserved for situations in which immediate conservation concerns for resources exist, or where opportunities to harvest an unusually abundant resource, surplus to reproductive needs, would be lost if no action was taken to modify existing regulations.

Emergency Orders and Regulations:

The Alaska State Legislature, in enacting AS 16.05.060, delegated to the Commissioner and his authorized designees the power to control certain aspects

of public utilization of fish and game. In 1990, ADF&G authority for emergency orders was expanded to include bag limits, and methods and means modifications as well as the usual authority for opening and closing fishing seasons or areas. Emergency regulations, via the Administrative Procedures Act ⁶, were required to change quotas, bag limits, size limits or gear restrictions prior to BOF action in the spring of 1990, which gave ADF&G expanded authority. Authority to change size limits by emergency order was not granted by the BOF however. Emergency orders and regulations have the same force and effect as regulations of the Boards of Fisheries and Game, or statutes enacted by the Legislature, and they carry the same criminal penalties if violated. An emergency order remains effective until rescinded, superseded by a subsequent emergency, and/or unless an expiration date is specified (ADF&G 1983). Use of emergency order authority allows ADF&G to take regulatory action in response to unexpected resource scarcity or abundance. Sport fisheries in the AYK Region generally proceed under the published basic regulations. Emergency regulations expire after 120 days unless made permanent via the process outlined in the Administrative Procedures Act.

A number of regulatory changes were enacted during the 1987 calendar year, some by the emergency regulation process, the majority through action by the Alaska Board of Fisheries. A full discussion of regulation development in 1987 may be found in Arvey et al. (1991). Action by the Board of Fisheries on finfish proposals for the AYK Region is scheduled to occur only in alternate years. The next scheduled Alaska Board of Fisheries meeting at which changes in AYK Region sport fish regulations were considered occurred in the winter of 1989-1990. Changes enacted at that time went into effect in the spring of 1990. No emergency regulations were enacted in 1990. Three emergency orders (Appendix B) were issued in 1990 for the AYK Region.

3-Z-S-8-90. Closed the Nome River to the taking of chum salmon on sport fishing gear, on July 2.

3-WF-01-90. Closed the Chatanika River to the taking of whitefish by sport fishermen, on October 11, 1990.

3-NP-02-90. Established 1) a 26-inch minimum length limit for northern pike harvested in Harding Lake, 2) a ban on the use of spears and bow and arrows in Harding Lake and, 3) a closure of Harding Lake to sport fishing for northern pike (during the spawning period) from 12:01 a.m. April 1, to 11:59 p.m. May 31.

News Releases, Announcements, and Articles:

Five news releases (also called field announcements) concerning AYK Region sport fisheries were issued by the Department in 1990. They are listed below.

1. June 1, 1990. Harding Lake Fishery Studies. This news release announced the initiation of the floating net pen fish rearing project. The project reared about 200,000 juvenile fish in the

⁶ Alaska Statute 44.62.180(3)

spring and summer for release in the lake in September. Studies focussed on growth and survival rates of lake trout, rainbow trout, sockeye salmon, Arctic grayling and Arctic char.

2. July 31, 1989. Interior Alaska Stocking Program. The stocking program and the Statewide Stocking Plan for Recreational Fisheries are discussed.
3. August 23, 1990. Harding Lake Fish Release. The release of pen reared fish is announced.
4. October 9, 1990. Fishery Closure. The closure of the Chatanika River to sport fishing for whitefish on October 11, 1990 is announced in this new release. Reason for the emergency closure is the dramatic decline in abundance of humpback whitefish
5. Harding Lake Fishery Restrictions. This news release announced that the northern pike fishery was being restricted by ADF&G emergency order.

Several news articles concerned with AYK Region sport fisheries appeared in major Alaska newspapers or in the Alaska Fish and Game magazine in 1990. They are listed below.

1. January 5-6, 1990. When its cold it's hot. By: Kelly Bostian, Fairbanks Daily News-Miner (FDNM). An article about ice houses in the Fairbanks area.
2. July 13-14, 1990. A better fish trap, biologists build weir for Chatanika whitefish. By: Kelly Bostian, Fairbanks Daily News-Miner (FDNM). An article about Chatanika River whitefish populations, and the new ADF&G research weir being constructed.
3. June 1-2, 1990. The big day is near. FDNM. Announced the upcoming Fishing is Fun Day on June 9, 1990.
4. May-June, 1990. Five Game Fishes of the North: Lake Trout. By John Burr. Alaska Fish and Game Magazine, Volume 22, No. 3. The distribution, and habits of lake trout as well as fishing information is presented.
5. May-June, 1990. Five Game Fishes of the North: Arctic char and Dolly Varden. By Fred DeCicco. Alaska Fish and Game Magazine, Volume 22, No. 3. The distribution and characteristics of both species are discussed.
6. May-June, 1990. Five Game Fishes of the North: Sheefish. By Kenneth Alt. Alaska Fish and Game Magazine, Volume 22, No. 3. Distribution of sheefish populations and fishing information is presented.

7. May-June, 1990. Five Game Fishes of the North: Northern pike. By Gary Pearse and Alan Burkholder. Alaska Fish and Game Magazine, Volume 22, No. 3. The distribution of northern pike in Alaska and fishing information is presented.

Preseason and Post-Season Regulatory Activities

The process of developing appropriate fishing regulations continues each year both during the primary fishing seasons, as well as before and after. The following section describes the salient features of that process in 1990.

Advisory Committees:

Public input concerning regulation changes is provided by several means, including direct testimony to the Board of Fisheries and by participation in local fish and game advisory committees. Local advisory committees have been established throughout the state to assist the Boards of Fish and Game in assessing fisheries and wildlife issues and proposed regulation changes in the areas that will be affected. Most active committees meet at least once a year, usually in the fall prior to Board meetings. Advisory meetings allow opportunity for direct public interaction with Department staff who are expected to attend, answer questions, and provide clarification concerning proposed regulatory changes.

The following local advisory committees for the AYK Region have been established: (1) in Western Alaska: Central Bering Sea, Lower Kuskokwim, Central Kuskokwim, Lower Yukon; (2) in Arctic Alaska: Norton Sound, Kotzebue, Northern Seward Peninsula, Upper Kobuk, Lower Kobuk, Noatak/Kivalina, Western Arctic, Eastern Arctic, St. Lawrence Island, Southern Norton Sound, and; (3) in Interior Alaska: Middle Nenana River, Delta, Eagle, Fairbanks, Lake Minchumina, Middle Yukon, Grayling/Anvik/Holy Cross, Koyukuk, McGrath, Ruby, Tanana/Rampart/Manley, Minto/Nenana, Upper Tanana/Forty Mile, Yukon Flats.

Several of the committees did not meet to discuss fisheries issues in 1990. Sport Fisheries Division staff participated in meetings of the Fairbanks and Delta committees in 1990. Sport fisheries issues in Nome, Kotzebue, and in the lower Yukon and Kuskokwim River areas were handled by Division of Commercial Fisheries staff attending those respective meetings.

Alaska Board of Fisheries:

The Board of Fisheries (BOF) met in February, 1990 to consider changes in sport fish regulations in the AYK Region. A total of 14 proposals were considered that concerned sport fishing in the AYK Region. The proposals were submitted by the public as well as ADF&G staff. The following is a summary of proposals submitted for Board of Fisheries action during 1990 that specifically concerned sport fisheries in the AYK Region. The proposal numbers listed are those assigned in the 1990 BOF proposal book. A designation following each proposal number identifies whether the proposal originated with ADF&G staff or from the public.

Proposal
Number

358. Public. Establish a sport fishing closure in the Unalakleet River that would coincide with weekly commercial fishing closures during the king salmon season. Proposal rejected.
359. Public. Open portions of the Tolovana River drainage to winter ice fishing. Proposal rejected.
360. Public. Remove size limit and seasonal closure for northern pike in the Tanana River drainage. Proposal rejected.
361. Public. Remove size limit and seasonal closure for northern pike in the Yukon River drainage. Proposal rejected.
362. Public. Provide a slot size limit for Arctic grayling in the Salcha River. Proposal rejected.
363. Public. Raise bag limits for Arctic grayling in the Nenana River drainage to 10 fish per day. Proposal rejected.
364. Public. Raise size limit for northern pike in the Dall River to 36 inches, and change the bag limit to 10 fish, of which two may be over 36 inches in length. Proposal rejected.
365. Public. Increase the bag limit and size limit for northern pike in the Dall River. Proposal rejected.
366. Public. Increase the minimum size limit for northern pike in the Dall River. Proposal rejected.
367. Public. Increase the minimum size limit for northern pike in the Dall River. Proposal rejected.
368. Public. Remove the 12 inch minimum size limit for Arctic grayling in the Chena River. Proposal amended and adopted to limit fishing to artificial lures and single hooks upstream of the Chena River Flood Control Dam, and change the daily bag limit to two fish, 12 inches or greater in length.
371. ADF&G Staff. Establish a size limit for Arctic char or two Dolly Varden in waters of the North Slope of Alaska. Daily bag limit would be 10 fish, of which only two may be 20 inches or greater. Proposal adopted.
372. ADF&G Staff. Allow the use of bait on hooks with a gap size larger than 3/4 inch in the Salcha River. Proposal adopted.
375. Public. Close sport fishing on the Unalakleet River above the mouth of the Chirokey River until August 1. Proposal rejected.

376. ADF&G Staff. Establish a guideline harvest range for the recreational chinook salmon fishery in the Chena River. The range shall be from 300 to 600 chinook salmon per year. Proposal adopted.

AYK Sport Fishing Regulations:

Published regulations for 1990 are reproduced as Appendix C.

AYK SPORT FISHERIES ENHANCEMENT

Interior Alaska Lake Stocking Program

Selected waters (lakes, ponds and Piledriver Slough) of the Tanana River Valley are stocked on a continuing basis with rainbow trout, coho salmon, Arctic grayling, chinook salmon, lake trout, sockeye salmon, and Arctic char. Resulting fisheries composed a minimum of 42% of the recreational angling effort and 61% of the fish harvested in the Tanana River drainage in 1990 (Table 4). By providing stocked fish in the Tanana River valley, the ADF&G hopes to:

1. ease harvest pressure on wild stocks;
2. provide increased angling opportunity for increasing numbers of anglers; and,
3. diversify angling opportunity.

Lake stocking in the Tanana River valley takes place over an approximate 130,000 km² area, mostly near communities and along the road system, but also in a number of remote locations accessible only by off-road vehicle (ORV), dog team, or airplane. About half of the yearly sport effort on stocked lakes takes place during the winter on larger accessible lakes.

ADF&G stocking in the Interior began in the mid-1950's when barren lakes along the road system were stocked with rainbow trout or salmon. Between 1968 and the early 1980's, 15 lakes (including Birch and Quartz lakes) were chemically treated to eradicate endemic fish populations (Doxey 1987).

Throughout the 1970's, hatchery ability to provide stocked fingerlings steadily increased as fisheries enhancement received growing emphasis. Native Alaska rainbow trout brood stocks were developed when the importation of eggs from outside the state was discontinued in the late 1970's. As suitable brood stocks were developed and new hatcheries came on line, numbers of available stocked fish increased so that by 1985, average annual harvest and effort levels for stocked waters had risen by more than 40% and 20%, respectively (Doxey 1987).

The growth and success of the interior Alaska stocking program has been largely due to the development of, and production from, Alaska state hatcheries, particularly the Clear Hatchery, located about 145 km south of Fairbanks at the Clear Air Force Station (Figure 8) and the Fort Richardson hatchery near Anchorage. The Clear Hatchery program began in 1977, with an

initial mission to experimentally incubate, rear and release chum salmon to determine whether large scale enhancement of salmon would be feasible under conditions found in interior Alaska. In recent years, production of sport fish species has taken precedence over anadromous salmonids, and a large proportion of its output consists of fingerling and sub-catchable rainbow trout, coho salmon, and Arctic grayling. The facility presently has a capacity of about 9.0 million eggs.

Success of the stocking program is evaluated annually. The level of evaluation varies according to the size and accessibility of the lake and the importance and intensity of the sport fishery. Minimal evaluations address the question of whether the stocked fish survived and are providing sport fishing. More comprehensive evaluations provide limnological data, growth rate data, and fishery statistics such as CPUE, population estimates, comparison of performance between species, and other parameters.

1990 Stocking Results

Approximately 2.037 million fish of seven species were stocked in different locations in 1990, of which 765,068 were rainbow trout. By comparison, 3.07 million fish were stocked in the area in 1989, of which 747,000 were rainbow trout.

Most of the stocked rainbow trout in 1990 were fingerling (approximately 1.0 g) or sub-catchable (20 to 30 g) sized fish reared at Clear Hatchery, but about 63,000 were of catchable size (65 to 135 g) reared at the Fort Richardson Hatchery in Anchorage (Table 11). All of the rainbow trout were of the Swanson River (Kenai Peninsula) strain.

Rainbow trout were stocked into Piledriver Slough for the first time in 1987, and subsequent observations and catch returns in 1988 indicated that at least some large rainbow trout stocked in 1987 overwintered in, or returned to Piledriver Slough (Doxey 1989). Natural reproduction in the slough was not documented. Stockings of only the catchable size cohort continued in 1990 when 20,000 were released at a size of about 112 g..

A total of seven stocking events for rainbow trout took place in Harding Lake alone in 1990. Four lots of fish (totalling approximately 161,000 rainbow trout) were stocked which had been previously reared in floating net pens anchored in the lake.

Approximately 309,000 fry- and fingerling-size Arctic grayling were stocked in area lakes, ponds, and gravel pits in 1990, all of Moose Lake (Susitna River drainage) brood stock. These fish were incubated and reared at the Clear Hatchery (Table 12). Approximately 32,000 were short-term reared in floating net pens at Harding Lake before release into Harding Lake.

Arctic char and lake trout stockings in 1990 (Table 13) were limited to only those into Harding Lake in contrast to 1989 when over 20 separate water bodies in the Tanana drainage were stocked with Arctic char and 19 locations for lake trout.

Table 11. Number and size of rainbow trout stocked in AYK Region waters in 1990.

Receiving Water	Number Stocked	Size (g)
Bathing Beauty Pond	700	1.6
Birch Lake	48,345	22.9
Chena Lake	23,092	97.1
Grayling Lake	500	1.6
Chena Lake	8,159	107.0
Harding Lake	150,000	1.7
Harding Lake	103,312	1.2
Harding Lake	49,912 ^a	3.2
Harding Lake	99,907 ^a	6.9
Harding Lake	1,019	110.0
Harding Lake	9,970 ^a	125.5
Harding Lake	1,000 ^a	177.0
Hidden Lake (EAFB)	3,600	1.6
Johnson Road Pit # 2	2,000	1.6
Little Harding Lake	1,000	1.6
Lost Lake	1,000	1.6
Manchu Lake	2,900	1.6
Piledriver Slough	20,000	111.9
Quartz Lake	150,632	1.2
Quartz Lake	52,914	2.4
Quartz Lake	33,843	22.9
Red Dog Mine	203	70.0
Red Dog Mine	250	92.8
Twenty-eight mile Richardson Hwy Pit	800	1.6
Total rainbow trout	765,068	

^a Reared in floating net pens before release.

Table 12. Number and size of Arctic grayling stocked in AYK Region waters in 1990

Receiving Water	Number Stocked	Size(g)
<u>Arctic grayling</u>		
Bolio Lake	20,000	0.02
Eighty-one mi. Pit (Richardson Hwy)	5,000	0.02
Grayling Lake	10,000	0.02
Harding Lake	51,200	0.02
Harding Lake	2,400 ^a	3.9
Harding Lake	29,972 ^a	5.2
Johnson Road Pit # 2	10,000	0.02
Long Pond	10,000	0.02
Lower Twin Lake	90,000	0.02
Otto Lake	30,000	0.02
Sansing Lake	40,000	0.02
Sansing Lake	200	5.6
Thirty-one mi. Pit (Richardson Hwy)	10,000	0.02
Subtotal Arctic grayling		308,772

^a Reared in floating net pens before release.

Table 13. Number and size of Arctic char, lake trout, sheefish, sockeye salmon, and coho salmon stocked in AYK Region waters in 1990.

Receiving water	Number Stocked	Size (g)
<u>Arctic char</u>		
Harding Lake	1,312	653.0
Harding Lake	49,900 ^a	20.1
Harding Lake	20,614	35.2
Harding Lake	15,159	35.9
Harding Lake	7,331	50.1
Harding Lake	11,230	56.6
Harding Lake	40,000	6.1
Harding Lake	10,000	5.7
Total Arctic char	155,546	
<u>Lake trout</u>		
Harding Lake	73,700 ^a	10.3
Harding Lake	72,000	4.6
Total Lake trout	145,700	
<u>Sheefish</u>		
Sansing Lake	7	6,100.0
<u>Sockeye</u>		
Harding Lake	400,000 ^a	0.20
Harding Lake	289	1.7
Total sockeye	400,289	
<u>Coho salmon</u>		
Birch Lake	131,000	2.7
Jan Lake	8,800	2.7
Little Harding Lake	3,600	2.7
Quartz Lake	150,000	2.7
South Jan Lake	21,000	2.4
Twenty-eight mi. Pit (Richardson Hwy)	500	2.7
Total coho	314,900	

^a Reared in floating net pens before release.

Harding Lake was stocked experimentally for the first time in 1988 with 500,000 sockeye salmon fry from the Gulkana Hatchery. It was again stocked in 1989 with 515,000 post-emergent (0.16 g) sockeye salmon fry. In 1990 400,000 emergent sockeye fry and 290 fed fry were released in Harding Lake (Table 13). A total of 314,900 coho salmon was stocked in Six Tanana Area lakes and ponds in 1990 (Table 14). Stocked coho salmon were all fingerling size (2.7 g). All released coho salmon were of Wood Creek (Nenana River) stock, and all were reared at Clear Hatchery.

Only a few large (7) Sheefish adults were released in 1990. These fish were released in Sansing Lake near Clear Hatchery, and the individuals represented hatchery broodstock, originally from the Koyukuk River.

Anadromous Fish Releases by State Hatcheries

All anadromous fish hatched and/or reared at Clear Hatchery in 1990 were stocked into landlocked lakes in the Tanana River drainage. No anadromous species were stocked into flowing waters of the drainage in 1990. Coho salmon of the Wood Creek (Nenana River) stock were released in 1989 in tributaries to the Nenana River. Released salmon in 1989 ranged in weight from about 2.5 to 4.2 g. and were progeny of the 1988 brood year.

The other state operated hatchery in the AYK Region was initiated in 1980 in the Kotzebue area at Sikusuilaq Creek, approximately 50 km upstream from the mouth of the Noatak River (Figure 18). The initial purpose of the facility was to examine the feasibility of large-scale chum salmon enhancement in a far northern environment. Hatchery expansion starting in 1987 is intended to bring the facility into larger scale production than in prior years of feasibility testing. Enhanced hatchery returns are intended to benefit the salmon gill net fishery which operates near Kotzebue in marine waters. The scale of salmon releases was relatively small, less than two million chum salmon fry prior to 1988. Returns from prior year releases have been identified both in the commercial fishery and at the hatchery. Releases of chum salmon fry in the lower Noatak River in 1990 totalled 6,364,003 fish, an increase of about 300,000 fry over the 1989 release. Most of the fry were released on June 13 at an average size of 0.45 g, but a later release was made on July 3 of 138,203 fry that had been reared to an average size of approximately 1.43g. Returns in 1990 of about 36,000 adult chum salmon were identified in the commercial fishery harvest and at the hatchery site from hatchery releases in 1985-1987. Expanded hatchery capability was initiated in 1987 with the installation of new egg and fry incubators in order to increase rearing capacity to 10 million eggs. In addition, new raceways were installed to allow short term rearing of fry before release.

LAND USE, HABITAT AND WATER QUALITY

Habitat-Related Fisheries Issues

Commercial development of minerals and timber, and construction of highways, can have significant impacts on watersheds as well as fish. The following is

Table 14. Gaging station records^a of mean and maximum discharge^b for 1990, and mean and maximum discharge for the period of record, for 11 rivers in the AYK Region.

River	1990		Period of Record		
	Mean	Max	Mean	Max	Years
Kuskokwim at Crooked Cr	44.9	175.0	41.0	392.0	1951-90
Yukon at Eagle	87.0	284.0	83.2	545.0	1950-90
Yukon at Pilot Station	235.4	900.0	224.3	1,100.0	1975-90
Tanana at Fairbanks	22.7	63.9	19.9	96.4 ^c	1973-90
Chena at Fairbanks	1.3	8.4	1.4	74.4	1948-90
Salcha near Salchaket	1.6	12.6	1.6	97.0	1948-90
Snake near Nome	0.2	1.9	0.2	4.2	1965-90
Kobuk near Kiana	13.3	130.0	15.2	152.0	1976-90
Wulik near Kivalina	1.3	19.7	1.0	31.4	1984-90
Kuparuk near Deadhorse	0.8	70.0	1.3	118.0	1971-90
Sagavanirktok, Pump 3	1.1	8.4	1.3	23.0	1982-90

^a Data from Lamke et al. 1991.

^b Cubic feet per second x 1,000.

^c Flood of Aug. 16, 1967 reached an approximate discharge of 125,000 ft³.

a brief description of the types of commercial development in the Tanana and AYK Areas, and known fisheries impacts.

Placer Mining:

The majority of Alaska's placer mining takes place in the AYK Region. Placer mining effluents, if not controlled, have the potential to significantly alter stream habitats and to impact fish populations. Elevated stream turbidity and sediment loads may reduce oxygen exchange rates through abrasion of gill tissues, prevent foraging by sight feeding fishes, limit aquatic plant growth by displacement or smothering, and generally reduce abundance and/or diversity of aquatic macroinvertebrates important for fish production (Weber 1986). Placer mining activities may also increase the toxic metal content (arsenic, mercury) of stream water (ADEC 1986). In addition to changes in water quality, placer mining can affect the physical characteristics of the streambed by altering channel flow and modifying riparian habitat. Tailing deposits can inhibit fish passage and decrease overwintering habitat.

In the AYK Region, in 1990, 2,879 new mining claims (on state and federal lands) were recorded by the Alaska Department of Natural Resources (ADNR), a receiving agency for state and federal mining permits (Swainbank et al. 1991). The number of claims recorded in 1990 represents a decrease of 29% from the number recorded in 1989. The total number of active claims on file in 1990 (40,149) was fewer than the number of claims on file in 1989 (44,126). The ADF&G issues permits for mining in streams supporting anadromous and resident fish per its Alaska State Statute Title 16 authority. In some cases, where development is within a stream supporting only non-anadromous fish and will not block fish passage, or where fish are not present, permits may not be required.

Gold production decreased in 1990 for the first year since 1987, from more than 8,852 kg (284,617 troy oz) in 1989 to 7,205 kg (231,700 troy oz) for the entire state (Swainbank et. al. 1991). Decreased production was due in large part to the poor production season on the 'BIMA' (owned by Western Gold Exploration and Mining Company) offshore dredge at Nome. The 'BIMA' dredge ceased operations permanently late in the dredging season of 1990. The majority of the gold operations were placer mines, with the remainder consisting of lode operations. Placer mining accounted for 83% of the gold produced in 1990.

The U.S. Environmental Protection Agency (EPA) issued placer mining effluent guidelines in 1988, which had been under consideration since 1986. Guidelines for water discharge permits went into effect in 1989 requiring 100% recycling of mine process waters, in addition to placing certain restrictions on the use of hydraulic mining technologies. Certain "best practice" mining methods are recommended in the guidelines. An NPDES (National Pollutant Discharge Elimination System) permit is required for discharge of mine waste into receiving streams. There were 64 such permits issued in 1990 (ADEC 1991).

The Alaska legislature implemented rental and royalty fees from all mining on state land in response to an Alaska Supreme Court decision that the Alaska

Statehood Act required collection of these fees from mining activity on state lands.

Gravel Mining:

There are few documented instances where gravel mining in AYK has affected fish populations. Gravel mining of a streambed has the potential to cause instream fanning, erosion, and deterioration of water quality and fish habitat (ADEC 1986). A complete listing of gravel mining sites and impacted streams is available in ADEC (1986). Swainbank et al. (1991) contains a listing of gravel mine operations in 1990. Reclamation of 19 gravel mine pits in the Prudhoe Bay area of the North Slope has increased overwintering habitat for fish inhabiting connecting streams, and increased fish production (Hemming 1988 and 1990).

Statewide production of sand and gravel in 1989 and 1990 was at the lowest level since the early 1970's due to declines in major construction activities.

Industrial Metals Development:

Offshore dredging for gold in Norton Sound by the 'Bima' (world's largest bucketline dredge owned by Western Gold Exploration and Mining Company) ceased in 1990 after five years of operation on its offshore leases.

Cominco Alaska Inc. mined 1,318,200 tons of zinc-lead-silver concentrate from the Red Dog deposit north of Kotzebue in its first year of operation (Swainbank 1991). There was concern in 1990 that heavy metal contamination of Red Dog and Ikalukrok creeks would occur both from natural leaching of the ore body as it was stripped for ore production and from discharge of impounded waters that were not treated to remove contaminants. Such contamination would affect fish resident to, and migrating through, the lower Wulik River, resulting in public health concerns for the river fisheries for char, Arctic grayling and salmon. Regulatory enforcement by the Alaska Department of Environmental Conservation, including fines imposed, in conjunction with mitigative and preventative measures taken by Cominco Inc. reduced danger of fish contamination in the Wulik River.

Oil and Gas Development:

Oil and gas extraction activities in the AYK Region are presently restricted to the North Slope. North Slope development has affected fish habitat on and near transportation corridors (such as the Dalton Highway and the Trans-Alaska Pipeline) and by the extraction of gravel for road and building construction and maintenance. There is evidence that construction of Prudhoe Bay's West Dock Causeway has disrupted east and west migratory movements as well as recruitment of Arctic cisco in the Colville and Sagavanirktok rivers (Gallaway et al. 1987; Moulton et al. 1986). Documentation of oil spills and resulting adverse impacts (if any) on fish populations from contamination is lacking in the AYK Area. A number of studies throughout the 1980's have attempted to determine or predict impacts from offshore gravel causeways, such as the Endicott Causeway, in the Beaufort Sea to fish, fisheries and fisheries habitat.

Timberland Development:

Logging had no significant impact to fisheries resources in the AYK Region in 1990. Little commercial logging presently occurs in the region. The largest commercial timber harvests in the Kuskokwim Basin occur from McGrath to Stony River (ADNR 1988). Some commercial logging of spruce currently occurs in the Tanana River basin, primarily on state owned land along the Tanana River.

Road construction associated with logging could impact fish populations by accelerating the rate of soil erosion and sedimentation in streams, however, because of the minimal level of industry development at the present time, no such problems have been identified in the region.

Highway Development:

Roads allow increased access to streams and lakes, thereby increasing the utilization of sport fishery resources. Improperly designed or constructed road culverts can create partial or complete barriers to fish migration. Major highways in the AYK Region include the Steese Highway which provided road access via Fairbanks to the Yukon River and Birch Creek; the Taylor Highway leading to the Forty-Mile and Yukon rivers; the Parks and Richardson highways which provide access to the Tanana River and many of its tributaries and lakes; the Dalton Highway to the North Slope which crosses the Yukon River, upper tributaries of the Koyukuk River and the Sagavanirktok and Kuparak rivers on the North Slope. A newly-constructed road north of Kotzebue connects the Red Dog Mine in the headwaters of the Wulik River to the Chukchi Sea coast south of Kivalina. Its purpose is to serve the Red Dog Mine site, and for transport of ore to the port facility on the coast. The Seward Peninsula has three major roads which cross more than a dozen rivers important for sport fishing. Of the road systems in the AYK Region, culvert crossing on the Dalton, Red Dog Mine and Seward Peninsula Highways have been of most concern to the Department. There are 40 - 50 culverts on the Seward Peninsula built in the 1950's of which many constitute partial barriers to tributary spawning and rearing. Habitat Division personnel, the Alaska Department of Transportation, and Bureau of Land Management (BLM) personnel have worked cooperatively to improve stream habitat by removing gravel berms in the Nome and Pilgrim rivers which will provide more rearing habitat for young Arctic grayling.

Commercial Utilization of Fisheries Resources

Commercial sport fishing activities, through establishment of lodges and guiding services, offer a source of revenue to residents in the Tanana and AYK Areas. A brief description of known commercial uses of sport fish species follows.

Wilderness Lodges and Guiding:

Lodges and sport fish guiding operations (including outfitters) are significant factors in the utilization of sport fishery resources in the AYK Region, although the extent of this influence has yet to be fully determined.

More information regarding the location and operation of lodges and guiding and outfitting operations is needed. A partial list of the type of operation at various locations can be found in Appendix D. The number of resident or nonresident clients served, species targeted, or types of fishing experiences offered at these locations is presently incompletely documented.

Commercial Fisheries:

Important commercial fisheries exist for finfish and shellfish species in the AYK Region. The largest is the commercial salmon fishery, followed by the herring and shellfish fisheries. Estimated wholesale values of the fisheries in 1989, (the latest year for which statistics have been compiled) were approximately \$16.4 million for salmon, \$3.5 million for herring and \$1.2 million for shellfish (Savikko and Page 1990).

Commercial fisheries for finfish species other than salmon, or herring, are sometimes allowed under authority of a permit issued by the Commissioner of the Department of Fish and Game or his designee, usually an area manager of the Division of Commercial Fisheries. Permits to commercially harvest whitefish, sheefish, northern pike, blackfish *Dallia pectoralis*, lamprey *Lampetra japonica*, Dolly Varden and burbot are issued occasionally for limited (usually local) commercial markets. In many cases, permits are issued by the Department, but harvests are either not made or are not reported. Only a small commercial harvest of non-anadromous freshwater finfish was reported for the AYK Region in 1990. The Division of Commercial Fisheries maintains data records of such harvests.

Commercial harvests of freshwater species in the Norton Sound and Kotzebue Area in 1990 are reported by Bue (pers. comm. ⁷). Historical harvests are summarized in Merkouris and Lean (1989). The harvest of sheefish from Hotham Inlet near Kotzebue takes place during winter months of two calendar years (October 1989-October 1990). Commercial harvest of sheefish during this period is estimated to total 852 fish. Five catcher-sellers participated in the fishery, harvesting 3,734 kg (8,224 lbs). This fishery is limited by quota to 11,350 kg. Adult Dolly Varden taken incidentally in the Kotzebue commercial chum salmon fishery are sometimes sold to commercial fish processors. A total of 604 such fish were harvested and sold in 1990. The mean weight of the commercially sold Dolly Varden was 3.2 kg (7.0 lbs) in 1990. Virtually no fishing activity under the authority of a freshwater fishing permit took place in either the Norton Sound or Kotzebue Areas in 1990 for the harvest of whitefish or other freshwater species such as burbot or northern pike.

Francisco et al. (1991) reports that in 1990, 11 fishermen harvested and sold 1,664 whitefish in the Kuskokwim Area.

A fishery has taken place in the Colville River since 1964 for broad whitefish, humpback whitefish, Arctic cisco and least cisco. Reported

⁷ Bue, Fred. 1991. Personal Communication. ADF&G, Division of Commercial Fisheries. P.O. Box 1148 Nome AK 99762.

harvests (Holder pers. comm.⁸) in this fishery in 1990 include: 5,694 humpback whitefish, 19,374 Arctic cisco, and 21,003 least cisco.

Freshwater fishery permits have been issued in various years for whitefish at Healy Lake, whitefish in Lake Minchumina, and whitefish and burbot in the Tanana River. Although several permits for commercial harvest of freshwater finfish were issued in 1990 for the Yukon River drainage, only two of the permit holders reported taking any fish under the permits issued. A reported 2,074 lbs of whitefish (species not specified) was harvested in the Yukon River downstream of Tanana (Holder 1991). A harvest of 809 whitefish (weight and species unspecified) was reported for the Tanana River in 1990.

Land Withdrawals, Status, and Planning

Land use designations by private owners, state, or federal agencies may affect fisheries management considerations within given land parcels. A brief description of various national land designations in the Tanana and AYK Areas and known influences on the use of sport fish within land units follows.

ANILCA:

The Alaska National Interest Lands Conservation Act (ANILCA), enacted into law in 1980, completed the implementation of the Alaska Native Claims Settlement Act (ANCSA) and addressed outstanding issues such as subsistence opportunity, energy development, economic growth and transportation planning. Legislative solutions to these issues included the creation or expansion of five national conservation systems in Alaska: national parks, wildlife refuges, wild and scenic rivers, wilderness preservation lands and national forests.

The purpose of ANILCA is to preserve for future generations certain lands and waters in Alaska, protect resources related to subsistence needs and the subsistence lifestyle for rural residents, and to protect those resources related to recreational opportunities, such as sport fishing and hunting (ANILCA 1980). The Act directs specific management guidelines for conservation system units within Alaska.

To maintain state responsibility for fish and game management on newly designated conservation system units, ANILCA required the state to distinguish between user groups and assign priority opportunities for subsistence uses of fish and game resources.

National Parks, Monuments and Preserves:

All National Park Service (NPS) managed lands in the AYK Region are discussed under the following section.

A memorandum of understanding between the State of Alaska and the National Park Service (NPS) stipulates that State fish and game regulations apply on NPS lands except when a more restrictive harvest approach is desired by NPS.

⁸ Holder, Russ. 1991. Personal Communication. ADF&G, Division of Commercial Fisheries, 1300 College Rd. Fairbanks, AK 99701.

NPS may promulgate regulations concerning consumptive uses of resources which are more restrictive than state laws. The ANILCA intends for NPS to provide opportunities for continued subsistence and traditional activities.

Park land designation has some influence on utilization of the sport fishery resource by restricting types of development within the parks. Large scale commercial development (i.e.. fishing lodges) is not allowed, but small lodge facilities for a few guests have been allowed in Kobuk National Park. Construction of temporary facilities (such as fish camps or tents) on park lands in Alaska was granted under ANILCA, but has been the subject of dispute with NPS. NPS goals include minimizing the sport fish take by encouraging release of captured fish or the taking of only small individuals of the more abundant species (NPS 1985a, 1986a). Motorized boat, snow machine, and airplane access is allowed for sport fishing on park lands in Alaska. Fish stocking or enhancement activities can be allowed if the purpose is to restore fish populations to "natural or healthy" levels.

Conservation system units within AYK (Figure 20) are as follows:

1. Kobuk Valley National Park is 688,000 ha (1,700,000 ac) in size and includes one wild and scenic river and 77,000 ha (190,000 ac) of wilderness. NPS has proposed an additional 168,000 ha (414,720 ac) be set aside as wilderness. Regional residents account for more than 90% of park use. NPS estimates that out-of-region recreational use is limited to about 25-75 users per year. Most out-of region visitors fly in their own planes to sport fish at the mouth of the Salmon River and other tributaries of the Kobuk River. Local boats can be chartered for fishing. Lodges in Ambler and Shungnak accommodate a small number of visitors. There are reported instances of subsistence and sport fish user conflicts on the Kobuk River (see NPS 1986a).
2. Gates of the Arctic National Park and Preserve consists of 2,939,000 ha (7,263,000 ac) of combined park and wilderness lands and 42,000 ha (103,932 ac) of park lands only. The park also includes six wild and scenic rivers. Recreational fishing is mostly for Arctic grayling, Dolly Varden and lake trout, with the most heavily used areas being Walker and Chandler lakes. Two lodges on Walker Lake, at the headwaters of the Alatna River, advertise sport fishing opportunities, and local air-taxi operators drop off anglers at other areas. Sport fishing mostly occurs in conjunction with other activities such as river running, hunting and backpacking (see NPS 1985a).
3. Cape Krusenstern National Monument was created in 1980 to protect archaeological sites, preserve prehistoric and historic Native cultures, protect habitat for fish and wildlife and protect the viability of subsistence resources. The NPS directs management of the monument which is 267,000 ha (659,807 ac) in size. Access and development restrictions of park lands apply to monument lands. Fishing for whitefish, ciscoes, Arctic char/Dolly Varden, chum salmon and northern pike is primarily by subsistence users.

Recreational use of the monument is extremely limited and occurs mostly in conjunction with subsistence activities (NPS 1985b).

4. Noatak National Preserve is 2,630,000 ha (6,500,000 ac) in size, and includes one Wild and Scenic River and 2,350,000 ha (5,800,000 ac) of designated wilderness land. It is also a UNESCO Biosphere Reserve, designed to maintain genetic pools. Recreational use is estimated at 2,000 to 2,500 visitors per year, who participate in river excursions, hunting and sport fishing. Arctic grayling and Dolly Varden are the most common sport fish. About 25 commercial operators provide air and guiding service. Popular drop off points for sport fishing include the Kelly and Cutler Rivers. Construction of a State hatchery (Sikusuilq Creek) in the Lower Noatak River has caused some concern on the part of NPS regarding maintenance of the chum salmon gene pool as expressed in the Biosphere Preserve philosophy (NPS 1986b).
5. Bering Land Bridge National Preserve consists of 1,127,000 million ha (2,800,000 ac), with 90% of use related to subsistence and local use activities. NPS has proposed that an additional 121,000 ha (299,520 ac) of the Preserve be designated as Wilderness. Very little sport fishing occurs in the Preserve because better fishing opportunities are available on the Seward Peninsula outside of the Preserve (NPS 1986c).
6. Yukon-Charley Rivers National Preserve is 1,023,000 ha (2,530,000 ac) in size and includes the Charley River and its main tributaries as a Wild and Scenic River. NPS has proposed that 442,380 ha (1,093,120 ac) be designated as Wilderness. Sport fishing is primarily for Arctic grayling, with northern pike found in lower tributary streams and Dolly Varden found in one tributary (NPS 1985c).

A General Management Plan for each unit (except the Yukon-Charley River National Preserve which was completed in 1985) was completed by the NPS in 1987 after taking public and state agency input. General Management plans are intended to establish management direction, determine public access policies, and allowable public uses including priorities for fisheries research within each park unit. It is intended that supplemental plans will be developed in subsequent years to deal with specific fisheries projects, public uses, and access problems. Many such planning efforts were underway in 1990.

National Wildlife Refuges:

Refuges (Figure 20) are mandated to conserve fish and wildlife habitat, fulfill international treaty obligations, provide for continued subsistence opportunities and ensure water quality. Each refuge has specific legislative purposes and although each is regulated by federal law, the USFWS recognizes a master memorandum of understanding with the State of Alaska which vests primary responsibility for fish and wildlife management with the state, unless subsistence opportunities are compromised. The Alaska Supreme Court in 1989 decided the McDowell case and found that allowing a priority for rural

residents for subsistence was unconstitutional. In 1990, as a result of this decision, and the fact that Alaska's fish and game regulations were out of compliance with ANILCA, the management of subsistence hunting and fishing on federal lands was transferred to federal authority. In practice, this has meant that subsistence hunting on federal land is managed by the federal government, under the auspices of a federal subsistence board, while subsistence fishing has largely remained a state responsibility because submerged lands that are deemed navigable are considered to be state lands under the Statehood Act. Refuge managers review and adopt ADF&G management plans unless the plans are formally determined to be incompatible with the purposes of the refuge. Different management goals exist for each refuge. Policy ranges from that of minimal interference with human use, to that of promotion of increased Wilderness and Wild and Scenic River designations. All guides and outfitters are required to have special use permits in addition to state licenses (for big game guides). There are eight National Wildlife Refuges (NWR) in the AYK Region, and a summary of each follows. Comprehensive Conservation Plans (available from U.S. Fish and Wildlife Service) were completed for all refuges except the Yukon Delta NWR and Arctic NWR in 1987. The plans contain detailed information on the environment, management alternatives, environmental consequences as well as maps for each unit. Step-down plans, more specific plans for fisheries, river management and other public uses of the lands and their resources, are scheduled for completion in subsequent years. Some refuge step-down plans were initiated in 1990.

1. Selawik NWR is 890,327 ha (2,200,000 ac) in size and includes one Wild and Scenic River and 97,126 ha (240,000 ac) of Wilderness lands. The preferred management alternative is for minimal interference. Mechanized travel to any in-holdings, oil and gas studies and recreational opportunities would be allowed. Recreational use levels are extremely low, with most sport fishing targeting on sheefish in the Kobuk River, adjacent to the NWR (USFWS 1986).
2. Yukon Delta NWR is the largest of Alaska's 16 refuges and consists of 10.52 million ha (26,000,000 ac) including two Wild and Scenic Rivers and 769,000 ha (1,900,000 ac) of Wilderness lands. The management plan permits oil and gas leasing on only 3% of the refuge. Habitat and population manipulation may be conducted on some of the lands. Most sport fishing occurs on the Kisaralik River, but increasing interest from Togiak River fishing guides in establishing commercial guiding on the Andreafsky and Kwethluk rivers and other refuge rivers has been expressed. Rainbow trout are found in the Kwethluk, Kasigluk, Kisaralik, Tuluksak, and Aniak rivers. Sport harvest of sheefish has increased and large numbers of northern pike are caught by locals in the winter. If sport fish guiding increases, the refuge staff envisions conflicts with subsistence users and plans to launch an extensive monitoring program (USFWS 1987a).
3. Yukon Flats NWR is 4,530,000 ha (11,200,000 ac) in size, has two Wild and Scenic Rivers, and borders the Trans-Alaska pipeline. The management plan directs minimal disturbance of habitat and increased

wilderness land designations. Limited fly-in sport fishing exists and is mostly incidental to hunting and river running. The Dall River receives the heaviest sport fishing pressure, due to access from the Dalton Highway (USFWS 1985). A fisheries management plan was published in April, 1990.

4. Koyukuk and Innoko NWR are 1,820,000 and 283,000 ha (4,500,000 and 700,000 ac) in size, respectively. The management plan calls for minimal management. Staff has little information on sport fishing, but believes some occurs in conjunction with hunting and river running (USFWS 1987c). A fisheries management plan was in preparation in 1990.
5. Nowitna NWR consists of 809,389 ha (2,000,000 ac) and one Wild and Scenic River. The management plan is for minimal management. Sport fishing for trophy sheefish is an established activity on the Nowitna River. Northern pike are also sought by anglers. It is believed that most sport fishing occurs in conjunction with hunting (USFWS 1987b). A fisheries management plan was in preparation in 1990.
6. Kanuti NWR is 647,511 ha (1,600,000 ac) in size. The management plan emphasizes the restoration of fish populations to natural and healthy levels. The plan also strives to increase fishing opportunities, but would designate no wilderness areas, and would allow some oil and gas studies (USFWS 1987d). A fisheries management plan was in preparation in 1990.
7. Arctic NWR consists of 7,900,000 ha (19,500,000 ac), four Wild and Scenic Rivers and 3,240,000 ha (8,000,000 ac) of wilderness lands. Section 1002 set aside 607,000 ha (1,500,000 ac) of land on the coastal plain of ANWR for future oil and gas exploration and development pending authorization by the U.S. Congress. The management plan maintains the existing range and intensity of management and recreational economic uses. Opportunities for fishing and other public uses would be maintained, as would scientific research. Most sport fishing for Arctic grayling, Dolly Varden, Arctic char, lake trout and northern pike occurs in conjunction with river trips and hunting (USFWS 1988). A fisheries management plan was in preparation in 1990, as was a river management plan.
8. Tetlin NWR, in the upper Tanana River valley is 284,000 ha (700,000 ac) in size. Under the management plan, fisheries would continue to be managed for harvests supported almost entirely by naturally reproducing stocks (USDI 1987). Only subsistence and sport fishing will be allowed on the refuge. A fishery management plan was published in April 1990.

Float trips on refuge rivers of both the north and south slope are a recognized and growing popular use. The Kongakut River on the north slope is considered most popular, followed by the Hulahula and Canning rivers. The Ivishak and Sagavanirktok rivers are also sometimes used by float parties.

The Sheenjek and Porcupine rivers are the most popular rivers on the Brooks Range south slope for recreation (USFWS 1988).

Wild and Scenic Rivers:

In the AYK Region, 23 rivers in national parks, preserves and refuges have been placed within the national wild and scenic river system. The Wild and Scenic Rivers Act of 1968 stipulates that these rivers shall be preserved in free flowing condition, generally free of impoundments, and have primitive shorelines and watersheds. The wild and scenic river designation positively impacts utilization of the sport fish resource by affording anglers the possibility of a pristine and uncrowded fishing experience. Access to rivers is controlled and facilities are restricted, thus potentially precluding the development of fishing lodges among other uses. Wild and scenic rivers in the AYK Area are listed in Appendix A.

Wilderness Land Designations:

The Wilderness Act of 1964 restricts modes of access and development on designated parcels of land. Wilderness land designation is intended to promote solitude and primitive recreational opportunities. Depending upon interpretation of the wilderness modifications in ANILCA, land managers may restrict the use of power chain saws, generators, and other similar motors. Stream clearance, weir construction, and field camp operations in wilderness areas in support of fisheries field research may be restricted, depending upon circumstances (Cunning pers. comm.⁹).

Natural Factors Affecting Sport Fisheries

The timing and severity of natural catastrophic events may affect sport fish habitat and life history. Known natural occurrences in 1990 are described and their impacts on sport fish are estimated in the following paragraphs.

Fires:

Fires and fire suppression measures by agencies such as the BLM and State of Alaska, Division of Forestry, are common during the summer months; and, during particularly dry and warm years, forest and tundra fires are a major feature in interior and northern Alaska. Fires in Alaska generally do not penetrate the duff layer to mineral soil and thus do not represent a great potential erosion problem. In addition, frozen ground in large areas of the Arctic and interior assists in curtailing fire-induced soil erosion. Major impacts of fire on fisheries can occur with the use of earth moving equipment by fire-fighters to prevent enlargement of a blaze, and aerial deployment of fire retardant. BLM has strict guidelines regarding fire retardant use near water bodies. BLM personnel state that the retardant presently in use is biodegradable and if mistakenly introduced into a water body, would have

⁹ Cunning, Artina. 1988. Personal Communication. ADF&G, Division of Wildlife Conservation, 333 Raspberry Rd. Anchorage AK 99502.

minimal and short-lived impact on fish populations (Jones pers. comm.¹⁰). Thus, it is the BLM position that fire retardant use in Alaska does not directly threaten fish populations.

The 1990 fire season was one of the most severe on record, with 932 unique fires recognized by the Alaska Fire Service (AFS). By the end of July, 1990 fire danger conditions, based on the Canadian Forest Fire Weather Index System, indicated that Alaska was under severe drought conditions equal to those influencing Yellowstone National Park at the end of July 1988 (U.S. Department of Interior/Bureau of Land Management Alaska Fire Service, 1991). A total of 3.2 million acres (1,295,000 hectares) burned in 1990, compared to the five-year average (1983-1988) of 253,964 ha. Lightning was the primary cause of all fires reported. The Beaver fire, (108,930 ha) started on June 27, and the Tok River fire (39,660 ha) started on July 1. The largest fire in the state, the Coleen River blaze on the Arctic National Wildlife Range (187,907 ha) started on June 29 (U.S. Department of Interior/Bureau of Land Management Alaska Fire Service, 1991).

Snow Pack Assessment:

Snowpack depth and duration impacts fish life history by influencing such factors as water level, sunlight penetration, and insulation of water bodies in periods of extreme cold. Snow survey data obtained from the Soil Conservation Service (USDA 1990) provided snowpack summaries for water year 1990 by region. (Water year 1990, hereafter referred to as WR 90, is the period of time from 1 October 1989 through September 30, 1990).

In the Arctic, winter snow accumulation was below average as of May, 1990, with accumulations ranging from 50% to 80% of normal. In the Upper Yukon Basin, the basin-wide snowpack at the headwaters of the Yukon River was near, to slightly below average for the year. Maximum snow water equivalent (SWE) was reached in April at about 16.5 cm, while SWE in the Central Yukon peaked in April at about 12.7 cm and was above normal. Breakup was earlier than normal in the Tanana Basin, where the SWE was about 13.2 cm in April compared to the average of about 10 cm. In the Koyukuk Basin SWE was below average. Kuskokwim area winter snowfall was well above normal, with SWE of 25.4 cm in April, compared to average values of 15.2 cm. Comparable snowfall data were not available for the Kotzebue/Norton Sound Basins.

Stream Discharge Assessment:

Stream flows have a significant impact on fish life history, especially maximum stream discharge events. Stream discharge records obtained from the U.S. Geological Survey (Lamke et al. 1991) provide monthly and annual mean and extreme discharge values by river (Table 14).

Of the streams in the AYK Region for which discharge data are collected, many of the sites throughout the region recorded mean discharge rates in WR 90 that exceeded the annual mean discharge of all years of record (Table 14).

¹⁰ Jones, Mark. 1988. Personal Communication. BLM, Alaska Fire Service, PO Box 35005, Ft. Wainwright, AK 99703.

However, maximum discharges did not approach the maximum recorded flows. Maximum discharges at all sites in WR 90 did not exceed maximum flow in the years of record. No serious flooding and streambed disruption was noted during the summer of 1990 in any AYK Region drainage.

Mean Air Temperature and Precipitation:

Average annual temperature and precipitation regimes influence the timing of stream freeze-up and break-up occurrences, and by affecting the duration and severity of the seasons, plays a major role in the annual water budget. Climatological data for four cities in the AYK Region (Fairbanks, Nome, Kotzebue and Barrow) were obtained from the US Weather Service (NOAA 1989). The mean monthly and yearly temperature (F) and precipitation (inches) for the period of record (1958-1990) are provided by this source. Climatological data from only four locations may not adequately represent micro-climatic conditions throughout the region, but may provide an indication of seasonal weather patterns in the AYK Region.

Mean monthly air temperatures in WY 1990 for the four cities were generally slightly elevated over the historic means, except for midwinter dips in January and February for all stations (Figure 25).

Mean monthly precipitation values for WY 1990 for the four cities are more variable than temperature means. The interior, as judged from the precipitation records from Fairbanks, experienced near average moisture from October 1989 to May 1990, and then experienced higher than average precipitation in June through August 1990. Higher than normal precipitation was recorded in July and August 1990 in Barrow and Nome (Figure 26).

ACKNOWLEDGEMENTS

The author thanks John H. Clark for his continued support and generous help in preparation of this report series. The author is grateful to members of the staff, including Mike Mills, Mike Doxey, Robert Clark, Margaret Merritt, Fred Andersen, Jerry Hallberg, John Burr, Fronty Parker and Fred DeCicco for responding generously to requests for assistance. Thanks also to Sara Case for final preparation of the report, and to Carol Hepler for preparation of several maps.






Figure 25. Monthly mean air temperature for the 1990 water year compared to monthly mean air temperature, 1958-1990, in four AYK locations.


FAIRBANKS



BARROW



KOTZEBUE



NOME




Figure 26. Mean monthly precipitation for water year 1990 compared to mean monthly precipitation, 1958-1990, in four AYK locations.

LITERATURE CITED

- Alaska Department of Environmental Conservation (ADEC). 1991. 1990 Annual Mining Report. Alaska Department Environmental Conservation, Juneau, Alaska. 39 pp.
- Alaska Department of Fish and Game (ADF&G). 1978. Alaska's fisheries atlas. Vol. 2 [R. F. McLean and K. J. Delaney, comps.]. 43pp. + 153 maps.
- _____. 1983. Emergency order standards of applicability. Alaska Department of Fish and Game, Commissioner's office. Juneau, Alaska. 18pp.
- _____. 1984. Sport Fish Survey. Booklet published by Alaska Department of Fish and Game, Division of Sport Fisheries. Juneau, Alaska.
- _____. 1986. Alaska Habitat Management Guide, Arctic Region. Vol. II: Distribution, abundance, and human use of fish and wildlife. Div. of Habitat, Alaska Department of Fish and Game, Juneau, AK.
- _____. 1991. Salmon Fisheries in the Yukon Area, Alaska. A report to the Alaska Board of Fisheries. Regional Information Report No. 3F91-02. Alaska Department of Fish and Game, Division of Commercial Fisheries, AYK Region, Anchorage, AK. 31 pp.
- Alaska Department of Labor (ADL). 1991. Alaska Population Overview 1990 Census and Estimates. Demographic Unit, Research and Analysis. Juneau, Alaska. 144 pp.
- Alaska Department Natural Resources (ADNR). 1988. Kuskokwim Area Plan. Juneau, Alaska.
- Alt, K. T. 1978. Inventory and cataloging of sport fish and sport fish waters of western Alaska. Alaska Department of Fish and Game, Federal Aid in Fish Restoration. Annual Report of Progress. 1977-1978, Project F-9-10, 19(G-I-P): 36-76.
- _____. 1981. A life history study of sheefish and whitefish in Alaska. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual report of progress, 1980-1981. Project F-9-13, 22(RII): 28 pp.
- _____. 1984. Inventory and cataloging of sport fish and sport fish waters of western Alaska. Part B: Fisheries resource investigations and sheefish adaptability studies. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1983-1984. Project F-9-16, 25(G-I-P-B).
- _____. 1987. Review of Sheefish (*Stenodus leucichthys*) studies in Alaska. Alaska Department of Fish and Game. Fishery Manuscript No. 2. Juneau, Alaska. 69 pp.
- Alaska National Interest Lands Conservation Act (ANILCA). 1980. Act of Congress. Public Law 96-487, December 2, 1980.

LITERATURE CITED (Continued)

- Arvey, W. D. and A. Burkholder. 1990. Stock assessment of northern pike in the vicinity of the Yukon River Haul Road crossing, 1988 and 1989. Alaska Department of Fish and Game, Fishery Manuscript No. 90-1, Anchorage.
- Arvey, W. D. 1991. Annual management report for sport fisheries in the Arctic Yukon and Kuskokwim Region, 1989. Alaska Department of Fish and Game. Fishery Management Report, No. 91-3, Anchorage.
- Arvey, W. D., M. F. Merritt, M. J. Kramer, and J. E. Hallberg. 1990. Annual management report for sport fisheries in the Arctic Yukon and Kuskokwim Region, 1986. Alaska Department of Fish and Game, Fishery Management Report No. 90-1, Anchorage.
- Arvey, W. D., M. J. Kramer, J. E. Hallberg, J. F. Parker, and A. L. DeCicco. 1991. Annual management report for sport fisheries in the Arctic Yukon and Kuskokwim Region, 1987. Alaska Department of Fish and Game, Fishery Management Report No. 90-1, Anchorage.
- Arvey, W. D., and J. F. Parker. 1991. Annual management report for sport fisheries in the Arctic Yukon and Kuskokwim Region, 1988. Alaska Department of Fish and Game, Fishery Management Report No. 91-2, Anchorage.
- Arvey, W. D., and M. J. Mills. 1991. Sport harvest of anadromous salmon in the Yukon River drainage, 1977-1989. Alaska Department of Fish and Game. Fishery Data Series No. 91-41, Anchorage.
- Baxter, R. 1977. Hoholitna River reconnaissance survey, 1977. Kuskokwim Resource Report #3. Arctic Yukon and Kuskokwim Region, Commercial Fisheries Division, Alaska Department of Fish and Game, Anchorage.
- Behnke, R. J. 1980. A systematic review of the genus *Salvelinus*. In Charrs: salmonid fishes of the genus *Salvelinus*, E.K. Balon, ed. Dr. W. Junk publishers, The Hague.
- Bendock, T. N.. 1980. Inventory and cataloging of arctic area waters. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1979-1980. Project F-9-12, 21(G-I-I).
- _____. 1982. Inventory and cataloging of arctic waters. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1981-1982. Project F-9-14, 23(G-I-I).
- _____. 1983. Inventory and cataloging of arctic area waters. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1982-1983. Project F-9-15, 20(G-I-I).

LITERATURE CITED (Continued)

- Bendock, T. N. and Burr, J. M. 1984. Inventory and cataloging of arctic area waters. Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual performance report, 1983-1984. Project F-9-16, 25(G-I-I).
- Bendock, T. N. and J. M. Burr. 1985. Unpubl. Freshwater fish distributions in the central arctic coastal plain (Topagoruk River to Ikpikpuk River). Alaska Department of Fish and Game, Division of Sport Fish, Fairbanks.
- Bernard, D. R. and A. L. DeCicco. 1987. Stock assessment of the Dolly Varden char of Kotzebue Sound. Alaska Department of Fish and Game, Fisheries Data Series No. 19, Juneau.
- Burkholder, A. 1991a. Abundance and composition of northern pike, Harding Lake, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-9, Anchorage.
- Burkholder, A. 1991b. Stock composition of northern pike captured in Minto Flats during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-23, Anchorage.
- Burkholder, A. 1991c. Abundance, egg production, and age-sex-length composition of the chinook salmon escapement in the Salcha River, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-5, Anchorage.
- Burr, J. M. 1987. Synopsis and bibliography of lake trout, *Salvelinus namaycush* in Alaska. Alaska Department of Fish and Game, Fishery Manuscript No. 5, Juneau.
- Burr, J. M. 1991. Lake trout population studies in interior Alaska, 1990, including abundance estimates of lake trout in Glacier, Sevenmile, and Paxson lakes during 1989. Alaska Department of Fish and Game, Fishery Data Series No. 91-7, Anchorage.
- Clark J. H. 1991. Abundance, length, and age composition of sockeye salmon, Arctic char, and least cisco in pelagic waters of Harding Lake, Alaska, 1989 and 1990.. Alaska Department of Fish and Game, Fishery Data Series No. 91-4, Anchorage.
- Clark J. H., Viavant T. R., Skaugstad C. and McKinley T. R. 1991. Growth, survival, and costs of rearing game fish in floating net-pens at Harding Lake, Alaska, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-2, Anchorage.
- Clark, R. A. 1987. Arctic grayling harvests, stock status, and regulatory concerns in the Arctic Yukon Kuskokwim region. A report to the Alaska Board of Fisheries, December, 1987. Alaska Department of Fish and Game, 1300 College Rd. Fairbanks, Alaska, 99701.

LITERATURE CITED (Continued)

- Clark, R. A. 1990. Stock status of Chena River Arctic grayling. Alaska Department of Fish and Game, Fishery Data Series No. 90-4, Anchorage.
- Clark R. A. 1991. Stock status of Chena River Arctic grayling during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-35, Anchorage.
- Clark R. A. 1991. Stock assessment of Arctic grayling in Fielding Lake. Alaska Department of Fish and Game, Fishery Data Series No. 91-11, Anchorage.
- Clark R. A. Fleming, D. M. and W. P. Ridder. 1991. Stock assessment of Arctic grayling in the Salcha, Chatanika, and Goodpaster rivers. Alaska Department of Fish and Game, Fishery Data Series No. 91-15, Anchorage.
- DeCicco, A. L. 1985. Inventory and cataloging of sport fish and sport fish waters of western Alaska with emphasis on Arctic char life history studies. Alaska Department of Fish and Game Federal Aid in Fish Restoration, Annual Performance Report, 1984-1985. Project F-9-17, Job G-I-P-A. Volume 26 pp 41-134.
- DeCicco A. L. 1991. Seward Peninsula Arctic grayling study, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-24, Anchorage.
- Doxey, M. 1987. Tanana drainage lake stocking evaluations, 1986. Alaska Department of Fish and Game, Fishery Data Series No. 31, Juneau.
- Doxey, M. 1989. Evaluation of stocked waters in the Tanana drainage, 1988. Alaska Department of Fish and Game, Fishery Data Series No. 106, Juneau.
- Evenson, M. J. 1990a. Movement, abundance, and length composition of burbot in rivers of interior Alaska during 1989. Alaska Department of Fish and Game, Fishery Data Series No. 90-3, Anchorage.
- Evenson M. J. 1990b. Age and length at sexual maturity of burbot in the Tanana River, Alaska. Alaska Department of Fish and Game, Fishery Manuscript No. 90-2, Anchorage.
- Evenson M. J. 1991a. Abundance, egg production, and age-sex-size composition of the chinook salmon escapement in the Chena River, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-6, Anchorage.
- Evenson M. J. 1991b. Abundance and size composition of burbot in rivers of interior Alaska during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-33, Anchorage.
- Fleming D. F. and W. P. Ridder 1991. Stock assessment of Arctic grayling in Mineral Lake Outlet, 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-22, Anchorage.

LITERATURE CITED (Continued)

- Francisco, R. K., K. Schultz, D. Schneiderhan, and D. Huttunen. 1987. Annual Management Report Kuskokwim Area 1985-1986. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bethel. Unpublished agency report.
- Francisco, R. K., C. Burkey Jr., D. B. Molyneaux, C. Anderson, H. H. Hamner, K. E. Hyer, M. W. Coffing, and C. Utermohle 1991. Annual Management Report Kuskokwim Area 1990. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 3B91-11. Bethel, AK. Unpublished report. 193 pp.
- Furniss, R. A. 1974. Inventory and cataloging of arctic area waters. Alaska Department of Fish and Game Federal Aid in Fish Restoration, Annual performance report, 1973-1974. Project F-9-6, 15(G-I-I).
- Gallaway, B., W. Gazey and L. Moulton. 1987. Population trends for arctic cisco in the Colville River of Alaska as reflected by the commercial fishery. Biological Papers, University of Alaska, Fairbanks.
- Hallberg, J. E. 1984. Evaluation of Interior Alaska waters and sport fish with emphasis on managed waters- Fairbanks District. Alaska Department of Fish and Game Federal Aid in Fish Restoration, Annual performance report, 1983-1984. Project F-9-16, 25(G-III-H).
- Hallberg, J. E. and R. A. Holmes. 1987. Abundance and size composition of Chatanika River least cisco and humpback whitefish with estimates of exploitation by recreational spear fishermen. Alaska Department of Fish and Game. Fishery Data Series No. 25. Juneau, Alaska. 26 pp.
- Hallberg, J. E. and A. E. Bingham. 1991. Creel surveys conducted in Interior Alaska during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-56, Anchorage.
- Hemming, C. 1988. Aquatic habitat evaluation of flooded North Slope gravel mine sites (1986-1987). Alaska Department of Fish and Game, Division of Habitat. Technical Report No. 88-1. Juneau 69 pp.
- Hemming, C. R. 1990. Fisheries investigations of flooded North Slope gravel mine sites, 1989. Alaska Department of Fish and Game, Division of Habitat. Technical Report No. 90-2. 38 pp.
- Holmes, R. and Pearse, G. 1987. Northern Pike stock status and regulatory concerns in the Arctic Yukon Kuskokwim Region. A report to the Alaska Board of Fisheries, December, 1987. Alaska Department of Fish and Game, 1300 College Rd., Fairbanks, Alaska, 99701.

LITERATURE CITED (Continued)

- Kretsinger, C. 1987. Fishery inventory of lakes and streams in the Kigluaik mountains and Imuruk Basin watershed (Seward Peninsula). Unpublished paper, U.S. Bureau of Land Management. Kobuk District Office, Fairbanks.
- Lafferty R., Parker, J. F., and D. R. Bernard. 1991. Stock assessment and biological characteristics of burbot in lakes of interior Alaska during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-57, Anchorage.
- Lamke, R. D., B. B. Bigelow, J. L. VanMaanen, R. T. Kemnitz, and K. M. Novcaski. 1991. Water Resources Data Alaska Water Year 1990. U.S. Geological Survey Water-Data Report AK-90-1. 252 pp.
- Lean, C. 1985. Unpubl. Agency rept. Unalakleet River escapement studies, 1984. Alaska Department of Fish and Game, Division of Commercial Fisheries. Nome, Alaska.
- Lean, C., S. Merkouris, H. Hamner, and M. Wyatt. 1986. Unpubl. Agency report. Annual Management Report, 1986, Norton Sound-Port Clarence-Kotzebue. Alaska Department Fish and Game, Division of Commercial Fisheries. Nome, Alaska.
- Merkouris, S.E. and C.F. Lean. 1989. Unpubl. Agency report. Annual Management Report, 1988 Norton Sound-Port Clarence-Kotzebue. Alaska Department of Fish and Game, Division of Commercial Fisheries, AYK Region. Regional Information Report No. 3N89-10. 187 pp.
- Mills, M. J. 1979. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1977-1978. Project F-9-11, 20 (SW-1): 112 pp.
- _____. 1980. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1979-1980. Project F-9-12, 21 (SW-1): 65 pp.
- _____. 1981. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1980-1981. Project F-9-13, 22 (SW-1): 78 pp.
- _____. 1982. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1981-1982. Project F-9-13, 23 (SW-1): 115 pp.
- _____. 1983. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1982-1983. Project F-9-14, 24 (SW-1): 118 pp.

LITERATURE CITED (Continued)

- _____. 1984. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1983-1984. Project F-9-16, 25 (SW-1): 122 pp.
- _____. 1985. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1984-1985. Project F-9-17, 26 (SW-1): 88 pp.
- _____. 1986. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1985-1986. Project F-9-18, 27 (SW-1): 137 pp.
- _____. 1987. Alaska statewide sport fish harvest studies. Alaska Department of Fish and Game. Federal Aid in Fish Restoration, Annual Report of Progress, 1986-1987. Project F-9-19, 28 (SW-1): 91 pp.
- _____. 1988. Alaska statewide sport fisheries harvest report 1987. Alaska Department of Fish and Game, Fishery Data Series No. 52, Juneau.
- _____. 1989. Alaska statewide sport fisheries harvest report 1988. Alaska Department of Fish and Game, Fishery Data Series No. 122, Juneau.
- _____. 1990. Harvest and participation in Alaska sport fisheries during 1989. Alaska Department of Fish and Game, Fishery Data Series No. 90-44, Anchorage.
- _____. 1991. Harvest, catch and participation in Alaska sport fisheries during 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-58, Anchorage.
- Morrow, J. E. 1980. Analysis of the Dolly Varden charr, *Salvelinus malma*, of northwestern North America and northeastern Siberia. In Charrs. Salmonid Fishes of the genus Salvelinus. ed. by Balon, E.K. Publ. by Dr. W. Junk, the Netherlands. pp 323-338.
- Moulton, L., B.J. Gallaway, M.H. Fawcett, W.B. Griffiths, K.R. Critchlow, R.G. Fechhelm, D.R. Schmidt, and J.S. Baker. 1986. 1984 central Beaufort Sea fish study. Prudhoe Bay waterflood project. Environmental monitoring program. Vol. 2, Chapt. 3. LGL, Woodward-Clyde consultants and U.S. Army Corps of Engineers, Anchorage, Alaska. 204 pp
- National Oceanic Atmospheric Administration (NOAA). 1989. Local climatological data for Fairbanks, Nome, Kotzebue and Barrow, Alaska. Annual summary for 1988 and 1989. Fairbanks, Alaska.
- National Park Service (NPS). 1984. Kobuk Valley National Park draft statement for management. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.

LITERATURE CITED (Continued)

- _____. 1985a. Gates of the Arctic National Park General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- _____. 1985b. Cape Krusenstern Monument General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- _____. 1985c. Yukon Charley Rivers National Preserve General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- _____. 1986a. Kobuk National Park General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- _____. 1986b. Noatak National Preserve General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- _____. 1986c. Bering Land Bridge National Preserve General Management Plan. USDI: NPS, Alaska Regional Office, Anchorage, Alaska.
- Peckham, R.D., and M. Doxey. 1983. Evaluation of Interior Alaska waters and sport fish with emphasis on managed waters- Delta District. Alaska Department of Fish and Game. Federal Aid in Fisheries Restoration, Annual report of progress 1982-1983. Project F-9-15, 24(G-III-I). 38 pp.
- Redick, R. R. 1967. A review of literature on lake trout life history with notes on Alaska management. Alaska Department of Fish and Game, Informational Leaflet 111. 19 pp.
- Ridder W. P. 1991. Summary of recaptures of Arctic grayling tagged in the middle Tanana River drainage, 1977 through 1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-34, Anchorage.
- Rue, S., N. Hemming and M. McGuiness. 1987. A method of evaluating the recreation potential of Alaskan rivers. Submitted under contract to the National Park Service, Alaska Regional Office by the Wildlife Federation of Alaska, the Alaska affiliate of the National Wildlife Federation. 118 pp.
- Russell, R. 1980. A fisheries inventory of waters in the Lake Clark Monument area. Alaska Department of Fish and Game, Division of Sport Fish, and United States Department of the Interior, National Park Service.
- Savikko, H., and T. Page 1990. 1989 preliminary Alaska commercial fisheries harvest and values. Regional Informational Report No. 5J90-07. Alaska Department of Fish and Game, Juneau, AK. 102 pp.
- Selkregg, L. L. 1976. Alaska regional profiles: Arctic region. Arctic Environmental Information and Data Center, University of Alaska, Anchorage.

LITERATURE CITED (Continued)

- Skaugstad C. and B. McCracken 1991. Fecundity of chinook salmon, Tanana River, Alaska. Alaska Department of Fish and Game, Fishery Data Series No. 91-8, Anchorage.
- Stokes, J. 1985. Natural resource utilization of four upper Kuskokwim communities. Technical Paper No. 86. Alaska Department of Fish and Game, Division of Subsistence.
- Swainbank, R. C., Bundtzen, T. K. and J. Wood. 1991. Alaska's Mineral Industry 1990. Special Report 45. Alaska Department of Natural Resources (ADNR). Division of Geological and Geophysical Surveys, Juneau Alaska. 78 pp.
- Timmons, S.L. 1990. Abundance and length, age, and sex composition of Chatanika River humpback whitefish and least cisco. Alaska Department of Fish and Game, Fishery Data Series No. 90-2, Anchorage.
- Timmons L. S. 1991. Opinions and regulatory preferences of holders of sport fishing licenses resident in the Arctic-Yukon-Kuskokwim area of Alaska in 1988. Alaska Department of Fish and Game, Fishery Data Series No. 91-1, Anchorage.
- Timmons L. S. 1991. Evaluation of the rainbow trout stocking program for Piledriver Slough. Alaska Department of Fish and Game, Fishery Data Series No. 91-25, Anchorage.
- Timmons L. S. and R. A. Clark 1991. Stock status of Piledriver Slough Arctic grayling. Alaska Department of Fish and Game, Fishery Data Series No. 91-37, Anchorage.
- Townsend, A.H. 1991. Distribution of fishes in Alaska's upper Birch Creek drainage during 1984 and 1990. Alaska Department of Fish and Game Division of Habitat. Technical Report No. 91-2. 14 pp.
- U.S. Army Corps of Engineers, Alaska District. 1967. Harbors and Rivers in Alaska. Interim Report No. 6. Northwestern Alaska.
- U.S. Department of Interior/Bureau of Land Management Alaska Fire Service. 1991. 1990 Alaska Fire Service fire season statistics. Intelligence section. FCC Fort Wainwright Alaska. 42 pp.
- U.S. Department of Agriculture (USDA). 1990. Alaska snow surveys, May 1, 1990. Soil Conservation Service, Anchorage, Alaska. 27 pp.
- U.S. Fish and Wildlife Service (USFWS). 1982. Arctic National Wildlife Refuge coastal plain resource assessment. Initial report baseline study of the fish, wildlife, and their habitats. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503. 507 pp.

LITERATURE CITED (Continued)

- _____. 1985. Yukon Flats National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1986. Selawik National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1987a. Yukon Delta National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1987b. Nowitna National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1987c. Koyukuk and Innoko National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1987d. Kanuti National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1988. Arctic National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- _____. 1987. Tetlin National Wildlife Refuge Plan. USDI: USFWS, Region 7. 1011 E. Tudor Rd., Anchorage, Alaska 99503.
- Viavant T. and J. H. Clark 1991a. Distribution and relative abundance of stocked species in Harding Lake, 1987-1990. Alaska Department of Fish and Game, Fishery Data Series No. 91-20, Anchorage.
- Viavant T. and J. H. Clark 1991b. Experimental hook and line fishing at Harding Lake, Alaska, 1989-1991. Alaska Department of Fish and Game, Fishery Data Series No. 91-38, Anchorage.
- Weber, P. K. 1986. Downstream effects of placer mining in the Birch Creek Basin, Alaska. Alaska Department of Fish and Game, Habitat Division. Technical Report no. 86-7. 21 pp.
- Wild and Scenic Rivers Act 1968. U.S. Public Law 90-542. Amended December, 1980 with Public Law 96-603-605.
- Whitmore, C., D. J. Bergstrom, and F. M. Andersen. 1987. Annual Management Report, 1987, Yukon Area. Regional Information Report No. 3A88-30. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage and Fairbanks. 167 pp.
- Whitmore, C., D. J. Bergstrom, F. M. Andersen, G. Sandone, J. Wilcock, L. Barton and D. Messiar. 1990. Annual Management Report, Yukon Area, 1988. Regional Information Report No. 3A90-28. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage and Fairbanks.

APPENDIX A

Appendix A. National Wild and Scenic Rivers in the Arctic Yukon Kuskokwim Region.

Streams Within The National Park System

Alatna River. The main stem within the Gates of Arctic National Park.

John River. The portion of the river within the Gates of the Arctic National Park.

Kobuk River. The portion within the Gates of the Arctic National Park and Preserve.

Noatak River. The river from its source in the Gates of the Arctic National Park to its confluence with the Kelly River in the Noatak National Preserve.

North Fork of the Koyukuk River. The portion within the Gates of the Arctic National Park.

Tinayguk River. The portion within the Gates of the Arctic National Park.

Salmon River. The portion with the Kobuk Valley National Park.

Charley River. The entire river, including its major tributaries, Copper Creek, Bonanza Creek, Hosford Creek, Derwent Creek, Flat-Orthmer Creek, Crescent Creek, and Moraine Creek, within the Yukon-Charley Rivers National Preserve.

Streams Within The National Wildlife Refuge System

Andreafsky River. The portion from its source, including all headwaters, and the East Fork, within the boundary of the Yukon Delta National Wildlife Refuge.

Ivishak River. The portion from its source, including all headwaters and an unnamed tributary from Porcupine Lake with the boundary of the Arctic National Wildlife Range.

Nowitna River. The portion from the point where the river crosses the west limit of township 18S, Range 22E, Kateel River meridian, to its confluence with the Yukon River within the boundaries of the Nowitna National Wildlife Refuge.

Selawik River. The portion from a fork of the headwaters in township 12N, Range 10E, Kateel River meridian to the confluence of the Kugarak River; within the Selawik National Wildlife Refuge .

Sheenjok River. The segment within the Arctic National Wildlife Refuge.

Wind River. The portion from its source, including all headwaters and one unnamed tributary in township 13 S, within the boundaries of the Arctic National Wildlife Refuge.

-continued-

Streams Located Outside National Parks and Refuges

Fortymile River. The mainstem within the State of Alaska; O'Brien Creek; South Fork; Napoleon Creek, Franklin Creek, Uhler Creek, Walker Fork downstream from the confluence of Liberty Creek; Wade Creek; Mosquito Fork downstream from the vicinity of Ketchumstuk; West Fork Dennison Fork downstream from the confluence of Logging Cabin Creek; Dennison Fork downstream from the confluence of West Fork Dennison Fork; Logging Cabin Creek; North Fork; Hutchison Creek; Champion Creek; the Middle Fork downstream from the confluence of Joseph Creek; and Joseph Creek.

Delta River. The segment from and including all of the Tangle Lakes to a point one-half mile north of Black Rapids.

Beaver Creek. The segment of the main stem from the vicinity of the confluence of the Bear and Champion Creeks downstream to its exit from the northeast corner of township 12N, Range 6E, Fairbanks meridian within the White Mountains National Recreation Area, and the Yukon Flats National Wildlife Refuge.

Birch Creek. The segment of the main stem from the south side of Steese Highway in township 7N, Range 10E, Fairbanks meridian, downstream to the south side of the Steese Highway in township 10N, Range 16E,.

Unalakleet River. The segment of the main stem from the headwaters in township 12S, Range 3W, Kateel River meridian extending downstream approximately 65 miles to the western boundary of township 18S, range 8W.

APPENDIX B

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH AND GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-Z-S-8-90

Issued at Nome, Alaska
11:00 p.m. ADT July 2, 1990

Effective Date: 6:00 p.m. ADT
Thursday, July 2, 1990

Expiration Date: December 31, 1990
unless superseded by subsequent
emergency order.

EXPLANATION:

This emergency order closes the Nome River to the taking of chum salmon on sport fishing gear. Effective 6:00 p.m. July 2 until further notice, all chum salmon caught on rod and reel in the Nome River must be immediately released.


REGULATION:

5 AAC 70.020. SEASONS AND BAG LIMITS, SEWARD PENINSULA WATERS (e). is amended to read:

5 AAC 70.020. SEASONS AND BAG LIMITS, SEWARD PENINSULA WATERS (e). The Nome River is closed to the taking of chum salmon.

Don W. Collinsworth
Commissioner

By delegation to :


Alfred L. Decicco
Area Management Biologist

JUSTIFICATION:

A boat survey on June 30, 1990 showed that only about 40 chum salmon had escaped into the Nome River. The 1985 and 1986 parent year escapements were below the escapement goal of 2,000 fish and the return of chum salmon spawners to the Nome River is expected to be below that goal. In only two of the last ten years has the escapement goal been reached. In order to rebuild depleted chum salmon runs in the Nome River, conservative measures are necessary at this time.

DISTRIBUTION:

The distribution list is on file at the Fairbanks and Nome Department of Fish and Game offices.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH AND GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-WF-01-90

Issued at Fairbanks, October 10, 1990

Effective Date 6:00 p.m.
Thursday, October 11, 1990

Expiration date December 31, 1990
unless superseded by subsequent
emergency order.

EXPLANATION:

This emergency order closes the Chatanika River to the taking of whitefish by sport fisherman.


REGULATION:

5 AAC 70.035. METHODS AND MEANS, is amended to read:

(i) the Chatanika River is closed to the taking of whitefish from 6:00 p.m. Thursday, October 11, 1990 until December 31, 1990 unless superseded by a subsequent emergency order.

Don W. Collinsworth
Commissioner

By delegation to:


Jerome E. Hallberg
Area Management Biologist

JUSTIFICATION:

- 1.) The estimated abundance of humpback whitefish in the Chatanika River in 1990 is less than 9,000 fish, approximately half of what it was in 1989 and only 25 percent of the 1988 estimate.
- 2.) Catch rates in the spear fishery are less than those in previous years.
- 3.) Weak year classes in 1985 and 1986 have drastically reduced the recruitment to the humpback whitefish population and have contributed to the declines in abundance.

Consequently, an emergency order closing the whitefish sport fishery is necessary to provide for conservation of the spawning stock and to prevent further depletion of the humpback whitefish population in the Chatanika River.

SPORT FISHING

Emergency Order

ALASKA DEPARTMENT
OF FISH AND GAME

Under Authority of AS 16.05.060

Emergency Order No. 3-NP-02-90

Issued at Fairbanks, December 6, 1990

Effective Date: 12:01 a.m.
Monday, December 10, 1990

Expiration Date: Until superseded by
regulation or by subsequent emergency
order.

EXPLANATION:

This emergency order establishes 1) a 26-inch minimum length limit for northern pike harvested in Harding Lake, 2) a ban on the use of spears and bow and arrows in Harding Lake and, 3) closes Harding Lake to sport fishing for northern pike (during the spawning period) from 12:01 a.m. April 1, to 11:59 p.m. May 31.

REGULATION:

5 AAC 70.010, 5 AAC 70.020, and 5 AAC 70.035 are therefore amended to read:

5 AAC 70.010. FISHING SEASONS: (d) northern pike may be taken in Harding Lake only from June 1 through March 31;


5 AAC 70.020. BAG LIMITS, POSSESSION LIMITS, AND SIZE LIMITS: (q) in Harding Lake, the minimum legal size limit for northern pike is 26 inches;

5 AAC 70.035. METHODS AND MEANS: (i) In the Arctic-Yukon-Kuskokwim Area suckers and burbot may be taken by spear, or bow and arrow from January 1, through December 31. Except as specified in (j) of this section, northern pike and whitefish (excluding sheefish) may be taken by spear, or bow and arrow from September 1, through April 30, and from January 1 through December 31, may be speared by persons completely submerged;

(j) Harding Lake is closed to the taking of fish by spear or by bow and arrow;

Don W. Collinsworth
Commissioner

By delegation to:


Jerome E. Hallberg
Area Management Biologist

JUSTIFICATION:

Available harvest information and biological data indicate that the Harding Lake northern pike population is being over-harvested. Harvest surveys indicate that removal by sport fishermen has totalled approximately 2,000 fish annually since 1987. Studies conducted at Harding lake during the spring of 1990, reveal that total abundance of northern pike (greater than 12 inches in length) has decreased to about 1,700 fish, that there has been poor recruitment of young fish in recent years and that few large fish remain in the population. Regulation changes are therefore required to prevent continuing over-harvest, to begin rebuilding the population, and to allow continuation of a limited recreational fishery.

The 26-inch minimum length limit and the elimination of spears and bow and arrows are necessary to reduce harvest to a sustainable level and assure that fish have the opportunity to spawn at least twice before being harvested. The spring closure will minimize disturbance of northern pike prior to and during the spawning period. Although the spearing or taking of burbot by bow and arrow in Harding Lake presents no threat to the conservation of that species, it is thought that prohibition of this gear type will enhance enforcement of regulations designed to protect the northern pike resource.

DISTRIBUTION:

The distribution list of this emergency order is available from the Department of Fish and Game office in Fairbanks.



APPENDIX C

ARCTIC-YUKON-KUSKOKWIM AREA

SEASON AND CATCH LIMIT

WATER AND SPECIAL REGULATIONS

	SALMON	DOLLY VARDEN ARCTIC CHAR/ LAKE TROUT/ RAINBOW/ GRAYLING	SHEEFISH	NORTHERN Pike BURBOT	HALIBUT	OTHER FISH SHELLFISH
Goodnews River drainage Special Regulations: 1. In all flowing waters of the Goodnews River drainage upstream of the Togiak National Wildlife Refuge wilderness area boundary only unbaited, single-hook artificial lures may be used. 2. No person may sport fish from a boat or the river bank within 300 feet of a legally operating subsistence set gillnet downstream of the Togiak National Wildlife Refuge Wilderness area boundary.	A,E	F,G,J,I	L	N,P		U
Kanektok River drainage: Special Regulations: 1. In all flowing waters of the Kanektok River drainage upstream of the Togiak National Wildlife Refuge wilderness area boundary only unbaited, single-hook artificial lures may be used. 2. No person may sport fish from a boat or the river bank within 300 feet of a legally operating subsistence set gillnet downstream of the Togiak National Wildlife Refuge Wilderness area boundary.	A,E	F,G,J,I	L	N,P		U
Kivalina River drainage:	A,C	G,K		N,P		U
Kobuk River drainage: upstream of the mouth of Mauneluk River remainder of the Kobuk River	A,C A,C	G,J,I G,J,I	M L	N,P N,P		U U
Kuskokwim Bay drainages: All waters that drain into Kuskokwim Bay (excluding the Kuskokwim River) from Cape Avinof to Cape Newenham	A,E	F,G,J,I	L	N,P		U
Kuskokwim River drainage:	B,E	F,G,J,I	L	N,P		U
Noatak River drainage:	A,C	G,K,I	L	N,P		U
North Slope waters: (All waters drainage into the Arctic Ocean and all waters draining into the Chukchi Sea north of Point Hope)	A,C	G,K,I	L	N,P		U
Seward Peninsula waters: (All waters draining into the Bering Sea from Cape Darby to Cape Prince of Wales on the Seward Peninsula) Special Regulations: *Salmon Lake, its tributaries, and the outlet stream 300 feet downstream from the lake outlet are closed to salmon fishing.	B*,D*	H,J		N,P		U
Trans-Alaska Pipeline (A corridor 5 miles wide on each side of the alignment.)	Closed	G,J,I	L	N,P		U
Unalakleet River drainage:	B,C	H,J		N,P		U
Wulik River drainage:	A,C	G,K		N,P		U
Yukon River drainage: from the mouth of the Tanana River upstream to, and including, the Hodzana River remainder of the drainage	A,C A,C	G,J,I G,J,I	L L	O,P N,P		U U

CODE KEY: ARCTIC-YUKON-KUSKOKWIM AREA Use these codes to determine open season, catch and length limits.

CODE	OPEN SEASON	BAG, POSSESSION, AND SIZE LIMITS
A KING SALMON	Entire Year	3 per day, 3 in possession, only 2 can exceed 28 inches
B KING SALMON	Entire Year	1 per day, 1 in possession, no size limit
C OTHER SALMON	Entire Year	10 per day, 10 in possession, no size limit
D OTHER SALMON	Entire Year	10 per day, 10 in possession, only 3 which may be chum salmon and coho salmon, in combination
E OTHER SALMON	Entire Year	5 per day, 5 in possession, no size limit
F RAINBOW TROUT	Entire Year	2 per day, 2 in possession, only 1 over 20 inches
G GRAYLING	Entire Year	10 per day, 10 in possession, no size limit
H GRAYLING	Entire Year	5 per day, 5 in possession, only 1 over 15 inches
I LAKE TROUT	Entire Year	4 per day, 4 in possession, no size limit
J ARCTIC CHAR/DOLLY VARDEN	Entire Year	10 per day, 10 in possession; no size limit

ARCTIC-YUKON-KUSKOKWIM AREA

K	ARCTIC CHAR/DOLLY VARDEN	Entire Year	10 per day, 10 in possession, only 2 over 20 inches
L	SHEEFISH	Entire Year	10 per day, 10 in possession, no size limit
M	SHEEFISH	Entire Year	2 per day, 2 in possession, no size limit
N	NORTHERN PIKE	Entire Year	10 per day, 10 in possession, no size limit
O	NORTHERN PIKE	Entire Year	5 per day, 5 in possession, only 1 over 20 inches
P	BURBOT	Entire Year	15 per day, 15 in possession, no size limit
Q	HALIBUT	Feb. 1—Dec. 31	2 per day, 4 in possession, no size limit
R	KING CRAB (Red and blue in combination)	In waters south of 60° north latitude: June 1—Jan. 31 In waters north of 60° north latitude: Entire Year	6 per day, 6 in possession, males only: Red King Crab 4½ inches or more; Blue King Crab 5½ inches or more 6 per day, 6 in possession, males only: Red King Crab 4½ inches or more; Blue King Crab 5½ inches or more
S	DUNGENESS CRAB	Entire Year	5½ inches or more: 12 per day, 12 in possession, males only
T	TANNER CRAB (C. Bairdi and C. Opilio in combination)	Entire Year	12 per day, 12 in possession, males only: C. Bairdi—5½ inches or more; C. Opilio—3 1/8 inches or more
U	OTHER FISH/SHELLFISH	Entire Year	No bag, possession or size limit

OTHER ARCTIC—YUKON—KUSKOKWIM AREA REGULATIONS

METHODS AND MEANS. 1. In all lakes, multiple hooks with gap between point and shank greater than one-half inch may be used for taking fish other than salmon.

2. Sucker and Burbot, may be taken by spear or bow and arrow from January 1 through December 31.

3. Northern Pike and Whitefish (excluding sheefish) may be taken by spear or bow and arrow from September 1 through April 30 and may be speared by persons completely submerged from January 1 through December 31.

SHELLFISH METHODS AND MEANS. (a) Shellfish may be taken only as follows:

(1) shrimp may be taken with pots and ring nets;

(2) crab may be taken with pots, ring nets, diving gear, dipnets, hooked or hookless lines either operated by hand or attached to a pole or rod, or by hand.

(3) clams may be taken with rakes, shovels, and manually operated clam guns.

(b) No more than five pots per person, regardless of type, and a maximum of ten pots per vessel, regardless of type, may be used to take shellfish at any time

(c) Each sport fisherman shall plainly and legibly inscribe his or her first initial, last name, and home address on a keg or buoy attached to each pot. A keg or buoy attached to a pot must also be inscribed with the name or the Coast Guard number of the vessel used to operate the pot.

(d) Escape mechanisms as described in the Personal Use Shellfish Regulations (page 56) must be provided for each pot.

(e) The bag, possession, and pot limits for shellfish in this chapter are not in addition to those allowed under subsistence fishing regulations, nor to those allowed under personal use fishing regulations.

(f) No person may mutilate or otherwise disfigure any crab in any manner which would prevent the determination of the minimum size restrictions until the crab has been processed for human consumption. No person may take or possess shellfish smaller than the minimum legal size limits.

(g) The operator of a commercially licensed and registered shrimp fishing vessel used in the sport fish taking of shrimp during a closed commercial shrimp season or within a closed commercial shrimp district, section, or subsection may not possess more than 500 pounds onboard the vessel.

HARVEST RECORDING FORM. A person must obtain a harvest recording form from the department prior to fishing for king crab in the Norton Sound section of the Northern District.

TANANA AREA

This is a summary of the official regulations codified in 5 AAC 70.001-060 (as they apply to waters of the Tanana River Drainage) which are available for inspection at libraries, department offices, and Department of Public Safety offices throughout the state.


INSTRUCTIONS:

1. Find the water (alphabetically listed) that you intend to fish. If the water, or any portion of it, is not listed, the regulations in the shaded entry apply.
2. Use the Code Key to determine open season, catch, and length limits. Read Special Regulations.
3. An asterisk (*) denotes Special Regulations apply.

SEASON AND CATCH LIMIT

WATER AND SPECIAL REGULATIONS

All waters not listed below	SALMON	DOLLY VARDEN/ ARCTIC CHUM/ LAKE TROUT/ RAINBOW/ GRAYLING	WHYTEFISH/ SHEEPFISH	NORTHERN PIKE/ BURBOT	OTHER FISH
	A,B	C,D,F,J,M	N,O	P,R,S	U
Chena River and its tributaries: downstream from a department marker 300 feet downstream from the Chena River flood control structure remainder of drainage Special Regulations: (Unless otherwise specified, each special regulation applies to the entire Chena River and its tributaries) 1. Upstream of the Chena River Dam, only unbaited, single-hook artificial lures may be used. 2. Downstream of the Chena River Dam, bait may be used only on hooks with a gap size larger than 3/4 inch. 3. *From April 1 to the first Saturday in June grayling may not be possessed or retained; all grayling caught between these dates must be released immediately. 4. The Chena River from the confluence of the South Fork (river mile 77) to the first bridge (river mile 88) is designated a catch and release water for grayling. Grayling may not be possessed or retained. All grayling caught must be released immediately.	A,B Closed	I* I*	N,O N,O	P,S P,S	U U
Delta Clearwater River and its tributaries: Special Regulations: 1. Only unbaited artificial lures or flies may be used. 2. *Grayling may not be possessed or retained from April 1 to the first Saturday in June; all grayling caught between these dates must be released immediately.	B	H*	N,O	P,S	U
Fielding Lake Special Regulations: *No set lines may be used.		F,K	O	T*	U
Goodpaster River drainage:	Closed	F	N,O	P,S	U
Harding Lake Special Regulations: *No set lines may be used.	A,B	C,F,K,M	N,O	P,T*	U
Kantishna River drainage Special Regulations: *From August 15 through December 31 the Toklat River and the Kantishna River downstream from the mouth of the Toklat River are closed to the taking of chum salmon. All chum salmon caught between these dates must be released immediately.	*A,B	C,D,F,J,M	N,O	P,S	U
Mineral Lake Outlet (From the outlet of Mineral Lake to its confluence with the Little Tok River) Special Regulations: 1. Only unbaited artificial lures or flies may be used. 2. *Grayling may not be possessed or retained from April 1 to the first Saturday in June; all grayling caught between these dates must be released immediately.		H*	N,O	P,S	U
Piledriver Slough upstream from its confluence with Moose Creek: Special Regulations: Only unbaited artificial lures or flies may be used.	B	D,G	N,O	P,S	U
Richardson Clearwater River drainage: Special Regulations: 1. Only unbaited artificial lures or flies may be used. 2. *Grayling may not be possessed or retained from April 1 to the first Saturday in June; all grayling caught between these dates must be released immediately.	B	H*	N,O	P,S	U



TANANA AREA	SEASON AND CATCH LIMIT					
	SALMON	DOLLY VARDEN/ ARCTIC CHAR/ LAKE TROUT/ RAINBOW/ GRAYLING	WHITEFISH/ SHEEPSKIN	NORTHERN PKE/ BURBOT	OTHER FISH	
WATER AND SPECIAL REGULATIONS						
Salcha River and its tributaries: downstream from a marker placed approximately 2½ miles upstream from the Richardson Hwy. Bridge: upstream from a marker placed approximately 2½ miles upstream from the Richardson Hwy. Bridge:	A,B Closed	H* H*	N.O N.O	P,S P,S	U U	
Special Regulations: 1. Only unbaited artificial lures or flies may be used; except that bait may be used only on hooks with a gap size larger than ¼ of an inch. 2. *Grayling may not be possessed or retained from April 1 to the first Saturday in June; all grayling caught between these dates must be released immediately. 3. Fishing from the Richardson Hwy. Bridge over the Salcha River is prohibited.						
Sensing Lake	A,B	E,F				
Shaw Creek and its tributaries: Special Regulations: 1. Only unbaited artificial lures or flies may be used upstream of the Richardson Hwy. Bridge. 2. Downstream of the Richardson Hwy. Bridge, bait may be used only on hooks with a gap size larger than ¼ inch. 3. *Grayling may not be possessed or retained from April 1 to the first Saturday in June in Shaw Creek drainage and the Tanana River from 2 miles upstream of the mouth of Shaw Creek to 2 miles downstream of the mouth of Shaw Creek; all grayling caught between these dates must be released immediately.	B	H*	N.O	P,S	U	
"T" Lake Special Regulations: *No set lines may be used.				O	P,T*	U
Tangle Lakes Special Regulations: *No set lines may be used.		F,L	O	T*	U	
Tolovana River drainage (Includes Minto Flats, Chatanika River, Tatalina River, and Goldstream Creek:	A,B	F	N.O	Q,S	U	

CODE KEY: TANANA AREA Use these codes to determine open season, catch and length limits.

CODE	OPEN SEASON	CATCH, POSSESSION, AND SIZE LIMITS
A	KING SALMON	
	18 inches or more	Entire Year
	Less than 18 inches	Entire Year
		1 per day, 1 in possession
		10 per day, 10 in possession
B	OTHER SALMON	
	18 inches or more	Entire Year
	Less than 18 inches	Entire Year
		3 per day, 3 in possession
		10 per day, 10 in possession
C	RAINBOW TROUT (In Lakes)	
	20 inches or more	Entire Year
	Less than 20 inches	Entire Year
		2 per day, 2 in possession
		10 per day, 10 in possession
D	RAINBOW TROUT (Flowing waters)	Entire Year
		5 per day, 5 in possession, no size limit
E	RAINBOW TROUT	Entire Year
		3 per day, no size limit
F	GRAYLING	Entire Year
		5 per day, 5 in possession, no size limit
G	GRAYLING	Entire Year
		5 per day, 5 in possession, 12 inch minimum length limit
H	GRAYLING	First Saturday in June through March 31
		5 per day, 5 in possession, 12 inch minimum length limit
I	GRAYLING	First Saturday in June through March 31
		2 per day, 2 in possession, 12 inch minimum length limit
J	LAKE TROUT	Entire Year
		2 per day, 2 in possession, no size limit
K	LAKE TROUT	Entire Year
		2 per day, 2 in possession, 18 inch minimum length limit
L	LAKE TROUT	Entire Year
		1 per day, 1 in possession, 18 inch minimum length limit

TANANA AREA

M	ARCTIC CHAR/DOLLY VARDEN	Entire Year	10 per day, 10 in possession, no size limit
N	SHEEPFISH	Entire Year	2 per day, 2 in possession, no size limit
O	WHITEFISH (Excluding Sheefish)	Entire Year	15 per day, 15 in possession, no size limit
P	NORTHERN PIKE	Entire Year	6 per day, 5 in possession, only 1 over 30 inches
Q	NORTHERN PIKE	June 1 through October 14	6 per day, 5 in possession, only 1 over 30 inches
R	BURBOT (in Lakes)	Entire Year	5 per day, 5 in possession, no size limit
S	BURBOT (Flowing Waters)	Entire Year	15 per day, 15 in possession, no size limit
T	BURBOT	Entire Year	2 per day, 2 in possession, no size limit
U	OTHER FISH	Entire Year	No bag, possession or size limit

OTHER TANANA AREA REGULATIONS

METHODS AND MEANS. (1) In all lakes, multiple hooks with gap between point and shank greater than one-half inch may be used for taking fish other than salmon.

(2) Suckers and burbot may be taken by spear or bow and arrow from January 1 through December 31.

(3) Northern pike and whitefish (excluding sheefish) may be taken by spear or bow and arrow from September 1 through April 30 and may be speared by persons completely submerged from January 1 through December 31.

(4) Burbot may be taken by set lines in all lakes in the Tanana River drainage (except for Fielding, Harding, "T", and Tangle Lakes) from October 15 through May 15.

(5) Burbot may be taken by set lines in rivers year round.

(6) The total aggregate number of hooks used on set lines, closely attended gear, and ice fishing gear may not exceed the daily bag limit for burbot in the water being fished.

(7) All ice houses not removed from the ice at the end of the day's fishing must be registered and a permit secured from the Department. Each registered ice house must have permit numbers displayed on its side and roof in distinguishable numbers not less than 12 inches in height.

KING SALMON HARVEST RANGES (1) The guideline harvest range for king salmon in the Chena River (downstream from a marker 300 feet downstream from the Chena flood control structure) recreational fishery is 300 to 600 fish. (2) The guideline harvest range for king salmon in the Salcha River (downstream from a marker placed approximately 2.5 miles upstream from the Richardson Highway bridge) recreational fishery is 300 to 700 fish.

STATEWIDE PROVISIONS—PERSONAL USE SHELLFISH

A sport fishing license is required to take shellfish for personal use in all areas of the state under the guidelines listed below. However, in the southeast and Yakutat areas nonresidents are required to abide by the bag and possession and methods and means regulations listed on pages 10 through 14.

Residents 60 years or more in age and all persons less than 16 years in age are exempt from these license requirements.

Harvest, bag and possession, and pot limits for shellfish allowed under personal use may not be added to those allowed for subsistence use.

Unless otherwise allowed in specified areas, gear that may be used in the taking of shellfish is as follows:

Shrimp - pots, ring nets, or trawl;

Crab - pots, ring nets, diving gear, dip nets, hooked or hookless hand lines and by hand; lines attached to rods or poles may only be used when fishing through the ice in the Bering Sea;

Clams - rakes, shovels, manually operated clam guns.

Abalone - abalone irons, diving gear, by hand;

Other Shellfish - all gear above and trawls, hand jigging, and hand operated dredges.

It is unlawful to buy or sell personal use caught shellfish, their parts, or their eggs.

Each personal use fisherman shall plainly and legibly inscribe his first initial, last name and address on a keg or buoy attached to unattended fishing gear. Kegs or buoys attached to pots must also be inscribed with the name or U.S. Coast Guard number of the vessel used to operate the pots.

No more than 5 pots of any type per person and 10 pots of any type per vessel may, in aggregate, be used to take personal use shellfish at any given time.

APPENDIX D

Appendix C. (Page 6 of 7).

TANANA AREA

WATER AND SPECIAL REGULATIONS

- Selkirk River and its tributaries:
downstream from a marker placed approximately 2 1/2 miles upstream
Richardson Hwy. Bridge;
 - upstream from a marker placed approximately 2 1/2 miles upstream
Richardson Hwy. Bridge;
 - Special Regulations: 1. Only unbaited artificial lures or
caps that bear a marker placed between these dates
in June, all grayling caught between these dates
 - 2. Grayling may not be used only on hooks with a gap size
of the Richardson Hwy. Bridge
 - 3. Fishing from the Richardson Hwy. Bridge
- Sansing Lake**
- Shore Creek and its tributaries:
of the Richardson Hwy. Bridge;
 - Special Regulations: 1. Only un-
baited artificial lures or
with a gap size larger
than 1/4 inch
 - 2. Downstream of the Ri-
chardson Hwy. Bridge
 - 3. Grayling may
in June in Shu-
of the mou-
all gray-
"T" L

Appendix D. Partial listing of wilderness lodges, guiding, and outfitting operations in the AYK Region.

Location	Operation	Remarks
<u>Seward Peninsula/Norton Sound:</u>		
White Mtn.	Fishing lodge, guiding	Opened in 1986
Niukluk, Fish R.	Outfitters	
Unalakleet R.	1 lodge	
<u>Kuskokwim:</u>		
Tonzona R.	2 lodges	Includes hunting
Holitna R.	2 lodges, 6 guides	Includes hunting
Hoholitna R.	1 lodge	
Aniak R.	6 guides, misc. outfitters	
<u>Arctic:</u>		
Kobuk R.	a few small lodges (20 guests) ^a	
Walker Lk.	1 lodge	
Alatna R.	1 lodge	

^a NPS encourages nonconsumptive use of Kobuk Valley Park, so these lodges may offer more for the sightseer than the angler.

Append

Loca

Sex

Wh

Ni

U