

Region SOUTHCENTRAL

USGS Quad Cordova C-7

DEC 18 1996

Anadromous Water Catalog Number of Waterway 221-40-10685-2006

REGION III
 HABITAT AND RESTORATION
 DIVISION
 Local Name

Name of Waterway _____

USGS Name

Addition Deletion Correction Backup Information

For Office Use

Nomination #	<u>97 028</u>	<u>[Signature]</u>	<u>12/8/91</u>
Revision Year:	_____	Regional Supervisor	Date
Revision to:	Atlas _____ Catalog _____	<u>[Signature]</u>	<u>3/18/97</u>
	Both <u>X</u>	AWC Project Biologist	Date
Revision Code:	<u>A-2</u>	<u>[Signature]</u>	<u>12/12/97</u>
		Drafted	Date

OBSERVATION INFORMATION

Species	Date(s) Observed	Spawning	Rearing	Present	Anadromous
Pink	5/21/96	Yes		Yes	<input checked="" type="checkbox"/>
Coho	5/21/96	Probable	Probable	Probable	<input type="checkbox"/>
					<input type="checkbox"/>
					<input type="checkbox"/>
					<input type="checkbox"/>


IMPORTANT: Provide all supporting documentation that this water body is important for the spawning, rearing or migration of anadromous fish, including: number of fish and life stages observed; sampling methods, sampling duration and area sampled; copies of field notes; etc. Attach a copy of a map showing location of mouth and observed upper extent of each species, as well as other information such as: specific stream reaches observed as spawning or rearing habitat; locations, types, and heights of any barriers; etc.

Comments: Site sampled during ADF&G, Habitat and Restoration Division's, participation in a Forest Practices inspection of Tatitlek Corporation's timber harvest operation. The current Anadromous Waters Atlas depicts one anadromous fish stream flowing into Snug Corner Cove. During our inspection, we discovered a tributary (Figure 1 and 2) to the currently documented stream (the newly discovered stream is actually the larger of the two). We observed several adult pink salmon carcasses along this stream from its mouth to the approximate location marked on the attached map. There is no obvious physical barrier where I designated the upper limit of pink salmon distribution. The gradient becomes gradually steeper, the substrate coarser, and no additional carcasses were observed. Although we did not sample for them, the lower reach of the stream provides excellent coho salmon spawning and rearing habitat.

Name of Observer (please print) Michael Wiedmer, Habitat Biologist
 Date: 12/18/96 Signature: [Signature]
 Address: Alaska Department of Fish and Game
333 Raspberry Road, Anchorage, AK 99518

This certifies that in my best professional judgment and belief the above information is evidence that this waterbody should be included in or deleted from the Catalog of Waters Important for Spawning, Rearing or Migration of Anadromous Fishes per AS 16.05.870.

Signature of Area Biologist: _____



PREVIOUSLY
CATALOGUED
STREAM

NEW
STREAM

ADD STREAM
2008-10-10-10685-
w/ P5

CORDOVA
C-8

CORDOVA
C-7

SOUND

KNOWLES
BAY

Red Head

Knowles Head

Porcupine

Goose Island

Gornet Cove

Shag Corner

Shag Corner