

State of Alaska  
 Department of Fish and Game  
 Nomination for Waters  
 Important to Anadromous Fish

ALASKA DEPT. OF  
 FISH & GAME

AWC Volume SE SC SW W AR IN USGS Quad Unalaska Feb 10 1997

Anadromous Water Catalog Number of Waterway 302-21-10850 REGION II  
 Name of Waterway Camp Creek (aka Laundry Creek) USGS name \_\_\_\_\_ Local name XX DIVISION  
 Addition XX Deletion \_\_\_\_\_ Correction \_\_\_\_\_ Backup Information \_\_\_\_\_

For Office Use

Nomination # <u>97 250</u>	Regional Supervisor _____	Date _____
Revision Year: _____	_____	_____
Revision to: Atlas _____ Catalog _____	_____	_____
Both _____	_____	_____
Revision Code: <u>E-9</u>	Drafted	Date _____

OBSERVATION INFORMATION

Species	Date(s) Observed	Spawning	Rearing	Migration	Anadromous
Dolly Varden	6/27/96		XXX		
Dolly Varden	6/28/96		XXX		

**IMPORTANT:** Provide all supporting documentation that this water body is important for the spawning, rearing or migration of anadromous fish, including: number of fish and life stages observed; sampling methods, sampling duration and area sampled; copies of field notes; etc. Attach a copy of a map showing location of mouth and observed upper extent of each species, as well as any other information such as: specific stream reaches observed as spawning or rearing habitat; locations, types, and heights of any barriers; etc.

Comments: Sample Site 1 - 6/27/96 - Hand sampled pool on south side, side channel, beneath undercut bank (downstream of a mainstem diversion weir). Captured 4 DV measuring 270, 120, 90, 85 mm.

Sample Site 2 - 2/27-28/96 - Minnow trap baited with cured salmon eggs set on north shore mainstem, fished overnight from 1930 hr. to 1030 hr. (15 hr. total); captured no fish. May have set trap in too high a velocity zone. Sample site 2 was located about 500 meters downstream of sample site 1.

Sample Site 3 - 6/28/96 - Beach seine sample consisting of 2 hauls, each covering about 75 feet of stream reach in a small, north side, side channel of the stream. Captured 4 DV measuring 165, 145, 140, 65 mm.

Several reaches of this stream have been impacted by World War II military activities including rechannelization, diversion, and road crossings. From on site ground surveys the more productive fish habitat appears to be found from the stream's mouth upstream about 2,800 meters. Marginal rearing habitat is found further upstream (including that reach at sample site 1) to a point about 4,000 meters upstream of the mouth.

[note: length measurements, above, are tip of snout to fork of tail]

Name of Observer (please print) C. Wayne Dolezal

Date: 2/12/97 Signature: *C. Wayne Dolezal*

Address: ADF&G - Habitat and Restoration Division  
333 Raspberry Road  
Anchorage, Alaska 99518-1599

This certifies that in my best professional judgement and belief the above information is evidence that this waterbody should be included in or deleted from the Catalog of Waters Important for Spawning, Rearing or Migration of Anadromous Fishes per AS 16.05.870.

Signature of Area Biologist: \_\_\_\_\_ Rev. 7/93

