Disappearance Creek
102-40-43
Westlund - Joyner

102-40-10430

13-14,000 chum in stream.
Many, many carcasses.
Spawning area - 65+%

Map: Craig A-1
Lat: 55° 07' 52"
Long. 132° 10' 51"



STREAM IN .ORY FORM

Name: Disappearance Creek
Number: 102-40-43

Observers:
Date: 10-5-73
Temperature:
Weather:

Station No.	1	2	3															
Pool Size/Type	3/2	1/2	1/2															
Riffle Type Width - Depth	40' 6"	65' 1"	18'															
Pool Riffle Frequency	3	2	2															
Bottom Type (Riffle)	4	4	4															
Color/Turbidity	1/1	1/1	1/1															
Velocity	2	1	2															
Flow (C.F.S.)	64.31	73.70	26.16															
Temp. (°F.)	44°	44°	44°															
pH	5	5	5															
Fry Abundance	0	0	0															
Benthos Sample No.	1	0	0[1]															
Higher Plant Class	1	1	1															
Aquatic Veg. Density	3	3	3															
Ident. (1) Mosses (2) Algae	2	2	2															

General Remarks (rehab., land use, barriers, log jams, etc.):

PEAK ESCAPEMENT RECORD

NAME: Disappearance Creek

STREAM CATALOG NUMBER: 102-40-43

DATE	PINK	CHUM	OTHER SPECIES	REMARKS
1961-Weir Count	1,365	20,285		Total accumulative Weir Count
1962-Weir Count	143	23,799		Total accumulative weir count
Sept. 18, 1963		9,000 (Weir)		2-3,000 chum at the mouth (F)
Sept. 24, 1964				10,000 chums schooled at the mouth (A)
Sept.16-Oct.12, 1965	488	25,590		Total accumulative Weir Count
Sept.14-Oct.5, 1966	295	25,005		Total accumulative Weir Count
Sept.13-Oct.5, 1967	68	24,947		Total accumulative Weir Count
Sept. 18, 1968		7,000		12-15,000 fish at mouth and 600 intertidally. Vis. excellent in stream and mouth. Few scattered jumps in South Arm. (A)
Aug.26-Oct. 10, 1969	1,169	12,411		Total accumulative Weir Count
Sept.9-Sept.30 1970	86	8,373		Total accumulative Weir Count
Sept.15-Oct.14, 1971	445	2,186		Total Accumulative Weir Count
Sept.12-Oct.17, 1972	1,829	63,752		Total Accumulative Weir Count
Sept. 6-Oct.18, 1973	3,102	55,306		Total Accumulative Weir Count
Sept.10-Oct.8, 1974	374	43,024		Total Accumulative Weir Count