


103-70-10290

103-70-29 6/10/77
MOUTH



103-70-29 6/10/77
MOUTH




103-70-29
6-10-77
Freitag

Ponds in forest off stream.

Too slow for good spawning area

Banks show erosion which exposed clam beds from a period when this area was below sealevel.


Bottom has brown algae cover.



- 300m
(984')

- 200m
(656')

- 100m
(328')



Name: _____
Latitude: 55° 33' ~~18"~~ 28"
Longitude: 133° 25' ~~22"~~ 10"
Geodetic Map No: Craig C5 - 9
Location: North shore of San Fernando
Island, east shore of Garcia Cove,
0.3 miles north of head.

Catalog No: 103-70-029
Former Stream No: _____

Work Area: Ketchikan
Watershed Length: 0.6 miles
Drainage Area: 1 sq. mile
Water Supply Type: runoff from low forested
hills and scattered muskegs.

Trails & Survey Routes: Easy walking up stream bed.

Aerial Survey Notes: Too small for aerial survey.

Anchorage: Off shore in Garcia Cove.

Tide Stage when Surveyed: Low

RESOURCES

COMMERCIAL FISHERIES (species, escapement, timing, spawning area):
Pinks and chums; escapement of 600 on 8/24/76.
Many good rearing areas for coho.

SPAWNING AREA: 100 meters²
Schooling Areas: Many pools up entire length of stream.

Spawning Areas: Most areas are not excellent spawning but ASA is
estimated at about 100m² for first 600 m of stream.

SHELLFISH POTENTIAL: None noted.

SPORT FISHERIES: None noted.

LAND USE (history, present, proposed): None known; presently in a natural state.

REHABILITATION POTENTIAL: None needed.

SOILS: several areas with overhanging banks and areas of soft sediments eroded by the
stream; generally banks are low and stable.

GAME RESOURCES (species, use, habitat): None noted; reportedly good furbearer
habitat.