

ALASKA DEPARTMENT OF FISH AND GAME
DIVISION OF COMMERCIAL FISHERIES
NEWS RELEASE



Sam Cotten, Commissioner
Forrest R. Bowers, Acting Director



Contacts:

Greg Buck, Bristol Bay Area Research Biologist
Jordan Head, Asst. Area Research Biologist
Phone: (907) 267-2355
Fax: (907) 267-2442

Anchorage Office
333 Raspberry Road
Anchorage, AK 99518
Date Issued: 11/9/2018

2019 BRISTOL BAY SOCKEYE SALMON FORECAST

FORECAST AREA: Bristol Bay

SPECIES: Sockeye Salmon

FORECAST OF THE 2019 RUN:

	Forecast	Forecast Range
TOTAL PRODUCTION:	(millions)	(millions)
Total Run	40.18	27.90 - 52.46
Escapement	12.58	
Commercial Common Property Harvest	27.60	
Bristol Bay Harvest	26.11	
South Peninsula Harvest	1.49	
Inshore Run	38.70	

METHODS

The 2019 Bristol Bay sockeye salmon forecast is the sum of individual predictions of nine river systems (Kvichak, Alagnak, Naknek, Egegik, Ugashik, Wood, Igushik, Nushagak, and Togiak rivers) and four age classes (ages 1.2, 1.3, 2.2, and 2.3, plus ages 0.3 and 1.4 for the Nushagak River). Adult escapement and return data from brood years 1972–2014 were used in the analyses.

Forecasts for each age class returning to a river system were derived from models based on the relationship between adult returns of that age class and either total returns or sibling returns from the same brood years. Models based on the most recent three and five years of returns were also evaluated. In general, models with statistically significant parameters and/or the best past performance (accuracy and precision) were chosen. Performance was evaluated using mean absolute deviation, mean absolute percent error, mean arctangent absolute percent error, and mean

percent error between forecasted and observed returns. These performance metrics were calculated and considered for each model across the most recent 3, 5, and 9-year time frames. In certain cases, competing models were averaged in a hybrid model approach.

The forecast range is the upper and lower values of the 80% confidence interval for the total run forecast. The confidence bounds were calculated from the deviation of actual runs and run forecasts from 2001 through 2018.

RESULTS

A total of 40.18 million sockeye salmon (range 27.90–52.46 million) are expected to return to Bristol Bay in 2019. This is 10% smaller than the most recent 10-year average of Bristol Bay total runs (44.4 million) while being 16% greater than the long-term (1963–2018) average of 34.2 million fish. All systems are expected to meet their spawning escapement goals.

Where practical, the department will manage escapements proportional to the run size and relative to the historical record (5AAC 06.355(d)(1)). Escapement is projected as the 75th quartile of the escapement range if the forecast is above the historical trend line (Egegik, Nushagak, and Togiak Rivers), as the midpoint (50th quartile) of the escapement range if the forecast is in line with the historical trend (Ugashik and Igushik Rivers), and as the 25th quartile of the escapement goal range if the forecast is below the historical trend line (Kvichak and Wood Rivers in 2019; Table 1). Because it is passively managed, the Alagnak River exploitation rate is assumed to be the same as the Kvichak River exploitation rate and therefore the escapement is projected to be the total run forecast minus expected harvest. Preseason harvest projections are provided to aid industry in planning. Once the run begins to develop, the department relies on catch and escapement data for management decisions.

A run of 40.18 million sockeye salmon would allow for a potential total harvest of 27.60 million fish—26.11 million fish in Bristol Bay and 1.49 million fish in the South Peninsula fisheries. A Bristol Bay harvest of this size is 8% smaller than the most recent 10-year harvest of 30.0 million which has ranged from 15.4 million to 41.9 million, and 23% greater than the long-term average harvest of 21.2 million fish (1963 to present).

The run forecast for each district and river system is as follows: 16.12 million to Naknek-Kvichak District (6.95 million to the Kvichak river, 3.97 million to the Alagnak river, and 5.21 million to the Naknek river); 9.07 million to the Egegik District; 3.46 million to the Ugashik District; 10.38 million to the Nushagak District (4.62 million to the Wood river, 4.18 million to the Nushagak river, and 1.58 million to the Igushik river); and 1.15 million to the Togiak District (Table 1).

We forecast that the 2019 run will consist of 15.16 million age-1.2 fish (38% of the total run), 5.49 million age-2.2 fish (14% of the total run), 17.05 million age-1.3 fish (42% of the total run), and 2.42 million age-2.3 fish (6% of the total run; Table 1).

DISCUSSION

Historically, sockeye salmon runs to Bristol Bay have been highly variable. The Bristol Bay total run has averaged 34.2 million from 1963 through 2018 and has averaged 44.4 million fish during the most recent 10-year period. Forecasting future salmon returns is inherently difficult and uncertain. We have used similar methods since 2001 to produce the Bristol Bay sockeye salmon forecast which have performed well when applied to Bristol Bay as a whole. Since 2001, our forecasts have, on average, under-forecast the run by 11% and have ranged from 44% below

actual run in 2014 to 19% above actual run in 2011. Forecasted harvests have had a mean absolute percent error of 14% since 2001.

Individual river forecasts have greater uncertainty compared to Bay-wide forecasts. Since 2001, on average, we have under-forecast returns to the Alagnak (-39%), Togiak (-12%), Kvichak (-22%), Wood (-17%), Nushagak (-21%), Ugashik (-0.2%), and Naknek (-12%) Rivers, and over-forecast returns to Igushik (13%) and Egegik Rivers (18%). Over-forecasting returns to some rivers while under-forecasting returns to other rivers means that the overall Bristol Bay forecast is often more accurate than the forecast to any individual river. The Nushagak District had another record return in 2018. These record returns have been driven by robust returns from the 2013 and 2014 brood years. Evidence regarding the strength of the 2015 brood year in the Nushagak District is mixed with a very high level of jacks (age-1.1 fish) in the Nushagak River and a very low level of jacks in the Wood River 2018 return.

The department would like to thank the Bristol Bay Fisheries Collaborative (BBFC) for funding assistance in 2018. The BBFC began in 2016 and is an agreement between ADF&G and the Bristol Bay Science and Research Institute (BBSRI) to work together with stakeholders to restore a world-class fishery management system and raise funds to support and maintain management. This agreement is supported by ADF&G, BBSRI, drift and set net fishermen, processors, municipalities, villages, support industries and other stakeholders. A list of organizations that committed financial support to the BBFC in 2018, as well as additional information about this agreement can be found at <https://www.bbsri.org/bbfc>.

Greg Buck and Jordan Head
Bristol Bay Research Staff

Table 1.—Forecast of total run, escapement, and harvest of major age classes of sockeye salmon returning to Bristol Bay river systems in 2019.

DISTRICT River	Millions of Sockeye Salmon								
	Forecasted Production by Age Class					Forecasted		South Peninsula ^a	BB Inshore
	1.2	2.2	1.3	2.3	Total	Escapement	Harvest		
NAKNEK-KVICHAK									
Kvichak	2.95	1.08	2.87	0.05	6.95	4.00	2.69	0.26	6.69
Alagnak	1.88	0.19	1.88	0.02	3.97	2.28	1.54	0.15	3.82
Naknek	2.18	0.58	2.00	0.45	5.21	1.40	3.61	0.19	5.01
Total	7.01	1.84	6.74	0.53	16.12	7.68	7.84	0.60	15.53
EGEGIK	2.51	3.04	1.81	1.72	9.07	1.70	7.04	0.34	8.74
UGASHIK	1.31	0.33	1.72	0.10	3.46	0.95	2.38	0.13	3.33
NUSHAGAK									
Wood	2.41	0.23	1.94	0.04	4.62 ^b	0.98	3.47	0.17	4.45
Igushik	0.62	0.01	0.94	0.01	1.58	0.28	1.25	0.06	1.52
Nushagak	1.12	0.02	2.95	0.02	4.18	0.77	3.26	0.15	4.02
Total	4.14	0.26	5.83	0.07	10.38	2.02	7.97	0.38	9.99
TOGIAK	0.18	0.01	0.95	0.01	1.15	0.23	0.87 ^c	0.04	1.10
BRISTOL BAY									
	15.16	5.49	17.05	2.42	40.18	12.58	26.11	1.49	38.70
	38%	14%	42%	6%	100%				

Note: This table is a summary. Slight differences may appear due to rounding.

^a Projected harvest is based on the current 5 year running average exploitation rate of 3.7%.

^b Nushagak River forecast total includes age-0.3 and age-1.4 fish.

^c Forecasts for Kulukak, Kanik, Osviak, and Matogak river systems are not included. These systems contribute approximately 50,000 sockeye salmon to Togiak District harvest each year.