

**Technical Report No. 19-04**

---

---

**Baseline Aquatic Biomonitoring for the Anarraaq and Aktigiruaq Prospects near the Red Dog Mine, 2014 – 2017, with comparisons to 2000 – 2002.**

by

**Alvin G. Ott, Audra L. J. Brase and Parker T. Bradley**



---

---

February 2019

Alaska Department of Fish and Game

Division of Habitat



## Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Habitat, Sport Fish, and Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, Technical Reports and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

<b>Weights and measures (metric)</b>		<b>General</b>		<b>Mathematics, statistics</b>	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H <sub>A</sub>
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	<i>e</i>
hectare	ha	at	@	catch per unit effort	CPUE
kilogram	kg	compass directions:		coefficient of variation	CV
kilometer	km	east	E	common test statistics	(F, t, $\chi^2$ , etc.)
liter	L	north	N	confidence interval	CI
meter	m	south	S	correlation coefficient	
milliliter	mL	west	W	(multiple)	R
millimeter	mm	copyright	©	correlation coefficient (simple)	r
		corporate suffixes:		covariance	cov
<b>Weights and measures (English)</b>		Company	Co.	degree (angular)	°
cubic feet per second	ft <sup>3</sup> /s	Corporation	Corp.	degrees of freedom	df
foot	ft	Incorporated	Inc.	expected value	<i>E</i>
gallon	gal	Limited	Ltd.	greater than	>
inch	in	District of Columbia	D.C.	greater than or equal to	≥
mile	mi	et alii (and others)	et al.	harvest per unit effort	HPUE
nautical mile	nmi	et cetera (and so forth)	etc.	less than	<
ounce	oz	exempli gratia	e.g.	less than or equal to	≤
pound	lb	(for example)		logarithm (natural)	ln
quart	qt	Federal Information Code	FIC	logarithm (base 10)	log
yard	yd	id est (that is)	i.e.	logarithm (specify base)	log <sub>2</sub> , etc.
		latitude or longitude	lat or long	minute (angular)	'
<b>Time and temperature</b>		monetary symbols (U.S.)	\$, ¢	not significant	NS
day	d	months (tables and figures): first three letters	Jan, ..., Dec	null hypothesis	H <sub>0</sub>
degrees Celsius	°C	registered trademark	®	percent	%
degrees Fahrenheit	°F	trademark	™	probability	P
degrees kelvin	K	United States (adjective)	U.S.	probability of a type I error (rejection of the null hypothesis when true)	$\alpha$
hour	h	United States of America (noun)	USA	probability of a type II error (acceptance of the null hypothesis when false)	$\beta$
minute	min	U.S.C.	United States Code	second (angular)	"
second	s	U.S. state	use two-letter abbreviations (e.g., AK, WA)	standard deviation	SD
<b>Physics and chemistry</b>				standard error	SE
all atomic symbols				variance	
alternating current	AC			population sample	Var
ampere	A			sample	var
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

***TECHNICAL REPORT NO. 19-04***

**BASELINE AQUATIC BIOMONITORING FOR THE ANARRAAQ AND  
AKTIGIRUQ PROSPECTS NEAR THE RED DOG MINE, 2014 - 2017,  
WITH COMPARISONS TO 2000 - 2002**

By

Alvin G. Ott, Audra L. J. Brase and Parker T. Bradley  
Division of Habitat, Fairbanks

Alaska Department of Fish and Game  
Division of Habitat  
1300 College Rd, Fairbanks, Alaska, 99701

February 2019

Cover: East Fork Ikalukrok Creek, August 2018, Photograph by Audra Brase

Technical Reports are available through the Alaska State Library, Alaska Resources Library and Information Services (ARLIS) and on the Internet: [http://www.adfg.alaska.gov/index.cfm?adfg=habitat\\_publications.main](http://www.adfg.alaska.gov/index.cfm?adfg=habitat_publications.main). This publication has undergone editorial and peer review.

*Note:* Product names used in the publication are included for completeness but do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

*Alvin G. Ott, Audra L.J. Brase and Parker T. Bradley  
Alaska Department of Fish and Game, Division of Habitat  
1300 College Rd., Fairbanks, AK 99701-1599, USA*

*This document should be cited as:*

*Ott, A.G., A.L.J. Brase and P.T. Bradley. 2019. Baseline Aquatic Biomonitoring for the Anarraaq and Aktigiruaq Prospects near the Red Dog Mine, 2014 -2017, with Comparisons to 2000 - 2002. Alaska Department of Fish and Game, Technical Report No. 19-04, Fairbanks, Alaska.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375

# Table of Contents

List of Figures .....	iii
Acknowledgements .....	vi
Executive Summary .....	vii
Introduction .....	1
Methods .....	2
Results and Discussion .....	4
Upper Competition Creek (Station 203) .....	4
Water Quality .....	4
Periphyton .....	5
Invertebrates .....	5
Fish .....	7
Sourdock Creek (Station 204) .....	7
Water Quality .....	7
Periphyton .....	7
Invertebrates .....	8
Fish .....	9
Lower Competition Creek (Station 202) .....	10
Water Quality .....	10
Periphyton .....	11
Invertebrates .....	12
Fish .....	13
West Fork Ikalukrok Creek (Station 205) .....	16
Water Quality .....	16
Periphyton .....	17
Invertebrates .....	17
Fish .....	19
Upper Ikalukrok Creek (Station 206) .....	19
Water Quality .....	19
Periphyton .....	19
Invertebrates .....	20
Fish .....	21
Ikalukrok Creek (Station 207) .....	22
Water Quality .....	22
Periphyton .....	22
Invertebrates .....	23
Fish .....	24
East Fork Ikalukrok Creek (Station 208) .....	24
Water Quality .....	24
Periphyton .....	25
Invertebrates .....	26
Fish .....	27
Grayling Junior Creek (Station 209) .....	28
Water Quality .....	28
Periphyton .....	28

Invertebrates .....	29
Fish.....	30
Noa Creek (Station 210) .....	34
Water Quality .....	34
Periphyton .....	35
Invertebrates.....	35
Fish.....	37
Moil Creek (Station 211) .....	37
Water Quality .....	37
Periphyton .....	38
Invertebrates.....	38
Fish.....	39
Sled Creek (Station 212).....	40
Water Quality .....	40
Periphyton .....	40
Invertebrates.....	41
Fish.....	42
Volcano Creek .....	42
Water Quality .....	42
Periphyton .....	43
Invertebrates.....	43
Fish.....	45
Literature Cited .....	49
Appendix 1. Total numbers and CPUE of Dolly Varden juveniles captured in minnow traps at various streams near the Red Dog Mine site.....	50
Appendix 1 continued.....	51
Appendix 1 continued.....	52

## List of Figures

Figure 1. Map showing sampling points and general location of Anarraaq Prospect. ....	1
Figure 2. Upper Competition Creek (left photo 7/31/15 and right photo 7/13/17).....	5
Figure 3. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Upper Competition Creek. ....	5
Figure 4. Average aquatic invertebrate density $\pm$ 1 SD, in Upper Competition Creek. ....	6
Figure 5. Percent Chironomidae and EPT, in Upper Competition Creek.....	6
Figure 6. Aquatic invertebrate taxa richness at Upper Competition Creek. ....	6
Figure 7. Sourdock Creek (left photo 7/11/2015 and right photo 7/3/2016). ....	7
Figure 8. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Sourdock Creek.....	8
Figure 9. Average aquatic invertebrate densities $\pm$ 1SD, in Sourdock Creek.....	8
Figure 10. Percent Chironomidae and EPT, in Sourdock Creek. ....	9
Figure 11. Aquatic invertebrate taxa richness, in Sourdock Creek.....	9
Figure 12. Catch per unit of effort for juvenile Dolly Varden in Sourdock Creek.....	10
Figure 13. Length frequency distribution of Dolly Varden in Sourdock Creek. ....	10
Figure 14. Lower Competition Creek (left photo 2000/2002 and right photo 7/30/2014). ....	11
Figure 15. Average concentration of chlorophyll-a ( $\pm$ 1SD) in lower Competition Creek. ....	11
Figure 16. Average aquatic invertebrate densities $\pm$ 1SD in lower Competition Creek.....	12
Figure 17. Percent Chironomidae and EPT in lower Competition Creek.....	12
Figure 18. Aquatic invertebrate taxa richness in lower Competition Creek.....	13
Figure 19. Catch per unit of effort for juvenile Dolly Varden in lower Competition Creek. ....	13
Figure 20. Length frequency distribution of Dolly Varden in Lower Competition Creek. ....	14
Figure 21. Average cadmium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17. ....	14
Figure 22. Average lead concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17. ....	15
Figure 23. Average selenium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17. ....	15
Figure 24. Average zinc concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17. ....	16
Figure 25. Average mercury concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17. ....	16
Figure 26. West Fork Ikalukrok Creek (7/12/2015), note white staining in right photo at confluence of West Fork and Ikalukrok.....	17
Figure 27. Average concentration of chlorophyll-a ( $\pm$ 1SD) in West Fork Ikalukrok Creek. ....	17
Figure 28. Average aquatic invertebrate densities $\pm$ 1SD in West Ikalukrok Creek. ....	18
Figure 29. Percent Chironomidae and EPT in West Ikalukrok Creek. ....	18
Figure 30. Aquatic invertebrate taxa richness in West Ikalukrok Creek. ....	18
Figure 31. Ikalukrok Creek immediately upstream of West Fork Ikalukrok Creek (left photo 7/3/2017) and Cub Creek (right photo 7/30/2014). ....	19
Figure 32. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Upper Ikalukrok Creek.....	20
Figure 33. Average aquatic invertebrate densities $\pm$ 1SD in Upper Ikalukrok Creek. ....	20
Figure 34. Percent Chironomidae and EPT in Upper Ikalukrok Creek. ....	21
Figure 35. Aquatic invertebrate taxa richness in Upper Ikalukrok Creek. ....	21
Figure 36. Ikalukrok Creek below Cub Creek on July 3, 2016 (left photo is the sample site and right photo is a rock scraped for a periphyton sample).....	22
Figure 37. Average aquatic invertebrate densities $\pm$ 1SD in Ikalukrok Creek. ....	23

Figure 38. Percent Chironomidae and EPT in Ikalukrok Creek. ....	23
Figure 39. Aquatic invertebrate taxa richness in Ikalukrok Creek. ....	24
Figure 40. Drift nets in East Fork Ikalukrok Creek. ....	25
Figure 41. Average concentration of chlorophyll-a ( $\pm$ 1SD) in East Fork Ikalukrok Creek. ....	25
Figure 42. Average aquatic invertebrate densities $\pm$ 1SD in East Fork Ikalukrok Creek. ....	26
Figure 43. Percent Chironomidae and EPT in East Fork Ikalukrok Creek. ....	26
Figure 44. Aquatic invertebrate taxa richness in East Fork Ikalukrok Creek. ....	27
Figure 45. Catch per unit of effort for juvenile Dolly Varden in East Fork Ikalukrok Creek. ....	27
Figure 46. Grayling Junior Creek (left photo drift nets on 7/3/2016 and right photo drift nets on 7/4/2017). ....	28
Figure 47. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Grayling Junior Creek. ....	29
Figure 48. Average aquatic invertebrate densities $\pm$ 1SD in Grayling Junior Creek. ....	29
Figure 49. Percent Chironomidae and EPT in Grayling Junior Creek. ....	30
Figure 50. Aquatic invertebrate taxa richness in Grayling Junior Creek. ....	30
Figure 51. Arctic grayling at the confluence of Grayling Junior Creek and Ikalukrok Creek. ...	31
Figure 52. Catch per unit of effort for juvenile Dolly Varden in Grayling Junior Creek. ....	32
Figure 53. Length frequency distribution of Dolly Varden in Grayling Junior Creek. ....	32
Figure 54. Average cadmium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks. ....	33
Figure 55. Average lead concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks. ....	33
Figure 56. Average selenium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks. ....	34
Figure 57. Average zinc concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks. ....	34
Figure 58. Average mercury concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks, note that samples were not tested for mercury in 2001. ....	34
Figure 59. Noa Creek (left photo 7/4/2016 and right photo 7/4/2017). ....	35
Figure 60. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Noa Creek. ....	35
Figure 61. Average aquatic invertebrate densities $\pm$ 1SD in Noa Creek. ....	36
Figure 62. Percent Chironomidae and EPT in Noa Creek. ....	36
Figure 63. Aquatic invertebrate taxa richness in Noa Creek. ....	37
Figure 64. Moil Creek (left photo 7/4/2016 is the confluence of Moil with Ikalukrok and right photo 7/4/2017). ....	37
Figure 65. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Moil Creek. ....	38
Figure 66. Average aquatic invertebrate densities $\pm$ 1SD in Moil Creek. ....	38
Figure 67. Percent Chironomidae and EPT in Moil Creek. ....	39
Figure 68. Aquatic invertebrate taxa richness in Moil Creek. ....	39
Figure 69. Sled Creek (left photo 7/5/2016 and right photo 7/5/2017). ....	40
Figure 70. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Sled Creek. ....	40
Figure 71. Average aquatic invertebrate densities $\pm$ 1SD in Sled Creek. ....	41
Figure 72. Percent Chironomidae and EPT in Sled Creek. ....	41
Figure 73. Aquatic invertebrate taxa richness in Sled Creek. ....	42
Figure 74. Volcano Creek (left photo minnow trap 7/29/2014 and right photo drift nets 7/29/2014). ....	43

Figure 75. Average concentration of chlorophyll-a ( $\pm$ 1SD) in Volcano Creek.....	43
Figure 76. Average aquatic invertebrate densities $\pm$ 1SD in Volcano Creek. ....	44
Figure 77. Percent Chironomidae and EPT in Volcano Creek. ....	44
Figure 78. Aquatic invertebrate taxa richness in Volcano Creek. ....	44
Figure 79. Catch per unit of effort for juvenile Dolly Varden in Volcano Creek. ....	45
Figure 80. Length frequency distribution of Dolly Varden in Volcano Creek.....	45
Figure 81. Average cadmium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks. ....	46
Figure 82. Average lead concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks. ....	46
Figure 83. Average selenium concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks. ....	47
Figure 84. Average zinc concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks. ....	47
Figure 85. Average mercury concentrations ( $\pm$ 1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks. ....	48

## **Acknowledgements**

We thank Teck Alaska Inc. for their financial and logistical support for aquatic biomonitoring in streams associated with the Anarraaq and Aktigiruaq Prospects. We specifically acknowledge the assistance provided by Teck Alaska Inc. employees: Wayne Hall, Robert Napier, Chris Eckert, Darren Jones, Joseph Diehl III, Dennis Sheldon, and former employees Jeff Clark and Karen Conitz.

Alaska Department of Fish and Game (ADF&G) Division of Habitat employees Bill Morris, Brad Wendling, Todd Nichols, Laura Jacobs, and Maria Wessel participated in field sampling and laboratory work.

Laura Jacobs and Nora Foster (NRF Taxonomic Services) were responsible for sorting and identification of aquatic invertebrates.

Robert Napier and Chelsea Clawson provided constructive reviews of this report.

## Executive Summary

This report summarizes the results of biomonitoring work performed in streams in the general vicinity of the Anarraaq and Aktigiruiq prospects located northwest of the Red Dog Mine. Biomonitoring included surveys of periphyton (measured by chlorophyll-a), aquatic invertebrates, and fish; these data were collected annually from 2014 to 2017. Biomonitoring data from 2000 to 2002 and reported in Weber Scannell and Ott (2006) are included here for comparisons. The purpose of this report is to document the existing aquatic environment and to provide a basis for monitoring as exploration continues and/or development of the ore bodies occurs.

Water quality in streams near the Anarraaq and Aktigiruiq prospects varies considerably. Creeks draining from the area where the orebody is located (West Fork Ikalukrok, Noa, Moil, Ikalukrok, and Competition creeks) exhibit naturally degraded water quality (e.g., high metals, low pH), low periphyton standing crop, low aquatic invertebrate density, and very few fish. No fish were found in Noa, Moil, West Fork Ikalukrok, and Ikalukrok creeks (upstream of its confluence with East Fork Ikalukrok Creek).

Water quality in Sourdock, upper Ikalukrok, East Fork Ikalukrok, Grayling Junior, and Sled creeks is of a higher quality (e.g., lower metals, neutral pH). Juvenile Dolly Varden and Arctic grayling were present in Competition Creek and juvenile Dolly Varden were present in Sourdock Creek. Arctic grayling and Dolly Varden were found in East Fork Ikalukrok Creek and in Ikalukrok Creek downstream of its confluence with the East Fork. Grayling Junior Creek contains both Arctic grayling and juvenile Dolly Varden, and in some years large numbers (about 300) of Arctic grayling may be found at the confluence of Grayling Junior Creek and Ikalukrok Creek. Sourdock Creek, a tributary to Competition Creek, supports rearing Dolly Varden. Sled Creek does not support fish and the absence of fish in Sled Creek is likely due to the fact that the stream goes subsurface during the ice-free season before entering Ikalukrok Creek.

Measurements of periphyton standing crop, aquatic invertebrates, and fish distribution also vary among the sample sites. Periphyton as measured by chlorophyll-a was highest in Sourdock and East Fork Ikalukrok creeks ( $\geq 4 \text{ mg/m}^2$ ), while aquatic invertebrate densities were highest in Lower Competition, Ikalukrok, East Fork Ikalukrok, and Sled creeks ( $\geq 10 \text{ invertebrates/m}^3$ ). The highest fish use by multiple species (Arctic grayling, Dolly Varden, and slimy sculpin) was documented in East Fork Ikalukrok and Grayling Junior creeks.

## Introduction

Teck Alaska Inc. has been conducting geotechnical drilling in the area of the Anarraaq Prospect since the mid-1990's and recently announced that there is a second prospect (Aktigiruaq) in the same general area. Both prospects are zinc (Zn) and lead (Pb) subsurface deposits located in Section 23, T32N, R19W (De Long Mountains A-2). The deposits are located about 16 km northwest of the Red Dog Mine (Figure 1).




Figure 1. Map showing sampling points and general location of Anarraaq Prospect.

Aquatic baseline data collection near the Anarraaq Prospect began in 2000 and continued through 2002 (Table 1). Alaska Department of Natural Resources (ADNR) technical reports summarize water quality, periphyton, aquatic invertebrate, and fish data collected in 2000, 2001, and 2002 (Weber Scannell and Ott, 2006).

From 2014 to 2017, sampling work focused on streams which flow to the west and east from the Anarraaq/Aktigiruaq ore bodies (Table 1). Volcano Creek, a tributary to Ikalukrok Creek, is a potential site for a new tailings dam and an aquatic biomonitoring station was established in the creek in 2014. Periphyton (chlorophyll-a concentrations), aquatic invertebrates (taxonomic richness and abundance), and fish (presence and use) data were collected at each site. Periphyton and aquatic invertebrates were collected in early July while fish sampling was done in late-July to early August.

The Anarraaq and Aktigiruaq Prospects may ultimately be developed as an underground mine located about 600 m below the ground surface. Details on mine development, operations, and closure are not available at this time, but such would be required prior to mine development, if initiated. Exploration access to the orebodies may be via an all-weather (exploration) road following Mainstem Red Dog Creek, crossing North Fork Red Dog, Grayling Junior, and Ikalukrok creeks, and then up Ikalukrok Creek to access the mine area. North Fork Red Dog, Grayling Junior, and Ikalukrok creeks are all anadromous waterbodies which support Arctic grayling, Dolly Varden, and slimy sculpin.

## Methods

Details of the methods used for the aquatic biomonitoring study are described in ADF&G Technical Report 17-09 *Methods for Aquatic Life Monitoring at the Red Dog Mine Site* (Bradley 2017). Location of the sample sites described in this report and the years they were sampled are shown in Table 1.

Periphyton was sampled directly from cobble on the streambed. The periphyton was collected from a specific area of cobble, following the rapid bioassessment techniques of Barbour et al. 1997, but with more replicates per site to increase sample precision. The concentrations of chlorophyll-a were determined to estimate periphyton standing crop.

Aquatic invertebrates were collected at each sample site using five drift nets installed in riffle habitat. The drift nets were set for one hour and the water depth and average water velocity through each net were measured. After one hour, the drift nets were pulled, materials (debris and

invertebrates) in the net were flushed to the cod end, transferred to a labeled sample container, preserved, and transported back to Fairbanks where they were sorted and identified.

Fish sampling typically consisted of setting ten minnow traps baited with salmon roe at each sample site for about 24 hours. Visual observations were made when appropriate. The minnow traps were pulled, the fish identified and measured (fork length, mm), and released. In some cases, juvenile Dolly Varden (between 90 and 140 mm long) were retained for whole body analyses of selected elements.

**Table 1. Location of Sample Sites and Years Sampled.**

Station No.	Stream/Site Name	Years Sampled
202	Lower Competition	2000-2002 and 2014-2017
203	Upper Competition	2000-2002 and 2014-2017
204	Sourdock	2000-2002 and 2014-2017
205	West Fork Ikalukrok	2000-2002 and 2015-2017
206	Ikalukrok (above West Fork)	2000-2002 and 2015-2017
207	Ikalukrok <sup>1</sup>	1997-1998, 2000-2002, and 2016-2017
208	East Fork Ikalukrok	1997-1998, 2000-2002, and 2016-2017
209	Grayling Junior	2000-2002 and 2016-2017
210	Noa	2000-2002 and 2016-2017
211	Moil	2000-2002 and 2016-2017
212	Sled	2000-2002 and 2015-2017
N/A	Volcano	2014-2017

<sup>1</sup> Sample site is located downstream of the Cub Creek seep.

## **Results and Discussion**

This section presents the biomonitoring results for each creek listed in Table 1. Biomonitoring data were collected from 2014-17 and are summarized here. Comparisons are made to prior work performed in 2000-02 and published in Weber Scannell and Ott (2006). Detailed data for fish catches can be found in Appendix 1. Additional detailed data (periphyton, aquatic invertebrates and fish whole body element concentrations) is available upon request<sup>2</sup>.

Periphyton attached microalgae biomass were collected in early July of each sample year and are presented as mg/m<sup>2</sup> chlorophyll-a.

Aquatic invertebrates were also collected in early July of each sample year and the densities are expressed as the average number of invertebrates/ m<sup>3</sup>, and as a comparison of the total percent of Ephemeroptera, Plecoptera and Tricoptera (EPT) vs. Chironomidae (CHIROs). In general, the higher the percentage of EPT at a site, the higher the water quality. Taxa richness, in this report, was defined as the total number of taxa found at a particular sample site.

Fish were sampled in early August of each sample year and densities are presented in Catch Per Unit Effort (CPUE) units as the number of fish caught per 24 hour period. Numbers of fish presented in the text are rounded to the nearest whole number.

### **Upper Competition Creek (Station 203)**

#### **Water Quality**

Upper Competition Creek (Figures 1 and 2) had moderately low pH and elevated concentrations of aluminum, cadmium, nickel, and zinc (Weber Scannell and Ott 2006). The substrate had a grayish-yellow precipitate in the early 2000s, but the precipitate in 2014 to 2017 varied from white to tan (Figure 2). Additionally, the water color in Upper Competition Creek has varied from an opaque white to orange.

---


<sup>2</sup> Submit detailed data requests to ADF&G Division of Habitat - 1300 College Rd, Fairbanks, Alaska 99708.



**Figure 2. Upper Competition Creek (left photo 7/31/15 and right photo 7/13/17).**

**Periphyton**


Average chlorophyll-a concentrations were low in Upper Competition Creek and ranged from 0.01 to 0.42 mg/m<sup>2</sup> over the sample years (Figure 3).




**Figure 3. Average concentration of chlorophyll-a (± 1SD) in Upper Competition Creek.**

**Invertebrates**


The average aquatic invertebrate density in Upper Competition Creek has varied from 1.2 to 14.9 invertebrates/m<sup>3</sup> (Figure 4). The percent Chironomidae was highest in 2014 and 2017, but in 2000 and 2016 the EPT was higher than Chironomidae (Figure 5). Taxa richness varied from 12 to 21 taxa per site over the sample years (Figure 6).



**Figure 4. Average aquatic invertebrate density  $\pm 1$  SD, in Upper Competition Creek.**



**Figure 5. Percent Chironomidae and EPT, in Upper Competition Creek.**



**Figure 6. Aquatic invertebrate taxa richness at Upper Competition Creek.**

## Fish

The catch per unit of effort of Dolly Varden in minnow traps at Upper Competition Creek was five fish in 2000 and three fish in 2002. Catches of these fish coincided with the two years that had the highest periphyton concentrations (Figure 3). Catches were zero for all other sample years. No fish were caught at the site from 2014-2017, which suggests that water quality has degraded to the point that fish are avoiding this stream reach. Upper Competition Creek appears to have changed over the time frame of our sampling effort with every indication that basic biological productivity has decreased.

## Sourdock Creek (Station 204)

### Water Quality

Sourdock Creek (Figures 1 and 7) had moderate alkalinity (as  $\text{CaCO}_3$ ), sulfate concentrations (2000-2002 median = 116 mg/L), and median hardness of 170 mg/L. The pH was neutral with slightly elevated concentrations of aluminum, cadmium, and zinc (Weber Scannell and Ott 2006). The large boulders were covered with a thick layer of moss from 2000 to 2002, but most of the moss was gone during the 2014 to 2017 sample period (Figure 7).




**Figure 7. Sourdock Creek (left photo 7/11/2015 and right photo 7/3/2016).**

### Periphyton

The average chlorophyll-a concentration in Sourdock Creek ranged from 3.57  $\text{mg/m}^2$  in 2015 to 12.44  $\text{mg/m}^2$  in 2002 (Figure 8). Chlorophyll-a was considerably higher each sampling year in


Sourdock Creek than it was in Upper Competition Creek. These two creeks merge, just downstream of the sample sites, to form Competition Creek.




**Figure 8. Average concentration of chlorophyll-a ( $\pm$  1SD) in Sourdock Creek.**

### Invertebrates


The average aquatic invertebrate density in Sourdock Creek was highest in 2017 (Figure 9). Average density ranged from a low of 3.0 to a high of 15.4 invertebrates/m<sup>3</sup>. The percent Chironomidae was much higher from 2014 to 2017 than from 2000 to 2002 (Figure 10). Taxa richness varied from 12 to 19 taxa per site over the sample years (Figure 11).



**Figure 9. Average aquatic invertebrate densities  $\pm$  1SD, in Sourdock Creek.**




**Figure 10. Percent Chironomidae and EPT, in Sourdock Creek.**



**Figure 11. Aquatic invertebrate taxa richness, in Sourdock Creek.**


## Fish

The catch per unit of effort of juvenile Dolly Varden at Sourdock Creek was highest in 2000 and has decreased over the sample period to zero fish caught (Figure 12). The element concentrations (e.g., metals) in Competition Creek may have increased, leading to a chemical barrier to the upstream movement of Dolly Varden juveniles from overwintering habitat in the Wulik River. Small streams around the Red Dog Mine area are typically used by rearing Dolly Varden during the ice-free season and the lower CPUE in Sourdock Creek indicates that this movement has been curtailed.



**Figure 12. Catch per unit of effort for juvenile Dolly Varden in Sourdock Creek.**

The length frequency distribution for all juvenile Dolly Varden caught in Sourdock Creek is presented in Figure 13. There appear to be at least two year classes present (most likely 1+ and 2+) which is consistent with data collected in other Red Dog Mine area streams.



**Figure 13. Length frequency distribution of Dolly Varden in Sourdock Creek.**

## Lower Competition Creek (Station 202)

### Water Quality

Lower Competition Creek (Figure 1 and 14) water quality appeared to be moderated by input from Sourdock Creek. Element concentrations (metals) at Lower Competition Creek were substantially lower than at the Upper Competition Creek sample site from 2000 to 2002 (Weber Scannell and Ott 2006). Unlike the Upper Competition Site, no samples from Lower Competition Creek


contained concentrations of iron, nickel, or lead that exceeded the chronic criteria for aquatic life (Weber Scannell and Ott 2006). It was apparent in the 2014 to 2017 sample period that water quality had changed from that observed from 2000 to 2002 (clear water) to red/orange staining and opaque water (Figure 14).



**Figure 14. Lower Competition Creek (left photo 2000/2002 and right photo 7/30/2014).**

### Periphyton

Average chlorophyll-a concentrations in Lower Competition Creek from 2014 through 2017 were substantially lower than those found from 2000 to 2002 (Figure 15) – an indication of degraded water quality.



**Figure 15. Average concentration of chlorophyll-a ( $\pm$  1SD) in lower Competition Creek.**

## Invertebrates

The average aquatic invertebrate density in Lower Competition Creek was highest in 2000 (Figure 16). Average density ranged from 1.3 in 2014 to 26.2 invertebrates/ m<sup>3</sup> in 2000. Aquatic invertebrate densities from 2014 to 2017 were substantially lower than the previous sample period. The percent Chironomidae also was higher from 2014 to 2017 than it was from 2000 to 2002 (Figure 17). Taxa richness varied from 13 to 22 taxa per site over the sample years and generally was higher during the 2014 to 2017 sample period (Figure 18).




Figure 16. Average aquatic invertebrate densities  $\pm$  1SD in lower Competition Creek.





Figure 17. Percent Chironomidae and EPT in lower Competition Creek.




**Figure 18. Aquatic invertebrate taxa richness in lower Competition Creek.**

### Fish


Juvenile Dolly Varden use lower Competition Creek as rearing habitat during the ice-free season (Bradley and Ott 2018). In 2000, fyke nets were used to catch fish moving either upstream or downstream in early July and late July. Catches yielded four juvenile Arctic grayling and 38 juvenile Dolly Varden (Weber Scannell and Ott 2006).

Minnow trap data collected from 2000 to 2002 and from 2014 to 2017 are presented in Figure 19. The CPUE has been highly variable with a low of two fish/day in 2000 and a high of 47 fish/day in 2016. Generally, the CPUE was higher from 2014 to 2017 and may reflect a higher number of fish using this section of the creek due to degraded water quality conditions in the upper part of the drainage (e.g., Upper Competition Creek).




**Figure 19. Catch per unit of effort for juvenile Dolly Varden in lower Competition Creek.**

The length frequency distribution of juvenile Dolly Varden in Lower Competition Creek is presented in Figure 20. There appear to be at least two year classes (most likely 1+ and 2+) which dominate the catch, and a small number of larger fish (multiple age classes). Data presented in Figure 20 include the minnow trap and the fyke net catches.




**Figure 20. Length frequency distribution of Dolly Varden in Lower Competition Creek.**


Juvenile Dolly Varden were retained from Lower Competition Creek from 2015-2017 for whole body element concentration (cadmium, lead, selenium, zinc, and mercury). These data were compared with Dolly Varden collected in Mainstem Red Dog and Anxiety Ridge creeks (Figures 21 to 25) during the same time frames. The cadmium, lead, and selenium concentrations were higher in Mainstem Red Dog Creek and the zinc and mercury concentrations were similar (Figures 21 to 25).




**Figure 21. Average cadmium concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17.**




**Figure 22.** Average lead concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17.



**Figure 23.** Average selenium concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17.



**Figure 24. Average zinc concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17.**



**Figure 25. Average mercury concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Competition, Anxiety Ridge, and Mainstem Red Dog creeks, 2015-17.**

## West Fork Ikalukrok Creek (Station 205)

### Water Quality


West Fork Ikalukrok Creek (Figures 1 and 26) had a relatively high hardness combined with low alkalinity and higher concentrations of sulfate, which indicates that this system was dominated by calcium sulfate rather than calcium bicarbonate (Weber Scannell and Ott 2006). From 2000 to 2002, the pH in this creek was low and ranged from 4.3 to 6.8. West Fork Ikalukrok Creek had high concentrations of most elements analyzed, especially aluminum, cadmium, copper, nickel, and zinc. Since sampling began in the area, a white precipitate (probably zinc and/ or aluminum) has been observed at the mouth of the creek as the waters mixed with Ikalukrok Creek (Figure 26).



**Figure 26. West Fork Ikalukrok Creek (7/12/2015), note white staining in right photo at confluence of West Fork and Ikalukrok.**

### Periphyton

In West Fork Ikalukrok Creek, the average chlorophyll-a concentrations from 2015 through 2017 were generally lower than those found from 2000 to 2002 (Figure 27). Average chlorophyll-a concentration varied from a low of 0.25 mg/m<sup>2</sup> in 2017 to a high of 3.45 mg/m<sup>2</sup> in 2002.



**Figure 27. Average concentration of chlorophyll-a ( $\pm$  1SD) in West Fork Ikalukrok Creek.**

### Invertebrates

The average number of aquatic invertebrates per m<sup>3</sup> of water in West Fork Ikalukrok Creek was low (<2 invertebrates/m<sup>3</sup>) from 2000 to 2002, but higher from 2015 to 2017 (Figure 28). In 2015, the aquatic invertebrate density was very high (27.7/m<sup>3</sup>) and was dominated by mayflies. The

percent Chironomidae exceeded the EPT in four of the six years (Figure 29). Taxa richness varied from 15 to 21 taxa per site over the sample years (Figure 30).




Figure 28. Average aquatic invertebrate densities ± 1SD in West Ikalukrok Creek.




Figure 29. Percent Chironomidae and EPT in West Ikalukrok Creek.




Figure 30. Aquatic invertebrate taxa richness in West Ikalukrok Creek.

## **Fish**

Fish have not been caught or been observed in West Fork Ikalukrok Creek, which may be due to degraded water quality in Ikalukrok Creek from various seeps, including the Cub Creek seep (located approximately 2 km downriver). These mineral seeps likely create a chemical barrier to fish passage.

## **Upper Ikalukrok Creek (Station 206)**

### **Water Quality**


Upper Ikalukrok Creek (upstream of West Fork Ikalukrok Creek) is a clear water system with fairly good water quality (Weber Scannell and Ott 2006). From 2000 to 2002, the pH was near neutral and ranged from 6.5 to 8.1. Concentrations of all metals in Upper Ikalukrok Creek are substantially lower than in Ikalukrok Creek downstream of the Cub Creek seep (Figure 31).



**Figure 31. Ikalukrok Creek immediately upstream of West Fork Ikalukrok Creek (left photo 7/3/2017) and Cub Creek (right photo 7/30/2014).**

### **Periphyton**


In Upper Ikalukrok Creek average chlorophyll-a concentration from 2000 to 2002 and from 2015 to 2017 were similar. Throughout the sample time frame, average chlorophyll-a concentration varied from a low of 0.3 mg/m<sup>2</sup> (2016) to a high of 3.48 mg/m<sup>2</sup> (2002) (Figure 32).




**Figure 32. Average concentration of chlorophyll-a ( $\pm$  1SD) in Upper Ikalukrok Creek.**

### **Invertebrates**


The average number of aquatic invertebrates in Upper Ikalukrok Creek was moderately high for all years except 2016 when it was extremely high (Figure 33). In 2016, the aquatic invertebrate density was 259.9/m<sup>3</sup> and was due to a substantial number of mayflies. The percent EPT in two of the six years of sampling greatly exceeded the chironomids (Figure 34), again due to abundant mayflies. The taxa richness varied from a low of 13 taxa in 2002 to a high of 28 taxa in 2017 (Figure 35).



**Figure 33. Average aquatic invertebrate densities  $\pm$  1SD in Upper Ikalukrok Creek.**



**Figure 34. Percent Chironomidae and EPT in Upper Ikalukrok Creek.**



**Figure 35. Aquatic invertebrate taxa richness in Upper Ikalukrok Creek.**

## Fish

Similar to West Fork Ikalukrok Creek, fish have not been caught or observed in Upper Ikalukrok Creek, even though there appears to be high quality fish habitat in the creek. This lack of fish may be due to degraded water quality downstream that likely creates a chemical barrier to fish passage.

## Ikalukrok Creek (Station 207)

### Water Quality

Ikalukrok Creek, upstream of East Fork Ikalukrok Creek and downstream of West Fork Ikalukrok Creek, is directly impacted by natural mineral seeps – the most visible being Cub Creek, located upstream of this section of Ikalukrok Creek (Weber Scannell and Ott 2006). The 2005-2018 pH of water samples from Cub Creek have ranged from 2.5 to 7.3, with a median value of 3.4 (Napier, 2019 pers comm). Substrate in this section of Ikalukrok Creek is stained red with iron flocculent (Figure 36) and in some years the staining extends downstream for several kilometers. Specific element concentrations in Ikalukrok Creek were high (aluminum, cadmium, copper, iron, nickel, lead, and zinc) and often exceeded the US EPA chronic criteria for aquatic life (Weber Scannell and Ott 2006). The pH was below the range for aquatic life in most of the water samples collected by Teck (Weber Scannell and Anderson 2000).




**Figure 36. Ikalukrok Creek below Cub Creek on July 3, 2016 (left photo is the sample site and right photo is a rock scraped for a periphyton sample).**

### Periphyton


In Ikalukrok Creek, the average chlorophyll-a concentrations in all samples were low. Additional samples were collected in 1997 and 1998 as a part of a Teck supplemental environmental project (Ott 1997), and in most years, chlorophyll-a was below the detection limit. The highest average chlorophyll-a concentrations were in 2001 (0.08 mg/m<sup>2</sup>), 2016 (0.06 mg/m<sup>2</sup>), and 2017 (0.05 mg/m<sup>2</sup>).

## Invertebrates


In Ikalukrok Creek aquatic invertebrate density varied from a low of 3.6 invertebrates/m<sup>3</sup> to a high of 28.9 invertebrates/m<sup>3</sup> (Figure 37). The percent chironomids generally exceeded the EPT, but in three samples (1998, 2016, and 2017) the percent EPT was higher (Figure 38). The higher percent in those years was due to both mayflies and stoneflies which may have been drifting from the upper reaches of the system. Taxa richness was highly variable, from a low of 10 taxa in 1998 to a high of 26 taxa in 2016 (Figure 39).



**Figure 37. Average aquatic invertebrate densities  $\pm$  1SD in Ikalukrok Creek.**



**Figure 38. Percent Chironomidae and EPT in Ikalukrok Creek.**



**Figure 39. Aquatic invertebrate taxa richness in Ikalukrok Creek.**

## Fish

During the 2000 sampling event, one lethargic adult Arctic grayling was observed in Ikalukrok Creek (Weber Scannell and Ott 2006), however since then no fish have been caught or observed in this section of Ikalukrok Creek. Similar to West Fork Ikalukrok and Upper Ikalukrok creeks, it is assumed that the degraded water quality from various seeps is limiting fish movement into high quality habitat particularly in Ikalukrok Creek upstream of West Fork Ikalukrok Creek.

## East Fork Ikalukrok Creek (Station 208)

### Water Quality


East Fork Ikalukrok Creek is a clear water system that joins with Ikalukrok Creek just downstream of the sample site (Figures 1 and 42). Only one spring water sample exceeded acute chronic criteria for cadmium, lead, and zinc (Weber Scannell and Ott 2006). The pH was near neutral and ranged from 6.6 to 8.5 with lower values in early spring during snowmelt. Water has moderately high hardness (median 130 mg/L) and alkalinity (median 117 mg/L), which is typical of a calcium-bicarbonate dominated system (Weber Scannell and Ott 2006). Extensive aufeis occurs in the canyon-like area both above and below the falls.



**Figure 40. Drift nets in East Fork Ikalukrok Creek.**

**Periphyton**

Chlorophyll-a concentrations were high in East Fork Ikalukrok Creek in most years, ranging from 1.21 mg/m<sup>2</sup> in 2016 to 7.36 mg/m<sup>2</sup> in 2002 (Figure 41). Generally, among the sample sites, East Fork Ikalukrok Creek was highly productive as measured by chlorophyll-a concentration.



**Figure 41. Average concentration of chlorophyll-a ( $\pm$  1SD) in East Fork Ikalukrok Creek.**

## Invertebrates

Aquatic invertebrate density in East Fork Ikalukrok Creek varied from a low of 0.7 invertebrates/m<sup>3</sup> to a high of 26.0 invertebrates/m<sup>3</sup> (Figure 42). Generally, the average aquatic invertebrate density was high during all sample events except July 2002. The percent chironomids exceeded the EPT in most of the samples, but EPT (both mayflies and stoneflies) was well represented in 1997 and 1998 (Figure 43). Taxa richness varied from 13 to 20 taxa per site over the sample years (Figure 44).




Figure 42. Average aquatic invertebrate densities  $\pm$  1SD in East Fork Ikalukrok Creek.






Figure 43. Percent Chironomidae and EPT in East Fork Ikalukrok Creek.



**Figure 44. Aquatic invertebrate taxa richness in East Fork Ikalukrok Creek.**

### Fish

Fish sampling with minnow traps and angling, including visual observations and aerial surveys, was conducted in East Fork Ikalukrok Creek. Fish sampling with minnow traps was done in East Fork Ikalukrok Creek in 1999, from 2000 to 2002 (two sampling events per year), and in 2016 and 2017 (Figure 45). The majority of catches were juvenile Dolly Varden, but two slimy sculpin (81 and 108 mm) were also caught in 2017.



**Figure 45. Catch per unit of effort for juvenile Dolly Varden in East Fork Ikalukrok Creek.**

Aerial surveys (helicopter) were conducted opportunistically and Arctic grayling were sampled by angling (Weber Scannell and Ott 2006). In some years (1999, 2000, and 2002) from 1997 to 2005, several hundred Arctic grayling were observed during aerial surveys in East Fork Ikalukrok Creek. Several Arctic grayling collected by angling in East Fork Ikalukrok Creek had been previously tagged in Mainstem Red Dog or North Fork Red Dog creeks.

In spring 2011, 14 adult Arctic grayling in North Fork Red Dog Creek were radio-tagged and tracked for one year. One of the adult Arctic grayling spent part of the summer in East Fork Ikalukrok Creek before moving to overwintering habitat in the Wulik River immediately downstream of Ikalukrok Creek (Ott and Morris 2012 and 2013). Five of the radio-tagged Arctic grayling returned to Mainstem Red Dog and North Fork Red Dog creeks in spring 2012 for spawning.

## Grayling Junior Creek (Station 209)

### Water Quality


Grayling Junior Creek is a clear water system that joins with Ikalukrok Creek just downstream of the sample site. It is the first major tributary entering Ikalukrok Creek from the east after the East Fork Ikalukrok and Ikalukrok creeks merge (Figures 1 and 48). Overall water quality was considered excellent with only a few samples exceeding the US EPA aquatic life criteria for aluminum and iron (Weber Scannell and Ott 2006). The pH was neutral to slightly basic and concentrations of zinc were slightly elevated and ranged from detection to 106  $\mu\text{g/L}$ .



**Figure 46. Grayling Junior Creek (left photo drift nets on 7/3/2016 and right photo drift nets on 7/4/2017).**

### Periphyton


Chlorophyll-a concentrations in Grayling Junior Creek varied from a low of 1.56  $\text{mg/m}^2$  in 2016 to a high of 4.63  $\text{mg/m}^2$  in 2002 (Figure 47). Generally, among the sample sites, Grayling Junior Creek was moderately productive as measured by chlorophyll-a concentration.




**Figure 47. Average concentration of chlorophyll-a ( $\pm$  1SD) in Grayling Junior Creek.**

### **Invertebrates**


Aquatic invertebrate density in Grayling Junior Creek varied from a low of 3.3 invertebrates/m<sup>3</sup> to a high of 11.6 invertebrates/m<sup>3</sup> (Figure 48). The EPT was composed of mayflies and stoneflies with very few caddisflies. In all sample years, the percent chironomids exceeded the percent EPT (Figure 49). Taxa richness was variable among sample events, varying from a low of 10 taxa in 2000 to a high of 24 taxa in 2017 (Figure 50).



**Figure 48. Average aquatic invertebrate densities  $\pm$  1SD in Grayling Junior Creek.**



**Figure 49. Percent Chironomidae and EPT in Grayling Junior Creek.**



**Figure 50. Aquatic invertebrate taxa richness in Grayling Junior Creek.**

## **Fish**


Aerial surveys (helicopter) were conducted opportunistically and Arctic grayling were sampled by angling (fish were tagged and recaptures recorded) and minnow traps (Weber Scannell and Ott 2006). In July 1999, we estimated about 300 adult Arctic grayling in Ikalukrok Creek at the mouth of Grayling Junior Creek (Figure 51). Mark-recapture sampling indicated that Arctic grayling moved between the Red Dog Creek drainage and Ikalukrok Creek drainage including Grayling Junior Creek. The very high numbers of fish seen at the mouth of Grayling Junior Creek was only documented in 1999.




**Figure 51. Arctic grayling at the confluence of Grayling Junior Creek and Ikalukrok Creek.**

Fish sampling with minnow traps was done in Grayling Junior Creek from 2000 to 2002 (two sampling events per year), 2004 (one sampling event), and in 2016 and 2017 (one sampling event) (Appendix 3). Dolly Varden juveniles, slimy sculpin, and age-0 Arctic grayling were captured in minnow traps. Arctic grayling age-0 fish were captured in late August 2004 indicating that spawning occurred there in spring 2004 ( $n = 5$ , 65 to 79 mm long, average 71.2 mm).

The CPUE for minnow traps varied from a low of two in 2017 to a high of 44 in 2002 (Figure 52). Length frequency distribution of Dolly Varden is shown in Figure 53. The majority of these fish, based on length, are likely age 1 and 2 with multiple age classes for the larger fish ( $\geq 120$  mm).



**Figure 52. Catch per unit of effort for juvenile Dolly Varden in Grayling Junior Creek.**




**Figure 53. Length frequency distribution of Dolly Varden in Grayling Junior Creek.**


In spring 2011, 14 adult Arctic grayling in North Fork Red Dog Creek were radio-tagged and tracked for one year. Two of the adult Arctic grayling spent part of the summer in Grayling Junior Creek. One of the radio-tagged fish in Grayling Junior was still present late in the fall and presumed to be dead (Ott and Morris 2012 and 2013). Five of the radio-tagged Arctic grayling returned to Mainstem Red Dog and North Fork Red Dog creeks in spring 2012 for spawning.

Juvenile Dolly Varden were retained in 2001 and 2004 for whole body element concentration (cadmium, lead, selenium, zinc, and mercury). These data were compared graphically with Dolly Varden collected in Mainstem Red Dog and Anxiety Ridge creeks (Figures 54 through 58) during the same time frame. The cadmium and lead concentrations were higher in Mainstem Red Dog Creek (Figures 54 and 55).


The selenium concentrations in whole body Dolly Varden were similar with the exception of fish from Mainstem Red Dog Creek in 2001 which had higher concentrations (Figure 56). Zinc concentrations were highest in fish from Grayling Junior Creek in 2001, but similar among fish from the three creeks for the remaining samples (Figure 57). Mercury concentrations were similar from all three creeks with concentrations near the detection limit for mercury (Figure 58).




**Figure 54. Average cadmium concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks.**




**Figure 55. Average lead concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 56. Average selenium concentrations ( $\pm 1SD$ ) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 57. Average zinc concentrations ( $\pm 1SD$ ) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 58. Average mercury concentrations ( $\pm 1SD$ ) in juvenile Dolly Varden collected from Grayling Junior, Anxiety Ridge, and Mainstem Red Dog creeks, note that samples were not tested for mercury in 2001.**

## Noa Creek (Station 210)

### Water Quality

Noa Creek, a tributary to Ikalukrok Creek, has degraded water quality with 95% of water samples exceeding the chronic aquatic life criteria for aluminum and cadmium, 90% exceeding the nickel


and zinc criteria, and 76% have a pH below the water quality criteria for aquatic life (Weber Scannell and Ott, 2006). Noa Creek is small and incised with breakup flows of about 20 cfs and has dense riparian vegetation (Figure 59).



**Figure 59. Noa Creek (left photo 7/4/2016 and right photo 7/4/2017).**

### **Periphyton**

Average chlorophyll-a concentrations in Noa Creek were low ranging from 0.17 mg/m<sup>2</sup> in 2002 to 0.32 mg/m<sup>2</sup> in 2000 (Figure 60).




**Figure 60. Average concentration of chlorophyll-a ( $\pm$  1SD) in Noa Creek.**


### **Invertebrates**

Aquatic invertebrate density varied from a low of 3.9 invertebrates/m<sup>3</sup> to a high of 38.1 invertebrates/m<sup>3</sup> and was dominated with aquatic diptera (Figure 61). EPT was virtually absent


from the samples (Figure 62). Taxa richness varied from 15 to 21 taxa per site over the sample years (Figure 63).



**Figure 61. Average aquatic invertebrate densities  $\pm$  1SD in Noa Creek.**



**Figure 62. Percent Chironomidae and EPT in Noa Creek.**



## **Figure 63. Aquatic invertebrate taxa richness in Noa Creek.**

### **Fish**

Fish sampling with minnow traps, including visual observations, was conducted in Noa Creek from 2000 to 2002 (two sampling days in 2000 and 2001 and one in 2002). Zero fish were caught and none were seen. Noa Creek is connected with surface flow to Ikalukrok Creek, so fish have access to the creek during the ice-free months

## **Moil Creek (Station 211)**

### **Water Quality**


Moil Creek, a tributary to Ikalukrok Creek, has degraded water quality with 95% of water samples exceeding the chronic aquatic life criteria for cadmium, copper, nickel, and zinc and 65% have a pH below the chronic criteria. Metals concentrations are high, especially copper, iron, aluminum, and zinc (Weber Scannell and Ott, 2006). Moil Creek is small with summer discharges typically ranging from three to five cfs (Figure 64).



**Figure 64. Moil Creek (left photo 7/4/2016 is the confluence of Moil with Ikalukrok and right photo 7/4/2017).**

## Periphyton


Average chlorophyll-a concentrations in Moil Creek were low, ranging from 0 mg/m<sup>2</sup> in 2002 to a high of 0.14 mg/m<sup>2</sup> in 2017 (Figure 65).




**Figure 65.** Average concentration of chlorophyll-a ( $\pm$  1SD) in Moil Creek.

## Invertebrates


Aquatic invertebrate density in Moil Creek varied from a low of 0.7 invertebrates/m<sup>3</sup> to a high of 6.5 invertebrates/m<sup>3</sup> and was dominated with aquatic diptera and miscellaneous aquatic species (Figure 66). EPT was virtually absent from the samples (Figure 67). Taxa richness varied from 13 to 21 taxa per site over the sample years (Figure 68).



**Figure 66.** Average aquatic invertebrate densities  $\pm$  1SD in Moil Creek.



**Figure 67. Percent Chironomidae and EPT in Moil Creek.**



**Figure 68. Aquatic invertebrate taxa richness in Moil Creek.**

## **Fish**

Fish sampling with minnow traps, including visual observations, was conducted in Moil Creek from 2000 to 2002 (two sampling days in 2000 and 2001 and one in 2002). Fish were not caught and none were seen. Moil Creek is connected with surface flow to Ikalukrok Creek so fish have access to the creek during the ice-free months. In some years, large adult Arctic grayling have been seen in Ikalukrok Creek near the mouth of Moil Creek, but have not been observed entering Moil Creek.

## Sled Creek (Station 212)

### Water Quality

Sled Creek, a tributary to Ikalukrok Creek, does not have a surface flow connection with Ikalukrok Creek. Overall, there was excellent water quality with only two samples exceeding the aquatic life criterion for aluminum and one sample exceeding the criteria for cadmium, copper, and zinc (Weber Scannell and Ott, 2006). Dense riparian vegetation is found throughout the sample reach (Figure 69).



Figure 69. Sled Creek (left photo 7/5/2016 and right photo 7/5/2017).

### Periphyton

Average chlorophyll-a concentrations in Sled Creek were relatively high ranging from 1.85 mg/m<sup>2</sup> in 2002 to 6.2 mg/m<sup>2</sup> in 2000 (Figure 70).






Figure 70. Average concentration of chlorophyll-a ( $\pm 1SD$ ) in Sled Creek.

## Invertebrates


Aquatic invertebrate density in Sled Creek varied from a low of 8.5 invertebrates/m<sup>3</sup> to a high of 59.3 invertebrates/m<sup>3</sup> (Figure 71). EPT were present in all sample years and twice exceeded the percentage of chironomids (Figure 72). In sample years where EPT was low there was a high number of aquatic diptera. Taxa richness was variable and ranged from a low of 14 to a high of 23 over the sample years (Figure 73).



**Figure 71. Average aquatic invertebrate densities  $\pm$  1SD in Sled Creek.**



**Figure 72. Percent Chironomidae and EPT in Sled Creek.**



**Figure 73. Aquatic invertebrate taxa richness in Sled Creek.**

## **Fish**

Fish sampling with minnow traps, including visual observations, was conducted in Sled Creek from 2000 to 2002 (two sampling days in 2000 and 2001 and one in 2002) and again in 2015. Fish were not caught and none were seen. Sled Creek is not connected with surface flow to Ikalukrok Creek during the ice-free season; therefore, fish have no access to the creek.

## **Volcano Creek**

### **Water Quality**


Volcano Creek was not sampled during the time frame from 2000 to 2002, but was sampled from 2014 to 2017. Visual observations and biological data collected indicate that Volcano Creek is a productive aquatic system (Figure 74).



**Figure 74. Volcano Creek (left photo minnow trap 7/29/2014 and right photo drift nets 7/29/2014).**

**Periphyton**


Average chlorophyll-a concentrations were high in Volcano Creek ranging from a low of 4.57 mg/m<sup>2</sup> in 2017 to a high of 6.32 mg/m<sup>2</sup> in 2014 (Figure 75). Chlorophyll-a concentrations in Volcano Creek were fairly consistent among the sample years.




**Figure 75. Average concentration of chlorophyll-a (± 1SD) in Volcano Creek.**

**Invertebrates**


Aquatic invertebrate density at Volcano Creek varied from a low of 3.7/m<sup>3</sup> to a high of 10.4/m<sup>3</sup> (Figure 76). EPT were present in all sample years with both mayflies and stoneflies represented (Figure 77). Samples were dominated by aquatic diptera, primarily chironomids. Taxa richness was variable and ranged from a low of 17 to a high of 25 over the sample years (Figure 78).



**Figure 76. Average aquatic invertebrate densities  $\pm$  1SD in Volcano Creek.**




**Figure 77. Percent Chironomidae and EPT in Volcano Creek.**




**Figure 78. Aquatic invertebrate taxa richness in Volcano Creek.**

## Fish

Fish sampling with minnow traps, including visual observations, was conducted in Volcano Creek from 2014 to 2017 (Appendix 1). In 2015 a second site (about 2 km further upstream) also was sampled. No fish were caught in 2014, but Dolly Varden were found in 2015, 2016, and 2017 (Figure 79) and slimy sculpin were caught in 2017. Length frequency distribution of all Dolly Varden caught is presented in Figure 80. Age 1 and 2 fish dominated the catch with a number of larger, older fish present.



**Figure 79.** Catch per unit of effort for juvenile Dolly Varden in Volcano Creek.




**Figure 80.** Length frequency distribution of Dolly Varden in Volcano Creek.


Juvenile Dolly Varden were retained in 2016 and 2017 for whole body element concentration (cadmium, lead, selenium, zinc, and mercury). A limited number of fish were retained in 2016 (n = 1) and 2017 (n = 2). These data were compared graphically with Dolly Varden collected in

Mainstem Red Dog and Anxiety Ridge creeks (Figures 81 through 84) during the same time frame. The cadmium concentrations in Volcano Creek were similar to those found in Mainstem Red Dog Creek (Figure 81).


Whole body lead and selenium concentrations in fish from Volcano Creek were similar to those found in Anxiety Ridge Creek but lower than fish collected in Mainstem Red Dog Creek (Figures 82 and 83). Zinc concentrations in juvenile Dolly Varden were slightly higher in Mainstem Red Dog Creek as compared to Anxiety Ridge Creek (Figure 84) and mercury concentrations were slightly higher in Anxiety Ridge Creek (Figure 85). All mercury concentrations were low.




**Figure 81. Average cadmium concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks.**




**Figure 82. Average lead concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 83. Average selenium concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 84. Average zinc concentrations ( $\pm$  1SD) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks.**



**Figure 85. Average mercury concentrations ( $\pm 1SD$ ) in juvenile Dolly Varden collected from Volcano, Anxiety Ridge, and Mainstem Red Dog creeks.**

## Literature Cited

- Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency; Office of Water; Washington, D.C.
- Bradley, P.T. 2017. Methods for Aquatic Life Monitoring at the Red Dog Mine Site. Alaska Department of Fish and Game Technical Report. No. 17-09.
- Bradley, P.T. and A.G. Ott. 2018. Aquatic biomonitoring at Red Dog Mine, 2017. A requirement under Alaska Pollution Discharge Elimination System Permit No. AK-0038652 – M1. Alaska Department of Fish and Game Technical Report. 18-06.
- Ott, A.G. 1997. July 6 Memo to J. Roberto at US Environmental Protection Agency. Alaska Department of Fish and Game Habitat and Restoration Division.
- Ott, A.G. and W.A. Morris. 2013. Aquatic biomonitoring at Red Dog Mine, 2012. National Pollution Discharge Elimination System Permit No. AK-003865-2. Alaska Department of Fish and Game Technical Report. 13-01.
- Ott, A.G. and W.A. Morris. 2012. Aquatic biomonitoring at Red Dog Mine, 2011. National Pollution Discharge Elimination System Permit No. AK-003865-2. Alaska Department of Fish and Game Technical Report. 12-02.
- Weber Scannell, P. and S. Anderson. 2000. Aquatic Taxa Monitoring Study at Red Dog Mine, 1997-1998. Alaska Department of Fish and Game Technical Report. 00-2.
- Weber Scannell, P. and A.G. Ott. 2006. Aquatic Baselines sampling, Wulik River Drainage. Volume I: Summary of Biological and Water Quality Information and Volume II: Appendices of Tabulated Data. Alaska Department of Natural Resources Technical Report. 03-05.

**Appendix 1. Total numbers and CPUE of Dolly Varden juveniles captured in minnow traps at various streams near the Red Dog Mine site.**

<b>Sample Location (Station)</b>	<b>Date Sampled</b>	<b>Hours Fished</b>	<b>Total Catch</b>	<b>CPUE</b>	<b>Other Fish</b>
Upper Competition Creek (203)	7/28/2000	23	5	5.2	
Upper Competition Creek (203)	7/7/2001	47	0	N/A	
Upper Competition Creek (203)	8/4/2001	28	0	N/A	
Upper Competition Creek (203)	8/3/2002	24	3	3.0	
Upper Competition Creek (203)	7/31/2014	19	0	N/A	
Upper Competition Creek (203)	7/31/2015	19.5	0	N/A	
Upper Competition Creek (203)	8/5/2016	23	0	N/A	
Upper Competition Creek (203)	8/6/2017	25.5	0	N/A	
Sourdock Creek (204)	7/9/2000	30	52	41.6	
Sourdock Creek (204)	7/28/2000	24	20	20.0	
Sourdock Creek (204)	7/7/2001	48	26	13.0	
Sourdock Creek (204)	8/4/2001	28	8	6.9	
Sourdock Creek (204)	7/9/2002	28.5	1	0.8	
Sourdock Creek (204)	8/3/2002	24	3	3.0	
Sourdock Creek (204)	7/31/2014	21	0	N/A	
Sourdock Creek (204)	7/31/2015	20	0	N/A	
Sourdock Creek (204)	8/5/2016	23.5	1	0.1	
Sourdock Creek (204)	8/6/2017	25	0	N/A	
Lower Competition Creek (202)	7/9/2000	32	4	3.0	
Lower Competition Creek (202)	7/29/2000	24	2	2.0	
Lower Competition Creek (202)	7/30/2000	24	2	2.0	
Lower Competition Creek (202)	7/6/2001	24	1	1.0	
Lower Competition Creek (202)	8/4/2001	28	11	9.5	
Lower Competition Creek (202)	7/9/2002	27	1	0.9	
Lower Competition Creek (202)	8/3/2002	24	33	33.0	
Lower Competition Creek (202)	7/31/2014	23	13	13.7	
Lower Competition Creek (202)	8/1/2015	26	35	32.3	
Lower Competition Creek (202)	8/5/2016	22.5	44	46.9	
Lower Competition Creek (202)	8/6/2017	27.25	22	19.4	

## Appendix 1 continued.

Sample Location (Station)	Date Sampled	Hours Fished	Total Catch	CPUE	Other Fish
West Fork Ikalukrok Creek (205)	7/8/2000	24	0	N/A	
West Fork Ikalukrok Creek (205)	7/28/2000	28	0	N/A	
West Fork Ikalukrok Creek (205)	7/7/2001	25	0	N/A	
West Fork Ikalukrok Creek (205)	8/3/2001	54	0	N/A	
West Fork Ikalukrok Creek (205)	7/11/2002	27	0	N/A	
West Fork Ikalukrok Creek (205)	8/2/2002	26	0	N/A	
West Fork Ikalukrok Creek (205)	8/1/2015	27	0	N/A	
West Fork Ikalukrok Creek (205)	8/5/2016	23	0	N/A	
West Fork Ikalukrok Creek (205)	8/7/2017	27	0	N/A	
Upper Ikalukrok Creek (206)	7/28/2000	28	0	N/A	
Upper Ikalukrok Creek (206)	7/7/2001	24	0	N/A	
Upper Ikalukrok Creek (206)	8/3/2001	54	0	N/A	
Upper Ikalukrok Creek (206)	7/11/2002	26.5	0	N/A	
Upper Ikalukrok Creek (206)	8/2/2002	26	0	N/A	
Upper Ikalukrok Creek (206)	8/1/2015	26.5	0	N/A	
Upper Ikalukrok Creek (206)	8/5/2016	23	0	N/A	
Upper Ikalukrok Creek (206)	8/7/2017	27	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	7/1999	24	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	7/8/2000	24.5	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	7/29/2000	23	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	7/7/2001	23.5	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	8/3/2001	53.5	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	7/12/2002	46.5	0	N/A	
Ikalukrok Creek (207) (DS of Cub Creek)	8/2/2002	25	0	N/A	
East Fork Ikalukrok Creek (208)	July of 1999	24	0	N/A	
East Fork Ikalukrok Creek (208)	7/8/2000	27	0	N/A	
East Fork Ikalukrok Creek (208)	7/29/2000	23	1	1.0	
East Fork Ikalukrok Creek (208)	7/8/2001	46	0	N/A	
East Fork Ikalukrok Creek (208)	8/3/2001	54	0	N/A	
East Fork Ikalukrok Creek (208)	7/12/2002	27	0	N/A	
East Fork Ikalukrok Creek (208)	8/1/2002	23	3	3.1	
East Fork Ikalukrok Creek (208)	8/5/2016	22.86	11	11.5	2 SS
East Fork Ikalukrok Creek (208)	8/7/2017	23.75	4	4.2	

## Appendix 1 continued.

<b>Sample Location (Station)</b>	<b>Date Sampled</b>	<b>Hours Fished</b>	<b>Total Catch</b>	<b>CPUE</b>	<b>Other Fish</b>
Grayling Junior Creek (209)	7/11/2000	23	14	14.6	
Grayling Junior Creek (209)	7/29/2000	28	8	6.9	
Grayling Junior Creek (209)	7/10/2001	42	5	2.9	
Grayling Junior Creek (209)	8/1/2001	46	12	6.3	
Grayling Junior Creek (209)	7/12/2002	26	0	N/A	
Grayling Junior Creek (209)	8/1/2002	24	44	44.0	
Grayling Junior Creek (209)	8/27/2004	24.02	20	20.0	5 AG, 2 SS
Grayling Junior Creek (209)	8/5/2016	27.75	4	3.8	3 SS
Grayling Junior Creek (209)	8/7/2017	24	2	2.0	1 SS
Noa Creek (210)	7/10/2000	28	0	N/A	
Noa Creek (210)	7/30/2000	23	0	N/A	
Noa Creek (210)	7/10/2001	22	0	N/A	
Noa Creek (210)	8/1/2001	46	0	N/A	
Noa Creek (210)	8/1/2002	22.5	0	N/A	
Moil Creek (211)	7/10/2000	27	0	N/A	
Moil Creek (211)	7/30/2000	23	0	N/A	
Moil Creek (211)	7/10/2001	22	0	N/A	
Moil Creek (211)	8/5/2001	24	0	N/A	
Moil Creek (211)	8/1/2002	24	0	N/A	
Sled Creek (212)	7/9/2000	26	0	N/A	
Sled Creek (212)	7/29/2000	25	0	N/A	
Sled Creek (212)	7/10/2001	24	0	N/A	
Sled Creek (212)	8/5/2001	24	0	N/A	
Sled Creek (212)	7/31/2002	23.5	0	N/A	
Sled Creek (212)	7/31/2015	22	0	N/A	
Volcano Creek (lower)	7/30/2014	26	0	N/A	
Volcano Creek (lower)	8/1/2015	22.5	32	34.0	
Volcano Creek (lower)	8/6/2016	22	3	3.3	
Volcano Creek (upper)	8/6/2016	22	4	4.4	
Volcano Creek (lower)	8/6/2017	21.75	2	2.2	2 SS