FINAL REPORT

Contract No. Research Unit #612 Reporting Period: 1 January 1980 -31 December 1982 Number of Pages:

Biological Investigations of Belukha Whales in waters of Western and Northern Alaska

Principal Investigators

John J. Burns, Kathryn J. Frost, Glenn A. Seaman, and Lloyd F. Lowry Marine Mammals Biologists Alaska Department of Fish and Game 1300 College Road Fairbanks, Alaska 99701

31 December 1982

Table of Contents

- Part I. Distribution, abundance and movements of belukha whales in western and northern Alaska by Glenn A. Seaman and Kathryn J. Frost
 - A. World Distribution
 - B. General Distribution in Alaska
 - C. Seasonal Distribution in Alaska
 - D. Regional Distribution and Abundance
 - 1. North Aleutian Basin
 - 2. Saint Matthew-Hall Basin
 - 3. Saint George Basin
 - 4. Navarin Basin
 - 5. Norton Basin
 - 6. Hope Basin
 - 7. Barrow Arch
 - 8. Diapir Field
 - E. Literature Cited

Part II. The biology and ecology of belukha whales in western and northern Alaska by John J. Burns and Glenn A. Seaman

- A. B. C.
- D.
- Ε.

Part III. Foods of belukha whales in western and northern Alaska by Lloyd F. Lowry, Kathryn J. Frost, and Glenn A. Seaman

- A. Introduction
- B. Methods and Materials
- C. Results
- D. Discussions
 - 1. Biases in stomach content analysis
 - 2. Feeding during spring migration
 - 3. Summer feeding in coastal area
 - 4. Sex and age related differences in foods
 - 5. Autumn and winter foods
 - 6. Trophic interactions
- E. Literature Cited

Part	I.	Distribution,	abundance,	and	movements	of	belukha	whales	in
		western and no	orthern Alas	ska					

1

by Glenn A. Seaman and Kathryn J. Frost

Α.	Worl	Distribution	•
в.	Gene	al Distribution in Alaska	•
c.	Seas	nal Distribution in Alaska	•
D.	Regi	nal Distribution and Abundance	• ·
	1.	North Aleutian Basin	•
	2.	Saint Matthew-Hall Basin	•
	3.	Saint George Basin	•
	4.	Navarin Basin	•
	5.	Norton Basin	•
·	6.	Hope Basin	•
	7.	Barrow Arch	•
	8.	Diapir Field	• .
Ε.	Lite	ature Cited	

A. World Distribution

Belukhas are widely though not uniformly distributed throughout most seasonally ice-covered waters of the northern hemisphere (Fig. 1). They are circumpolar, occurring off North America, Europe, and Asia (Kleinenberg et al. 1964). Based on a knowledge of seasonal patterns of movement and concentration areas, the presence of major though not complete geographical barriers, and differences in size of adult animals in different areas, it is likely that the population can be divided into a number of somewhat discrete stocks in some parts of their range (Sergeant and Brodie 1969; Gurevich 1980).

2

In general, belukhas spend the winter in ice-covered offshore waters. They are unable to make and maintain breathing holes in ice more than 8 cm thick so are found in areas where geographic, oceanographic, or meteorologic factors cause ice motion and the formation of openings (Kleinenberg et al. 1964; Burns et al. 1981). In spring, as soon as the ice begins to break up and move offshore belukhas move toward the coast, some making extensive migrations in excess of 2000 km and some moving relatively short distances toward shore. Most belukhas appear to spend the majority of the summer in coastal waters, especially in shallow bays or estuaries of large rivers, although an unknown proportion of some populations may remain associated with offshore pack ice. In late summer to late autumn they move generally away from the coast, ahead of or with advancing pack ice (Kleinenberg et al. 1964). In the eastern hemisphere belukhas occur regularly and in substantial numbers in the White, Barents, Kara, Laptev, East Siberian, and Okhotsk seas (Kleinenberg et al. 1964). They are sometimes present off the coasts of Norway, Holland, Denmark, and West Germany, and in cold winters have been sighted as far south as Great Britain (Tomilin 1957; Gurevich 1980).

Belukhas regularly occur throughout the north Atlantic and eastern Canadian Arctic north to 82°30'N near Ellesmere Island, western Greenland, and Spitsbergen and south to the Gulf of Saint Lawrence. They are occasionally present near the coast of Nova Scotia in the Bay of Fundy, and are rare off Labrador and Newfoundland. They are most abundant in Davis Strait, Baffin Bay, Ungava Bay, Hudson Bay, Hudson Strait, Foxe Basin, Lancaster Sound, Prince Regent Inlet, Barrow Strait, Peel Sound, Cumberland Sound, and Jones Sound and have also been observed near Iceland and Jan Mayen (Kleinenberg et al. 1964). The most southern extra-limital record along the east coast of North America is from Avalon, New Jersey (38°55'N). Locations of other extra-limital sightings from the east coast include Maine, Massachusetts, and Long Island (Reeves and Katona 1980).

In the western arctic belukhas are found in the Beaufort Sea, Amundsen Gulf, and M'Clure Strait. The western arctic stocks are apparently separated from those to the east by heavy pack ice which occurs in the western Canadian arctic islands (Sergeant and Brodie 1975). Belukhas are found in the Beaufort, Chukchi, and Bering seas, the latter including the Gulf of Anadyr, and Bristol Bay. A small apparently separate

stock occurs in Cook Inlet (Seaman and Burns 1981). They also are residents in the Sea of Okhotsk. During summer belukhas occur north of the Chukchi Peninsula and move as far west as the East Siberian Sea (Kleinenberg et al. 1964). In the eastern North Pacific region, extra-limital occurrences have been reported from as far south as Tacoma, Washington (47°15'N) (Scheffer and Slipp, 1948).

B. General Distribution in Alaska

Belukhas in Alaska are considered to comprise 2 stocks. One has a center of abundance in Cook Inlet where they are numerous throughout the year (Klinkhart 1966). They are known to range into the northern Gulf of Alaska from at least Kodiak Island to Yakutat Bay (Harrison and Hall 1978). Seasonal movements are poorly known, however, concentrations occur each summer near mouths of rivers flowing into Cook Inlet from the north and east. This project has not dealt with the Cook Inlet stock and it will not be considered in the remainder of this report.

The second, much larger, stock of belukhas ranges seasonally through the Bering, Chukchi, Beaufort, and at least parts of the East Siberian seas. During winter these whales occur throughout the ice fringe and front from the Alaska coast to Siberia, as well as in more northerly regions of the Bering and Chukchi sea pack ice where open water regularly occurs (Kleinenberg et al. 1964; Fay 1974; Seaman and Burns 1981). As the ice recedes in spring, a large segment of the population moves north, some of them passing Point Hope and Point Barrow during April to June (Braham and Krogman 1977; Fraker 1979). Those belukhas are thought to mostly

migrate eastward through offshore leads in the Beaufort Sea, then south along the west coast of Banks Island to Amundsen Gulf, then west to the Mackenzie River delta where they appear in late June (Sergeant and Hoek 1974; Fraker et al. 1978; Fraker 1980). Ice conditions allow late spring migrants to utilize a more direct route to the estuary. Other belukhas migrate less extensively and are seen in coastal waters of the Bering and Chukchi seas shortly after ice breakup in spring. During the summer months belukhas occur in the Bering, Chukchi, Beaufort, and East Siberian seas, primarily in coastal waters and the broad margin of pack ice. Major concentrations in western North American waters occur in Mackenzie Bay, Kugmallit Bay, Kasegaluk Lagoon, Kotzebue Sound, Norton Sound (including the Yukon River delta), and Bristol Bay. They have been recorded in major river systems several hundred kilometers from the ocean (Kleinenberg et al. 1964; Gurevich 1980; ADF&G, unpubl.). Belukhas leave the coastal zone in late summer to late autumn. Animals in the northern part of their range move southward ahead of and with the advancing ice pack, most of them passing through Bering Strait and into the Bering Sea (Fay 1974; Seaman and Burns 1981).

C. Seasonal Distribution in Alaska

We have compiled all available distribution information for belukhas in the Bering and Chukchi seas, the Beaufort Sea including Mackenzie Bay, and the eastern part of the East Siberian Sea (see also Gurevich 1980, for a review of the seasonal distribution of belukhas in Siberian waters). Data have been compiled by 2-month periods beginning in

January and are summarized in Fig. 3 through 8. Major locations are shown in Fig. 2.

б

January-February (Fig. 3)

Belukhas overwinter in both the Bering and southern Chukchi seas. Most midwinter sightings have been made from coastal villages south of 69°N latitude. Belukhas probably do not regularly overwinter in the Beaufort or northern Chukchi seas since the ice cover there is heavy, without extensive leads, polynyas, or other areas of predictably open water (Fay 1974). They may occasionally become entrapped by ice, however, and forced to remain in unsuitable regions (Freeman 1968). Mortality in such instances is probably high (Porsild 1918; Freeman 1968).

During January-February in the southeastern Chukchi Sea these whales have been observed from Point Hope to Bering Strait. Sightings have been by residents of Point Hope, Shishmaref, Wales, and Diomede. Winter distribution in the Chukchi Sea is probably variable depending on annual severity of ice conditions. Along the southwestern Chukchi coast they have been reported during winter from Cape Dezhnev (East Cape) and Serdtse Kamen' Cape (Kleinenberg et al. 1964). In the Bering Sea they occur in Mechigmen Gulf and Provideniya Bay on the Siberian coast, south and west of Saint Lawrence Island, and occasionally along the Alaskan coast from Norton Sound to Bristol Bay. In Bristol Bay belukhas are rarely seen by coastal residents during the coldest winter months (Brooks 1954; ADF&G, unpubl.). They generally occur in the outer regions of Bristol Bay and the Bering Sea at that time (Lensink 1961).

March-April (Fig. 4)

Observations from March and April indicate that belukhas are widely distributed in the Bering and Chukchi seas. They are present along the southern edge of the seasonal sea ice from Bristol Bay westward (Seaman and Burns 1981). Although sightings are widely dispersed throughout ice-covered regions of Bristol Bay and the Bering Sea, the greatest number of sightings has been in western Bering Sea from the ice edge to Bering Strait, including southeast of Saint George Island, south and southwest of Saint Matthew Island, around Saint Lawrence Island, and around the Diomede Islands and Cape Prince of Wales (Kenyon 1972; Braham et al. 1982; ADF&G, unpubl.). On the Siberian side they have been observed from Cape Navarin, Mechigmen Gulf, Serdtse Kamen' Cape and Cape Dezhnev (Kleinenberg et al. 1964; Seaman and Burns 1981).

7.

Belukhas in large numbers are first seen in nearshore waters of Bristol Bay in April as areas become ice free, frequently congregating at or near the mouths of large rivers to feed (Brooks 1956; Frost et al. 1982). They appear north of Bristol Bay along the coast in Etolin Strait and Hazen Bay in April, and are also commonly sighted off the shore ice in Norton Sound near Saint Michael, Shaktoolik and Cape Denbigh, near Cape Nome, and near the city of Nome.

Sightings in the Chukchi Sea in March and April occur mainly near the coast from Bering Strait to Cape Schmidt on the Siberian side and Point Barrow on the Alaskan side (Seaman and Burns 1981). The first sightings of belukhas off Point Hope are in March with larger numbers observed in April and May (Marquette 1976, 1977, and 1979; Braham and Krogman 1977; ADF&G, unpubl.). Belukhas first appear off Barrow in early to mid-April but most pass by in May (Braham and Krogman 1977; ADF&G, unpubl.).

May-June (Fig. 5)

In May and June belukhas are still reported throughout the northern Bering Sea. However, there are fewer offshore sightings and the majority of sightings are relatively nearshore. On the Siberian coast sightings occur along the coast from Cape Navarin to Cape Dezhnev (Tomilin 1957; Kleinenberg et al. 1964). In Alaskan waters belukhas occur from southeastern Bristol Bay to Bering Strait, including a few which pass by Saint Lawrence Island moving northward. Most sightings are from Bristol Bay and Norton Sound (Brooks 1954, 1955; Lensink 1961; Klinkhart 1966; Seaman and Burns 1981; Frost et al. 1982).

By June many belukhas have moved northward into the Chukchi Sea and arrived in the eastern Beaufort Sea where they congregate near Banks Island and Amundsen Gulf before moving into the Mackenzie River delta. Most sightings in the Chukchi Sea are from the Alaskan side, extending from Kotzebue Sound well into the Beaufort Sea northeast of Barrow (Childs 1969; Seaman and Burns 1981; Braham et al. 1982). Sightings during this period have also been made on the Siberian side near Serdtse Kamen' and along the coast as far west as Cape Schmidt (Kleinenberg et al. 1964).

July-August (Fig. 6)

July and August are the months during which peak use of coastal waters occurs in most areas. Along the Siberian coast belukhas are apparently rare in Karaginski Bay and common in the Gulf of Anadyr, western Bering Strait, and along the northern coast of the Chukchi Peninsula to the vicinity of Long Strait. There are few sightings during these months from the East Siberian Sea (Tomilin 1957; Kleinenberg et al. 1964). The distribution in Alaska during this period is generally continuous from Bristol Bay to the western Beaufort Sea and into Canadian waters of the eastern Beaufort Sea (Seaman and Burns 1981).

The largest number of sightings, and generally the largest groups of belukhas, are seen in inner Bristol Bay, particularly in Nushagak and Kvichak Bays (Brooks 1955; Lensink 1961; Frost et al. 1982); in Norton Sound near the Yukon River delta, Stebbins, Unalakleet, Shaktoolik, Koyuk, and Elim; in Kotzebue Sound and between Cape Lisburne and Point Barrow (mainly in and adjacent to Kasegaluk Lagoon); north of Barrow in late August and September (Seaman and Burns 1981; ADF&G, unpubl.); and in Canadian waters of the eastern Beaufort Sea (Fraker 1977). Groups of whales have also been sighted along the margin of the pack ice from Barrow southwest to Icy Cape and east to Barter Island (Harrison and Hall 1978; Braham et al. 1982; ADF&G, unpubl.).

September-October (Fig. 7)

The pattern of whale distribution changes markedly in September and October. Fewer whales are observed in coastal waters, and there is a general increase in offshore sightings. In the far north, animals from Siberia move east and seaward, while those from the eastern Beaufort move westward. Consequently, most sightings in September and October have been from the northern Chukchi Sea between Wrangell Island and northeast of Point Barrow (Seaman and Burns 1981).

Very large aggregations of belukhas have been seen at this time of year; sightings of 500 to more than 1,000 whales were made northeast of Barrow in September 1978 and October 1979 (John Bitters and Lloyd Zimmerman, pers. comm.) and of several thousand (perhaps more than 5,000) in the central Chukchi Sea in September 1974 (G.C. Ray and T. Dohl, pers. comm.).

Some sightings have also been made in the area from south of the pack ice to Bering Strait. Coastal residents of Bering Strait report belukhas moving southward in advance of the ice in October (Kleinenberg et al. 1964; ADF&G, unpubl.). Sightings along the Alaskan coast from Cape Prince of Wales to Bristol Bay become increasingly less common as winter approaches.

November-December (Fig. 8)

There are few sightings of belukhas whales in November and December. Most have been in the Bering Sea with a few in the Chukchi Sea from Point Hope southward (Seaman and Burns 1981). In general, sightings have been by coastal hunters and commercial airline pilots since survey efforts have been minimal during these months. Sightings of belukha whales from villages in Bering Strait indicate a predominently southward movement (Kleinenberg et al. 1964; F. Fay, pers. comm.; ADF&G, unpubl.) The southward movement characteristically peaks in November and early December with or in advance of the appearance of seasonal pack ice. This southward movement continues through midwinter (Kleinenberg et al. 1964).

It appears that belukhas maintain a close association with sea ice in winter, and that the timing of their southward migration is closely related to the timing of freeze-up and southward advance of the pack ice. Their distribution in March and April suggests that they are widely distributed throughout the Bering Sea in winter from the ice front to Bering Strait and the southern Chukchi Sea.

D. Regional Distribution and Abundance

1. North Aleutian Basin

For the purpose of this discussion the North Aleutian Basin is defined as the waters of Bristol Bay from Cape Newenham to Unimak Pass (Fig. 9).

12

Our information on the belukha whales of Bristol Bay comes from a variety of sources. From 1954 to 1958, J. Brooks conducted extensive studies on the distribution, movements, and feeding of belukhas in inner Bristol Bay. Lensink (1961) summarized Brooks' work and added information for areas north of Bristol Bay. More recent studies include the work of Harrison and Hall (1978) primarily in the Bering Sea, and Fried et al. (1979) and Lowry et al. (1982<u>a</u>) in Nushagak Bay. Other information is from the authors' observations in the area, ADF&G unpublished data, interviews with area residents, and correspondence with biologists working in the area.

Belukhas utilize the Bristol Bay area throughout the year. They are most common and occur in the largest concentrations in nearshore waters during ice-free months (Frost et al. 1982). Although small groups are occasionally observed nearshore in inner Bristol Bay during winter, they are considered uncommon there at that time (Brooks 1954, 1955; Lensink 1961; ADF&G, unpubl. data).

In general, during winter-early spring belukhas are widely distributed in outer Bristol Bay and the southeastern Bering Sea (Lensink 1961; ADF&G, unpubl.) and are believed to occur in close association with seasonal sea ice. They are probably more common during relatively heavy ice years when the seasonal ice extends south into the Bay than in years of less extensive ice cover. Most sightings at this time of year are of groups of 1 to 5 whales. A notable exception occurred on 13 April 1976 when over 300 whales were sighted northwest of Port Moller (Braham and Krogman 1977). Examination of satellite imagery indicates that this sighting occurred close to the southern edge the seasonal pack ice which extended unusually far to the south of that time (Burns et al. 1981).

In April as the seasonal ice starts to disintegrate and recede northward, belukha whales begin to move into coastal regions (Brooks 1956; Frost et al. 1982). Whales are found both offshore and nearshore at this time with sightings recorded from Hagemeister Island, Togiak Bay, and eastern Bristol Bay (Kenyon 1972). In April and May, concentrations of several hundred animals occur at the river mouths in Kvichak Bay (Brooks 1956; Frost et al. 1982). The first concentrations usually occur in and at the mouth of the Naknek River where the whales feed on smelt (Osmerus mordax). Belukhas, in groups of up to several hundred, ascend the Naknek River as soon as the ice goes out, moving at least as far upstream as King Salmon (30 river km from the mouth). When the ice in the Kvichak River breaks up (usually several weeks after breakup in the Naknek) belukhas move to the Kvichak River, apparently showing a preference for that area. They remain in Kvichak Bay during the sockeye salmon (Oncorhynchus nerka) smolt outmigration, often ascending the river on flood tides. Belukhas also occur in Nushagak Bay but are not common there until late June (Brooks 1956; Frost et al.

1982). In Nushagak Bay whales are most commonly seen near the mouth of the Snake River and in the northern part of the Bay near the junction of the Wood, Little Muklung, and Nushagak Rivers (Lowry et al. 1982a).

Prior to the mid-1960's belukhas moved into several of the major rivers of Bristol Bay from breakup until mid-June when an increase in boat traffic associated with the salmon fishery displaced them out of the rivers (Lensink 1961). Beginning in 1965 tape recorded sounds of killer whales (Orcinus orca) were used to repel belukha whales from the mouths of the Naknek River and later the Kvichak River (Fish and Vania 1971; N. Steen and D. Bill, pers. comm.). This effort was designed to reduce belukha whale predation of out migrating sockeye salmon smolt by keeping the whales away from areas with the highest smolt concentrations. The belukha "spookers" were normally in operation from the end of May through the first 2 weeks of June and effectively displaced belukhas from the Naknek and Kvichak Rivers during that period. When the use of spookers was discontinued in late June, belukhas again ascended these rivers but in low numbers, apparently due to heavy boat traffic. Attempts were made to extend the program to the Nushagak River but tides and other hydrological conditions prevented the establishment of a permanent program. After 1978 the belukha spooker program was discontinued, and belukhas have since resumed use of these river systems during the smolt outmigration (D. Bill and R. Randall, pers. comm.).

Belukhas are abundant in inner Bristol Bay through the remainder of the summer, but become progressively less common in autumn (Brooks 1954; Frost et al. 1982). They are observed there with some degree of

frequency until October when the whales are presumed to move offshore and westward. They have been reported east of Hagemeister Island in September (G. C. Ray, pers. comm.) and near Ugashik Bay in October (Harrison and Hall 1978). Local fishermen suggest they frequent the outer portions of the Bay. An October sighting near the Pribilof Islands confirms that belukhas do occur offshore over the continental shelf at this time (Harrison and Hall 1978). The degree to which belukhas utilize these offshore waters during summer and autumn is unknown. Sightings and changes in coastal abundance suggest that offshore habitats are not utilized extensively during the summer, but that they may be utilized during autumn. These changes correspond with the sharp decrease in abundance of anadromous fish in coastal waters.

The spring and summer movements of belukhas in inner Bristol Bay are reported to be closely tied with tidal movements. Lensink (1961) found that belukhas generally moved up the rivers and over the flats on flood tide. They usually returned to the Bay on ebb tide, although they occasionally remained in the deeper portions of the rivers through the tidal cycle. Fried et al. (1979 and pers. comm.) observed belukhas a considerable distance up the Nushagak River on all phases of the tidal cycle. S. Weston (pers. comm.) found that movements from Nushagak Bay up the Snake River occurred mainly on the flood tide and downstream movements on the ebb tide. Recent observations (Frost and Lowry, unpubl.) indicate that in the lower portions of the Snake River movements of whales are not closely correlated with tides although the direction of tidal flow has a major influence on river currents.

Although the general distribution of belukhas in inner Bristol Bay is similar at present to their distribution earlier in the century, the summer use patterns have changed in response to increased commercial fishing activity and the corresponding increased use of motorized boats. The principal salmon fishing season in Bristol Bay extends from about the 10th of June to the first week of August with the peak around the 4th of July. Adult red salmon start to appear in early to mid-June and at that time become the primary prey of belukha whales (Brooks 1955; Lensink 1961). With the commencement of the salmon fishery most whales avoid areas of intensive boat activity in the rivers and river mouths, particularly in the Naknek and Kvichak rivers, and tend to move to areas with less boat traffic. In Kvichak Bay, many belukhas appear to use the north side of the Bay, the area with the least fishing activity (R. Randall, pers. comm.).

During periods of intensive fishing in Nushagak Bay belukhas are commonly seen at the mouth of and ascending the Snake River where boat activity is relatively low (Fried et al. 1979; S. Weston, pers. comm.; Lowry et al. 1982<u>a</u>). Fishermen report that belukhas have used the Snake River area heavily for many years and many people believe that predation by the whales is the cause of poor salmon runs in the system. The Snake River originates from Lake Nunavagaluk which in Yupik Eskimo means "the poor lake" or, the lake with no fish. In 1979, smolt counts and escapement in the Snake River were much lower than in the surrounding stream and lake systems of similar size and characteristics (S. Fried, pers. comm.). Although the reason for the reduced productivity of this system

is not clearly understood, belukha predation on salmon smolt and adults may prevent the establishment of a larger salmon run.

Belukha whales calve in Bristol Bay in June and July (ADF&G, unpubl.). No large concentration areas associated with calving have been described in the literature. Fishermen and local residents have suggested that the lower Snake River and its mouth may be a calving site (Fried et al. 1979). A large number of belukhas (80-120) were observed in the mouth of the Snake River on 28 June 1979 but no newborn calves were seen (Fried et al. 1979). Neonates may have been present but due to their small size, dark coloration, and poor survey conditions, they may not have been observed (S. Weston, pers. comm.). Neonates were observed in a group of 400-600 whales near the mouth of the Snake River in early July 1982 (Lowry et al. 1982<u>a</u>). Belukhas may also calve in northern Kvichak Bay.

It is difficult to determine the abundance of whales in Bristol Bay; survey conditions are poor due to turbid water and dark colored juveniles are particularly difficult to see. Belkovich (1960) found that foraging belukhas in turbid waters were very sensitive to aircraft and tended to dive as soon as the aircraft was overhead. Aircraft at altitudes less than 200 m caused belukhas to dive deeper, remain longer underwater, spend less time at the surface, or move away from the area. However, Fraker (1980) flying at 300 m found that his survey aircraft did not disturb the whales. Sergeant (1973) in Hudson Bay and Fraker (1980) in the Mackenzie Estuary surveyed belukhas under similar hydrological conditions. Sergeant found that belukhas spend a third of

the time at the surface and the remainder of time underwater and thus multiplied his actual counts by 3 to account for the unseen animals underwater. Fraker assumed, since his view of an area was not instantaneous but lasted over 15 sec, that he would see a higher proportion; he multiplied the number of whales sighted by 2 to obtain the total number present. Most of the counts by Sergeant and Fraker were of nonfeeding whales which apparently remain at the surface longer than foraging whales.

There have been no systematic surveys of belukhas in inner Bristol Bay and the associated rivers. The best available counts were made by Brooks (1955, 1956) (Table 1). He estimated that at least 1,000 belukhas were present in Bristol Bay in 1954, and approximately half that number in 1955 and 1956. In the late 1970's fishermen reported belukhas to be moderately abundant (R. Baxter and M. Nelson, pers. comm.). Concentrations of several hundred whales still occur in the river mouths in spring and early summer. Sightings of up to 100-200 whales are regularly made in the Naknek River in April and May (D. Bill and N. Steen, pers. comm.). In late June 1979, during a flight over the north side of Kvichak Bay, R. Randall (pers. comm.) counted at least 250 belukhas and estimated that half the animals present were counted. During a single flight several years ago he counted 400 to 500 whales in Kvichak and Nushagak bays. Applying the sightability factor derived by Fraker (1978) this observation would account for at least 800 whales. Lowry et al. (1982a) estimated that 400-600 whales were in the vicinity of the Snake River mouth in early July 1982.

Table 1.	Estimated numbers of belukhas whales in inner Bristol Bay in 1954 and 1955 (Brooks 1955). Estimates were based on surface and aerial observations and interviews with fishermen and local residents.							
1954	May	June	July	August				
Kvichak Bay	250	250-400	?	600				
Nushagak Bay	?	250-400	400	400				
Total, both ba	ays, about 1,000			1997). 1997)				
1955								
Kvichak Bay	100	150-250	?	50-100				
Nushagak Bay	?	250	250-500	450				
Total, both ba	iys, about 525							

T2

2. Saint Matthew-Hall Basin

The proposed Saint Matthew-Hall OCS lease area, as discussed here, includes the coastal region of western Alaska from the southern Yukon Delta to Cape Newenham and westward to 174°W longitude (Fig. 10). Use of the Yukon River estuary by belukha whales is discussed in detail in the section dealing with the Norton Basin.

In winter and early spring belukhas occur throughout the Saint Matthew-Hall lease area except the immediate nearshore region where they may be excluded by shorefast ice. During occasional episodes of strong easterly winds which may break up the shorefast ice and move it offshore, belukhas have been seen near the mouths of the Yukon River, off Cape Romanzof, near Hooper Bay and in Kuskokwim Bay (Nelson 1887; Seaman, unpubl.). Residents of Hooper Bay report that such sightings occur during most winters (Seaman, unpubl.). Nelson (1887) reported that large numbers of belukha whales utilized the coastal regions south of Cape Vancouver during winter. Recent interviews with residents of coastal villages in that area generally confirm Nelson's observations but suggest that fewer whales may be present now (ADF&G, unpubl.).

Most winter and early spring observations of belukhas in the offshore portion of this basin during March and April have been west of 170°W. This is at least partially a result of the distribution of survey efforts. Belukhas are quite abundant in the large polynya and pack ice west and south of Saint Lawrence Island and are commonly sighted along the west and south shores of Saint Lawrence Island (see section on

Norton Basin). They have also been frequently sighted in polynyas south and southwest of Saint Matthew Island in March and April (ADF&G, unpubl.).

From breakup in May or June until freeze-up in October or November belukhas occur throughout coastal waters between the Yukon River delta and Cape Newenham. Their appearance and abundance is frequently associated with the availability and movements of various anadromous and marine fishes. In recent years their appearance in Kuskokwim Bay is reported by local residents to be irregular and of short duration. Sightings have been reported from Quinhagak, Tooksook Bay, and Kipnuk (ADF&G, unpubl.). Belukhas were considerably more common in Kuskokwim Bay earlier in the century (R. Baxter, pers. comm.). The last year belukhas were reported to be seen in large numbers near Quinhagak was around 1955. Belukhas have not been seen for many years in Goodnews Bay where they were previously very common (ADF&G, unpubl.). Formerly belukhas regularly entered the shallow waters of the Bay during the summer and were hunted by local residents.

Belukhas were formerly very abundant in the shallow waters of Jacksmith Bay (R. Baxter, pers. comm.). A village was located near there which depended to a large extent on an annual summer belukha hunt. It is said that in the early 1920's a large vessel came to Kuskokwim Bay and traded motor boats for king salmon (<u>Oncorhynchus tshawytscha</u>). The next year, about 1925, a very large belukha hunt took place in Jacksmith Bay in which it was reported that "all" the whales were killed. Belukhas failed to return to Jacksmith Bay in subsequent years and the settlement

there was abandoned (R. Baxter, pers. comm.). This may have been a cause and effect situation, but there is also the possibility that, as in other parts of Kuskokwim Bay, there was a general consolidation of many small settlements during this period.

Belukhas frequent the coastal waters between Cape Vancouver and the Yukon River delta during the spring, summer, and autumn. Observations by residents of Tanunak and local pilots indicate that belukhas are common in the Hazen Bay area where small groups are sighted every year during late spring and early summer. They are seen in the same area although somewhat less commonly during autumn. Belukhas are also present around Nunivak Island during the ice-free months but the degree of use at different times of the year is unclear. Historically, residents of Nunivak Island caught belukhas in nets during the autumn (Curtis 1930).

Belukhas are occasionally observed near and inside of Hooper Bay during the ice-free period, particularly during the late spring and early summer (Frost et al. 1982). The number of whales in the area varies greatly from year-to-year. Their presence is closely tied to runs of king and chum (<u>Oncorhynchus keta</u>) salmon (T. Ponaganuk, pers. comm.). Belukhas are also common just north of Hooper Bay near Cape Romanzof where they are often seen in May in association with schools of herring (<u>Clupea harengus</u>) (Frost et al. 1982). By early summer most whales leave this area and are believed to move to the Yukon River delta where they are very commonly seen during summer and autumn.

Based on available information about seasonal movements of belukhas in Norton Sound, Bering Strait, and the Saint Lawrence Island region it appears that a large portion of the whales that seasonally migrate through the Bering Strait to summer in the Arctic Ocean spend the winter in the Saint Matthew-Hall lease area. We estimate that from 40 to 75% of these whales either winter in the area or pass through it during migration.

3. Saint George Basin

The Saint George Basin lease area encompasses a large portion of the southeastern Bering Sea (Fig. 11). The northern portion of this area is on the continental shelf. The southern portion is off the shelf, with depths ranging from 200 m to as much as 1,500 m or more. The extent and characteristics of seasonal ice cover are highly variable in the lease area from year-to-year (Burns et al. 1981). During the "average" year, ice is usually present in the northern and northeastern portions, generally north of the Pribilofs. During cold winters or "heavy" ice years the ice may extend southward to approximately the continental shelf break, while in light ice years it may be entirely absent from the lease area.

It is difficult to assess the distribution and abundance of belukha whales in the Saint George Basin due to the scarcity of surveys and belukha sightings in the area. Most sightings have been made in conjunction with aerial and ship surveys directed at other species such as bowhead whales, walruses, and ice-associated seals. Since these

surveys are frequently restricted to particular habitats (ice front, shelf edge, etc.) where the target species are more likely to occur, they may not provide a reliable indication of the use of the area by belukhas.

We know of no sightings of belukha whales in the Saint George Basin during the summer. Harrison and Hall (1978) and Braham et al. (1982) surveyed a large portion of the area during summer and saw no belukhas. It is possible that they occur in small numbers since they have been seen in the area during the spring and autumn (Harrison and Hall 1978; ADF&G, unpubl.). When present, belukMas are probably restricted to the relatively shallow waters overlying the continental shelf and may be somewhat more common in the northeastern portion of the lease area which is closest to the coast.

In late summer and early autumn, belukhas start to leave the coastal areas of the Bering Sea and by mid- to late-autumn there is a clear decrease in abundance in some coastal areas, including in Bristol Bay. This decrease frequently parallels a decrease in the abundance of primary prey species. It is unclear where these whales go at this time, but, since there is little evidence of a shift to neighboring coastal areas, it is likely that at least some utilize the more offshore regions of the Bering Sea including the northern portion of the Saint George Basin. Harrison and Hall (1978) observed 2 belukhas on 11 October 1976, approximately 70 miles southeast of Saint George Island. Several species of suitable prey species are abundant in this area (Pereyra et al. 1976).

Belukhas apparently occur in greatest abundance in the lease area during winter and spring when seasonal ice excludes them from many nearshore regions. They are probably most common in Saint George Basin during heavy ice years when they have been observed in March and April near the Pribilof Islands, in western Bristol Bay, and south of Nunivak Island. They are probably less common when seasonal ice in Saint George Basin is minimal or absent. A significant but unknown proportion of the whales that winter in the Saint George Basin lease area probably summer in the coastal waters of the eastern Bering Sea and Bristol Bay.

We cannot presently estimate the number of belukhas utilizing the Saint George Basin. Based on limited sightings, the availability of apparently suitable habitat, and the area's proximity to coastal areas of high use by belukhas, the Saint George Basin lease area may be important to a large number of whales. We expect that the use of the lease area varies annually with peak use during winter and spring when ice is present in that portion of the area overlying the continental shelf.

4. Navarin Basin

The Navarin Basin includes a large portion of the central Bering Sea west of the Saint Matthew-Hall and northwest of the Saint George Basin lease areas (Fig. 11). The northern portion is on the continental shelf, while the southern part occurs over very deep water. Navarin Basin is remote; the closest land masses are Saint Matthew Island to the

east, Saint Lawrence Island to the northeast, and the coast of the USSR to the west.

It is difficult to assess utilization of the Navarin Basin by belukha whales due to the lack of settlements in the area and the near absence of sightings from any months except March, April, and early May. Aerial sightings of belukhas here have been opportunistic having been made in conjunction with surveys for other species.

It appears that the portion of the Navarin Basin overlying the continental shelf is an important part of belukha winter range. Similarly, the adjacent offshore waters to the east (western Saint Matthew-Hall lease area) and west (Gulf of Anadyr and westcentral Bering Sea) also appear to be important during winter. In the autumn and early winter large numbers of belukhas are consistently observed moving south into these regions through the Anadyr Strait between Saint Lawrence Island and eastern Siberia (Kleinenberg et al. 1964; Fay, pers. comm.; Seaman, unpubl. data). In March and April belukhas have frequently been sighted during the course of survey flights over the Navarin Basin (H. Braham, pers. comm.; ADF&G, unpubl.). They are also common east and west of the Basin during the same period. Although there are no sightings available for January and February their distribution then is probably similar to that in March and April although generally more northerly. As winter progresses the whales move southward with the advancing pack ice. Belukhas are thought to be rare or uncommon south of the continental

shelf because they are generally shallow feeders and the ice with which they are usually associated in winter rarely extends south of the shelf break (Burns et al. 1981).

As in other offshore areas, belukhas appear to move inshore or northward out of the Navarin Basin in spring. Residents of Gambell see these whales passing through Anadyr Strait in March and April, with the numbers diminishing in May (Seaman, unpubl.). Kleinenberg et al. (1964) observed several hundred whales in late May moving northward by Cape Navarin into the Gulf of Anadyr. In early May 1976, Soviet scientists observed numerous belukhas in this area as far south as the shelf break. These whales may have been associated with the ice remnant which predictably occurs in the northern Navarin Basin each year (Burns et al. 1981). The belukhas that summer in the Gulf of Anadyr may remain in association with the ice remnant until June, then move west to coastal areas of the Gulf of Anadyr where they are common until freeze-up (Tomilin 1957; Kleinenberg et al. 1964). Belukhas are probably rare or uncommon in the ice-free waters of the Navarin Basin in summer and early autumn when they are abundant in coastal areas. They probably return to the region when coastal areas such as the Gulf of Anadyr freeze over. Kleinenberg et al. (1964) indicate that new ice appears in the nearshore protected waters of the Gulf of Anadyr in October.

Many of the belukhas which summer in the Chukchi, Beaufort, and East Siberian seas probably occur seasonally in the Navarin Basin. Some may utilize the area for a major portion of those months when ice is present, while others may occur there for only a few days. The abundance of belukhas in the area is probably highly variable depending

on ice and feeding conditions, but is likely to be greatest during years of heavy ice.

5. Norton Basin

For the purpose of this discussion Norton Basin includes Norton Sound, the southwest coast of the Seward Peninsula, and the Chirikof Basin including Saint Lawrence Island (Fig. 12).

Belukha whales are uncommon during the coldest winter months in Norton Sound due to the usual presence of an extensive, comparatively unbroken ice cover. Hunters from Elim have reported sightings and occasional entrapment of belukhas in openings in the ice south of that village (Seaman, unpubl.) but such sightings are uncommon since leads in the ice are not regularly present in areas accessible by local hunters.

Belukhas utilize the coastal areas of Norton Sound including the Yukon River delta during the entire ice free period from breakup in May or June until freeze-up in October or November. Belukhas have often been sighted as early as April off the shorefast ice near Shaktoolik and Cape Denbigh. They are most common near the eastern Norton Sound villages of Stebbins, Saint Michael, Unalakleet, Shaktoolik, Koyuk, and Elim from late May through June and from September until November, although they are present throughout the summer (Frost et al. 1982). People from Stebbins, Saint Michael, and Elim believe that belukhas seen in spring frequent the mouths and nearshore waters off the Yukon River during the summer. Ray (1964 and 1975) identified the historically important belukha hunting areas in Norton Sound as Pastol Bay, the mouth of the Inglutalik River (Norton Bay), and Golovnin Bay. Nelson (1887) found that belukhas were very common in southern Norton Sound near Saint Michael and near the mouths of the Yukon River. Ford (no date) reported that they were historically common in Golovnin Bay and Golovnin Lagoon. Residents of Golovin and White Mountain confirm Ford's observations.

Belukhas begin to utilize the coastal areas of Norton Sound at a time which corresponds with the arrival of migratory and anadromous fishes. Herring spawning coincides closely with breakup, commencing in May in the southern Bering Sea and late May or early June in Norton Sound (Barton 1979). Belukhas are regularly seen following schools of herring throughout the coastal regions of Norton Sound, particularly near Golovnin Bay, Cape Denbigh, Point Dexter, and near Saint Michael (Nelson 1887; Giddings 1967; Barton, pers. comm.; Frost et al. 1982). Pilots in the region have observed belukhas feeding on herring in mid-June near Besboro Island. In 1981, belukhas were seen chasing and eating herring off Klikitarik (east of Saint Michael) in late April and Cape Stephens in mid-May (ADF&G, unpubl.). At least 100 were present and feeding on herring in the shallows near Point Dexter in late May 1981. Runs of herring are followed slightly later by capelin (<u>Mallotus villosus</u>) and salmon.

Throughout the summer and autumn belukhas are found near and in the mouths of the Yukon River where they feed on salmon. In July 1981 over 100 belukhas were seen feeding just off the northern mouth of the Yukon River and another smaller group was sighted to the east in outer Pastol

Bay (D. Ljungblad, pers. comm.). King, chum, sockeye, and silver (<u>Oncorhynchus kisutch</u>) salmon enter the Yukon River from late May to early September (Geiger and Andersen 1978). There are numerous historical accounts of belukhas ascending the Yukon River several hundred miles above tidal waters, probably following salmon. They have been reported from Nulato and Koyukuk, over 800 km from the river mouth, as recently as 1981 (Nelson 1887; Collins 1945; Lensink 1961; ADF&G, unpubl.). Residents of Tanana remember seeing belukhas near their village in the early 1900's. A group of 4 or 5 belukhas was reported several km upriver from Tanana (1200 km from the river mouth) in June 1982 and at the same general time a single large adult was reported 130 km further upriver above Rampart (F. Andersen, pers. comm.). In recent years belukhas have been observed occasionally at Mountain Village, 110 km upriver; historically they were seen there much more commonly.

In general, belukhas appear to move up the Yukon River less frequently than they did 50-75 years ago. They are still very common, however, around the mouths of the river where they feed in the shallows. Although the use of those waters may have been altered to some extent by increased fishing and related motorboat activity the changes are not reported to be great (ADF&G, unpubl.).

Near Saint Michael from midsummer to freeze-up Nelson (1887) found that belukhas fed extensively on the abundant saffron cod (<u>Eleginus gracilis</u>) found there. He observed that feeding occurred mainly at night and in the early morning in the bay near Saint Michael and in the many tidal

creeks south to Kuskokwim Bay. In late September and October of 1976 and 1981 groups of 30-60 belukhas were feeding during daytime on schools of saffron cod near Cape Darby and Rocky Point at the entrance of Golovnin Bay (Lowry et al. 1982<u>b</u>). About 150 belukhas were seen between Topkok and Bluff in early September 1981 (R. Nelson, pers. comm.).

There have been no systematic surveys directed toward determining the abundance of belukhas in Norton Sound or adjacent areas. The best available information on abundance is based on the observations of local residents and biologists working in the area. In combination, those sources suggest that the number of belukhas utilizing Norton Sound and the coastal regions south to Kuskokwim Bay at least equals and probably exceeds the number in Bristol Bay. A conservative estimate may be 1,000-1,200 whales, possibly as many as 2,000. Although calving probably occurs in Norton Sound, specific calving areas have not been. identified.

Along the coast of the Seward Peninsula from Cape Nome to Wales, belukhas are seen from spring through autumn. Some move north through leads in the ice off Wales in late March and early April. These early sightings are probably of whales migrating to the Beaufort Sea. Belukhas are most common near Wales from mid-March through mid-May when movement is generally northward and in October and November when they are reported to be moving south. Whales sighted during the summer may be moving in either direction. References by Curtis (1930), Thornton (1931), Van Valin (1941), and Ray (1964) indicate that belukhas were historically common in the coastal waters north and south of Wales

during autumn and early winter. Recent sightings suggest this is true today, and that at that time of year the whales are mostly moving southward.

Belukhas are sometimes seen in the pack ice off Cape Nome and the city of Nome as early as April. They were seen in early May of 1979 by Eskimos hunting walrus between Nome and Sledge Island. Cape Woolley and Cape Nome were once productive hunting sites for belukhas, with whales present throughout the ice-free periods but most common in early summer and autumn (Ray 1964; Seaman, unpubl.). In November 1977, 150-200 belukhas were seen moving by Cape Nome, and in November 1979, 75-100 whales were observed feeding there (R. Nelson, pers. comm.).

During spring and summer, belukhas appear to move through the area from Cape Nome to Wales, sometimes foraging along the way, but not forming any major local concentrations. Near Cape Nome in spring and early summer they feed on schools of saffron cod and later have been observed following schools of herring (L. Barton, pers. comm.). The relationship between the belukhas of Norton Sound and those seen along the outer coast between Cape Nome and Wales is unknown, but they may be the same whales moving back and forth or through the area. Historically large numbers of belukhas occurred in Port Clarence and Grantley Harbor, but today they are seen only occasionally and in small numbers (Ray 1964 and 1975; Seaman, unpubl.). In previous years when belukhas were common in Port Clarence and Grantley Harbor their appearance coincided with the appearance of spawning herring.

Near Saint Lawrence Island belukhas are seen commonly in the spring, and occasionally in autumn and winter, particularly when saffron and arctic (<u>Boreogadus saida</u>) cods are abundant (Seaman, unpubl.). They are rarely observed during the summer, usually as single animals or in very small groups of both gray and white individuals (Fay, pers. comm.; Seaman, unpubl.).

In some years large numbers of belukhas are seen along the north and west shores of Saint Lawrence Island prior to freeze-up. Occasional whales are seen in late October but most arrive from the north in November and December. Local residents report that belukhas are seen more often in the autumn at Gambell than at Savoonga; whales seen at Savoonga are usually following the coast of the island towards the west, occasionally remaining in the area for several days. Either before or shortly after the ice appears, belukhas move southward, at least some of them moving into the Gulf of Anadyr (Kleinenberg et al. 1964). F. Fay (pers. comm.) reported at least a thousand animals north of Saint Lawrence Island in late November and early December 1957. This group followed the north coast past Gambell, headed toward the Gulf of Anadyr. Sightings of equal or greater numbers were made north and west of Saint Lawrence Island in November of 1974 and 1976 (Seaman, unpubl.). Smaller groups (15-25) are seen nearly every year in December-March along the western and southern shores in areas of open water created by strong ocean currents and prevailing northwesterly winds. Groups of up to 250-300 have been reported near Southwest Cape (Seaman, unpubl.). In April, groups of belukhas (some comprised of over 100 whales) have been observed moving north by Gambell, Southwest Cape, Southeast Cape, and

East Cape. C. Ray (pers. comm.) and Braham et al. (1982) also reported many belukhas in this area in spring, particularly north and northwest of the Island.

6. Hope Basin

Hope Basin includes the southeastern Chukchi Sea from Bering Strait north to Cape Lisburne (Fig. 13). Most of our information on the distribution and movements of belukha whales in the coastal regions of the Hope Basin is based on field studies undertaken by the Alaska Department of Fish and Game at Point Hope and in Kotzebue Sound, and on interviews and conversations with long-time residents of coastal villages. There is little published information regarding belukhas in this area.

During winter, belukhas are not uncommon in the southern Chukchi Sea. Eskimo hunters from Wales see them in nearshore leads throughout the winter. In the 1950's when seal hunting was still a major winter occupation, belukhas were occasionally seen near Shishmaref, by hunters traveling to the shore lead. On 5 March 1976, hunters reported a group of 35 belukhas trapped in the ice about 45 km southwest of Shishmaref (ADF&G, unpubl.).

According to older residents of Shishmaref and Wales, belukhas were once common along the northern Seward Peninsula from Ikpek to Cape Espenberg during breakup and throughout the summer (Seaman, unpubl.). In the early 1900's, reindeer herders from Shishmaref and Wales saw belukhas
inside Lopp and Arctic lagoons in late June and July with the number of animals present varying from a few to several hundred. If left undisturbed whales would remain in the lagoons for extended periods. Belukhas occasionally entered Shishmaref Lagoon in July during periods of high water, and occurred along the nearby coast until freeze-up when they were sometimes caught in nets set in the drifting ice near the village (Seaman, unpubl.). At one time some of the people from Kotzebue Sound spent their summers fishing and hunting seals, caribou, and belukhas along the coast from Shishmaref to Cape Prince of Wales (Nelson 1887; Curtis 1930; Hall 1975; ADF&G, unpubl.).

Belukhas have been infrequently sighted near Shishmaref in recent years. One group of about 20 was sighted in the ice 7 km west of the village on 4 June 1979 (ADF&G, unpubl.). This change in occurrence of whales near Shishmaref apparently took place over a period of few years which coincided with the introduction and increased use of outboard-powered boats. Undoubtedly large numbers of belukhas pass along the north side of the Seward Peninsula in spring on their way to Kotzebue Sound and locations further to the north, but this migration is probably far enough offshore to pass unnoticed by coastal residents.

Belukhas have been reported as common summer residents of Kotzebue Sound for as long as there are published records for the area (Nelson 1887; Curtis 1930; Foote and Cook, unpubl.; Ray 1964; Foote 1965; Foote and Williamson 1966; Saario and Kessel 1966; Hall 1975; Ray 1975; Giddings 1967; Seaman and Burns 1981). Belukhas first appear in Kotzebue Sound in late May to mid-June, usually during or shortly after breakup when

ice is still present but is broken and scattered. They are often first seen in pockets of open water in northern Kotzebue Sound from Sheshalik to Cape Blossom. In 1978 the first confirmed sighting in Kotzebue Sound was made on 11 June southeast of Chamisso Island by a Kotzebue hunter enroute to Elephant Point. In 1979, a pilot from Kotzebue reported a group of about 30 whales on 1 June south of Cape Blossom. A group of 80-100 was seen at the same location on 6 June, and was observed approaching Sheshalik spit from the southwest shortly thereafter (G. Barr, pers. comm.). These first sightings in 1978 and 1979 were probably somewhat earlier than usual, since in both years the winters and springs were unusually warm and breakup occurred early. Foote and Cook (unpubl.) found that the first belukhas usually appeared near Sheshalik in mid- to late-June.

Eschscholtz Bay is a large shallow bay in the southeastern corner of Kotzebue Sound about 85 km southeast of Kotzebue. It is presently the most productive belukha hunting site in the Kotzebue Sound area. Belukhas normally appear in Eschscholtz Bay in mid-June, slightly later than in northern Kotzebue Sound. In 1978, hunters from Deering sighted a group of at least 50 on 12 June, 6 km west of Elephant Point, and in 1979 over 200 were seen on 8 June along the northwest shore (N. Lee, pers. comm.). Belukhas appeared somewhat later in 1982 with the first whales seen on 21 June. When in the area, belukhas normally move into Eschscholtz Bay each day on the flood tide and leave on the ebb tide, but sometimes remain in the bay through the tidal cycle. They follow a deep channel which extends from Chamisso Island and parallels the north shore towards the Buckland River. On high tide and the first part of

ebb tide, the whales commonly disperse along the north and east shores of the bay. On some flood tides they do not deeply penetrate the bay but concentrate in the shallow waters along the northeast shore. This may be due in part to avoidance of boat traffic near Elephant Point and the Buckland River. In June they are usually intercepted by hunters who herd or drive them into shallow waters of the inner bay. In June 1978, 1979, 1981, and 1982 belukhas continued to move in and out of the bay for a week to 10 days, presumably until hunting activity disturbed them to the the point they would no longer enter (Seaman and Burns, unpubl.). In some years after hunting ceases and all hunters leave some whales return to Eschscholtz Bay and remain until at least mid-July. (Seaman, unpubl.)

There appears to be considerable local movement of belukhas in Kotzebue Sound. The whales seen near Sheshalik, Kotzebue, and Cape Blossom are almost certainly part of the same group seen in Eschscholtz Bay. In 1979, during times when belukhas were not seen in Eschscholtz Bay, many whales were seen off Sheshalik, in Kotzebue Sound proper, and in Spafarief Bay just west of Eschscholtz Bay. On several occasions when there was much boat activity near both Sheshalik-Kotzebue and Eschscholtz Bay, belukhas were seen near Cape Blossom and seaward of Sheshalik. The residents of Deering say that belukhas are not seen near their village. Historically as well as in recent years the whales seem to have preferred the northern and eastern parts of the Sound.

The distribution and movements of belukhas in Kotzebue Sound appear to be markedly different today than in the early 1900's (Seaman and Burns

1981; ADF&G, unpubl.). Residents of Noatak and Kotzebue have noted that the greatest change occurred shortly after the introduction of outboardpowered boats in the 1920's and early 1930's. Foote and Cook (unpubl.) stated that before motorboats were used belukhas came very close to shore and often entered the shallows behind Sheshalik spit as well as Hotham Inlet. In the 1940's and 1950's there was a large increase in both the number and size of motorboats near Kotzebue and Sheshalik. Hunting became more difficult as fewer belukhas came into the shallows near these sites. By the 1960's boat traffic in northern Kotzebue Sound was heavy, traditional hunting methods gave way to less organized hunts, and fewer belukhas were seen in these shallow areas. Today belukhas are even less common near Sheshalik than in the 1960's although they are still common offshore. Many people from Kotzebue believe that the noises associated with modernization, such as electrical generation, construction, barge traffic, and low flying aircraft have compounded the problem,

Noticeable changes in utilization patterns and movements of belukhas have also occurred in Eschscholtz Bay. Traditionally only the people from the small village of Buckland and occasionally Deering hunted belukhas in Eschscholtz Bay. In the early 1900's the village was located on the lower Buckland River, and residents seasonally moved down river to Eschscholtz Bay for the belukha hunt. The whales, which were present in large but variable numbers every year, were hunted from umiaks and kayaks for 1 or 2 weeks in late June or July, or until enough meat was obtained. After the boat hunt was over (usually mid-July) belukhas returned to the bay and frequently stayed for days at a time

moving over the tidal flats on flood tide and to the deep water at ebb tide. The older people remember that very large numbers of whales were present after the hunt in July in the shallows east of Elephant Point and along the north shore. Belukhas frequented these areas until early August, after which time they were more commonly seen in western Eschecholtz Bay, near Chamisso Island and the Choris Peninsula, or in Spafarief Bay.

In the early 1920's a reindeer processing plant was established at Elephant Point. About the same time the Buckland people moved their summer hunting camp from the north side of the bay to this location and by the late 1930's the village of Buckland was situated at Elephant Point year-round. With the increase in noise and activity, belukhas spent more time on the northern side of the bay and came less frequently into the shallows east of Elephant Point. However, boat traffic was generally moderate prior to the 1950's since the village was located very near the hunting area. Boat traffic increased somewhat about 1954 when the village was relocated up the Buckland River to above tidewater and people began moving regularly back and forth. In the late 1960's a few hunters from other areas began to come to Eschscholtz Bay to hunt and by 1975 there were many additional boats, particularly from Kotzebue. Hunters are of the opinion that uncontrolled boat traffic in June and early July, particularly during flooding tides, acts to reduce the number of belukhas entering Eschscholtz Bay and to decrease hunting success.

Belukhas are known to both feed and calve in Kotzebue Sound. As in Norton Sound the whales probably follow local movements of fish, feeding on species which are particularly abundant at certain times (Seaman and Burns 1981; Seaman et al. 1982). In Eschscholtz Bay there are known to be substantial runs of herring, smelt, char (<u>Salvelinus alpinus</u>), and salmon, in addition to large numbers of saffron cod (Barton 1979; Burns, Frost, and Seaman, pers. obs.).

Calving has been reported in all coastal regions of the Sound; however, it is unknown whether calves are born only in shallow coastal regions or whether calving also occurs offshore. Most observations of calving are from near Sheshalik and from the eastern end of Eschscholtz Bay. The former area may be of lesser importance at present due to avoidance by whales as discussed above.

The actual number of whales using the Kotzebue Sound area during the ice-free months is poorly known. Our estimate of abundance is based on our field studies, interviews with local residents, and occasional observations of local pilots and biologists working in the area. In July 1962, Burns (unpubl.) saw 900-1,200 north of Chamisso Island, moving northward along the Choris and Baldwin peninsulas. On 8 July 1978 a resident of Buckland (N. Lee) saw an estimated 900-1,000 belukhas scattered in the shallows along the northwest shore of Eschscholtz Bay. At least 500 whales were seen from boats in Eschscholtz Bay on the first hunt in June 1978 and, based on hunting success, that is a very conservative estimate of the numbers of belukhas in the area that year. Local hunters reported that belukhas were also very abundant in 1977.

40.

In 1979 and 1981 hunters reported low numbers of whales which was reflected in very low harvests. In 1982 belukhas were very abundant in southeastern Kotzebue Sound and were also common near Sheshalik. Considering all observations we estimate that the peak number of whales in Kotzebue Sound during summer ranges from 500 to 1,800 with considerable year-to-year variability which cannot at present be explained. This estimate is based primarily on observations made in southeastern Kotzebue Sound and may poorly reflect whale abundance in other portions of the Sound. Systematic surveys of the area are needed in order to refine these estimates.

Belukhas appear off Kivalina and Point Hope, which are along the migratory route of whales headed to the eastern Beaufort Seas, much earlier in spring than they do in Kotzebue Sound. The northward spring migration past Point Hope has been documented by Foote (1960), Fiscus and Marquette (1975), Marquette (1976, 1977, and 1979), and Braham and Krogman (1977). At Point Hope belukhas are seen moving north through leads in the ice as early as March. The earliest recent sighting was on 21 March 1976 when 2 groups of approximately 80 and 120 whales were seen moving north through a lead southeast of Point Hope (Seaman, unpubl.). In late March 1978 more than 100 were seen moving through the leads near Point Hope and about 1,000 were seen on 19 May 1980 (D. Smullin, pers. comm.). Belukhas are commonly seen and sometimes hunted throughout April and May, although hunting for belukhas takes place primarily when bowhead whales (Balaena mysticetus) are not available (Marquette 1977; Braham and Krogman 1977; ADF&G, unpubl.). During spring most belukhas are seen swimming north, although in May 1976 several small groups were

seen swimming south. Most sightings near Kivalina are in April and May and again in early July (Seaman, unpubl).

Hunters from Point Hope frequently see belukhas while hunting seals among the ice floes in late June and early July. During July, August, and early September, many belukhas are sometimes seen along the coast between Kotzebue Sound and Point Hope (Nelson 1887; Foote 1960; Seaman, unpubl.). Residents of Kivalina commonly see belukhas during the first part of September, usually swimming northwest along the coast toward Point Hope; they are rarely seen after that time. Seal hunters from Point Hope report seeing belukhas moving southward by the village during September and October. Belukhas are uncommon off Point Hope in January and February, following periods of strong northerly winds that form leads and polynyas in the ice.

7. Barrow Arch

The Barrow Arch area includes the Chukchi Sea coast from Cape Lisburne to Point Barrow (Fig. 14). Most of the information presented below is based on our field studies conducted from 1978 to 1981, including interviews with local residents of Barrow, Wainwright, and Point Lay, and aerial surveys of the coast between Barrow and Cape Sabine. There is little published information on belukhas in this area, especially for the summer open water period.

There are 2 peak periods of abundance of belukhas along the northern Chukchi Sea coast. The first is comprised of whales migrating northward through leads in the pack ice during March-June. The second comprises whales that move into the coastal zone after the ice moves offshore in June-July. The timing of breakup is variable from year-to-year, with the ice moving out of the southern regions such as Ledyard Bay earlier than Peard Bay and Point Barrow to the north. On the average the shore ice leaves Cape Sabine and Cape Beaufort regions in mid-June and Point Barrow about 1 month later.

In spring, northward migrating belukhas have been reported moving by Wainwright through leads in the ice as early as March (Nelson 1969; ADF&G, unpubl.). However the peak of the spring migration past Wainwright and Barrow occurs in April and May (Braham and Krogman 1977; Seaman and Burns 1981). Groups of from 10 to several hundred whales have been seen in the flaw zone between Cape Lisburne and Barrow. The spring migration is largely complete by late May with most whales moving into the eastern Beaufort Sea to summer in coastal waters off the Mackenzie River delta (Fraker 1979).

The coastal area of the northern Chukchi Sea used most intensively by belukhas in June and July is in and adjacent to Kasegaluk Lagoon. Belukhas characteristically appear in the southern part of this region near Ledyard Bay in mid- to late-June then move gradually northward following the retreat of seasonal ice. Childs (1969) reported a group of 50 or more belukhas near the Pitmegea River on June 24, 1958.

Residents of Point Lay regularly see whales near Cape Beaufort prior to their arrival at Kasegaluk Lagoon.

Belukhas first appear near the village of Point Lay in late June or early July. In 1978 the people of Point Lay saw the first "summer" belukhas (as opposed to spring migrants to the Mackenzie River delta which may be seen in April and May) of the year in Naokok Pass on 2 July. At least 100 whales were moving northward close to the shore. In 1979 the first "summer" whales, a group of at least 100, were seen at Kukpowruk Pass on 22 June. This is one of the earliest recorded sightings in this area and was probably due to a very early breakup that year.

After their arrival belukhas may be found both outside the barrier islands and in deeper portions of Kasegaluk Lagoon, though nearshore waters outside the lagoon are used most extensively (Fig. 15). They are usually concentrated in and outside of major passes, particularly Kukpowruk, Utukok, Icy Cape, and Akoliakatat, and to a lesser extent Akunik. At a given time, most of the whales in an area usually concentrate in the same pass, rather than dispersing to several passes. In concentrations we observed the greatest number of animals were in or just outside of a pass, and most of the remainder were in the downstream plume of lagoon water. Most were within 1/2-3/4 km from shore, usually with a few small groups or solitary animals (all adults) farther offshore in deeper water. On 10 July 1978 a large group of whales observed near Kukpowruk Pass was concentrated in and south of the pass; the nearshore current was moving south that day. Whales seen at

Akoliakatat Pass on 13 and 15 July 1979 were in and to the northeast of the pass; the current was moving to the northeast. Water temperatures in these lagoon plumes were as much as 2°C warmer than up current from the passes.

Belukhas also occur in the deeper channels inside Kasegaluk Lagoon. Point Lay residents report that whales only enter the lagoon when water is moving out a pass. They often enter by one pass and leave by another. In some years many whales enter the lagoon while in others very few do so. The reason for this variability is unknown.

Sightings from 1979, the year for which our data are most complete, illustrate the movement of belukhas within the area. By late June the shore ice had moved offshore along the southern third of the lagoon; the pack ice remained close to shore near Utukok Pass and Solivik Island. The first belukhas were seen and hunted on 22 June at Kukpowruk Pass, after which they reportedly moved northward and offshore. They were next seen on 24 June moving southward along the beach near Point Lay and at the same location on 28 June moving northward. A pilot estimated 400-500 whales near Kukpowruk Pass on 30 June. None were seen on 1 July but many were present nearshore and inside of the lagoon near Kukpowruk Pass on 2 July. There appeared to be some movement of whales between the coast and the edge of the ice pack a short distance to the north.

On 3 July the coast was surveyed from Naokok Pass to Point Barrow. The pack ice extended south parallel to the coast to approximately midway between Utukok and Icy Cape passes. Several small groups of belukhas

were seen along the ice edge north of Utukok Pass. At approximately the same time over 500 were reported by a hunter about 125 km to the south at Cape Beaufort. No whales were seen between the 2 points. On 4 July a southwest wind blew the ice to just north of Icy Cape and a pilot saw approximately 200 whales near Icy Cape Pass. On 8 July over 500 whales were seen at Utukok Pass and on 9 July sightings of 300-500 animals were made in the vicinity of both Akunik and Naokok Passes.

The last large sighting of belukhas made at Point Lay in 1979 was on 13 July. Over 100 whales were moving northward from Kukpowruk Pass where they were seen the day before. East of Icy Cape, at Akoliakatat Pass, approximately 1,600-2,400 belukhas were present from 13 to 18 July, at which time they moved further east and north. They were sighted at Pingorarok Pass (1,000+ animals) on 19 July moving northward, and at Wainwright, 40 kilometers to the northeast between 17 and 20 July, with peak numbers late on the 19th (>500) and on the 20th (400). No whales were seen at Akoliakatat Pass on 19 July. Based on the timing of observations, the belukhas seen at Wainwright on the 19th and 20th were the same animals that were observed at Akoliakatat a few days earlier.

According to the residents of Point Lay, belukhas left the Kasegaluk Lagoon region unusually early in 1979. They are usually seen at Point Lay until at least the end of July and sometimes, as in 1978, until the middle of August. In 1981, however, no belukhas were seen on aerial surveys flown after 15 July. The whales usually depart to the north, occasionally following the coast where they are seen at Wainwright and

less commonly at Barrow. The factors affecting the timing of belukha movements are poorly known, but may include ice conditions, water temperature, food availability, human disturbance in the form of hunting or aircraft overflights, and the presence of killer whales (Belkovich 1960; Fish and Vania 1971; Sergeant 1973; ADF&G, unpubl.).

Belukhas are known to calve in the coastal region off Kasegaluk Lagoon. On aerial surveys in July 1978 and 1979, neonates were observed among the adult and subadult whales. Twice, on 8 and 10 July 1978, belukhas were observed giving birth in the lagoon (Seaman, unpubl.).

Our best estimates of the abundance of belukhas along this sector of the coast come from aerial photographic counts of whales at concentration areas. At Kukpowruk Pass on 10 July 1978, 703 whales were counted. At Akoliakatat Pass in 1979, 1,104 were counted on 13 July and 1,601 on 15 July. There are many problems associated with deriving an estimate of the total number of whales in concentration areas from aerial counts (Brodie 1971; Sergeant 1973; Fraker 1977). Due to their small size and dark coloration neonates and yearlings are difficult to see. Some of the animals are underwater at any given time, and, depending on turbidity of the water and the depth to which the belukhas dive, may or may not be visible to an observer. Animals outside the main concentration area (farther offshore) are more widely dispersed, appear to remain underwater longer and therefore are not seen in aerial photographs. By applying correction factors to account for the above problems in sightability we estimate the total numbers of belukhas at Kukpowruk Pass on 10 July 1978 as 1,200, at Akoliakatat Pass on 13 and

15 July, 1979 as 1,650 and 2,400 (Table 2). Based on the above observations we estimate that 2,000-3,000 belukhas may occur near Kasegaluk Lagoon in most years, although in some years the abundance of whales in the area may be considerably less.

Belukhas appear only occasionally at Wainwright and Barrow during the ice-free periods. Van Valin (1941) described a belukha hunt in "late spring" (presumably July) at Wainwright after the ice had gone out. Belukhas were seen and hunted there on 17 and 18 July, 1979. Nelson (1969) noted that during the ice-free season, belukhas were most commonly seen in late July and August, and were usually moving northeast along the coast. Informants from Wainwright confirmed that information and added that "long ago" belukhas in moderate numbers sometimes congregated at the mouth of Wainwright Inlet and moved into the Kuk River during summer. They are rarely seen near Wainwright in September. They were reported to be numerous off Wainwright on 3 September, 1975 and small groups were seen off the coast between Wainwright and Barrow on 11 and 13 September (Fiscus et al. 1976).

Hunters near Barrow occasionally see belukhas moving along the shore in summer and early autumn. Murdoch (1885) reported that in 1881, 1882, and 1883 large groups of belukhas passed by Barrow as soon as there was open water near the beach, and appeared again a week to 10 days later. He saw 100 or more whales pass by Barrow within 200 yards of the shore on 28 September 1881 but noted that September sightings were uncommon. Several older residents of Barrow indicated that belukhas were once commonly seen near the village every summer. Boat traffic and noise

	Kupowruk Pass	Akoliakatat Pass	
	10 July 1978	13 July 1979	15 July 1979
Photographic count	703	1104	1601
Correction factor for whales underwater and therefore not observed (1)	(140) 20%	(221) 20%	(320) 20%
Correction factor for areas where whales were not included in photos (2)	(211) 25%	(133) 10%	(192) 10%
Correction factor for neonates not observed (3)	(53) 5%	(73) 5%	(106) 5%
Correction factor for yearlings not observed (4)	(84) 8%	(117) 8%	(169) 8%
Total estimated number of belukhas in concentration	1191	1648	2388

Table 2. Counts from photographs, correction factors and total estimated numbers of belukhas seen near Kasegaluk Lagoon on survey flights in 1978 and 1979.

- 1. Brodie (1971) working in clear water in Cumberland Sound estimated that he missed counting 40% of the animals because they were underwater and too deep to see. He did not use aerial photographs. Sergeant (1973) believed that he saw only 33% of the total animals while working in the murky waters of Hudson Bay. Seaman is of the opinion that he was able to count a much greater proportion of the total animals present in and adjacent to Kasegaluk Lagoon through the use of aerial photographs. On the average, whales were observed for 15-20 seconds and appeared on as many as 3 or 4 frames.
- 2. Outside of the main concentration areas--the area covered by photographs--there were widely scattered individuals or small groups of whales that could not be included in photos taken on a single pass of the airplane. We assume that the age composition of those whales was similar to the main concentration.
- 3. Brodie (1971) estimated that neonates comprised 10% of the population. Since the peak of calving may not occur until several weeks after our survey flights we assummed only half of the calves had been born.

from large generators, vehicles, aircraft, etc., or other unknown factors may have discouraged whales from passing by nearshore in recent years.

There is little information on the distribution and numbers of belukhas in the Barrow Arch portion of the northwestern Chukchi Sea. Burns (unpubl.) recorded 13 whales in 5 scattered locations from 74° 20'N to 74' 41'N and 160° 54'W to 167° 24'W on 12 September, 1974. On aerial survey flights conducted 10 through 20 September, 1980, belukhas were sighted at 4 locations from 72° 35'N to 73° 00'N and 164° 00'W to 169° 00'W (Burns, unpubl.).

8. Diapir Field

The Diapir Field planning area includes the Alaskan Beaufort Sea and a portion of the northeastern Chukchi Sea east of 162°W longitude (Fig. 16). The continental shelf is quite limited in the Beaufort Sea generally extending offshore less than 100 km. Nearly continuous ice cover persists through much of the winter with a few offshore leads developing in the spring (Fraker 1979; Burns et al. 1981). Shorefast ice usually persists through June. In most years the pack ice retreats northward in mid- to late summer leaving the coastal waters ice free until freeze-up in late September to early November (Burns et al. 1981). In some years the ice never leaves the coastal waters of the Beaufort Sea while in other years the southern edge may be a hundred kilometers or more north of the continental shelf break.

Belukha whales are absent from the Diapir Field during most of the winter, from late November through March. Ice and weather conditions do not produce areas with predictable open water and favorable wintering conditions for belukhas generally do not occur. Small numbers of whales may become entrapped by ice during the autumn migration, but the incidence of this is probably low. Observations of entrapped belukhas in the eastern Canadian Arctic (Porsild 1918; Freeman 1968) suggest that attempts to overwinter under these conditions frequently result in high mortality.

Belukha whales are common, and at times very abundant, in the Diapir Field during the spring, summer, and autumn. The majority of belukhas that seasonally occur in the Beaufort Sea are part of a group of 6,000 to perhaps as many as 11,500 whales that summer in the Canadian Beaufort Sea and overwinter in the Bering and southern Chukchi seas (Braham and Krogman 1977; Fraker 1979; Davis and Evans 1982; ADF&G, unpubl.). The spring migration of belukha whales in the Chukchi Sea past Point Hope commences in mid- to late-March (see Hope Basin and Barrow Arch planning area discussions). The earliest recorded sighting of belukha whales passing Point Barrow was on 2 April 1977 when a Barrow hunter, Arnold Brower (pers. comm.), sighted over 60 animals moving through a narrow lead off the shorefast ice. Four days later several hundred whales were seen. It is possible that belukhas occasionally pass by Barrow as early as late March. Belukhas are known to utilize offshore leads during the spring migration (Braham et al. 1982) and it is likely that some pass Point Barrow unnoticed by local hunters.

The peak of the spring migration past Point Barrow occurs from late April to the third week of May and varies in relation to ice conditions (Braham and Krogman 1977; G. Carroll, pers. comm.). The general north and east migration past Point Barrow continues through at least early July. Braham et al. (1982) observed belukhas north of Barrow up until their last aerial surveys in late June. Harrison and Hall (1978) observed 2 groups of 9 and 23 belukhas north of Barrow the first week of July and Murdoch (1885) reported whales along the coast at Barrow in the middle of July immediately following breakup.

The proposed migration route of belukha whales from Point Barrow to the eastern Beaufort Sea is described in detail by Fraker (1979). Observations by Ljungblad (1981) confirm that in late May and June migrating whales utilize offshore leads in pack ice which extend northeast from Barrow, into the Beaufort Sea. Many whales appear to congregate in the leads, polynyas, and open water west of Banks Island and in the Amundsen Gulf in May and early June. By late June most of these whales have moved to the shallow, warmer waters of the Mackenzie River estuary where concentrations in excess of 2,000 whales are observed in some of the bays (Fraker et al. 1978). However, some belukhas have been seen moving eastward in Alaskan waters as late as 27 June (Ljungblad 1981).

Belukha whales have never been observed in large numbers during the summer in the coastal waters of the Alaskan Beaufort Sea. This is in marked contrast to the Canadian portion of the Beaufort Sea where up to 7,000 whales have been observed in the Mackenzie River estuary during July and August (Fraker 1980). Long-time residents of the Alaskan

mid-Beaufort region, Jim and Harmon Helmericks (pers. comm.), indicate that belukhas are common off the shorefast ice until it moves away from the coast which usually occurs between late June and the middle of July. During the remainder of the summer, belukhas are rarely observed in the ice-free coastal waters of the Alaskan mid-Beaufort Sea.

During summer, belukhas are occasionally observed in the pack ice or ice-free waters near Barter Island. In July, these whales are typically observed close to shore in small groups of 5 to 15 whales moving eastward. During August belukhas occur in groups of similar size but their direction of movement is less pronounced as whales are observed moving in both east and west parallel to the coast. Residents of Kaktovik on Barter Island killed several belukhas on 19 August 1980.

There have been very few sightings of belukha whales in the offshore waters and pack ice of the Beaufort Sea during July and early August. Harrison and Hall (1978) saw 4 whales on 18 August 1976, approximately 75 km north of Cape Halkett, and Ljungblad et al. (1982) made 3 sightings of 26 animals at 96, 220, and 270 km north of Prudhoe Bay on 22 August 1981 (Fig. 17). Considering the extensive effort that has been devoted to offshore aerial surveys in this region (e.g. Harrison and Hall 1978; Ljungblad et al. 1980; Ljungblad 1981), it appears that belukhas are uncommon in the open water areas of the Alaskan Beaufort Sea during July and early August.

Sightings of belukha whales in the eastern Beaufort Sea become increasingly infrequent in late August and September. Whales are seen headed

west past Herschel Island during September; a group of 2,000 was sighted near there on 21 September 1972 (Fraker et al. 1978). It appears that most whales move offshore prior to entering the Alaskan Beaufort Sea since they are rarely reported in nearshore waters. Johnson (1979) reported 2 sightings of whales swimming westward just offshore of the Jones Islands; a group of 75-100 was seen on 15 September 1977 and approximately 35 were seen on 23 September 1978. Cummings (pers. comm.) heard vocalizations of belukha whales near Prudhoe Bay in September 1980. No belukha whales were seen during marine mammal research done from a small boat in the nearshore Alaska Beaufort Sea from 20 August to 21 September 1980 and 16 August to 11 September 1981 (Frost and Lowry 1981; Lowry et al. 1981). The majority of belukha whales seem to remain well offshore throughout their westward migration out of the Beaufort Sea (Fig. 17). Large numbers of whales have been seen on several occasions in September and early October in the region 35-220 km north of Point Barrow (NARL, unpubl.; Burns, unpubl.). During September they may be dispersed over a wide area. For example on 21 September 1977, 100 were seen north of Prudhoe Bay and 2,000 were seen near Herschel Island. On 20 September 1980, Burns (unpubl.) observed numerous belukhas in a 140 km long band of pack ice north and east of Barrow. As discussed in the previous section whales are only rarely seen near Point Barrow during autumn.

E. Literature Cited

Barton, L. H. 1979. Finfish resource surveys in Norton Sound and Kotzebue Sound. Pages 75-313 in Environmental Assessment of the

Alaskan Continental Shelf, Final Reports of Principal Investigators, Vol. 4. Env. Res. Lab., Boulder, Colo.

_____, I. M. Warner, and P. Shafford. 1977. Herring spawning surveys--southern Bering Sea. Pages 1-112 <u>in</u> Environmental Assessment of the Alaskan Continental Shelf, Annual Reports of Principal Investigators, Vol. 7. Env. Res. Lab., Boulder, Colo.

- Belkovich, V. M. 1960. Some biological observations on the white whale from the aircraft. Zool. Zh. 39(9):1414-1422. (Translated from Russian by U.S. Naval Oceanographic Office, Transl. No. 403. 16pp.)
- Braham, H. W., and B. D. Krogman. 1977. Population biology of the bowhead (<u>Balaena mysticetus</u>) and beluga (<u>Delphinapterus leucas</u>) whale in the Bering, Chukchi, and Beaufort Seas. Northwest and Alaska Fisheries Center Processed Report. 29pp.

______, B. D. Krogman, and G. M. Carroll. 1982. Bowhead whale (<u>Balaena mysticetus</u>) migration, distribution, and abundance in the Bering, Chukchi, and Beaufort seas, 1975-1978, with notes on the distribution and life history of white whales (<u>Delphinapterus</u> <u>leucas</u>). NOAA Tech. Memo. NMFS SSRF-000 (in press).

Brodie, P. F. 1971. A reconsideration of aspects of growth, reproduction and behavior of the white whale (Delphinapterus

<u>leucas</u>), with reference to Cumberland Sound, Baffin Island population. J. Fish. Res. Board Can. 28:1309-1318.

Brooks, J. W. 1954. Beluga. Annual Report for 1954. Alaska Department of Fisheries. pp. 51-57.

_____. 1955. Beluga Investigation. Annual Report for 1955. Alaska Department of Fisheries. pp. 98-106.

_____. 1956. Marine predator control and investigation. Annual Report for 1956. Alaska Department of Fisheries. pp. 50-59.

- Burns, J. J., L. H. Shapiro, and F. H. Fay. 1981. Relationship of marine mammal distribution, densities, and activities to sea ice conditions. Pages 489-670 <u>in</u> Environmental Assessment of the Alaskan Continental Shelf, Final Reports of Principal Investigators, Vol. 11. Env. Res. Lab., Boulder, Colo.
- Childs, H. E., Jr. 1969. Birds and mammals of the Pitmegea River Region, Cape Sabine, northwestern Alaska. Biol. Pap. No. 10, Univ. of Alaska, Fairbanks. 76pp.
- Collins, H. B. 1945. The Aleutian Islands: their people and natural history. Smithsonian Institute War Background Studies, No. 21. Washington, D.C.

- Curtis, E. S. 1930. The North American Indian. Vol. 20. The Alaskan Eskimo. University Press, Cambridge, Mass. 320pp.
- Davis, R. A., and C. R. Evans. 1982. Offshore distribution and numbers of white whales in the eastern Beaufort Sea and Amundsen Gulf, summer 1981. Report by LGL Ltd., Toronto, Ontario for SOHIO Alaska Petroleum Co., Anchorage, Alaska and Dome Petroleum Ltd., Calgary, Alberta. 78pp.
- Fay, F. H. 1974. The role of ice in the ecology of marine mammals of the Bering Sea. Pages 383-399 <u>in</u> D. W. Hood and E. J. Kelly, eds. Oceanography of the Bering Sea. Occas. Publ. No. 2, Inst. Mar. Sci., Univ. of Alaska, Fairbanks.
- Fiscus, C. H., and W. M. Marquette. 1975. National Marine Fisheries Service field studies relating to the bowhead whale harvest in Alaska, 1974. Processed Rep. Natl. Mar. Fish Serv., Seattle, Wash. 23pp.
- , W. M. Marquette, and H. W. Braham. 1976. Abundance and seasonal distribution of bowhead and belukha whales - Beaufort Sea, northeastern Chukchi Sea. Pages 159-182 <u>in</u> Environmental Assessment of the Alaskan Continental Shelf, Annual Report of Principal Investigators, Vol. 1. Env. Res. Lab., Boulder, Colo.
- Fish, J. F., and J. S. Vania. 1971. Killer whale, <u>Orcinus orca</u>, sounds repel white whales, <u>Delphinapterus</u> <u>leucas</u>. Fish. Bull. 69:531-535.

Foote, D. C. 1960. The Eskimo hunter at Point Hope, Alaska, September 1959 to May 1960. U.S. Atomic Energy Comm. Contract No. AT(04-3)-315.

______. 1965. Exploration and resource utilization in northwestern arctic Alaska before 1855. Ph.D. Thesis. McGill Univ., Montreal, Quebec, Canada. 400pp.

, and H. A. Williamson. 1966. A human geographical study. Pages 1041-1107 <u>in</u> N. J. Wilimovsky and J. N. Wolfe, eds. Environment of the Cape Thompson Region, Alaska. U.S. Atomic Energy Comm., Oak Ridge, Tenn.

- Ford, J. P. No date. Hunting the white whale. The Wide World Magazine. Wikersham File No. 6705, Univ. of Alaska, Fairbanks. 4pp.
- Fraker, M. A. 1977. The 1977 whale monitoring program, Mackenzie estuary, N.W.T. F. F. Slaney and Co., Ltd., Vancouver, British Columbia, Canada. 53pp.

______. 1978. The 1978 whale monitoring program, Mackenzie estuary, N.W.T. F. F. Slaney and Co., Ltd., Vancouver, British Columbia, Canada. 28pp.

. 1979. Spring migration of bowhead (<u>Balaena mysticetus</u>) and white whales (<u>Delphinapterus leucas</u>) in the Beaufort Sea. Fish. Mar. Serv. Tech. Rep. No. 859, Dep. Fish. and Environ., Canada. 36pp.

_____. 1980. Status and harvest of the Mackenzie stock of white whales (<u>Delphinapterus leucas</u>). Rep. Int. Whal. Comm. 30:451-458, SC/31/SM8.

, D. E. Sergeant, and W. Hoek. 1978. Bowhead and white whales in the southern Beaufort Sea. Beaufort Sea Project, Tech. Rep. No. 4.

- Freeman, M. M. R. 1968. Winter observations on beluga (<u>Delphinapterus</u> leucas) in Jones Sound, N.W.T. Can. Field-Nat. 82:276-286.
- Fried, S. M., J. J. Laner, and S. C. Weston. 1979. Investigation of white whale (<u>Delphinapterus leucas</u>) predation upon sockeye salmon (<u>Oncorhynchus nerka</u>) smolts in Nushagak Bay and associated rivers: 1979 aerial reconnaissance surveys. Alaska Dep. Fish and Game, Unpubl. Rep., Proj. 11-41-6-340. 15pp.
- Frost, K. J., and L. F. Lowry. 1981. Feeding and trophic relationships of bowhead whales and other vertebrate consumers in the Beaufort Sea. Final Rep., Contract No. 80-ABC-00160, Natl. Marine Mammal Lab., NMFS, NOAA. 106pp.

_____, ____, and J. J. Burns. 1982. Distribution of marine mammals in the coastal zone of the Bering Sea during summer and autumn. Final Rep., NOAA, Outer Continental Shelf Environmental Assessment Program, Contract NA-81-RAC-00050. 166pp.

- Geiger, M. F., and F. M. Andersen. 1978. Annual management report, 1980, Yukon area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Juneau, Alaska.
- Giddings, J. L. 1967. Ancient men of the arctic. Alfred A. Knopf, New York. 391pp.
- Gurevich, V. S. 1980. Worldwide distribution and migration patterns of the white whale (beluga), <u>Delphinapterus</u> <u>leucas</u>. Rep. Int. Whal. Comm. 30:465-480, SC/31/SM 14.
- Hall, E. S., Jr. 1975. The Eskimo Storyteller. Univ. of Tennessee Press, Knoxville. 491pp.
- Harrison, C. S., and J. D. Hall. 1978. Alaskan distribution of the beluga whales (Delphinapterus leucas). Can. Field-Nat. 92:235-241.
- Johnson, S. R. 1979. Fall observations of westward migrating white whales (<u>Delphinapterus leucas</u>) along the central Alaskan Beaufort Sea coast. Arctic 32:275-276.

- Kenyon, K. W. 1972. Aerial surveys of marine mammals in the Bering Sea, 6-16 April 1972. Unpubl. Rep., Bur. Sport Fish. Wildl., U.S. Fish Wildl. Serv., Seattle, Wash. 79pp.
- Kleinenberg, S. E., A. V. Yablokov, B. M. Belkovich, and M. N. Tarasevich. 1964. Beluga (<u>Delphinapterus leucas</u>) - Investigation of the species. 376pp. (Translated from Russian by Israel Program Scientific Translations, Jerusalem, 1969.)
- Klinkhart, E. G. 1966. The beluga whale in Alaska. Alaska Department of Fish and Game, Juneau, Alaska. 11pp.
- Lensink, C. J. 1961. Status report: Beluga studies. Unpubl. ms., Div. of Biol. Res., Alaska Dep. Fish and Game, Juneau, Alaska. 20pp.
- Ljungblad, D. K., M. F. Platter-Reiger, and F. S. Shipp, Jr. 1980. Aerial surveys of bowhead whales, North Slope, Alaska. Final Report: Fall 1979. Naval Ocean Systems Ctr., Tech. Doc. 314, San Diego, Calif. 181pp.

. 1981. Aerial surveys of endangered whales in the Beaufort Sea, Chukchi Sea, and northern Bering Sea. Final Report: Fall 1980. Naval Oceans Systems Ctr., Tech. Doc. 449, San Diego, Calif. 211pp.

Lowry, L. F., K. J. Frost, and R. R. Nelson. 1981. Trophic relationships among ice-inhabiting phocid seals and functionally related marine mammals. Pages 1-41 in J. J. Burns, L. F. Lowry, and K. J. Frost. Trophic relationships, habitat use and winter ecology of ice-inhabiting phocid seals and functionally related marine mammals in the Arctic. Annual Report, Contract No. NA-81-RAC-00045, Outer Continental Shelf Environmental Assessment Program, Juneau, Alaska.

_____, _____, and ______. 1982a. Investigations of belukha whales in coastal waters of western and northern Alaska: marking and tracking of whales in Bristol Bay. Annual Report, Contract No. NA-81-RAC-00049, Outer Continental Shelf Environmental Assessment Program, Juneau, Alaska. 43pp.

______, _____, and J. J. Burns. 1982<u>b</u>. Investigations of marine mammals in the coastal zone of western Alaska during summer and autumn. Annual Report, Contract No. NA-81-RAC-00050, Outer Continental Shelf Environmental Assessment Program, Juneau, Alaska. 37pp.

Marquette, W. M. 1976. Bowhead whale field studies in Alaska, 1975. Mar. Fish. Rev. 38(8):9-17.

. 1977. The 1976 catch of bowhead whales (<u>Balaena</u> <u>mysticetus</u>) by Alaskan Eskimos, with a review of the fishery, 1973-1976, and a biological summary of the species. Processed Rep. Natl. Mar. Fish. Serv., Seattle, Wash. 80pp.

______. 1979. The 1977 catch of bowhead whales (<u>Balaena</u> <u>mysticetus</u>) by Alaskan Eskimos. Rep. Int. Whal. Comm. 29:281-289, SC/30/Doc. 35.

Murdoch, J. 1885. Report of the international polar expedition to Point Barrow, Alaska. U.S. Gov. Printing Off., Washington, D.C.

- Nelson, E. W. 1887. Report upon natural history collections made in Alaska between the years 1887 and 1881. Arctic Series Publ. No. 3, Signal Service, U.S. Army, Washington, D.C.
- Nelson, R. R. 1969. Hunters of the northern ice. Univ. of Chicago Press, Chicago, Illinois. 429pp.
- Pereyra, W. T., J. E. Reeves, and R. G. Bakkala. 1976. Demersal fish and shellfish resources of the eastern Bering Sea in the baseline year 1975. Northwest and Alaska Fisheries Ctr. Processed Report, NMFS, NOAA, Seattle, Wash.
- Porsild, M. P. 1918. On "Savssats": a crowding of arctic animals of holes in the sea ice. Geogr. Rev. 6:215-228.
- Ray, D. J. 1964. Nineteenth century settlement and subsistence patterns in Bering Strait. Arctic Anthropology 2(2):61-94.

. 1975. The Eskimos of Bering Strait, 1650-1898. Univ. of Washington Press, Seattle, Wash. 305pp.

Reeves, R. R., and S. K. Katona. 1980. Extra-limital records of white whales (<u>Delphinapterus leucas</u>) in eastern North American waters. Can. Field-Nat. 94:239-247.

- Saario, D. J., and B. Kessel. 1966. Human ecological investigations at Kivalina. Pages 969-1039 in N. J. Wilimovsky and J. N. Wolfe, eds. Environment of the Cape Thompson Region, Alaska. U.S. Atomic Energy Comm., Oak Ridge, Tenn.
- Scheffer, V. B., and J. W. Slipp. 1948. The whales and dolphins of Washington State with a key to the cetaceans of the west coast of North America. Am. Midl. Nat. 39:257-337.
- Seaman, G. A., and J. J. Burns. 1981. Preliminary results of recent studies of belukhas in Alaskan waters. Rep. Int. Whal. Comm. 31:567-574, SC/32/SM 13.

_____, L. F. Lowry, and K. J. Frost. 1982. Foods of belukha whales (<u>Delphinapterus leucas</u>) in western Alaska. Cetology 44:1-19.

Sergeant, D. E. 1973. Biology of white whales (<u>Delphinapterus leucas</u>) in western Hudson Bay. J. Fish. Res. Board Can. 30:1065-1090.

_____, and P. F. Brodie. 1969. Body size in white whales, Delphinapterus leucas. J. Fish. Res. Board Can. 26:2561-2580.

_____, and _____. 1975. Identity, abundance, and present status of populations of white whales, Delphinapterus leucas, in North America. J. Fish. Res. Board Can. 32:1047-1054.

______, and W. Hoek. 1974. Seasonal distribution of bowhead and white whales in the eastern Beaufort Sea. Pages 705-719 <u>in</u> J. C. Reed and J. E. Sater, eds. The coast and shelf of the Beaufort Sea. Arctic Inst. North Am., Arlington, Va.

Thornton, H. R. 1931. Among the Eskimos of Wales, Alaska: 1890-93. The Johns Hopkins Press, Baltimore, Md. 235pp.

Tomilin, A. G. 1957. Cetacea. <u>In</u> V. G. Heptner, ed. Mammals of the USSR and adjacent countries. (Translated from Russian by Israel Program for Scientific Translations, 1967.)

Van Valin, W. B. 1941. Eskimoland speaks. The Caxton Printers, Ltd., Caldwell, Idaho. 242pp.



Figure 1.

Current world distribution of belukha whales, not including extralimital occurrences.



.


















the text.













Banks Island CHUKCHI BEAUFORT SEA SEA Point Barrow Cape Halkett Jones Islands Prudhoe Bay Barter Island Herschel Island ALASKA Mackenzie River CANADA Figure 18. Map of the Diapir Field shaving locations mentioned in the text.