Alaska Department of Fish and Game Division of Wildlife Conservation

> Federal Aid in Wildlife Restoration Research Final Report Survey-Inventory Activities 24 April 1997-23 April 1998

Abundance, Timing, and Demography of Neotropical Migratory Birds During Migration

John M Wright



James L Davis

Grant SE-3-5 March 1998

STATE OF ALASKA Tony Knowles, Governor

DEPARTMENT OF FISH AND GAME Frank Rue, Commissioner

DIVISION OF WILDLIFE CONSERVATION Wayne L. Regelin, Director

Persons intending to cite this material should receive permission from the author(s) and/or the Alaska Department of Fish and Game. Because most reports deal with preliminary results of continuing studies, conclusions are tentative and should be identified as such. Please give authors credit.

Free copies of this report and other Division of Wildlife Conservation publications are available to the public. Please direct requests to our publications specialist.

> Mary Hicks Publications Specialist ADF&G, Wildlife Conservation P.O. Box 25526 Juneau, AK 99802 (907) 465-4190

The Alaska Department of Fish and Game administers all programs and activities free from discrimination on the basis of race, religion, color, national origin, age, sex, marital status, pregnancy, parenthood, or disability. For information on alternative formats for this and other department publications, please contact the department ADA Coordinator at (voice) 907-465-4120, (TDD) 1-800-478-3648, or FAX 907-586-6595. Any person who believes she/he has been discriminated against should write to ADF&G, PO Box 25526, Juneau, AK 99802-5526 or O.E.O., U.S. Department of the Interior, Washington DC 20240.

RESEARCH FINAL REPORT

STATE:	Alaska GRANT NO.: SE-3-5	
COOPERATOR:	Alaska Bird Observatory	
STUDY TITLE:	Abundance, Timing, and Demography of Neotropcial Migratory Bird During Migration	ls
AUTHOR:	John M Wright	
PERIOD:	24 April 1997–23 April 1998	

SUMMARY

In 1997 monitoring of migratory landbirds continued for the sixth year at the Creamer's Refuge migration station in a cooperative project with the Alaska Bird Observatory. Mist nets were used to capture birds on 42 days between 25 April and 14 June and 66 days between 15 July and 30 September. A new standard array of nets was established in 1997 using a subset of previous years' nets to reduce variations in net effort. Comparisons of capture rates in the original and reduced set with data from fall 1994–1996 found no significant differences. Spring 1997 capture rates (4.3 birds/100 net hours; 354 birds captured) were low, comparable to 1993, the previous low year. Fall 1997 capture rates (33.4 birds/100 net hr; 4890 birds captured) were the second highest recorded since netting began in 1992. Most birds captured (83%–93% of individuals and 71%–81% of species) in both spring and fall were neotropical migrants. In spring the most commonly captured species were the yellow-rumped (myrtle) warbler, American robin, dark-eyed (slate-colored) junco, Swainson's thrush, northern waterthrush and orange-crowned warbler. In fall, the yellow-rumped (myrtle) warbler, American tree sparrow, orange-crowned warbler, dark-eyed (slate-colored) junco, Lincoln's sparrow, ruby-crowned kinglet, yellow warbler, Swainson's thrush and Wilson's warbler were most common. The proportion of juveniles (hatch-year birds) in the fall captures was 0.89, slightly higher than average.

Key words: Alaska, migration monitoring, mist netting, neotropical migratory birds.

i

SUMMARY	i
BACKGROUND	1
OBJECTIVE	2
STUDY AREA	2
METHODS	2
RESULTS	2
Spring 1997	2
Fall 1997	3
EDUCATION	4
CONCLUSIONS AND RECOMMENDATIONS	4
ACKNOWLEDGMENTS	4
LITERATURE CITED	5
Table 1 Mist netting effort at Creamer's Refuge migration station, 1992–1997	6
Table 2 Numbers of birds captured at Creamer's Refuge migration station, 1997	7
Table 3 Capture rates (birds/100 net hr) of common species and total for all species,	
Creamer's Refuge migration station, spring 1997	.9
Table 4 Capture rates (birds/100 net hr) of common species and total for all species,	
Creamer's Refuge migration station, fall 1997	10
Table 5 Proportion of juvenile (HY) birds in total captures of common species	
and for all individuals, Creamer's Refuge migration station, fall 1997	11
Table 6 Difference in fall capture rates using all nets and reduced set of 30 nets	
(used in 1997)	12
Table 7 Educational programs and visitors, Creamer's Refuge banding station, 1997	12

CONTENTS

BACKGROUND

Declines in populations of migrant landbirds have been well documented in North America. Long distance migrants, including flycatchers, thrushes, and wood warblers that breed in North America and winter in neotropical Central and South America, are among the species of concern (Askins et al. 1990; Sauer and Droege 1992). Alaska is an important part of the breeding range of several species of boreal forest landbirds known to be declining in other portions of their North American breeding range.

A concerted international conservation effort, *Partners In Flight* Neotropical Migratory Bird Conservation Program, was developed in the early 1990s to address this problem. In Alaska, the *Boreal Partners in Flight* working group provides coordination and direction for local projects so that a comprehensive statewide monitoring and research program can be formed from the individual efforts of the many partners.

The Creamer's Refuge migration station is operated by the Alaska Bird Observatory in cooperation with ADF&G and a host of other contributors. The migration station is part of the statewide monitoring network, providing information on abundance and productivity on a broad scale to complement Breeding Bird Survey, off-road point counts, and MAPS

(Monitoring Avian Productivity and Survival) studies. Approximately 150 migration monitoring sites are located throughout North America. Creamer's is the northernmost migration monitoring station and the longest-running station in Alaska.

OBJECTIVE

• Gather information on the abundance and productivity of migratory songbirds as part of a long-term monitoring project in the boreal forest of central Alaska.

STUDY AREA

The migration station is located on Creamer's Field Migratory Waterfowl Refuge in Fairbanks and encompasses about 15 ha of boreal forest, shrub, and wetland habitats.

METHODS

An array of mist nets was used to capture birds. Nets were opened about 7 hours each day, from sunrise to early afternoon, weather permitting. The spring 1997 netting season extended from 25 April to 14 June, and fall season 15 July to 30 September. In the first 5 years of this project, 4 to 51 nets were operated daily at the station. In 1997 a standard array of mist nets was established using a subset of previous years' nets so that all nets could be operated consistently, regardless of bird abundance or number of staff available. This reduced set of nets will be the standard array operated in future years. To examine the effect this change in nets might have on capture rates, we used data collected in fall 1994 to 1996 to compare total captures in all nets used those years to the number caught in the reduced set of nets in use those same years. Birds were banded with standard aluminum leg bands, and we collected data on age, sex, wing chord, tail length, fat index, breeding condition, and molt.

We also provided public education programs at the banding station. A 30–45 minute banding demonstration with information about bird migration, biology, and conservation was provided to 2–3 classes of local school children each school day in May and September. Structured programs were also provided to groups from summer science camps and tours. We supplied scheduled visitors with activity guides and individual booklets before their visit to the banding station. Impromptu presentations were provided to unscheduled drop-in visitors.

RESULTS

Spring 1997

Twenty-six nets were operated on 42 days in spring 1997 for a total of 7548 net hours, down 36% from the average netting effort of the previous 4 years (Table 1). Three hundred fifty-four individual birds were captured, representing 27 species (Table 2). This was the lowest absolute number of birds captured in the 6 years of spring netting, but the capture rate of 4.3 birds per 100 net hours was equivalent to 1993 and only slightly lower than 1995 and 1996 (Table 3). As in previous springs, most species and individuals captured were neotropical migrants; 81% of the species were Type A (44%; species wintering primarily south of the US/Mexico border)

or Type B (37%; species with some populations wintering south of the US/Mexico border). Individuals captured comprised 93% Type A (38%) or Type B (55%).

Among the 16 most commonly captured species, the American robin was captured in spring 1997 at the highest rates ever recorded for the species in the 6 years of netting, and the graycheeked thrush was captured at the lowest rates (Table 3). When 1997 capture rates for these 16 species are compared with the average rate of capture for the first 5 years (1992–1996), 9 species were captured less frequently than the previous 5-year average and just 4 were captured at rates greater than the average. Three species were captured at rates in spring 1997 nearly equal to their 5-year average.

FALL 1997

In fall 1997 nets were operated for 66 days for a total of 14,617 net hours. The reduction in nets to 30 from a maximum of 51 in prior years was not accompanied by a decline in net hours because in previous years it was common to open only a portion of the total number of nets on a given day and nets were usually operated for approximately 10 fewer days than in 1997 (Table 1). A total of 4890 individual birds were captured in fall 1997, including 35 species. The overall capture rate of 33.4 birds/100 net hours was the second highest recorded in the 6 years of fall netting, exceeding the average rate (27.7) for the first 5 years. Most (71%) species captured were Type A (40%) or Type B (31%) neotropical migrants, as were most (83%) individuals (Type A, 38%; Type B, 45%).

Among the 19 commonly captured species, 4 (alder flycatcher, Swainson's thrush, American robin, and fox sparrow) were captured at the highest rates recorded in the 6 years of fall netting, and no species set a new minimum record (Table 4). Thirteen species were caught at higher rates than their 1992–1996 average, 3 were captured at lower rates than average, and 3 were captured at average rates.

In fall 1997, 89% of birds captured were young of the year (HY, hatch year) (Table 5). This falls within the range (84–90%) observed in 4 of 5 previous years, excluding the exceptionally low (75%) year of 1993. The proportion of young in 3 of the most commonly captured species, Wilson's warbler, American tree sparrow, and white-crowned (Gambell's) sparrow was higher in 1997 than in any prior year, while a new low was recorded for only 1 species, the Fox sparrow.

The effect of reducing the overall number of nets was examined using data from fall 1994 to 1996 by comparing the capture rates using all nets with capture rates in the reduced set of 30 nets now used as the standard array (Table 6). The absolute difference in capture rates between the full and reduced arrays for the 19 most common species averaged 8% to 9% for the 3 years. There was no significant difference between capture rates using the full and reduced set of nets (paired-sample *t*-tests; 1994, t = -1.49, df = 18, P > 0.15; 1995, t = -1.85, df = 18, P > 0.05; 1996, t = -0.91, df = 18. P > 0.35). Only 3 of 19 species had average absolute differences in capture rate greater than 10%. The blackpoll warbler and Wilson's warbler were the only species captured at rates averaging >10% higher in the reduced set, and only the Savannah sparrow was captured an average of >10% less in the reduced set of nets. Overall, 7

species were captured at higher rates (\bar{x} difference = +10%) in the reduced set of nets all 3 years, while 3 species were consistently caught at lower rates (\bar{x} difference = -8%) and capture rates for 9 species were not consistently higher or lower (\bar{x} absolute difference = 7%).

EDUCATION

The number of participants in banding demonstrations more than doubled from 1996 to 1997. In 1997, 95 groups with 2237 individuals (primarily school classes) attended educational programs at Creamer's Refuge (Table 7). More than 1900 activity booklets were distributed to these groups prior to their arrival at the station. Another 400 unscheduled individuals received informal presentations at the banding station.

CONCLUSIONS AND RECOMMENDATIONS

This year was the seventh year of the long-term monitoring effort at Creamer's Refuge. This cooperative project conducted by the Alaska Bird Observatory has proven capable of sustaining the consistent effort necessary to maintain a project requiring 15–20 years or more of data collection. Recent analyses and reviews have confirmed the usefulness of standardized netting stations for study of the abundance and productivity of migratory bird populations (Dunn and Hussell 1995; Johnson and Geupel 1996; Dunn et al. 1997). In addition to its value in monitoring species of concern, this project is also gathering important information on the timing of migration, reproduction, molt, juvenile dispersal, changes in body condition, and other life history events and has become a significant educational opportunity for Fairbanks school children and others.

ACKNOWLEDGMENTS

This project was supported by Endangered Species funding through Federal Aid in Wildlife Restoration. Funding was also provided by the Alaska Bird Observatory and its members, US Fish and Wildlife Service (Cost Share Challenge Grant), ARCO Alaska, Exxon Company USA, Bureau of Land Management, Alaska Conservation Foundation, MAPCO Alaska Petroleum, Alyeska Pipeline Service Company, ABR, Inc., and Fairbanks Curling Lions Club.

The dedicated board of directors and staff of the Alaska Bird Observatory are responsible for the success of this long-term project. I would especially like to thank Lori Quakenbush, Board President; Anna-Marie Barber, Chief Biologist; Nancy DeWitt, Executive Director; Steve Springer, Banding Biologist; and Sara McDaniel, Education Coordinator for the key roles they played in this effort. Thanks also to the interns and volunteers who regularly got up before dawn to run the nets. Mark Ross, ADF&G, assisted with materials for the banding demonstrations and developed complementary educational programs for visiting school children.

LITERATURE CITED

- ASKINS RA, JF LYNCH, AND R GREENBERG. 1990. Population declines in migratory birds in Eastern North America. Pages 1–57 in DM Power, editor. Current Ornithology Volume 7, Plenum Press, New York.
- DUNN EH AND DT HUSSELL. 1995. Using migration counts to monitor landbird populations; review and evaluation of current methods. Pages 43-88 *in* DM Power, editor. Current Ornithology Volume 12, Plenum Press, New York.
- DUNN EH, DT HUSSELL, AND RJ ADAMS. 1997. Monitoring songbird population change with autumn mist netting. *Journal of Wildlife Management* 61: 389–396.
- JOHNSON MD AND GR GUEPEL. 1996. The importance of productivity to the dynamics of a Swainson's Thrush population. *Condor* 98: 133–141
- SAUER JR AND S DROEGE. 1992. Geographic patterns in population trends of neotropical migrants in North America. Pages 26-42 *in* JM Hagan and DW Johnston, editors. Ecology and Conservation of Neotropical Migrant Landbirds, Smithsonian Institution Press, Washington, DC.

PREPARED BY:

John M Wright Wildlife Biologist III

SUBMITTED BY:

Kenneth R Whitten Research Coordinator **APPROVED BY:**

Wayne L Regelin, Director

Division of Wildlife Conservation

Steven R Peterson, Senior Staff Biologist Division of Wildlife Conservation

1992-1997
uge migration station,
at Creamer's Refu
Mist netting effort
Table 1

				IC	ar		
Season	Description	1992	1993	1994	1995	1996	1997
Spring	Number of days nets open	42	43	41	45	44	42
	Net hours	6903	10,552	11,252	12,731	12,411	7548
	Number of nets per day	16–29	16–33	27-47	36-47	33-45	26
	(range)						
Fall	Number of days nets open	46	53	52	58	57	99
	Net hours	5890	13,711	13,934	14,156	14,985	14,617
	Number of nets per day	4-35	11–47	21–51	16-49	18-49	30
	(range)						

Species	Winter ^a	Spring 1997	Fall 1997
Sharp-shinned Hawk (Accipter striatus)	B	1	4
American Kestrel (Falco sparverius)	N		1
Lesser Yellowlegs (Tringa flavipes)	Α	1	
Solitary Sandpiper (Tringa solitaria)	Α	4	
Downy Woodpecker (Picoides pubescens)	R	1	1
Hairy Woodpecker (Picoides villosus)	R ·		1
Northern Flicker (Colaptes auratus)	В	3	2
Western Wood-Pewee (Contopus	Α		3
Yellow-bellied Flycatcher (Empidonax	Α		2
flaviventris)			
Alder Flycatcher (Empidonax alnorum)	Α	3	126
Hammond's Flycatcher (<i>Empidonax</i> hammondi)	Α	16	74
Gray Jay (Perisoreus canadensis)	R	1 .	
Tree Swallow (<i>Tachycineta hicolor</i>)	B	-	1
Cliff Swallow (Hirundo pyrrhonota)	Ā		1
Black-capped Chickadee (Poecile	R	5	35
atricapillus)		e e	
Boreal Chickadee (Poecile hudsonicus)	R		6
Ruby-crowned Kinglet (Regulus calendula)	В	3	197
Arctic Warbler (Phylloscopus sibilatrix)	Р		5
Gray-cheeked Thrush (Catharus minimus)	Α	5	41
Swainson's Thrush (Catharus ustulatus)	Α	28	188
Hermit Thrush (Catharus guttatus)	В	1	18
American Robin (Turdus migratorius)	В	43	39
Varied Thrush (Ixoreus naevius)	Ν		6
Orange-crowned Warbler (Vermivora celata)	Α	20	646
Yellow Warbler (Dendroica petechia)	Δ	10	180
Yellow-rumped Warbler (Dendroica	B	74	1234
Townsend's Warbler (Dendroica	Α		10
townsendi)			100
Blackpoll Warbler (Dendroica striata)	A	3	102
Northern Waterthrush (Seiurus	Α	25	49
Wilson's Warbler (Wilsonia pussilla)	А	13	186
American Tree Sparrow (Spizella arborea)	N	1	763
Savannah Sparrow (Passerculus	B	17	106
sandwichensis)		±1	100
Fox Sparrow (Passerella iliaca)	В	2	66
Lincoln's Sparrow (Melospiza lincolnii)	Α	8	246

Table 2 Numbers of birds captured at Creamer's Refuge migration station, 1997

Table 2 Continued

Species	Winter ^a	Spring 1997	Fall 1997
White-crowned Sparrow (Zonotrichia	В	13	70
leucophrys)			
Dark-eyed Junco (Junco hyemalis)	В	37	466
Rusty Blackbird (Euphagus carolinus)	Ν		1
Common Redpoll (Carduelis flammea)	Ν	15	5
Totals		354	4890
Dates nets open	•	25 Apr–14 Jun	15 Jul-30 Sep
Number of days nets operated		42	66
Number of net hours		7548	14,617

^a Winter range/migration: A = primarily south of US/Mexico border; B = some populations south of US/Mexico border; P = Pacific/Eurasia/Africa; N = North America; R = Resident.

Species	1992	1993	1994	1995	1996	1997
Hammond's Flycatcher	0.17	0.12	0.14	0.17	0.06	0.12
Ruby-crowned Kinglet	0.29	0.01	0.04	0.05	0.06	0.04
Gray-cheeked Thrush	0.23	0.13	0.19	0.09	0.18	0.07
Swainson's Thrush	0.41	0.45	0.49	0.38	0.63	0.33
American Robin	0.45	0.34	0.28	0.35	0.29	0.50
Orange-crowned Warbler	0.58	0.47	0.40	0.36	0.24	0.25
Yellow Warbler	0.67	0.26	0.36	0.06	0.17	0.13
Yellow-rumped Warbler	1.58	0.93	0.48	0.70	0.65	0.94
Northern Waterthrush	0.33	0.13	0.58	0.19	0.44	0.32
Wilson's Warbler	0.48	0.51	0.51	0.46	0.14	0.17
American Tree Sparrow	0.51	0.01	0.08	0.07	0.19	0.01
Savannah Sparrow	0.83	0.14	0.29	0.25	0.42	0.20
Lincoln's Sparrow	0.09	0.05	0.11	0.16	0.10	0.11
White-crowned Sparrow	0.20	0.08	0.16	0.08	0.07	0.17
Dark-eyed Junco	0.42	0.09	0.21	0.29	0.73	0.45
Common Redpoll	1.17	0.18	1.99	0.68	0.08	0.20
Total (all species)	10.46	4.27	6.91	4.63	4.87	4.29

Table 3 Capture rates (birds/100 net hr) of common species and total for all species, Creamer's Refuge migration station, spring 1997

Species	1992	1993	1994	1995	1996	1997
Alder Flycatcher	0.58	0.55	0.44	0.47	0.48	0.86
Hammond's Flycatcher	0.14	0.10	0.29	0.35	0.59	0.50
Black-capped Chickadee	0.58	0.18	0.21	0.48	0.31	0.23
Ruby-crowned Kinglet	0.88	0.65	0.91	0.87	1.56	1.36
Gray-cheeked Thrush	0.63	0.18	0.20	0.35	0.11	0.28
Swainson's Thrush	1.06	0.67	0.73	1.24	0.59	1.29
American Robin	2.73	0.15	0.27	0.24	0.24	0.27
Orange-crowned Warbler	7.35	1.71	4.23	6.12	3.06	4.41
Yellow Warbler	2.17	0.43	0.64	1.38	0.76	1.29
Yellow-rumped Warbler	13.40	0.41	7.18	4.15	7.77	8.44
Blackpoll Warbler	1.38	0.15	0.35	0.36	0.33	0.70
Northern Waterthrush	0.69	0.22	0.12	0.20	0.22	0.36
Wilson's Warbler	2.15	0.79	0.83	1.19	0.50	1.27
American Tree Sparrow	1.81	2.88	2.46	4.66	1.83	5.22
Savannah Sparrow	0.75	0.52	0.20	1.23	0.55	0.72
Fox Sparrow	0.26	0.23	0.30	0.23	0.25	0.45
Lincoln's Sparrow	2.73	1.09	1.27	2.56	1.56	1.68
White-crowned Sparrow	0.69	0.31	0.43	0.30	0.80	0.48
Dark-eyed Junco	2.33	1.18	3.84	2.92	3.86	3.17
Total (all species)	43.16	12.58	25.32	30.78	26.73	33.41

đ

ð

Table 4 Capture rates (birds/100 net hr) of common species and total for all species, Creamer's Refuge migration station, fall 1997

Species	1992	1993	1994	1995	1996	1997
Alder Flycatcher	0.89	0.61	0.76	0.64	0.72	0.68
Hammond's Flycatcher	0.89	0.85	0.76	0.96	0.89	0.90
Black-capped Chickadee	0.94	0.75	0.69	0.90	0.93	0.94
Ruby-crowned Kinglet	0.82	0.90	0.89	0.91	0.87	0.87
Gray-cheeked Thrush	0.79	0.76	0.75	0.78	0.88	0.76
Swainson's Thrush	0.82	0.79	0.86	0.91	0.96	0.90
American Robin	0.42	0.50	0.89	0.76	0.86	0.85
Orange-crowned Warbler	0.83	0.63	0.86	0.86	0.84	0.79
Yellow Warbler	0.91	0.58	0.88	0.84	0.94	0.94
Yellow-rumped Warbler	0.91	0.75	0.94	0.91	0.95	0.94
Blackpoll Warbler	0.94	0.71	0.86	0.94	0.96	0.92
Northern Waterthrush	0.93	0.87	0.94	0.93	0.97	0.90
Wilson's Warbler	0.90	0.67	0.84	0.89	0.85	0.96
American Tree Sparrow	0.67	0.72	0.87	0.81	0.85	0.90
Savannah Sparrow	0.87	0.92	0.96	0.96	0.98	0.92
Fox Sparrow	0.69	0.77	0.79	0.81	0.78	0.68
Lincoln's Sparrow	0.92	0.79	0.93	0.95	0.97	0.93
White-crowned Sparrow	0.86	0.90	0.90	0.91	0.93	0.94
Dark-eyed Junco	0.84	0.88	0.96	0.90	0.91	0.89
Total (all individuals captured)	0.84	0.75	0.90	0.87	0.90	0.89

Table 5 Proportion of juvenile (HY) birds in total captures of common species and for all individuals, Creamer's Refuge migration station, fall 1997

	Difference				
Species	1994	1995	1996		
Alder Flycatcher	-0.14	-0.13	0.00		
Hammond's Flycatcher	-0.10	-0.03	-0.03		
Black-capped Chickadee	-0.10	0.04	-0.16		
Ruby-crowned Kinglet	0.04	0.15	0.11		
Gray-cheeked Thrush	0.10	0.06	0.00		
Swainson's Thrush	0.01	0.02	-0.17		
American Robin	0.00	0.04	-0.04		
Orange-crowned Warbler	0.08	0.10	0.06		
Yellow Warbler	0.03	-0.04	0.13		
Yellow-rumped Warbler	0.05	0.07	0.13		
Blackpoll Warbler	0.17	0.19	0.18		
Northern Waterthrush	-0.08	-0.10	0.00		
Wilson's Warbler	0.16	0.16	0.08		
American Tree Sparrow	0.03	0.12	-0.15		
Savannah Sparrow	-0.25	-0.11	-0.11		
Fox Sparrow	0.03	0.04	0.08		
Lincoln's Sparrow	0.09	0.07	0.08		
White-crowned Sparrow	0.07	-0.17	0.01		
Dark-eyed Junco	-0.05	-0.04	-0.04		
\overline{x} Absolute Difference	0.08	0.09	0.08		
Paired-sampled t-test	<i>P</i> >0.15	<i>P</i> >0.05	<i>P</i> >0.35		

\$

9

Table 6 Difference in fall capture rates using all nets and reduced set of 30 nets (used in 1997)

Table 7 Educational programs and visitors, Creamer's Refuge banding station, 1997

Season	Number of groups	Number of individuals in groups	Number of drop-in visitors
Spring	39	939	
Summer	17	294	400
Fall	39	1004	
Totals	95	2237	400

Alaska's Game Management Units



The Federal Aid in Wildlife Restoration Program consists of funds from a 10% to 11% manufacturer's excise tax collected from the sales of handguns, sporting rifles, shotguns, ammunition, and archery equipment. The Federal Aid program allots funds back to states through a formula based on each state's geographic area and number of paid hunting license holders. Alaska receives a maximum 5% of revenues collected each year. The Alaska Department of Fish and Game uses federal aid funds to help restore, conserve, and manage wild birds and mammals to benefit the

public. These funds are also used to educate hunters to develop the skills, knowledge, and attitudes for responsible hunting. Seventy-five percent of the funds for this report are from Federal Aid.



James L Davis