

*Gulf of Alaska - Marine Mammals*

**Organochlorine, Pesticides and Polybrominated Diphenyl Ether Contaminant Concentrations in Multiple Tissue Matrices of Live Steller Sea Lions (*Eumetopias jubatus*) in Alaska**

**Kimberlee B. Beckmen**, Alaska Department of Fish & Game, kimberlee.beckmen@alaska.gov  
Kathy Burek, Alaska Veterinary Pathology Services, fnkab1@uaf.edu  
Ken Pitcher, Alaska Department of Fish & Game, kwpitcher@yahoo.com  
Gina Ylitalo, Northwest Fisheries Science Center, National Marine Fisheries Service,  
gina.ylitalo@noaa.gov  
Brian Fadely, Alaska Fisheries Science Center, National Marine Fisheries Service,  
brian.fadely@noaa.gov

Blood, blubber, milk, and feces were collected from 53 free-ranging and 3 captive Steller sea lions (*Eumetopias jubatus*) in Alaska from 1998-2003 to assess exposure to selected organochlorine (OC) contaminants (e.g., dioxin-like PCBs, DDTs). Organochlorine contaminant relationships among multiple matrices of individuals were used to determine the appropriate tissue for exposure monitoring in live animals and thus minimize invasive sampling techniques. Concentrations of certain OC contaminants in blubber, milk and blood were highly correlated within individuals; however fecal concentrations were only correlated with those measured in blood. Thus a blood sample may be the best alternative as a less-invasive indicator of relative contaminant exposure in lieu of surgical blubber biopsy while feces may be used as a non-invasive monitoring tool of relative OC exposure without direct handling of animals. Regional OC contaminant exposure was compared in blubber samples of pups through sub-adults of the stable eastern stock in Southeast Alaska (n=48) as compared to the endangered western stock of the Gulf of Alaska (n=55) and Aleutian Islands (n=43). Pesticides and polybrominated diphenyl ethers were detected in 25 and 15 animals respectively, including 4 individuals that were sampled at 5 month intervals. Transplacental transfer of OCs was extremely low. Concentrations of OCs peaked in pups sampled between 2 - 6 weeks of age, declined by midway through the suckling period, and increased again through the first year of the presumed dependent suckling period though the weaning period. These data suggests that exposure to OCs is at a level of concern especially in young pups in portions of the range of the endangered western stock of Steller sea lions.



# Alaska Marine Science SYMPOSIUM 2008

## **Book of Abstracts for Oral Presentations and Posters**

**January 20-23, 2008**

**Hotel Captain Cook, Anchorage, Alaska**

### **Sponsoring Organizations**

Alaska Department of Fish and Game

Alaska Fisheries Science Center

Alaska Ocean Observing System

Alaska Pacific University

Alaska Sea Grant

Alaska SeaLife Center

Alliance for Coastal Technologies

Exxon Valdez Oil Spill Trustee Council

Kachemak Bay Research Reserve

Minerals Management Service

National Ocean Service

National Park Service

North Pacific Fishery Management Council

North Pacific Research Board

North Slope Science Initiative

Oceans Alaska Science and Learning Center

Oil Spill Recovery Institute

Pollock Conservation Cooperative Research Center

Prince William Sound Science Center

University of Alaska Fairbanks

US Arctic Research Commission

USGS Alaska Science Center

[www.alaskamarinescience.org](http://www.alaskamarinescience.org)