

Tracking the movements of surf and white-winged scoters with satellite telemetry.

*DANIEL H. ROSENBERG & MICHAEL J. PETRULA, Alaska Dept. Fish & Game, Anchorage. We studied the distribution, seasonal movements, and migration patterns of surf scoters (*Melanitta perspicillata*) and white-winged scoters (*M. fusca*) with satellite telemetry. Using floating mist-nets, we captured ducks in Prince William Sound, Alaska from 1998-2000. A veterinarian surgically implanted satellite transmitters in a total of 43 surf scoters and 23 white-winged scoters. Post-surgical mortality rates in 1998 and 1999 were 52%. In 2000, we focused efforts on identifying the causes and reducing mortality of implanted birds. Movements of scoters were monitored every two to four days. For surf scoters we identified nesting areas in the Yukon and Northwest Territories of Canada, and the Yukon Flats National Wildlife Refuge, Alaska. Molting areas were identified in Norton and Kotzebue Sound and Kuskokwim Bay (mouth of the Kuskokwim River) in western Alaska. Wintering areas were identified in Prince William Sound and British Columbia. For white-winged scoters we identified nesting areas in the Northwest Territories; molting areas in the Beaufort Sea (Canada), lower Cook Inlet, and Bristol Bay, Alaska; and wintering areas in Prince William Sound and the Alaska Peninsula. We also obtained information on the timing and routes of migration.

SECOND NORTH AMERICAN DUCK SYMPOSIUM & WORKSHOP

*OCTOBER 11-15, 2000,
Saskatoon, Saskatchewan
Canada*

