

Fishery Management Report No. 17-04

**Area Management Report for the Sport Fisheries of
Anchorage, 2011–2015**

by

Jay Baumer

and

Brittany Blain

February 2016

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the *Système International d'Unités* (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H_A
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	e
hectare	ha	at	@	catch per unit effort	CPUE
kilogram	kg	compass directions:		coefficient of variation	CV
kilometer	km	east	E	common test statistics	(F, t, χ^2 , etc.)
liter	L	north	N	confidence interval	CI
meter	m	south	S	correlation coefficient (multiple)	R
milliliter	mL	west	W	correlation coefficient (simple)	r
millimeter	mm	copyright	©	covariance	cov
		corporate suffixes:		degree (angular)	$^\circ$
Weights and measures (English)		Company	Co.	degrees of freedom	df
cubic feet per second	ft ³ /s	Corporation	Corp.	expected value	E
foot	ft	Incorporated	Inc.	greater than	>
gallon	gal	Limited	Ltd.	greater than or equal to	\geq
inch	in	District of Columbia	D.C.	harvest per unit effort	HPUE
mile	mi	et alii (and others)	et al.	less than	<
nautical mile	nmi	et cetera (and so forth)	etc.	less than or equal to	\leq
ounce	oz	exempli gratia (for example)	e.g.	logarithm (natural)	ln
pound	lb	Federal Information Code	FIC	logarithm (base 10)	log
quart	qt	id est (that is)	i.e.	logarithm (specify base)	log ₂ , etc.
yard	yd	latitude or longitude	lat or long	minute (angular)	'
		monetary symbols (U.S.)	\$, ¢	not significant	NS
Time and temperature		months (tables and figures): first three letters	Jan, ..., Dec	null hypothesis	H_0
day	d	registered trademark	®	percent	%
degrees Celsius	°C	trademark	™	probability	P
degrees Fahrenheit	°F	United States (adjective)	U.S.	probability of a type I error (rejection of the null hypothesis when true)	α
degrees kelvin	K	United States of America (noun)	USA	probability of a type II error (acceptance of the null hypothesis when false)	β
hour	h	U.S.C.	United States Code	second (angular)	"
minute	min	U.S. state	use two-letter abbreviations (e.g., AK, WA)	standard deviation	SD
second	s			standard error	SE
Physics and chemistry				variance	
all atomic symbols				population	Var
alternating current	AC			sample	var
ampere	A				
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

FISHERY MANAGEMENT REPORT NO. 17-04

**AREA MANAGEMENT REPORT FOR THE SPORT FISHERIES OF
ANCHORAGE, 2011–2015**

by
Jay Baumer
and
Brittany Blain

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1565

February 2017

The Fishery Management Reports series was established in 1989 by the Division of Sport Fish for the publication of an overview of management activities and goals in a specific geographic area, and became a joint divisional series in 2004 with the Division of Commercial Fisheries. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.adfg.alaska.gov/sf/publications/>. This publication has undergone regional peer review.

*Jay Baumer,
Alaska Department of Fish and Game, Division of Sport Fish,
333 Raspberry Road, Anchorage, AK 99518-1599, USA*

and

*Brittany Blain,
Alaska Department of Fish and Game, Division of Sport Fish,
333 Raspberry Road, Anchorage, AK 99518-1599, USA*

This document should be cited as follows:

Baumer, J., and B. Blain. 2016. Area management report for the sport fisheries of Anchorage, 2011–2015. Alaska Department of Fish and Game, Fishery Management Report No. 17-04, Anchorage.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	iii
LIST OF FIGURES.....	iv
LIST OF APPENDICES.....	iv
ABSTRACT.....	1
INTRODUCTION.....	1
AREAWIDE OVERVIEW.....	3
Knik Arm Streams.....	3
Chester Creek.....	3
Eagle River.....	3
Eklutna River.....	4
Fish Creek.....	4
Ship Creek.....	4
Sixmile Creek.....	5
Peters Creek.....	5
Turnagain Arm Streams.....	5
Bird Creek.....	6
California and Glacier Creeks.....	6
Campbell Creek.....	6
Indian Creek.....	7
Ingram Creek.....	7
Rabbit Creek.....	7
Twentymile River.....	7
Placer River.....	8
Portage River.....	8
Anchorage Lakes.....	8
Stocking Program.....	10
Fisheries Management and Objectives.....	11
Fisheries Performance.....	11
CHINOOK SALMON FISHERIES.....	15
Fishery Description.....	15
Campbell Creek.....	16
Eagle River.....	16
Ship Creek.....	17
Fishery Management and Objectives.....	20
Campbell Creek.....	20
Eagle River.....	21
Ship Creek.....	21
Stocking Program.....	21
Ship Creek.....	21
Eagle River.....	21
Fishery Performance.....	21
Campbell Creek.....	22
Eagle River.....	22
Ship Creek.....	22

TABLE OF CONTENTS (Continued)

	Page
CHUM SALMON FISHERIES.....	23
Fishery Description.....	23
Fishery Management and Objectives.....	24
Stocking Program	24
Fishery Performance.....	25
COHO SALMON FISHERIES	25
Fishery Description.....	25
Bird Creek.....	27
Campbell Creek	27
Ship Creek	28
Turnagain Arm Streams.....	29
Fishery Management and Objectives.....	29
Campbell Creek	29
Ship Creek	30
Turnagain Arm Streams.....	30
Stocking Program	30
Bird Creek.....	31
Campbell Creek	32
Ship Creek	32
Turnagain Arm Streams.....	32
Fishery Performance.....	32
Bird Creek.....	33
Campbell Creek	34
Ship Creek	35
Turnagain Arm Streams.....	35
PINK SALMON FISHERIES	36
Fishery Description.....	36
Fishery Management and Objectives.....	38
Stocking Program	38
Fishery Performance.....	38
SOCKEYE SALMON FISHERY	39
Fishery Description.....	39
Fishery Management and Objectives.....	40
Stocking Program	40
Fishery Performance.....	40
DOLLY VARDEN–ARCTIC CHAR FISHERIES.....	42
Fishery Description.....	42
Fishery Management and Objectives.....	42
Stocking Program	44
Fishery Performance.....	44

TABLE OF CONTENTS (Continued)

	Page
STEELHEAD–RAINBOW TROUT FISHERIES	45
Fishery Description.....	45
Fishery Management and Objectives.....	45
Fishery Performance.....	45
LAKE FISHERIES	47
Fishery Description.....	47
Arctic Char–Dolly Varden.....	48
Arctic Grayling	48
Chinook Salmon	48
Rainbow Trout.....	49
Fishery Management and Objectives.....	49
Invasive Species.....	49
Stocking Program	52
Arctic Char–Dolly Varden.....	53
Arctic Grayling	54
Landlocked Chinook Salmon.....	54
Rainbow Trout.....	54
REFERENCES CITED	56
APPENDIX A: SPORT FISHING REGULATIONS.....	57
APPENDIX B: STOCKING IN THE ANCHORAGE MANAGEMENT AREA	63
APPENDIX C: ESCAPEMENT ESTIMATES.....	75

LIST OF TABLES

Table	Page
1 Sport fishing effort expended in the Anchorage Management Area (AMA) compared to Southcentral Alaska and statewide, 1996–2015.	12
2 Saltwater, lake, and stream sport fishing effort (angler-days), Anchorage Management Area (AMA), 1996–2015.....	13
3 Sport fish catch and harvest of anadromous salmon, Anchorage Management Area, 1996–2009.....	14
4 Angler effort for streams in the Anchorage Management Area, 1996–2015.....	18
5 Chinook salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.	19
6 Chum salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.	24
7 Coho salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.....	26
8 Pink salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.....	37
9 Sockeye salmon sport catch (C) and harvest (H), Anchorage Management Area, 1996–2015.....	41
10 Dolly Varden sport fish catch (C) and harvest (H), Anchorage Management Area, 1996–2015.	43
11 Rainbow trout sport fish catch and harvest, Anchorage Management Area, 1996–2015.....	46
12 Angler lake sport fishing effort (lakes with names starting A–G), Anchorage Management Area, 1996–2015.....	50
13 Angler lake sport fishing effort (lakes with names starting H–Z and total AMA effort), Anchorage Management Area, 1996–2015.	51
14 Lake sport fish catch (C), harvest (H), and percent released (%), Anchorage Management Area, 1996–2015.....	55

LIST OF FIGURES

Figure		Page
1	Map of the Anchorage Management Area	2
2	Anchorage Management Area streams and stocked lakes.....	9
3	Sport fish catch and harvest of anadromous salmon, Anchorage Management Area, 2006–2015.....	15
4	Map of lower section of Ship Creek.....	20
5	Chum salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.....	25
6	A map of lower section of Campbell Creek showing areas open to coho salmon sport fishing and season dates.....	28
7	Coho salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.....	31
8	Pink salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.....	38
9	Sockeye salmon sport catch and harvest, Anchorage Management Area, 2006–2015.....	42
10	Dolly Varden sport catch and harvest, Anchorage Management Area, 2006–2015.....	44
11	Rainbow trout sport fish catch and harvest, Anchorage Management Area, 2006–2015.....	47
12	Angler fishing effort for lakes in the Anchorage Management Area, 2006–2015.....	52
13	Lake sport fish catch and harvest, Anchorage Management Area, 2006–2015.....	53

LIST OF APPENDICES

Appendix		Page
A1	Sport fishing regulations for Ship Creek, 1957–2015.....	58
A2	Sport fishing regulations for Eagle River, 1957–2015.....	60
A3	Sport fishing regulations for Campbell Creek, 1957–2015.....	62
B1	Chinook salmon stocking in the Anchorage Management Area by year (1996–2015) and site.....	64
B2	Coho salmon stocking in Anchorage Management Area by year (1996–2015) and site.....	65
B3	Rainbow trout stocking in the Anchorage Management Area by year (1996–2005) and site.....	66
B4	Rainbow trout stocking in the Anchorage Management Area by year (2006–2015) and site.....	69
B5	Arctic char stocking in the Anchorage Management Area by year (1996–2015) and lake.....	72
B6	Arctic grayling stocking in the Anchorage Management Area by year (1996–2015) and site.....	73
C1	Ship Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.....	76
C2	Eagle River salmon escapement estimates, Anchorage Management Area, 1996–2015.....	77
C3	Campbell Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.....	78
C4	Bird Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.....	79
C5	Rabbit Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.....	80

ABSTRACT

This report provides a summary of sport fisheries in the Anchorage Management Area from 2011 through 2015. Included are a description and historical overview of each sport fishery, how each fishery is currently managed, and the performance and escapement for each fishery for 2011–2015.

Key words: Anchorage Management Area, Alaska Board of Fisheries, sport fisheries , stocked fish, Chinook salmon, *Oncorhynchus tshawytscha*, coho salmon, *Oncorhynchus kisutch*, sockeye salmon, *Oncorhynchus nerka*, pink salmon, *Oncorhynchus gorbuscha*, chum salmon, *Oncorhynchus keta*, Dolly Varden, *Salvelinus malma*, groundfish, sharks

INTRODUCTION

The Anchorage Sport Fish Management Area (AMA) is part of the Southcentral Sport Fish Region and consists of all waters flowing into eastside Knik Arm and northside of Turnagain Arm from the Eklutna River drainage in the north to Ingram Creek in the south (Figure 1). The AMA includes the brackish waters on the south side of Turnagain Arm and the northeast side of Kink Arm and includes part of the Chugach Mountain Range. Even though the AMA is coastal and includes Turnagain Arm, the majority of fishing effort and opportunity is in fresh water. Turnagain Arm is muddy in nature and has extreme daily tidal fluctuations making it difficult for anglers to catch fish and difficult for boats to navigate. Most streams, except Ingram Creek, are fed from high mountain lakes, snow melt, or glaciers in the Chugach Mountain Range. These streams have year-round resident sport fish species as well as anadromous salmon that return to them. Many of these streams are enhanced with hatchery fish to provide additional sport fishing opportunity. A majority of the lakes, except those located in the Chugach Mountains (Symphony and Rabbit lakes), are relatively shallow in nature and are often man-made. Most of the lakes are stocked with fish to provide sport fishing opportunity and if not stocked would contain very few, if any, sport fish species.

Local communities within the area include Anchorage, Eagle River, Chugiak, Birchwood, Peters Creek, Eklutna, Indian, Bird, Girdwood, Portage, and the Joint Base Elmendorf-Richardson (JBER). Of Alaska's 738,432 residents (2015 U. S. Census Bureau Population Estimate data, https://factfinder.census.gov/faces/nav/jsf/pages/community_facts.xhtml, accessed December 2016), 40% or 298,695 people reside in the AMA. Anchorage is commonly referred to as the hub for the rest of the state and has the Ted Stevens International Airport acting as the gateway to connect travelers to smaller in-state flights or quickly access Alaska's primary highways. Even though there are 3 lakes dedicated as float-plane lakes and numerous local small plane runways, a majority of anglers access the AMA sport fisheries by road.

AMA land managers include private individuals, Municipality of Anchorage (MOA), Alaska Railroad Corporation (ARRC), Alaska Department of Natural Resources (ADNR), U.S. Forest Service (USFS), U.S. Bureau of Land Management (BLM), U.S. Department of Defense (DOD), and Alaska Native organizations. Management and research functions for AMA sport fisheries are conducted by the Alaska Department of Fish and Game (ADF&G), Division of Sport Fish (SF) staff from the Anchorage regional office.




Figure 1.—Map of the Anchorage Management Area.

All 5 species of salmon are found in the AMA: Chinook (*Oncorhynchus tshawytscha*), chum (*O. keta*), coho (*O. kisutch*), pink (*O. gorbuscha*), and sockeye (*O. nerka*) salmon. In addition, resident and stocked species in the AMA include Arctic char (*Salvelinus alpinus*) and grayling (*Thymallus arcticus*), Dolly Varden (*Salvelinus malma*), and rainbow trout (*Oncorhynchus mykiss*). There are also unconfirmed reports of steelhead (anadromous *O. mykiss*) and confirmed reports of invasive species such as northern pike (*Esox lucius*), Alaska blackfish (*Dallia pectoralis*), and domestic goldfish (Family Cyprinidae). Data from the ADF&G Statewide Harvest Survey (SWHS) used in this report groups Dolly Varden and Arctic char as 1 fish although they are 2 separate species, and both are found in the AMA.

AREAWIDE OVERVIEW

The AMA has approximately 1,575 miles of flowing waters and 371 lakes (ADF&G GIS Staff calculated using ArcView). All of the watersheds eventually flow into 2 arms of Cook Inlet (Knik Arm and Turnagain Arm). A majority of creeks support anadromous fish and many support resident species as well. Four creeks and 30 lakes are stocked with hatchery fish to improve and provide additional sport fishing opportunities in the AMA.

KNIK ARM STREAMS

Knik Arm streams are the streams that flow into Knik Arm range from Fish Creek, near Point Woronzof located at the edge of the Ted Stevens International Airport, to the Eklutna River and include Chester Creek, Eagle River, Ship Creek, Sixmile Creek, and Peters Creek.

Chester Creek

Chester Creek, also known as *Chanshtnu*, or “Grass Creek” (Kari et. al. 2003), travels approximately 16 miles from the Chugach Mountains and flows into Knik Arm of Cook Inlet. Some of the headwaters of Chester Creek are on DOD property. The mainstem of Chester Creek is fed by a tributary from Reflection Lake as well as tributaries formed from rain and snowmelt from the mountains. After the waters of Reflection Lake and a mountain tributary combine near Lee Street, they then flow into University Lake, also known as “University Lake” or Behm Lake near the Alaska Pacific University (APU) campus. After exiting University Lake, Chester Creek flows through or on the edge of many neighborhoods for approximately 5 miles until it enters Chester Lagoon. Chester Lagoon is a man-made lagoon separated into 2 sections (Westchester Lagoon and Chester Lagoon) by the Walter J. Hickel Parkway. The water flows out of Westchester Lagoon for approximately one-tenth of a mile to the mud flats of Knik Arm. Coho and sockeye salmon, rainbow trout, and Dolly Varden (ADF&G Anadromous Waters Catalog; <https://www.adfg.alaska.gov/sf/SARR/AWC/index.cfm?ADFG=main.home>, accessed January 2017) are present at times in Chester Creek and its tributaries. It is also believed that Chinook and chum salmon and steelhead trout are also present in Chester Creek but in very low numbers and they are not well documented. Rainbow trout are also enhanced into sections of Chester Creek. To protect wild rainbow trout and give them an opportunity to spawn, the entire Chester Creek drainage is closed to sport fishing April 15 through June 14.

Eagle River

The Eagle River drainage originates in the Chugach Mountains with most of its flow contributed by Eagle Glacier and 2 lakes (Eagle Lake and Symphony Lake). The lower portion of the river flows through flats on Fort Richardson; these flats were historically used as a large-weapon test

firing range and impact area. Access to Eagle River from the mouth upstream to Bravo Bridge, approximately 2 miles, is restricted by the military due to the presence of unexploded ordinance, and this reach of river is closed to all sport fishing year-round. The portion of Eagle River upstream from Bailey Bridge to the Glenn Highway Bridge is accessed through Fort Richardson. Upstream of the Glenn Highway, the river meanders through dedicated greenbelt as part of Chugach State Park. Developed public access points on Eagle River are limited. These access sites include 1) the Glenn Highway campground located immediately upstream of Glenn Highway, 2) a day use area upstream of Briggs Bridge, and 3) a parking area and unimproved small boat launch site located at Mile 7.4 of Eagle River Road. The Eagle River drainage is largely used for hiking, camping, and white-water float trips. According to the ADF&G Anadromous Waters Catalog, Eagle River contains all 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon). Dolly Varden are also present in Eagle River. Arctic grayling and rainbow trout have been reportedly seen and caught but these reports have not been confirmed yet. Mainstem Eagle River was enhanced with Chinook salmon in the early 1990s and with 1,010 rainbow trout in 1990. Symphony Lake, one of the headwater lakes of the Southfork of Eagle River was stocked with Arctic grayling in 2001 and 2003.

Eklutna River

Eklutna River is located 17 miles northeast of Anchorage and is the northern boundary to the AMA. Eklutna River is approximately 22 miles long and has as its origin the Eklutna Glacier as well as rain runoff and snowmelt from the Chugach Mountains. The Eklutna River was first dammed in the 1920s to provide power for the growing area. The dam was replaced after the 1964 earthquake. The dam on Eklutna Lake separates and limits flow into Eklutna River. According to the Anadromous Waters Catalog, the Eklutna River contains all 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon). Dolly Varden are also present in the Eklutna River (ADF&G Anadromous Waters Catalog). Eklutna Lake was stocked with rainbow trout in the early 1990s (ADF&G Alaska Lakes Database [ALDAT] <http://www.adfg.alaska.gov/index.cfm?adfg=fishingsportstockinghatcheries.lakesdatabase>, accessed January 2017).

Fish Creek

Fish Creek is a small creek located near Point Woronzof. Much of Fish Creek has been urbanized and runs through culverts through parts of Anchorage. No sport fishing takes place on Fish Creek although according to the Anadromous Waters Catalog, coho salmon can be present at times. Dolly Varden are also likely to be present in Fish Creek along with illegal introductions of goldfish into Cuddy Pond, which is an artificial pond that eventually flows into Fish Creek. It is unknown if the goldfish survive over the winter in the artificial pond.

Ship Creek

Ship Creek is located north of downtown Anchorage at the Port of Anchorage. The source of this creek is Ship Lake as well as rain runoff and snowmelt from the Chugach Mountains. This drainage is approximately 27 miles long and flows into Kink Arm of Cook Inlet. The majority of Ship Creek flows through U.S. Department of Defense (DOD) property. Most fishing occurs in the lower three-quarters of a mile section before entering Cook Inlet. Prior to the official designation of Anchorage in 1915, several indigenous names existed for the creek. Ship Creek was also called *Dgheyaytnu*, translated to “Stickleback Creek,” and *Dgheyay Leht*, which is “Where Stickleback Run” (Kari et al. 2003).

Four dams were constructed in the lower 11 miles of the creek during the 1940s and 1950s for power generation and as a water source for both the Municipality of Anchorage (MOA) and the military bases (Elmendorf and Fort Richardson, currently combined). These dams reduced Ship Creek wild salmon runs (Bosch 2010). All 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon) as well as Dolly Varden and rainbow trout have been observed in Ship Creek (ADF&G Anadromous Waters Catalog). Hatcheries run by ADF&G SF have been stocking Chinook salmon, coho salmon, and rainbow trout (only in 1969) into Ship Creek since 1966. The hatchery facility currently used to stock fish into Ship Creek and currently located on Ship Creek is the William Jack Hernandez Sport Fish Hatchery (WJHSF Hatchery).

In the fall of 2005, ADF&G received funding from U.S. Fish and Wildlife Service (USFWS) for a study evaluating the feasibility of restoring fish passage to Ship Creek upstream of the Elmendorf and Ft. Richardson dams. The report was completed in February 2007 and lists alternative actions for each dam site ranging from “do nothing” to “total removal.” Each alternative was described in terms of impacts and pros and cons for issues such as effectiveness of restoring fish passage, effects to water tables, sediment, creek shoreline stability, and costs. The report also identified issues outside the scope of this initial study (such as social issues) and issues that may require further study. This report was presented to the DOD as the landowner, and to stakeholder agencies, organizations, and the public in order to help choose a course of action that would achieve the goal of restoring fish passage on Ship Creek. DOD stated concerns about allowing fish passage upstream of the WJHSF Hatchery citing BASH (bird air strike hazard), safety issues related to new housing built along the banks of Ship Creek, and potential groundwater issues.

Sixmile Creek

Sixmile Creek is located in Southcentral Alaska on JBER and flows approximately 1 mile into Cook Inlet. The Sixmile Creek drainage includes Sixmile Creek and Lower and Upper Sixmile lakes, and occupies a valley created by an old channel of the Eagle River. The source of the Sixmile drainage is primarily by springs entering the south side of Upper Sixmile Lake. The creek is now flooded for most of its length by the waters of Upper and Lower Sixmile lakes, which were created by damming Sixmile Creek in 1951. Improvements made in 1966 to Talley Avenue crossing the upper portion of the system included the addition of a fish ladder–culvert easing fish access into Upper Sixmile Lake. Much of the old creek channel is still visible in the lakes. All 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon) as well as Dolly Varden and rainbow trout are present in Sixmile Creek (ADF&G Anadromous Waters Catalog). The WJHSF Hatchery stocks rainbow trout into Upper Sixmile Lake; however, these fish have the ability to move freely into Lower Sixmile Lake.

Peters Creek

Peters Creek is located approximately 21 miles to the northeast of Anchorage. The source of Peters Creek is rain and snowmelt from the Chugach Mountains. Chinook, coho, and pink salmon are present at times in Peters Creek. Dolly Varden are also present in Peters Creek (ADF&G Anadromous Waters Catalog).

TURNAGAIN ARM STREAMS

Turnagain Arm streams are the streams that flow into Turnagain Arm range from Campbell Creek, south of Point Woronzof located at the edge of the Ted Stevens International Airport, to

Ingram Creek and include Bird Creek, California Creek, Glacier Creek, Indian Creek, Rabbit Creek, Placer River, and Twenty Mile River. These drainages are prone to flash floods due to the steep nature of the Chugach Mountain Range.

Bird Creek

Bird Creek is southeast of Anchorage approximately 25 miles from downtown Anchorage. This is a popular stream for coho salmon anglers. Bird Creek is fed by high, small, mountain lakes as well as rain and snowmelt. Salmon can only travel up Bird Creek approximately 1.5 miles before reaching an unpassable waterfall. Prior to the falls, there is 1 tributary, Penguin Creek, that is about 0.5 miles from the mouth of Bird Creek. Sections of Bird Creek are popular for salmon fishing. All 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon) as well as Dolly Varden are present in Bird Creek (ADF&G Anadromous Waters Catalog). The WJHSF Hatchery stocks coho salmon into Bird Creek. The entire Bird Creek drainage is closed to fishing January 1 through July 13.

California and Glacier Creeks

California and Glacier creeks are located in Girdwood, Alaska. The community of Girdwood is located approximately 41 miles southeast from downtown Anchorage and near the end of Turnagain Arm. California and Glacier creeks combine 0.36 miles from Turnagain Arm of Cook Inlet. The lower sections of these streams are popular fisheries for locals in Girdwood. Both of the creeks are fed from rain, snowmelt and small lakes in the Chugach Mountains. All 5 species of Pacific salmon (Chinook, chum, coho, pink, and sockeye salmon) as well as Dolly Varden are present in California and Glacier creeks (ADF&G Anadromous Waters Catalog).

Campbell Creek

Campbell Creek travels approximately 21 miles from the Chugach Mountains and flows into Turnagain Arm of Cook Inlet. Campbell Creek was named after Sir Joseph Campbell who was an explorer in Alaska in the late 1700s (Anchorage Park Foundation, <http://anchorageparkfoundation.org/directory/campbell/>, accessed December 2016). Campbell Creek has also historically been known as *Qin Cheghi* (Kari et al. 2003). *Qin Cheghi*, or “Crying Ridge,” refers to Tanaina Peak to the east in the Chugach Mountains and the ridge along the north side of upper Campbell Creek (Kari et al. 2003). North Fork Campbell Creek and South Fork Campbell Creek are both fed by small mountain lakes, rain, and snowmelt. The North Fork and South Fork Campbell Creek combine near Piper Street to form the mainstem Campbell Creek which flows through many neighborhoods. From the junction near Piper Street, Campbell Creek flows approximately 7.5 miles into the artificial Campbell Lake. It exits Campbell Lake and flows about one-quarter mile to the mud flats of Turnagain Arm. Four species of Pacific salmon (Chinook, coho, pink, and sockeye salmon), as well as Dolly Varden and rainbow trout, are present in Campbell Creek at times (ADF&G Anadromous Waters Catalog). It is also believed that chum salmon and steelhead trout—*anadromous rainbow trout*, which were stocked with steelhead (Bosch 2010)—are present in Campbell Creek but in very low numbers and are not well documented. Rainbow trout and coho salmon are also enhanced into sections of Campbell Creek. Campbell Creek, the largest free-flowing stream in the Anchorage metropolitan area, supports a small wild Chinook salmon run. To protect the wild spawning rainbow trout, the entire drainage is closed to fishing April 15 through June 14.

The upper reach of Campbell Creek is composed of 2 tributaries, the North and South Forks, which drain from the Chugach Mountains east of Anchorage. Both forks flow through canyons in their upper reaches that are impassable to upstream fish migration. Downstream of the canyons, these tributary streams flow approximately 10 miles through the largely undeveloped forests and wetlands of Chugach State Park and Far North Bicentennial Park before converging near Piper Street. Campbell Creek flows through MOA greenbelt and private property from the confluence of the forks downstream to Cook Inlet. The greatest effects from urbanization have occurred in this reach of Campbell Creek.

MOA has made an effort to obtain and preserve the riparian habitat of Campbell Creek from Lake Otis Parkway downstream to Campbell Lake and to improve water quality. In 1981, the Bureau of Land Management (BLM) transferred title to the 4,000-acre Campbell Tract (Bicentennial Park) to the MOA. This area makes up the primary spawning and rearing habitat for salmon. Campbell Creek is stocked annually with coho salmon and rainbow trout.

Indian Creek

Indian Creek is a small stream located approximately 24 miles southeast of Anchorage near the small community of Indian. Indian Creek is fed from rain and snowmelt from the steep but south facing slopes of the Chugach Mountain Range before flowing into Turnagain Arm of Cook Inlet. According to the ADF&G Anadromous Waters Catalog, Indian Creek has Chinook, coho, and pink salmon present at times of the year. Chum salmon and Dolly Varden may also be present.

Ingram Creek

Ingram Creek is the southernmost stream in the AMA. Its headwaters are in Turnagain Pass in the Kenai Peninsula Mountain Range. Its water source is rain and snowmelt. According to the ADF&G Anadromous Waters Catalog, chum, coho, and pink salmon are present. Dolly Varden are also present in Ingram Creek. Ingram Creek was stocked in the late 1980s with coho and pink salmon (ADF&G Mark-Tag Lab Hatchery Release Database; <http://mtalab.adfg.alaska.gov/CWT/reports/hatcheryrelease.aspx>, accessed December 2016).

Rabbit Creek

Rabbit Creek is located south of Anchorage. Water sources include Rabbit Lake as well as rain and snowmelt. It flows through many developed neighborhoods and eventually flows through Potter Marsh Wildlife Refuge before entering Turnagain Arm of Cook Inlet. Rabbit Creek has small runs of Chinook, chum, coho, and sockeye salmon as well as Dolly Varden (ADF&G Anadromous Waters Catalog). Rabbit Lake is stocked with rainbow trout. Rabbit Creek downstream of the Old Seward Highway is closed to all sport fishing.

Twentymile River

Twentymile River is located approximately 47 miles southeast of Anchorage at the head of Turnagain Arm. Twentymile River divides into South Fork, Twentymile, North Fork, and Glacier River. Glacier River is fed from Carmen Lake and Twentymile Glacier. The sources of water feeding the Twentymile drainage, including Glacier River and Carmen Creek, are rain, Carmen Lake, snowmelt, and Twentymile Glacier. The lower sections of the Twentymile River drainage are often traveled by jet boats, but airboats are often used to reach the upper section. According to the ADF&G Anadromous Waters Catalog, Chinook, coho, chum, and sockeye salmon are present. Dolly Varden and eulachon (*Thaleichthys pacificus*) have also been

documented in Placer River. It is suspected that pink salmon and rainbow trout–steelhead may also be present, but these have not been documented.

Placer River

The Placer River is a glacial river located approximately 52 miles southeast of Anchorage at the head of Turnagain Arm of Cook Inlet. The sources of the water feeding the Placer River are rain, snowmelt and glaciers (Bartlett, Skookum, and Spencer glaciers). The Placer River is constantly changing channels and it is difficult for jet boats to navigate because of the shallowness. According to the ADF&G Anadromous Waters Catalog, coho salmon are present. Dolly Varden and sockeye salmon have also been documented in Placer River. It is suspected that chum salmon may also be present, but these have not been documented.

Portage River

Portage River is approximately 45 miles southeast of Anchorage and is located at the head of Turnagain Arm. Portage River is fed primarily by Portage Lake and Placer Creek, which flows out of Bear Valley on the north side of Portage Lake, which is glacial. Chum, coho, pink, and sockeye salmon have been documented in the Portage drainage (ADF&G Anadromous Waters Catalog).

ANCHORAGE LAKES

Anchorage Management Area (AMA) lakes are currently stocked with catchable-sized fish. Before the 1960s and the initiation of stocking efforts, few AMA lakes supported resident fish populations of recreational interest. Most lakes are landlocked and threespine stickleback (*Gasterosteus aculeatus*) was the only fish species present. In the 1960s, ADF&G began a rainbow trout stocking program to increase sport fishing opportunities within the AMA. Alaska blackfish were illegally introduced into AMA lakes and can now be found in almost all area lakes.

In order to evaluate the stocking program, a creel survey was conducted during 1986 on 4 AMA lakes (Havens et al. 1987). Results of this survey indicated that youths and adult males were the primary recreational anglers. The main objective of the survey was to determine if a single annual spring release of a large number of rainbow trout was suitable for anglers using AMA lakes. Data indicated that catch rates remained high for 2 to 6 weeks after stocking, and then dropped to below 1 fish per angler-hour. It was recommended, and adopted, that initial stocking would occur after ice-out and then a second stocking would be repeated after 4 to 6 weeks. Multiple stocking of high use lakes appears to increase fishing success throughout the open water season.

The AMA stocked lakes program has increased sport fishing opportunities for the general public, including the development of educational fishing classes for youths and adults, and an annual ice fishing jamboree for disabled and underprivileged anglers. Fish species stocked from 2011 to 2015 into AMA lakes included rainbow trout, Arctic char, grayling, and land-locked Chinook salmon. Locations of stocked AMA lakes can be found in Figure 2.



Figure 2.—Anchorage Management Area streams and stocked lakes.

STOCKING PROGRAM

The Alaska Department of Fish and Game produces a Statewide Stocking Plan, which outlines plans for stocking sport fish throughout Alaska every 5 years with annual updates as needed. The Division of Sport Fish stocking program is primarily funded with Federal Aid in Sport Fish Restoration funds and license sales revenue. Fish stocking plans go through a formal process and sites are reviewed prior to initiating stocking. The stocking plan for future years can be found online at <http://www.adfg.alaska.gov/index.cfm?adfg=fishingSportStockingHatcheries.stockingPlan>. Historical sport fish stockings can be found in the stocked fish database: <http://www.adfg.alaska.gov/index.cfm?adfg=SportStockingHatcheriesSearch.main>.

Historically, stocking in the AMA has been primarily by 2 fish hatcheries (Fort Richardson Hatchery and Elmendorf Hatchery), both located on Ship Creek since the late 1950s. The Fort Richardson hatchery was built in 1958 by the U.S. Army to provide fish for lakes on the DOD property. The Alaska Department of Fish and Game became involved in the early 1960s and assumed full operation of the Fort Richardson Hatchery facility by the late 1960s. The Fort Richardson Hatchery sits on the banks of Ship Creek, just downstream of the Glenn Highway (ADF&G website Fort Richardson State Fish Hatchery; <http://www.adfg.alaska.gov/index.cfm?adfg=fishingSportStockingHatcheries.ftrichardson>, accessed December 2016) and approximately 5 miles upstream from the Elmendorf Hatchery. The Elmendorf State Hatchery started in 1965. Elmendorf State Hatchery was located in Anchorage on the north bank of Ship Creek near the intersection of Reeve Boulevard and Post Road (ADF&G website Elmendorf State Fish Hatchery; <http://www.adfg.alaska.gov/index.cfm?adfg=fishingSportStockingHatcheries.elmendorf>, accessed December 2016). The permanent loss of heated effluent from the Fort Richardson (2003) and Elmendorf (2005) power plants resulted in a reduction in fish size and number of catchable-sized fish released from 2005 to 2011. The catchable rainbow trout program shifted from producing 1-year-old catchable fish to producing 3-year-old catchable fish and 2-year-old undersized catchable or subcatchable-sized fish. The closure of Elmendorf Hatchery in 2010 resulted in the release of rainbow trout fingerling into local lakes. The catchable Chinook salmon program saw reductions in fish size, number of fish released, and number of lakes stocked starting in 2006 when an outbreak of Bacterial Kidney Disease, also known as BKD, in the Chinook salmon at Fort Richardson Hatchery resulted in the release of fingerling instead of catchable Chinook salmon that year. In addition, from 2007 to 2009, only category 1 (landlocked) lakes were permitted for releases of catchable-sized fish from Elmendorf Hatchery after an Oregon State University graduate student reported the presence of DNA from the parasite that causes whirling disease. Catchable-sized Chinook salmon were not released in 2010–2011. Due to disease concerns, catchable-sized Arctic grayling production ceased in 2002. Arctic grayling fingerling and subcatchable-sized fish were released from 2003 to 2012.

In June 2011, construction of the new William Jack Hernandez Sport Fish Hatchery (WJHSF Hatchery) was completed. With over 100 rearing tanks, there is the capacity to produce over 6 million sport fish each year to stock throughout permitted areas of Alaska. The rainbow trout and Chinook salmon catchable production programs returned to historical stocking levels when the first catchable-sized fish were released from the WJHSF Hatchery in 2012. In addition, the first release of catchable-sized Arctic grayling from WJHSF Hatchery occurred in 2013; however, due to budget restrictions, the Arctic grayling stocking program was suspended indefinitely after stocking in 2015 to reduce operating expenses (ADF&G Statewide Stocking Plan for Sport Fish

2016; <http://www.adfg.alaska.gov/static/fishing/pdfs/hatcheries/16intro.pdf>, accessed December 2016).

FISHERIES MANAGEMENT AND OBJECTIVES

ADF&G has a constitutional mandate to manage on the principle of sustained yield. Within the sustained yield principle, SF goals seek to optimize social and economic benefits, and where possible, expand opportunity to participate in diverse fisheries on these stocks.

Fisheries management and objectives are implemented at the species level within a particular drainage, although other species present in the drainage may be impacted by species-specific objectives. Drainages in the AMA are dynamic, and often more than one species is present at a time, which must be considered when management objectives and actions are being assessed.

FISHERIES PERFORMANCE

Overall, there has been a slight decrease in sport fishing effort in the AMA, although the area sport fishing efforts mirror the statewide trends in sport fishing effort. According to the ADF&G Statewide Harvest Survey (SWHS), the AMA represents approximately 75,982 angler-days (2006–2015 average), which represents approximately 3% of the statewide effort and 5% of the effort for Southcentral Alaska (Table 1). Most (97% on average for 2006–2015) of the effort is in the freshwaters of the AMA (Table 2). Considering just freshwater effort in the AMA, 59% or 44,456 angler-days (2006–2015 average) are spent fishing on streams (as opposed to lakes), annually (calculated from Table 2). In 2011 and 2015, the freshwater effort was higher in streams (60.5% and 57.5%, respectively) versus lakes. However, from 2012 to 2014, effort was more evenly distributed between streams (2012–2014 average: 48.4%) and lakes (2012–2014 average: 49.5%) (calculated from Table 2).

In streams, anglers primarily focused on anadromous salmon versus trout (Dolly Varden or rainbow trout). The recent catch of anadromous salmon in streams has decreased, although catches have increased since the historical low of 2012 when 17,425 salmon were caught (Table 3). From 2013 to 2015, the salmon catch ranged from 22,196 to 43,615 salmon (Table 3) but have not improved to the levels observed in 1998 and 2000 when annually over 100,000 salmon were caught. In AMA streams, approximately 46,746 salmon are caught annually and 15,323 (33%) of those salmon were harvested on average during the past 10 years (2006–2015). Many of the local streams are stocked with anadromous salmon to support the growing population of Anchorage. Overall quality of smolt released in enhanced streams has improved since 2012 with the new WJHSF Hatchery, although catch of anadromous salmon has not reached pre-2010 levels on both wild and stocked streams, which could indicate poor marine survival (Figure 3). However, a detailed investigation at a scale that includes more areas, commercial and subsistence harvest, and an age-class analysis of fish would be needed to further support this claim. Over the last 10 years (2006–2015), coho salmon were harvested more frequently than any other species of salmon in the AMA, followed by pink, Chinook, chum, and sockeye salmon, in that order (Table 3).

Table 1.—Sport fishing effort expended in the Anchorage Management Area (AMA) compared to Southcentral Alaska and statewide, 1996–2015.

Year	Statewide effort	Southcentral effort	AMA		
			Effort	Percent of statewide	Percent of Southcentral
1996	2,006,528	1,434,943	126,325	6	9
1997	2,079,514	1,400,983	139,549	7	10
1998	1,856,976	1,258,782	125,513	7	10
1999	2,499,152	1,659,966	146,789	6	9
2000	2,627,805	1,844,824	167,499	6	9
2001	2,261,941	1,560,562	135,359	6	9
2002	2,259,091	1,569,513	111,694	5	7
2003	2,219,398	1,535,501	104,004	5	7
2004	2,473,961	1,709,671	101,943	4	6
2005	2,463,929	1,712,610	101,041	4	6
2006	2,297,961	1,605,852	103,800	5	6
2007	2,543,674	1,799,352	91,881	4	5
2008	2,315,601	1,622,920	111,121	5	7
2009	2,216,445	1,522,345	79,743	4	5
2010	2,000,167	1,371,470	61,704	3	4
2011	1,919,313	1,326,950	48,187	3	4
2012	1,885,692	1,252,263	45,407	2	4
2013	2,202,957	1,488,383	64,037	3	4
2014	2,309,853	1,571,650	79,306	3	5
2015	2,212,331	1,470,381	74,631	3	5
Average					
2011–2015	2,106,029	1,421,925	62,314	3	4
2006–2015	2,190,399	1,503,157	75,982	3	5

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Note: “Effort” means number of angler-days.

Table 2.—Saltwater, lake, and stream sport fishing effort (angler-days), Anchorage Management Area (AMA), 1996–2015.

Year	Saltwater		Freshwater						AMA Total Effort	
	Effort	Percent	Lake		Stream		Total freshwater			
			Effort	Percent	Effort	Percent	Effort	Percent		
1996	870	1	54,699	43	70,756	56	125,455	99	126,325	
1997	1,449	1	64,331	46	73,769	53	138,100	99	139,549	
1998	2,921	2	43,905	35	78,687	63	122,592	98	125,513	
1999	2,916	2	66,312	45	77,561	53	143,873	98	146,789	
2000	2,197	1	69,607	42	95,695	57	165,302	99	167,499	
2001	2,277	2	47,384	35	85,698	63	133,082	98	135,359	
2002	3,493	3	40,201	36	68,000	61	108,201	97	111,694	
2003	3,243	3	40,552	39	60,209	58	100,761	97	104,004	
2004	1,251	1	47,539	47	53,153	52	100,692	99	101,943	
2005	2,670	3	36,833	36	61,538	61	98,371	97	101,041	
2006	1,540	1	35,741	34	66,519	64	102,260	99	103,800	
2007	5,542	6	28,833	31	57,506	63	86,339	94	91,881	
2008	2,977	3	35,984	32	72,160	65	108,144	97	111,121	
2009	2,616	3	27,910	35	49,217	62	77,127	97	79,743	
2010	1,675	3	24,152	39	35,877	58	60,029	97	61,704	
2011	2,230	5	16,793	35	29,164	61	45,957	95	48,187	
2012	702	2	22,265	49	22,440	49	44,705	98	45,407	
2013	1,085	2	32,095	50	30,857	48	62,952	98	64,037	
2014	2,123	3	39,333	50	37,850	48	77,183	97	79,306	
2015	921	1	30,738	41	42,972	58	73,710	99	74,631	
Average										
2011–2015	1,412	2	28,245	45	32,657	53	60,901	98	62,314	
2006–2015	2,141	3	29,384	40	44,456	57	73,841	97	75,982	

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Note: “Effort” means number of angler-days.

Table 3.—Sport fish catch and harvest of anadromous salmon, Anchorage Management Area, 1996–2009.

Year	Chinook salmon			Chum salmon			Coho salmon			Pink salmon			Sockeye salmon			Total	
	Catch	Harv.	% RIs	Catch	Harv.	% RIs	Catch	Harv.	% RIs	Catch	Harv.	% RIs	Catch	Harv.	% RIs	Catch	Harv.
1996	11,187	4,456	60	4,522	607	87	27,726	17,795	36	19,189	2,862	85	2,349	884	62	64,973	26,604
1997	9,799	4,616	53	1,588	248	84	30,192	20,578	32	5,120	843	84	2,445	1,023	58	49,144	27,308
1998	6,022	2,156	64	5,154	513	90	64,522	42,219	35	28,138	3,205	89	2,100	1,271	39	105,936	49,364
1999	15,118	5,462	64	2,304	129	94	17,834	12,266	31	5,462	721	87	1,507	542	64	42,225	19,120
2000	11,848	4,752	60	3,936	340	91	46,888	28,191	40	38,236	3,123	92	1,182	537	55	102,090	36,943
2001	11,843	4,452	62	4,631	470	90	63,865	40,693	36	12,988	783	94	2,292	894	61	95,619	47,292
2002	7,070	2,421	66	6,540	472	93	41,219	26,260	36	11,651	1,168	90	694	330	52	67,174	30,651
2003	9,480	3,678	61	4,121	313	92	20,762	13,375	36	9,461	1,600	83	1,791	943	47	45,615	19,909
2004	7,713	3,160	59	2,185	306	86	25,474	13,447	47	7,897	1,272	84	1,012	286	72	44,281	18,471
2005	9,202	4,329	53	2,376	234	90	25,937	15,063	42	10,739	677	94	997	551	45	49,251	20,854
2006	6,857	3,165	54	4,427	242	95	35,854	19,863	45	23,926	2,345	90	341	91	73	71,405	25,706
2007	6,142	3,106	49	2,968	97	97	17,806	10,692	40	34,318	3,278	90	595	172	71	61,829	17,345
2008	5,464	2,647	52	6,033	283	95	26,124	17,996	31	30,004	2,032	93	1,719	223	87	69,344	23,181
2009	2,655	1,027	61	5,945	386	94	17,736	10,805	39	62,996	6,426	90	663	192	71	89,995	18,836
2010	2,141	1,130	47	3,852	335	91	5,983	4,466	25	14,625	2,578	82	618	193	69	27,219	8,702
2011	1,322	616	53	4,350	397	91	10,327	7,405	28	15,241	893	94	535	244	54	31,775	9,555
2012	334	113	66	1,749	137	92	6,535	4,187	36	8,587	1,488	83	220	64	71	17,425	5,989
2013	1,304	824	37	3,511	2,053	42	7,641	6,190	19	9,568	2,447	74	172	28	84	22,196	11,542
2014	1,382	882	36	3,737	541	86	13,939	9,430	32	12,941	1,473	89	656	288	56	32,655	12,614
2015	3,077	1,820	41	1,742	194	89	19,218	15,099	21	19,236	2,382	88	342	266	22	43,615	19,761
Average																	
2011–2015	1,484	851	47	3,018	664	80	11,532	8,462	27	13,115	1,737	85	385	178	57	29,533	11,892
2006–2015	3,068	1,533	50	3,831	467	87	16,116	10,613	32	23,144	2,534	87	586	176	66	46,746	15,323

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Note: Column labels are “catch,” “harvest,” and “percent released.”




Figure 3.—Sport fish catch and harvest of anadromous salmon, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

CHINOOK SALMON FISHERIES

FISHERY DESCRIPTION

Chinook salmon return to AMA streams from May through early July. Chinook salmon typically spend 2 to 4 years feeding in the ocean before returning to their natal stream to spawn (Healey 1991) so in any given year, the run will be made up of multiple year classes. Wild Chinook salmon runs are found in creeks or rivers across the area including Bird, California (a tributary to Glacier Creek in Girdwood), Carmen (a tributary of Glacier River in Twentymile River drainage), Campbell, Eagle, Eklutna, Glacier, Indian, Peters, Placer, Portage, Rabbit, Ship, and Twentymile creeks. Most of these streams support annual Chinook salmon runs of less than 100 fish each (Bosch 2010). As a result, sport fishing for Chinook salmon in streams has been closed with few exceptions.

Chinook salmon fisheries in the AMA include Campbell Creek, Eagle River, and Ship Creek. The largest Chinook salmon sport fishery is hatchery produced and occurs on Ship Creek. The Ship Creek sport fishery is centrally located in the heart of downtown Anchorage, providing a unique opportunity for anglers in an urban setting. Wild Chinook salmon fisheries include Eagle River and a youth-only fishery on Campbell Creek. ADF&G performs survey counts of Chinook salmon returning to sections of Ship, Rabbit, Bird, and Penguin creeks, as well as the south fork of Eagle River and Meadow Creek (tributary of Eagle River; Appendices C1–C5).

Campbell Creek

Prior to 2005, Chinook salmon sport fishing had not been permitted in Campbell Creek since statehood in 1959. However, in 2005, the Alaska State Legislature gave the Alaska Board of Fisheries (BOF) the authority to create youth-only fisheries (youth, under age 16, are not required to obtain a sport fishing license). The BOF created the Campbell Creek Youth-only Fishery to give area kids, 15 or younger, access to a Chinook salmon fishery where they would not have to compete with adult anglers. Kids (youth), under the age of 16 may fish for Chinook salmon on Campbell Creek between Dimond Boulevard and the Old Seward Highway between 6:00 AM and 10:00 PM daily on the last Saturday and Sunday of June of each year. Campbell Creek is closed to fishing for Chinook salmon the rest of the year. After the youth-only fishery, a combination of foot and float escapement surveys are performed to count spawning Chinook salmon in Campbell Creek. The current bag and possession limits for Chinook salmon during the Campbell Creek Youth Fishery are 1 per day, 1 in possession for Chinook salmon 20 inches or longer; and 10 per day, 10 in possession for Chinook salmon less than 20 inches long. After taking a Chinook salmon 20 inches or longer, a person may not fish for any species that same day in waters open to Chinook salmon sport fishing. Chinook salmon 20 inches or longer harvested in Campbell Creek must be immediately recorded on the back of the youth's harvest record card and count towards the annual limit of 5 Chinook salmon harvested from the fresh waters of Cook Inlet. Chinook salmon less than 20 inches in length do not need to be recorded on the back of the sport fishing license or harvest record card, and do not count toward the Cook Inlet seasonal limit. The small wild run of Chinook salmon has not been enhanced with hatchery releases and cannot provide a sustainable harvest in such a heavily urbanized location.

In 2006, the Campbell Lake Homeowners Association and MOA were permitted to lower Campbell Lake levels for 3 months in order to winterkill invasive aquatic vegetation species in the lake that were interfering with float plane travel. ADF&G staff expressed concern that Campbell Lake, as the only significant deep water body in the Campbell Creek drainage, probably provides significant habitat for stocks of wild Chinook and sockeye salmon, hatchery-enhanced runs of coho salmon, and resident species in the watershed. Permit requests to extend the refill date of May 15, 2007, by a week were filed but denied. In 2010, the Campbell Lake Homeowners Association and MOA were permitted to draw down Campbell Lake in order to finish the dredging project started in 2006.

Eagle River

Before the Chinook salmon fishery opening in 1992 in the Eagle River, angler effort targeting other salmon and resident species averaged about 2,300 angler-days from 1982 through 1991 (Stratton and Cyr 1997). In 1992, the first year of the Chinook salmon fishery, estimated effort was 4,908 angler-days for all species (Mills 1993). The estimated Chinook salmon catch was 109 fish and harvest was 48 fish (Mills 1993). Approximately 300 wild Chinook salmon and 1,000 hatchery Chinook salmon were projected to be available to sport anglers. Estimated effort for Eagle River was 3,396 angler-days in 1993; estimated catch and harvest were 88 and 47 Chinook salmon, respectively (Mills 1994). The low harvest in 1993 was surprising because over 2,300 hatchery Chinook salmon were expected to be available. Because the hatchery run failed to materialize and because the estimated angler-days of effort were at prestocking levels, the stocking program was eliminated in 1995. The fishery was reconfigured by the BOF in spring 1996 and the area open to Chinook salmon fishing was restricted to the portion of the Eagle River near the Glenn Highway campground. This small area was only open for four 3-day

weekends (Saturday, Sunday, Monday) beginning Memorial Day weekend. Effort was estimated at only 108 angler-days in 2009 and no catch was reported (Tables 4 and 5). It is thought that this limited fishery does not impact natural Chinook salmon runs.

From 2011 to 2015, Eagle River was only open to fishing for Chinook salmon from the Bailey Bridge on Fort Richardson upstream to ADF&G markers in the Chugach State Park Eagle River Campground on Saturdays, Sundays and Mondays for the 4 consecutive weeks beginning on the Saturday before Memorial Day. The current bag and possession limits for Chinook salmon in those waters of Eagle River open to salmon fishing are 1 per day, 1 in possession for Chinook salmon 20 inches or longer; and 10 per day, 10 in possession for Chinook salmon less than 20 inches in length. After taking a Chinook salmon 20 inches or longer, a person may not fish for any species that same day in waters open to Chinook salmon sport fishing. Chinook salmon 20 inches or longer harvested in Eagle River must be immediately recorded on the back of the angler's sport fishing license or harvest record card and count towards the annual limit of 5 Chinook salmon from the fresh waters of Cook Inlet. Chinook salmon less than 20 inches in length do not need to be recorded on the back of the sport fishing license or harvest record card, and do not count towards the Cook Inlet seasonal limit.

Ship Creek

Ship Creek was open to Chinook salmon sport fishing from 1957 through 1959 but was closed from 1960 through 1969. Chinook salmon fishing was allowed during selected periods in Ship Creek downstream of the Chugach Power Plant Dam from 1970 through 1972. From 1973 through 1986, the creek was closed to Chinook salmon sport fishing due in part to low Chinook salmon abundance throughout northern Cook Inlet. Beginning in 1987, as returns increased from annual stocking efforts, the lower portion of Ship Creek downstream of the Chugach Power Plant Dam was reopened to Chinook salmon sport fishing 2 days per week for 5 consecutive weeks in June and July. The season was expanded to 7 days per week, January 1 through July 13, in 1990. Nighttime closures from 11:00 PM to 6:00 AM were issued by emergency order to help address enforcement issues, and passed into regulation in 2001. The fishery now occurs during late May through early July in the lower 1 mile of Ship Creek, downstream of the Chugach Power Plant Dam (Figure 4). Fishing is also closed within 100 ft of the Chugach Power Plant Dam. The shoreline of the area open to Chinook salmon fishing is owned and managed by Alaska Railroad Corporation (ARRC) and the MOA. The Ship Creek Chinook Salmon Derby, which was first held in 1993, has become an annual event currently sponsored by Grace Alaska to benefit their Downtown Soup Kitchen.

The current bag and possession limits for Chinook salmon in those waters of Ship Creek open to salmon fishing are 1 per day, 1 in possession for Chinook salmon 20 inches or greater; and 10 per day, 10 in possession for Chinook salmon less than 20 inches. After taking a Chinook salmon 20 inches or longer, a person may not fish for any species that same day in waters open to Chinook salmon sport fishing. Chinook salmon 20 inches or longer harvested in Ship Creek must be immediately recorded on the back of the angler's sport fishing license or harvest record card and counted towards the annual limit of 5 Chinook salmon from the salt or fresh waters of Cook Inlet north of Bluff Point.

Table 4.—Angler effort for streams in the Anchorage Management Area, 1996–2015.

Year	Bird Creek		Campbell Creek		Eagle River		Ship Creek		Twentymile R.		Other		Total stream effort
	Effort	Percent of total	Effort	Percent of total	Effort	Percent of total	Effort	Percent of total	Effort	Percent of total	Effort	Percent of total	
1996	12,003	17	5,225	7	3,499	5	42,454	60	3,823	5	3,752	5	70,756
1997	12,136	16	5,897	8	2,059	3	47,826	65	3,170	4	2,681	4	73,769
1998	20,797	26	4,834	6	1,023	1	44,670	57	2,805	4	4,558	6	78,687
1999	13,033	17	4,446	6	2,096	3	52,294	67	3,370	4	2,322	3	77,561
2000	17,550	18	3,918	4	1,998	2	62,101	65	3,620	4	6,508	7	95,695
2001	13,662	16	6,222	7	1,214	1	56,402	66	4,161	5	4,037	5	85,698
2002	5,540	8	4,561	7	1,538	2	46,955	69	2,869	4	6,537	10	68,000
2003	3,691	6	4,937	8	1,382	2	40,380	67	3,418	6	6,401	11	60,209
2004	2,239	4	5,674	11	1,289	2	35,524	67	3,826	7	4,601	9	53,153
2005	8,365	14	6,933	11	1,130	2	39,610	64	2,554	4	2,946	5	61,538
2006	15,220	23	6,235	9	1,591	2	34,557	52	2,299	3	6,617	10	66,519
2007	13,247	23	5,779	10	955	2	30,676	53	2,749	5	4,100	7	57,506
2008	10,866	15	6,267	9	1,541	2	40,605	56	4,874	7	8,007	11	72,160
2009	13,605	28	2,774	6	108	0	23,663	48	1,729	4	7,338	15	49,217
2010	7,464	21	2,209	6	515	1	18,136	51	1,738	5	5,815	16	35,877
2011	5,027	17	2,897	10	474	2	12,473	43	1,843	6	6,450	22	29,164
2012	4,222	19	2,183	10	393	2	8,763	39	1,304	6	5,575	25	22,440
2013	4,142	13	4,271	14	463	2	17,287	56	1,323	4	3,371	11	30,857
2014	9,171	24	4,295	11	182	0	16,780	44	1,899	5	5,523	15	37,850
2015	8,953	21	4,760	11	300	1	24,253	56	989	2	3,717	9	42,972
Average													
2011–2015	6,303	19	3,681	11	362	1	15,911	48	1,472	5	4,927	16	32,657
2006–2015	9,192	20	4,167	10	652	1	22,719	50	2,075	5	5,651	14	44,456

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Table 5.—Chinook salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.

Year	Eagle River		Ship Creek		Salt water		Other freshwater		Area total	
	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest
1996	586	309	9,354	3,774	21	0	1,226	373	11,187	4,456
1997	306	140	9,045	4,456	30	20	418	0	9,799	4,616
1998	64	19	5,382	2,099	121	38	455	0	6,022	2,156
1999	48	22	14,275	5,204	201	167	594	69	15,118	5,462
2000	132	109	11,090	4,593	60	20	566	30	11,848	4,752
2001	132	58	10,656	4,286	262	108	793	0	11,843	4,452
2002	162	34	5,967	2,287	164	94	777	6	7,070	2,421
2003	76	25	8,667	3,588	78	52	659	13	9,480	3,678
2004	51	24	6,840	2,790	104	58	718	288	7,713	3,160
2005	25	25	7,578	4,081	183	108	1,416	115	9,202	4,329
2006	251	60	5,464	3,060	32	32	1,057	13	6,804	3,165
2007	125	47	4,888	2,615	695	366	434	78	6,142	3,106
2008	46	0	4,279	2,540	92	92	1,047	15	5,464	2,647
2009	0	0	1,869	884	209	143	577	0	2,655	1,027
2010	0	0	1,918	1,095	0	0	223	35	2,141	1,130
2011	17	0	1,171	600	16	16	118	0	1,322	616
2012	0	0	154	113	0	0	180	0	334	113
2013	0	0	1,265	824	0	0	39	0	1,304	824
2014	0	0	1,245	882	0	0	137	0	1,382	882
2015	0	0	3,002	1,761	16	16	59	43	3,077	1,820
Average										
2011–2015	3	0	1,367	836	6	6	107	9	1,484	851
2006–2015	44	11	2,526	1,437	106	67	387	18	3,063	1,533

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.



Figure 4.—Map of lower section of Ship Creek.

The sport fishing bag and possession limits for Chinook salmon less than 20 inches are currently 10 per day, 10 in possession. The limits for Chinook salmon 20 inches or longer in length from Ship Creek are currently 1 per day, 1 in possession. A signed Chinook salmon stamp, or a Permanent Identification Card, is required to fish for Chinook salmon. Harvest must be immediately recorded in ink on the back of the angler’s sport fishing license or harvest record card, and counts toward the Cook Inlet seasonal limit of 5 Chinook salmon. Chinook salmon less than 20 inches in length do not need to be recorded on the back of the sport fishing license and do not count toward the Cook Inlet seasonal limit.

FISHERY MANAGEMENT AND OBJECTIVES

In the majority of streams in the AMA, Chinook salmon runs are too small to allow a harvestable surplus. The management goal and objectives are to maintain historical Chinook salmon escapement levels, continue natural production, and provide viewing opportunities. Campbell Creek is the only creek that has a sustainable escapement goal (SEG) threshold in the AMA.

Campbell Creek

The management objectives for Campbell Creek Chinook salmon are to manage and protect the wild Chinook salmon run and to achieve an SEG threshold of 380 Chinook salmon. The SEG threshold for Campbell Creek Chinook salmon was developed in 2010 and includes the 2010

data in the analysis to develop this goal. Campbell Creek was previously managed for an SEG with a range of 50 to 700 fish (Munro and Volk 2010). In 2016, the SEG was reevaluated and determined to still be a suitable goal with the data available.

Eagle River

Eagle River is managed to allow small levels of opportunity for Chinook salmon fishing while ensuring wild populations are not impacted. In addition, the fishery is managed to maintain historical Chinook salmon escapement levels, continue natural production, and provide viewing opportunities. Eagle River no longer has an SEG for Chinook salmon (Munro and Volk 2010).

Ship Creek

This fishery began to develop in 1987, with fishing open for 2 days each week (Appendix A1) to allow stocked Chinook salmon returns to build to harvestable levels. The fishery was expanded to 7 days per week in 1991, resulting in a catch of over 1,600 Chinook salmon and a harvest of over 1,100 Chinook salmon during that year (Bosch 2010). Small saltwater Chinook salmon fisheries occur near the mouths of Ship and Bird creeks.

The 3 management objectives for the Ship Creek Chinook salmon fishery are as follows: 1) to maintain or increase current angler effort through smolt stocking, 2) to generate at least 50,000 angler-days of annual sport fishing opportunity directed at stocked Chinook and coho salmon in Ship Creek, and 3) to produce a return to Ship Creek of 6,000 to 9,000 adult Chinook salmon to assure 750 adult salmon are available (having passed above the Chugach Power Plant Dam) for natural spawning, fish viewing, and meeting hatchery egg-take needs.

STOCKING PROGRAM

Ship Creek

Currently, the only anadromous Chinook salmon stocking in the AMA occurs in Ship Creek. Attempts to enhance Ship Creek salmon runs occurred from 1966 through 1980 when Chinook salmon of Alaska and Oregon origin were stocked (Miller 1990; Stratton and Cyr 1995). During this period, eggs obtained from these stocks were incubated at Fire Lake Hatchery. The fry were reared to smolt in the Fort Richardson Hatchery before release. These releases were generally unsuccessful because consistent numbers of returning adults could not be established. A more successful hatchery enhancement program was established in 1987 using smolt releases from the Elmendorf Hatchery and Ship Creek Chinook salmon brood stock.

The Chinook salmon run is the result of an annual release of smolt raised at the WJHSF Hatchery. From 2011 through 2015, an average of 338,237 Chinook salmon smolt were released into Ship Creek annually (Appendix B1).

Eagle River

In 1990, an annual stocking program was initiated in Eagle River with approximately 105,000 Chinook salmon smolt of Ship Creek origin (Stratton and Cyr 1995). Due to poor returns and difficult fishing conditions, the stocking program was discontinued in 1995.

FISHERY PERFORMANCE

The AMA Chinook salmon fisheries are largely driven by the Ship Creek fishery. On average (2006–2015), the Ship Creek Chinook salmon fishery accounts for 50% of the total angler effort

in Anchorage streams each year (Table 4). Chinook salmon catches in the AMA have averaged 1,484 for the 5-year period from 2011 through 2015 and the average harvest for this period is 851 Chinook salmon, or about 57% of the estimated catch (Table 5).

Campbell Creek

The Youth-only Fishery began in 2005; however, because it is only open to youths 15 years old and younger (who do not require Alaska sport fishing licenses), none are eligible to receive a Statewide Harvest Survey, from which catch and harvest are estimated. Furthermore, even if other members of the household received a survey, it is likely the Youth-only Fishery effort was underreported. In 2005, the first year of the fishery, an estimated 750–1,000 youth anglers and adult chaperons participated in the fishery and harvested 150–200 Chinook salmon. No annual estimates of participation have been made for this Youth-only Fishery. Participation, effort, and success may reflect run strength and weather conditions during the fishery. Anecdotal information indicates that illegal harvest is probably greater than the legal harvest.

Escapement of Chinook salmon in Campbell Creek is assessed using a combination of foot and float surveys. The accuracy of these surveys varies greatly depending on timing, water levels, and water clarity. Chinook salmon survey escapements have ranged from 1,119 fish (1997) to 260 fish (2011) (Appendix C3). These estimates are based on foot surveys conducted along the length of the creek after the Youth-only fishery occurred. No surveys were conducted in 2012–2013 due to reductions in staff. The average observed escapement using data from surveys conducted in the last 10 years (2006–2015) has been approximately 514 Chinook salmon.

Eagle River

In Eagle River, the average effort from 2011 to 2015 was 362 angler-days or approximately 1% of angling effort for AMA streams (Table 4); much of the Eagle River effort is probably targeting Dolly Varden (Bosch 2010). In 2011, only 17 Chinook salmon were caught and 0 were harvested according to the SWHS. No Chinook salmon were reported caught or harvested in Eagle River from 2012 through 2015 (Table 5).

A small run of Chinook salmon and poor conditions did not allow for reliable Chinook salmon surveys (Appendix C2). The failure to enhance the fishery with hatchery releases, and typically poor fishing conditions with high, fast water during the season open to Chinook salmon fishing, all probably contributed to low angler effort and success. Anecdotal information and observations of fishery performance inseason suggest that the catch and harvest estimates will continue to remain low, but evidence indicates that illegal harvest is probably greater than the legal harvest.

Ship Creek

By far the largest Chinook salmon fishery in the AMA is the stocked Ship Creek fishery (Appendix B1). Angling effort targeting all species in Ship Creek peaked at 62,101 angler-days in 2000 (Table 4). From 2006 to 2015, the Ship Creek sport fishery produced an annual average catch and harvest of 2,526 and 1,437 Chinook salmon, respectively (Table 5). However, the most current 5-year average (2011–2015) of catch and harvest is now only 1,367 and 836 fish, respectively. Runs to Ship Creek are predicted to average about 5,000 Chinook salmon annually (Bosch 2010). Sufficient brood of 750 fish has been collected in most years since 2000 to meet egg-take goals and to produce not only the 315,000 smolt released specifically for Ship Creek, but to also provide for other Southcentral Chinook salmon smolt releases. Present regulations

combined with emergency order authority have allowed ADF&G to achieve these management objectives. In years where adequate hatchery fish are not returning, ADF&G will take action under Alaska Statute (16.05.060) to issue an emergency order allowing more fish to escape through the Ship Creek fishery and make it to the WJHSF Hatchery.

Broodstock needs were very minimal for 2010 because the new WJHSF Hatchery was not due to come on line until 2011. The new hatchery is able to once again produce 0-check smolt, and move away from rearing smolt for 2 years before release. The salmon not used for broodstock spawn naturally near the hatchery and provide an opportunity for the public to view spawning Chinook salmon. Historically, unofficial tallies at Elmendorf Hatchery (now the William Jack Hernandez Sport Fish Hatchery) have counted more than 60,000 viewers each year watching salmon in Ship Creek adjacent to the hatchery (WJHSF Hatchery Staff, personal communication).

An emergency order (EO 2-KS-2-29-12) was issued June 29, 2012, closing sport fishing on Ship Creek for the remainder of the Chinook salmon season in the sections open to the taking of Chinook salmon. The 2012 estimated final escapement in Ship Creek was 796 Chinook salmon (Appendix C1). In 2013, an emergency order (EO 2-KS-2-28-13) was issued on June 28 closing sport fishing on Ship Creek for the remainder of the Chinook salmon season in the sections open to the taking of Chinook salmon. In 2014, an emergency order (EO 2-KS-2-34-14) was issued on July 3 closing sport fishing on Ship Creek for the remainder of the Chinook salmon season in the sections open to the taking of Chinook salmon. In 2014, the escapement of Chinook salmon was estimated at 1,471 fish (Appendix C1). No emergency orders were issued in 2015 and the escapement of Chinook salmon in Ship Creek was 2,447 fish. This was the highest escapement since 2003 (2,733 fish). The average escapement the previous 10 years (2006–2015) was 1,379 Chinook salmon. Adding the average escapement (including brood stock; Appendix C1) and the estimated harvest (Table 5) for the previous 5 years (2006–2015), gives a conservative estimated annual run of about 2,816 (1,437 harvest + 1,379 escapement = 2,816 run) adult Chinook salmon returning to Ship Creek.

CHUM SALMON FISHERIES

FISHERY DESCRIPTION

Chum salmon do not return in significant numbers to AMA streams. Anglers targeting pink and coho salmon harvest most of the chum salmon. In 2015, the AMA estimated chum salmon sport catch was 1,742 fish of which 194 were harvested (Table 6). The majority of the catch and harvest were from Bird Creek. Chum salmon are also harvested in California, Fish, Glacier, Indian, Peters, and Ship creeks, and Eagle, Eklutna, Placer, and Twentymile rivers. Although chum salmon harvests have remained relatively low, catches hit a peak of 6,540 in 2002, followed by 6,033 caught in 2008, which was significantly higher than the most recent 5-year average (2011–2015) of 3,018 fish (Table 6, Figure 5). Chum salmon are counted during Chinook salmon escapement surveys although no directed chum salmon counts are conducted by ADF&G staff (Appendices C1–C5). ADF&G does not currently monitor chum salmon escapements.

Table 6.—Chum salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.

Year	Bird Creek		Ship Creek		Twentymile R.		Other streams		Saltwater		Area total	
	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest
1996	2,416	232	497	155	384	33	926	187	299	0	4,522	607
1997	1,014	209	229	24	87	0	113	15	145	0	1,588	248
1998	3,628	236	508	93	342	17	474	111	202	56	5,154	513
1999	1,156	99	354	16	81	14	244	0	469	0	2,304	129
2000	2,549	158	853	73	48	0	389	73	97	36	3,936	340
2001	1,489	87	1,347	218	357	35	1,270	130	168	0	4,631	470
2002	3,056	406	807	66	1,400	0	1,238	0	39	0	6,540	472
2003	1,857	155	1,014	67	357	25	349	66	544	0	4,121	313
2004	1,233	117	516	44	103	70	305	75	28	0	2,185	306
2005	1,548	116	338	100	0	0	420	18	70	0	2,376	234
2006	2,952	203	677	26	253	0	545	13	0	0	4,427	242
2007	2,027	61	425	22	102	0	295	0	119	14	2,968	97
2008	2,505	239	351	0	926	0	2,236	29	15	15	6,033	283
2009	4,205	316	517	40	69	11	1,154	19	0	0	5,945	386
2010	1,345	148	288	57	903	32	1,316	98	0	0	3,852	335
2011	1,006	111	827	0	22	0	2,265	286	230	0	4,350	397
2012	894	98	140	0	342	0	373	39	0	0	1,749	137
2013	1,366	449	1,885	1,512	58	0	202	92	0	0	3,511	2,053
2014	2,166	309	491	199	0	0	1,067	20	13	13	3,737	541
2015	925	97	331	97	19	0	467	0	0	0	1,742	194
Average												
2011–2015	1,271	213	735	362	88	0	875	87	49	3	3,018	664
2006–2015	1,939	203	593	195	269	4	992	60	38	4	3,831	467

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

In those freshwaters of the AMA open to fishing for salmon, the bag and possession limits for salmon (other than Chinook salmon) 16 inches or longer is 3 per day and 3 in possession, only 2 of which may be coho salmon (see regulations for exceptions). The limits for salmon (other than Chinook salmon) under 16 inches are 10 per day, 10 in possession. In saltwater, the limits for salmon, (other than Chinook salmon) are 6 per day, 6 in possession. Only 3 per day, 3 in possession may be coho salmon.

FISHERY MANAGEMENT AND OBJECTIVES

There are no formal management objectives for chum salmon in the AMA. ADF&G has a constitutional mandate to manage on the principle of sustained yield. Within the sustained yield principle, SF goals seek to optimize social and economic benefits, and where possible, expand opportunity to participate in diverse fisheries on these stocks.

STOCKING PROGRAM

There is currently no chum salmon stocking program in the AMA.




Figure 5.—Chum salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

FISHERY PERFORMANCE

Chum salmon catch and harvest has varied year to year. Specifically, sport caught chum salmon numbers have been on the decline since 2008 with 2012 and 2015 seeing nearly identical low catches of chum salmon (Figure 5). Chum salmon harvest has been steady over the last 10 years; however, a significant harvest of chum salmon was observed in 2013, specifically in Ship Creek. On average (2006–2015), 13% of chum salmon caught in AMA streams are harvested; however, in 2013, chum salmon harvest was 58% in AMA streams (Table 3).

COHO SALMON FISHERIES

FISHERY DESCRIPTION

Coho salmon fisheries in the AMA include the centrally located urban opportunities for anglers at Ship and Campbell creeks to remote, boat accessible angling opportunities at Twentymile and Placer rivers in Turnagain Arm. The largest coho salmon sport fisheries in the AMA include the 3 hatchery enhanced streams (Bird, Campbell, and Ship creeks) and the wild run returning to Twentymile River. Other drainages support smaller runs of wild coho salmon including the Placer River. Ship and Bird creeks contribute significantly to the overall AMA average catch and harvest of coho salmon (Table 7).

Table 7.—Coho salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.

Year	Bird Creek		Campbell Creek		Ship Creek		Twentymile River		Saltwater		Other freshwater		Area total	
	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest
1996	9,675	6,934	3,474	1,458	7,710	5,369	3,767	2,597	325	238	2,775	1,199	27,726	17,795
1997	9,097	6,771	3,006	1,651	13,448	9,413	2,664	1,332	479	243	1,498	1,168	30,192	20,578
1998	33,546	22,406	2,624	1,167	21,733	14,049	3,354	2,541	1,267	860	1,998	1,196	64,522	42,219
1999	6,284	4,611	1,880	1,341	7,064	4,649	1,457	1,051	170	158	979	456	17,834	12,266
2000	15,799	10,741	1,873	555	20,890	11,858	5,025	3,094	591	288	2,710	1,655	46,888	28,191
2001	11,563	8,449	2,748	813	39,615	26,419	5,724	2,742	732	463	3,483	1,807	63,865	40,693
2002	1,504	1,053	2,998	1,144	24,699	16,751	4,101	2,672	1,012	587	6,905	4,053	41,219	26,260
2003	1,117	776	2,873	1,457	8,831	6,094	3,039	2,116	867	441	4,035	2,491	20,762	13,375
2004	1,064	611	3,468	1,056	10,543	6,110	5,048	3,012	336	143	5,015	2,515	25,474	13,447
2005	5,331	3,281	4,552	1,989	10,922	6,830	1,632	1,334	473	473	3,027	1,156	25,937	15,063
2006	9,530	5,889	3,622	1,767	14,881	8,079	2,299	1,739	316	220	5,172	2,169	35,820	19,863
2007	7,461	3,287	1,051	758	5,845	3,934	998	719	993	915	1,458	1,079	17,806	10,692
2008	3,817	3,030	2,164	1,155	8,755	6,735	7,336	4,116	0	0	4,052	2,960	26,124	17,996
2009	6,020	3,296	577	364	4,014	2,974	2,052	1,329	284	207	4,789	2,635	17,736	10,805
2010	1,381	974	392	249	1,038	743	1,358	1,214	44	15	1,770	1,271	5,983	4,466
2011	2,024	1,324	784	502	3,748	2,443	1,281	1,087	55	55	2,435	1,994	10,327	7,405
2012	1,120	722	425	306	2,289	1,312	939	639	305	46	1,457	1,162	6,535	4,187
2013	1,251	980	168	136	3,733	3,264	1,304	895	0	0	1,185	915	7,641	6,190
2014	3,936	2,751	2,409	1,303	4,235	2,949	1,116	784	198	198	2,045	1,445	13,939	9,430
2015	4,712	4,097	2,993	1,908	9,309	7,232	685	503	125	104	1,394	1,255	19,218	15,099
Average														
2011–2015	2,609	1,975	1,356	831	4,663	3,440	1,065	782	137	81	1,703	1,354	11,532	8,462
2006–2015	4,125	2,635	1,459	845	5,785	3,967	1,937	1,303	232	176	2,576	1,689	16,113	10,613

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Once coho salmon leave the freshwater as smolt, they spend 1 winter out in the ocean prior to returning to their natal streams (Sandercock 1991). Coho salmon return to AMA streams from mid-July through mid-October. Hatchery-stocked stream runs peak in mid-August, whereas wild Turnagain Arm runs peak in mid-September.

In the fresh waters open to fishing for coho salmon, bag and possession limits for salmon other than Chinook salmon 16 inches or greater in length are 3 per day and 3 in possession, only 2 of which may be coho salmon except in stocked streams. In stocked streams like Bird, Campbell, and Ship creeks, the bag and possession limits for salmon other than Chinook salmon are 3 per day and 3 in possession of which all 3 can be coho salmon.

Beginning in 2011, a coho salmon 16 inches or longer that is removed from fresh water must be retained and becomes part of the bag limit of the person who originally hooked the fish.

Bird Creek

Little historical information is available regarding Bird Creek coho salmon. The first foot surveys were conducted in 1986 and 3 coho salmon were observed. Foot surveys from 1990 through 1992 indicated escapements ranging from 10 to 100 coho salmon (Stratton and Cyr 1995). The well-developed access, proximity to Anchorage, and lack of natural coho salmon production makes Bird Creek an ideal candidate for coho salmon enhancement. Bird Creek is open to sport fishing from the ADF&G markers approximately 500 yards upstream of the Seward Highway Bridge downstream to the mouth of the creek. A marker approximately 400 yards upstream of the Seward Highway Bridge marks the public-private property boundary. The open area was expanded in 1993 by approximately 100 yards as a result of a pending land exchange between the Alaska Department of Natural Resources (DNR), Municipality of Anchorage (MOA), and private landowners. To date, this land exchange has not occurred. Once the land swap is completed, the marker will be moved upstream. Bird Creek upstream of this reach is closed to all salmon fishing. The area open to salmon sport fishing in Bird Creek is within the intertidal reach, and coho salmon are harvested from late July through mid-September. The remainder of the drainage is closed year-round to salmon fishing.

In 2005, construction of the new parking and access project was completed. The enhanced coho salmon fishery had grown in popularity since the initial return in 1993 and quickly outgrew existing facilities for anglers. This project provided off-road parking for over 125 vehicles and developed camping, safe access for anglers and spectators, and sanitation facilities.

Campbell Creek

Although wild coho salmon historically returned to Campbell Creek in August and September, the number of returning adults was insufficient to support a viable sport fishery. Currently, coho salmon start to enter Campbell Creek in mid-July and are observed during annual ADF&G Chinook salmon surveys. This component of the Campbell Creek coho salmon run does not stay long in the area of Campbell Creek below Piper Street (the only area open to salmon fishing). These fish all appear to go up the North Fork of Campbell Creek above Campbell Airstrip Road (Bosch 2010). Coho salmon that return in August tend to hold in Campbell Lake, which is closed to all fishing, before moving into the creek. Most fish migrate upstream of Lake Otis Parkway and spawn in both the North and South Forks of Campbell Creek. From 1986 to 1992, before the returns of hatchery fish commenced, Campbell Creek coho salmon escapement surveys averaged 159 fish annually (Stratton and Cyr 1995). The reduction of Campbell Creek coho salmon runs

was probably a result of urbanization and development along the creek that reduced the number and size of wetlands and associated rearing habitat, an influx of pollutants and silt from storm drain runoffs, and poaching. A combination of foot and float escapement surveys are performed to count spawning coho salmon in Campbell Creek.

Campbell Creek was opened to coho salmon fishing in 1993 for the first time since 1971 (Appendix A3). A map of lower Campbell Creek showing areas open to coho salmon sport fishing can be found in Figure 6. Campbell Creek, from Dimond Boulevard to Shelikof Street, and Lake Otis to the Forks located near Piper Street, is open to coho salmon fishing from July 14 through September 30 each year. The bag and possession limits on Campbell Creek for salmon (other than Chinook salmon) 16 inches or greater in length are 3 per day, 3 in possession, of which all 3 may be coho salmon.




Figure 6.—A map of lower section of Campbell Creek showing areas open to coho salmon sport fishing and season dates.

Ship Creek

Prior to the start of World War II in 1939, Ship Creek had a wild run of coho salmon that supported sport, personal use, and subsistence fisheries (Bosch 2010). The dams constructed in the lower 11 miles of the creek were built for power generation and as a water source for the Municipality of Anchorage (MOA). The construction of these dams as well as the development of military installments during the 1940s and 1950s coincided with reduced Ship Creek salmon

runs. To rebuild the coho salmon runs, smolt were stocked annually into Ship Creek (ADF&G SF hatchery records).

Ship Creek was open to coho salmon sport fishing from 1957 through 1959, and again from 1964 to present (Appendix A1). Currently, only the reach downstream of the Chugach Power Plant Dam is open to salmon fishing Figure 4. Fishing is also closed within 100 ft of the Chugach Power Plant Dam. The bag and possession limits on Ship Creek for salmon (other than Chinook salmon) 16 inches or greater in length are 3 per day, 3 in possession of which all 3 may be coho salmon.

Turnagain Arm Streams

Turnagain Arm streams produce the largest wild stock coho salmon runs in the AMA; coho salmon return to several streams from late July through mid-September. In some systems, fresh fish are available into October.

The Twentymile River drainage supports the largest and most popular recreational coho salmon fishery in Turnagain Arm. Only about the first 10 miles of the Twentymile River are open to fishing after July 13. The Glacier River is only open to its confluence with Carmen River. The upper reaches of Twentymile River, Glacier River, and Carmen River are closed by regulation to sport fishing after July 14, and are always closed to salmon fishing. These rivers are heavily influenced by the summertime glacial runoff and fishing typically takes place in or near freshwater sloughs that feed these streams.

In the Placer River drainage, and Skookum and Lower Explorer creeks, sport fishing effort is minimal. Spawning areas in Lower Explorer and Skookum creeks are closed to sport fishing by regulation after July 14. Coho salmon are also harvested in California, Glacier, Ingram, Peterson, and Placer creeks and several Portage Valley streams.

In waters of Turnagain Arm open for fishing for coho salmon, the bag and possession limits for salmon (other than Chinook salmon), 16 inches or greater in length are 3 per day, 3 in possession. Only 2 may be coho salmon.

FISHERY MANAGEMENT AND OBJECTIVES

The Alaska Board of Fisheries has no established management plans or objectives for AMA coho salmon. The management objective and goal for wild stocks of coho salmon in systems considered too small to support a harvestable surplus is to maintain historical escapement levels, continue natural production, and provide viewing opportunities. Angler opportunity should be provided in those streams only when there is a harvestable surplus. Escapement goals have not been set for these streams. Angler participation and harvest of wild stocks have increased in many systems in recent years and should be monitored to ensure sustainability.

Campbell Creek

Campbell Creek was closed to all salmon fishing prior to 1993 when it was opened to fishing for coho salmon. A biological escapement goal of 200 fish was set for Campbell Creek coho salmon (it was changed to a sustainable escapement goal (SEG) of 100 to 500 wild coho salmon in 2001), and a weir was operated on Campbell Creek from 1993 to 1994 to count returning adult salmon. There is currently no escapement goal for coho salmon on Campbell Creek.

The Campbell Creek coho salmon fishery was established to provide additional angler opportunities in Anchorage by attempting to produce run of 3,500 adult coho salmon to Campbell Creek while maintaining historical levels of natural coho spawning. In addition, coho salmon enhancement was established to provide 5,000 angler-days of annual sport fishing opportunity directed at stocked coho salmon in Campbell Creek. The fishery will continue to be managed to maintain historical escapement levels and to provide continued natural production and viewing opportunities (ADF&G Statewide Stocking Plan; <http://www.adfg.alaska.gov/index.cfm?adfg=fishingSportStockingHatcheries.stockingPlan>, accessed January 2017).

Ship Creek

The Ship Creek coho salmon fishery management objectives are identical to the Statewide Stocking Plan. The objectives are to produce a return of 12,000 adult coho salmon to Ship Creek while assuring approximately 1,000 coho salmon are available at Ship Creek for natural spawning, fish viewing, and egg-take needs and to generate at least 35,000 angler-days of annual sport fishing opportunity directed at stocked Chinook and coho salmon in Ship Creek. (ADF&G Statewide Stocking Plan).

Present regulations provide for the harvest of coho salmon in excess of spawning and viewing requirements and allow optimum utilization of Ship Creek coho salmon. In 2014, an emergency order (EO 2-SS-2-48-14) was issued to increase harvest to 5 salmon, of which all 5 may be coho salmon in the sections open to salmon fishing. No other emergency orders were issued to liberalize or restrict the coho salmon fishery in 2011–2013 or 2015.

Turnagain Arm Streams

The management objective for the Turnagain Arm coho salmon fisheries is to provide angler opportunities while ensuring adequate spawning escapement. No escapement goals have been set for these systems. Large areas of streams at the head of Turnagain Arm are closed to all sport fishing after July 13 to protect spawning salmon.

STOCKING PROGRAM

Stocking of coho salmon into anadromous streams in the Anchorage Management Area is referred to as the *Urban Coho Salmon Stocking Program*. The primary purpose of the *Urban Coho Salmon Stocking Program* is to maintain or increase coho salmon sport fishing opportunities in Anchorage on a sustainable basis by supplementing natural runs with hatchery fish.

Prior to the *Urban Coho Salmon Stocking Program*, which was initiated in 1991, the greatest total coho salmon harvest (6,730 fish) occurred in 1988 (Mills 1989). On average (2006–2015), an estimated 16,113 coho salmon were caught annually in AMA sport fisheries and 10,613 of those fish were harvested (Table 7). The peak catch between 2006 and 2015 was 35,820 coho salmon caught in 2006 (Table 7, Figure 7). Approximately 420,000 smolt are stocked in Bird, Campbell, and Ship creeks annually. Coho salmon returning to Ship Creek are used for the broodstock although their origin is most likely from Little Susitna River coho salmon (Mark Tag Lab Database; <http://mtalab.adfg.alaska.gov/CWT/reports/hatcheryrelease.aspx>, accessed January 2017). In 2012, excessive mortality occurred during early rearing of coho salmon at the hatchery. As a result, Ship Creek received more smolt than Bird or Campbell creeks to ensure returning broodstock goals were met at the hatchery.

The increase in the AMA coho salmon sport harvest from 2013 to 2015 is directly related to increased sport fishing effort on Ship, Campbell, and Bird creeks, which had been enhanced with hatchery stocking, and due to increased effort on Twentymile River, which has natural stocks.




Figure 7.—Coho salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Bird Creek

Bird Creek historically produced few coho salmon, leaving no genetic concerns with stocking, and no attempts were made to collect wild Bird Creek coho salmon for broodstock. Thus, Little Susitna River coho salmon broodstock (now collected at WJHSF Hatchery site) have been and are currently used for Bird Creek stocking. Stocking of coho salmon smolt was initiated at Bird Creek in 1992 to increase the number of coho salmon for sport anglers because natural production is very low. The annual stocking of about 150,000 coho salmon smolt of Little Susitna River origin was initiated to increase the number of coho salmon for sport anglers. Smolt released in 1992 returned as adults to the Bird Creek fishery in 1993. Due to hatchery surplus, nearly 300,000 coho salmon smolt were released in 1997. Coho salmon smolt releases were suspended from 2001 through 2003 while the Bird Creek parking area was under construction. In the past 5 years (2011–2015), the average number of stocked coho salmon smolt was 108,132 (Appendix B2) but stocking varied between 136,047 smolt in 2011 and 70,004 smolt in 2012 due to an excessive die-off of coho salmon smolt at the hatchery. This stocking is part of the *Urban Coho Salmon Stocking Program* aimed at increasing coho salmon angling opportunities in the AMA.

Campbell Creek

Hatchery-produced coho salmon runs in Campbell Creek provide a unique opportunity for sport anglers to participate in quality fisheries in an urban setting. The annual stocking of 150,000 coho salmon smolt of Little Susitna River origin was initiated in 1992 to increase coho salmon runs to Campbell Creek. This stocking is part of the *Urban Coho Salmon Stocking Program* aimed at increasing coho salmon angling opportunities in the AMA. The number of coho salmon smolt stocked annually was reduced to about 76,000 in 1996, and 75,000 remains the current stocking goal. Between 2011 and 2015, excluding 2012, the average number of stocked coho salmon smolt was 59,047. In 2012, no smolt were released into Campbell Creek due to an excessive die-off coho salmon smolt at the hatchery.

Ship Creek

Hatchery-produced coho salmon runs in Ship Creek also provide a unique opportunity for sport anglers to participate in quality fisheries in an urban setting. The primary purpose of the stocking program is to maintain or increase coho salmon sport fishing opportunities in Anchorage on a sustainable basis by supplementing Ship Creek's natural run with hatchery fish.

The construction of dams in the 1940s and 1950s on Ship Creek made it difficult for wild coho salmon runs. To enhance and reestablish the coho salmon runs after the construction of the dams, the creek was stocked annually with coho salmon smolt from 1968 through 1977 (ADF&G SF hatchery records). These efforts proved unsuccessful in providing consistent numbers of returning adults. Nine different broodstocks from Ship Creek, Bear Lake (near Seward), Kodiak, Washington, and Oregon were used (Miller 1990). Eggs obtained from these stocks were incubated at Fire Lake Hatchery (located on Upper Fire Lake in Eagle River) and the resultant fry were reared to smolt at Fort Richardson Hatchery. No coho salmon smolt were released in Ship Creek from 1978 through 1986. From 1987 through 1994, ADF&G stocked coho salmon smolt in Ship Creek using fish of Ship Creek origin reared at Elmendorf Hatchery. Although these efforts have provided consistent coho salmon runs, these runs tend to enter the system slowly throughout the fall. Ideally, coho salmon runs that appeal to sport anglers exhibit a compressed run timing with large numbers of fish available in a relatively short time period. The decision was made to change broodstock for Ship Creek to Little Susitna River origin fish, which exhibit the preferred condensed run timing. The first release of Little Susitna River origin coho salmon smolt occurred in 1995. The current coho salmon run is the result of an annual release of about 240,000 smolt raised at the WJHSF Hatchery.

Turnagain Arm Streams

With the exception of the Bird Creek coho salmon enhancement program described above, there are currently no coho salmon stocking programs in Turnagain Arm.

FISHERY PERFORMANCE

Poor runs of coho salmon in many systems in Upper Cook Inlet from 1997 until 2000 resulted in closures of the Upper Cook Inlet commercial fisheries, reductions to bag and possession limits, bait restrictions to the sport fisheries, and a conservative approach to coho salmon management regionwide. These years of weak runs were followed by record runs from 2000 through 2002. The 2000 through 2002 estimated catches of coho salmon in the AMA are 3 of the 4 highest years ever recorded. The 2003 through 2009 estimated catches remained relatively good,

averaging a catch and harvest of 24,237 and 14,463, respectively; however, the areawide catch of 5,983 coho salmon in 2010 was the lowest on record since the early 1990s (Table 7). From 2011 to 2013, the coho salmon runs continued to be weak, although from 2013 to 2015, the coho salmon runs have been steadily increasing (Figure 7). The average annual catch in the last 5 years (2011–2015) was 11,532 coho salmon with 2014 and 2015 having the highest catches of those 5 years.

Bird Creek

The angler effort in Bird Creek in the last 5 years (2011–2015) has averaged 6,303 angler-days (Table 4), which is 2,889 angler-days lower than the 10-year (2006–2015) average of 9,192 angler-days. Catch and harvest of coho salmon have been low as a result of decreased angling effort (Figure 7).

The first returns from hatchery stockings in 1993 resulted in a catch and harvest of 7,799 and 6,195 coho salmon, respectively (Bosch 2010). In 1998, the fishery peaked with a catch and harvest of 33,546 and 22,406 coho salmon, respectively (Table 7). Fishery performance is linked closely to the number of properly-sized hatchery-reared smolt that are released. Nearly 300,000 coho salmon smolt were released in 1997 (Appendix B2) resulting in high catch rates the following year. The fishery continued to do well until 2002, when the catch dropped to 1,504 coho salmon of which 1,053 were harvested. This decline in fishery performance was due to Bird Creek not being stocked in 2001. Construction of the new parking area north of Bird Creek was scheduled to begin in 2002, and in order to conduct a safe and orderly fishery, Bird Creek was not stocked from 2001 through 2003. The loss of the stocking program for those 3 years corresponded to poor fishery performance from 2002 through 2004 (Table 7).

Between 2011 and 2015, the number of stocked coho salmon smolt ranged from 70,004 (2012) to 136,047 (2011) and the 5-year average (2011–2015) was 108,132 coho salmon smolt. However, the sport catch and harvest did not materialize as expected during these years. In 2010, 157,534 coho salmon smolt were released and the catch and harvest on those fish returning as adults in 2011 was only 2,024 and 1,324, respectively (Table 7). In addition, angler effort in 2011 was low at 5,027 angler days (Table 4). The stocking level in 2010 was similar to 1998 (Appendix B2), but catch and harvest the following year, 1999, was 6,284 and 4,611, respectively, and effort was 13,033 angler days (Tables 4 and 7). The coho salmon smolt from the new WJHSF Hatchery ranged from 18 g to 23 g at release (Mark Tag Lab Database, <http://mtalab.adfg.alaska.gov/CWT/reports/hatcheryrelease.aspx>) and were in good health.

The release location on Penguin Creek, a tributary of Bird Creek, was considered a possible cause for increased mortality. Prior to the new hatchery, fish were stocked approximately 0.5 miles up Penguin Creek on a small access road. A new stocking release site was utilized for fish from the new hatchery, about another river mile upstream on Penguin Creek. This new location was selected to provide smolt more time to imprint in the fast flowing freshwater of Penguin Creek. In 2014, a foot survey was conducted immediately after the coho salmon smolt were released, and it was determined that this additional upper section of Penguin Creek was very steep and may have contributed to increase mortality; however, no actual mortalities were observed on the foot survey. In 2015, stocked fish were divided into 3 different release locations and observed closely to try to determine the best release location for the future. One group of coho salmon smolt was released at the highest location, which was approximately 1.39 miles up Penguin Creek from the confluence with Bird Creek. The second group of coho salmon smolt

was released at more traditional (2009 and earlier) site, which is about 0.5 mile up Penguin Creek, and the final release location and the smallest group of smolt was released at a bridge about a 0.25 miles above the fishery on the mainstem of Bird Creek but just below the confluence of Bird and Penguin creeks. These 3 locations are the only locations where the stocking truck can realistically intersect with Penguin or Bird creeks. The lowest stocking location on Bird Creek was determined to be too close to the tidally influenced saltwater and the fish were swept out by the tide within an hour. It appears that the middle stocking location, which is approximately 0.5 miles up Penguin Creek, is the most suitable stocking location and will continue as the traditional stocking location until further assessments can be made. This section has less of a gradient than the upper stocking location and released smolt were seen holding in this section by observers.

The Bird Creek coho salmon fishery relies largely on hatchery enhancement, and attempts to estimate escapement during the previous 5 years (2011–2015) have only been successful twice (45 adult coho salmon were estimated in 2011 and 629 in 2014; Appendix C4) making it difficult to rely on these surveys as a meaningful index. This is a very short system and is subject to poor water clarity, high swift water, and high bear densities that can make survey conditions difficult and often unsafe.

Campbell Creek

The proximity of Campbell Creek within Anchorage makes it easy to access and fish. Even when hatchery runs are anticipated to be low, fishing effort remains fairly steady. The effort on Campbell Creek in the last 10 years has averaged 4,167 angler-days (Table 4) and is 486 angler-days higher than the average effort from the last 5-years. Effort decreased slightly in 2012, dropping to a historical low of 2,183 angle-days; however, a decrease in effort was common in other AMA streams that year and was followed by 3 years (2013–2015) of increased effort of at least 4,000 angler-days.

In Campbell Creek, the average catch and harvest of coho salmon for the last 5 years (2011–2015) was 1,356 and 831, respectively, both of which were less than the 10-year averages (1,459 and 845, respectively) (Table 7). Stocking in Campbell Creek has ranged from 0 coho salmon smolt in 2012 to 83,088 coho salmon smolt in 2013. Coho salmon released as smolt will spend one winter in the ocean prior to returning to spawn as adults.

In 2013, angler effort on Campbell Creek increased by 2,088 angler-days from the prior year (Table 4), although catch and harvest of coho salmon reflected the absence of stocking in 2012, with only 168 fish caught and 136 fish harvested (Table 7). In 2013, 83,088 coho salmon smolt were stocked (Appendix B2), and in 2014, these returned to spawn as adults. In 2014, the effort (4,295 angler-days) remained similar to the previous year (Table 4), but catch and harvest (2,409 and 1,303, respectively), reflected the increase in stocked coho salmon from 2013 (Table 7; Appendix B2). In 2014, stocking was reduced to 29,028 coho salmon smolt (Appendix B2). The 2015 catch and harvest (reflective of the 2014 stocked fish) was 2,993 and 1,908 fish, respectively (Table 7) and higher than anticipated. The harvest and catch levels in 2014 were similar to the levels in 2015, although the smolt released in 2014 (29,028) was a 33% of the smolt released in 2013 (83,088). In 2015, 52,110 coho salmon smolt were released into Campbell Creek (Appendix B2). The smolt released in 2015 will be returning as adults in 2016.

In 2011 and 2012, no streams surveys were performed due to reductions in staff and poor sampling conditions. From 2013 to 2015, stream surveys were resumed. In 2013, only 146 fish

were counted on the annual survey (Appendix C3). In 2014, 3,316 adult coho salmon were counted and in 2015, 1,178 coho salmon were counted (Appendix C3). Estimates from these surveys are consistent with angler reports of coho salmon abundance during that time and probably reflect the escapement of coho salmon in Campbell Creek.

Combining the average estimated harvest (2006–2015) of 845 fish (Table 7) and average stream surveys estimates (2006–2015) of 930 fish (excluding 2007, 2011, and 2012, when no surveys were performed; Appendix C3), the annual total run size of adult coho salmon for Campbell Creek would be estimated at 1,775 fish (845 harvest + 930 escapement = 1,775 run).

Ship Creek

Effort, and catch and harvest of coho salmon in Ship Creek remain the highest out of any stream in the AMA even though overall effort has decreased since the late 2000s and early 1990s (Tables 4 and 7). On average during the past 10 years (2006–2015), anglers have spent 22,719 angler-days of effort in Ship Creek (Table 4). This is slightly higher than the recent 5-year average (2011–2015) of 15,911 angler-days. This decrease was similar to other AMA streams and reached a historical low in 2012.

The average (2006–2015) catch and harvest of coho salmon in Ship Creek was 5,785 fish and 3,967 fish, respectively (Table 7). From 2011 to 2015, coho salmon catch and harvest was 4,663 and 3,440 fish, respectively. This was 1,122 and 527 fish, respectively, lower than the 10-year average (2006–2015). The catch and harvest of coho salmon in 2009 through 2014 were historical lows and averaged 3,176 fish and 2,281, respectively (calculated from Table 7).

The 10-year (2006–2015) average annual number of coho salmon smolt stocked in Ship Creek (254,116) has been fairly constant and has ranged from a low of 243,499 (2012) to high of 287,825 (2009) fish (Appendix B2). Even with consistent stocking numbers back to 1996 (Appendix B2), the average catch and harvest of coho salmon in the past 5 years are much less than most years in the late 1900s and early 2000s (Table 7).

It is difficult to estimate the entire coho salmon run size for Ship Creek. Stream surveys and hatchery counts do not always coincide. The stream survey counts need good visibility and suitable water conditions. Some fish counted in stream surveys may be counted again at the hatchery. If the hatchery needs additional fish to meet broodstock goals, they may catch fish near the hatchery, or if the hatchery has too many fish, they may return fish to the creek. These are all potential issues with stream foot survey data on Ship Creek. In recent years, SF biologists for the AMA have been working more closely with the hatchery staff to try to get a better end-of-season escapement estimate by combining both hatchery-counted fish and fish observed on the foot survey. Although rain is common in the fall, causing increased water levels and turbidity making accurate stream surveys difficult. The 10-year (2006–2015) average annual estimated escapement for Ship Creek is 2,076 adult coho salmon (Appendix C1). If the 10-year average annual harvest of 3,967 coho salmon (Table 7) is added to the estimated 10-year average annual escapement, the conservatively estimated annual drainage escapement is 6,043 coho salmon (3,967 harvest + 2,076 escapement = 6,043 run size).

Turnagain Arm Streams

Besides Bird Creek (See Bird Creek coho salmon section) and the Twentymile River drainage, a majority of the sportfishing effort for coho salmon on Turnagain Arm streams is small and not reported in the annual Statewide Harvest Survey (SWHS). Twentymile River and other

Turnagain Arm streams are wild coho salmon runs. The only river that does show up consistently in the SWHS results is Twentymile River. Twentymile River includes the entire Twentymile River drainage of which only the lower portion is open to coho salmon fishing (see Turnagain Arm Streams fishery description for more detail). The average annual effort in Twentymile River makes up 5% of the overall effort in the AMA (Table 4). It is difficult to participate in this coho salmon fishery without a boat. The average annual effort for the last 10 years (2006–2015) was 2,075 angler-days. This is 603 angler-days greater than the 5-year average (2011-2015) of 1,472 angler-days. This decrease in effort on Twentymile River is similar to other AMA streams. In 2015, a historically low angler effort was reported in the SWHS with only 989 angler-days spent fishing on Twentymile River (Table 4). In 2015, catch and harvest of coho salmon also hit a historical low with only 685 fish and 503 fish, respectively (Table 7). The 10-year (2006–2015) average annual catch and harvest of coho salmon was 1,937 fish and 1,303 fish, respectively.

Currently no escapement surveys are performed on the Twentymile River drainage. Historical aerial streams survey numbers and reasons for discontinuing the surveys are detailed in the 2010 Anchorage Management Report (Bosch 2010).

PINK SALMON FISHERIES

FISHERY DESCRIPTION

Pink salmon return annually to AMA streams in July and August, although the largest runs tend to occur in even-numbered years. However, since 2004, every odd-numbered year has observed a higher catch than the previous even-numbered year. In some cases, the differences were very significant. For example, in 2008, 30,004 pink salmon were caught followed by a catch of 62,996 in 2009 (Table 8). The 10-year average (2006–2015) catch was 23,143 pink salmon and 2,534 were harvested (Table 8). Historically, pink salmon fisheries had the second highest annual salmon catch in the AMA after coho salmon; however, a shift began in 2007 and annual catch of pink salmon has been greater than coho salmon in almost every year since then (Table 3). In 2015, the catch was nearly the same for both pink and coho salmon (19,236 and 19,218 fish, respectively (Table 3).

The majority of pink salmon caught in AMA streams are released (74–94% since 1996) and Bird Creek supports the largest pink salmon sport fishery in the AMA with an average (2006–2015) catch and harvest of 13,772 and 1,532 pink salmon, respectively (Table 8). Bird Creek flows into Turnagain Arm approximately 25 miles south of Anchorage and supports the primary AMA pink salmon sport fishery. Improvements in parking areas and access trails have increased Bird Creek's popularity as a fishing destination for both local and nonresident anglers. Pink salmon return to Bird Creek from mid-July to mid-August each year. Although the pink salmon stocks in Bird Creek are wild, beginning in 1992, ADF&G began releasing hatchery-reared coho salmon smolt, making Bird Creek even more popular with most anglers. Bird Creek is open to sport fishing from ADF&G markers located approximately 500 yards upstream of the Seward Highway Bridge and downstream to the mouth of the creek. The open area was expanded in 1993 by approximately 100 yards as a result of a pending land exchange between the Alaska Department of Natural Resources (DNR), the Municipality of Anchorage (MOA), and private landowners. To date, this land exchange has not occurred. Bird Creek upstream of this reach is closed to all salmon fishing.

Table 8.—Pink salmon sport fish catch and harvest, Anchorage Management Area, 1996–2015.

Year	Bird Creek		Ship Creek		Twentymile R.		Saltwater		Other freshwater		Area total	
	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest
1996	14,218	2,419	1,258	214	1,464	91	460	59	1,789	79	19,189	2,862
1997	3,038	535	1,484	80	72	32	128	9	398	187	5,120	843
1998	18,595	1,941	5,152	454	301	34	1,654	253	2,436	523	28,138	3,205
1999	3,913	507	789	80	80	0	185	40	495	94	5,462	721
2000	20,055	1,335	6,841	853	297	10	963	348	10,080	577	38,236	3,123
2001	7,662	333	2,815	190	234	23	321	64	1,956	173	12,988	783
2002	5,931	758	2,724	155	709	29	53	7	2,234	219	11,651	1,168
2003	6,152	1,033	1,055	291	109	0	997	150	1,148	126	9,461	1,600
2004	2,677	751	1,668	61	163	16	884	81	2,505	363	7,897	1,272
2005	8,624	433	839	108	93	46	85	24	1,098	66	10,739	677
2006	16,182	1,574	1,755	288	384	0	108	0	5,482	483	23,911	2,345
2007	25,861	1,717	3,559	405	299	88	1,287	743	3,312	325	34,318	3,278
2008	16,205	1,088	1,492	191	2,043	63	140	140	10,124	550	30,004	2,032
2009	37,299	3,812	7,620	785	185	0	159	74	17,733	1,755	62,996	6,426
2010	6,956	1,433	914	377	450	27	80	14	6,225	727	14,625	2,578
2011	5,366	339	5,268	21	292	56	85	85	4,230	392	15,241	893
2012	2,834	508	1,387	92	432	134	0	0	3,934	754	8,587	1,488
2013	3,657	672	4,085	1,734	60	0	0	0	1,766	41	9,568	2,447
2014	9,314	944	1,532	99	287	0	73	73	1,735	357	12,941	1,473
2015	14,047	1,435	1,932	378	58	0	0	0	3,199	569	19,236	2,382
Average												
2011–2015	7,044	780	2,841	465	226	38	32	32	2,973	423	13,115	1,737
2006–2015	13,772	1,352	2,954	437	449	37	193	113	5,774	595	23,143	2,534

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Most AMA streams support annual pink salmon runs, but run size is often small and doesn't support a recreational harvest. Other area streams with significant pink salmon catch and some harvest include Ship Creek and Twentymile River. Reported harvests from California, Fish, Glacier, Indian, Ingram, and Peters creeks, and Eagle, Eklutna, and Placer rivers are included in "other freshwater" on Table 8.

In those waters of the AMA open to fishing for salmon, the bag and possession limits for salmon (other than Chinook salmon) 16 inches or longer is 3 per day, 3 in possession, only 2 of which may be coho salmon (see regulations for exceptions). The limits for salmon (other than Chinook salmon) under 16 inches are 10 per day, 10 in possession. Rabbit and Sixmile creeks are closed to all salmon fishing and Campbell Creek is closed to pink salmon fishing.

FISHERY MANAGEMENT AND OBJECTIVES

There are no formal management objectives for pink salmon in the AMA. ADF&G has a constitutional mandate to manage on the principle of sustained yield. Within the sustained yield principle, SF goals seek to optimize social and economic benefits, and where possible, expand opportunity to participate in diverse fisheries on these stocks.

STOCKING PROGRAM

There is currently no pink salmon stocking program in the AMA.

FISHERY PERFORMANCE

Bird, Ship, and Twentymile creeks provide the only pink salmon fisheries of any significance in the AMA. All other pink salmon fisheries are included in "other freshwater" in Table 8. Fishery performance for all AMA pink salmon fisheries has improved since a significant drop in 2012. In addition, the majority of the catch continues to be released (Table 8, Figure 8).




Figure 8.—Pink salmon sport fish catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

The 5-year (2011–2015) average catch and harvest of pink salmon in the AMA was 13,115 and 1,737, respectively (Table 8). This is a significant drop from the 10-year catch and harvest averages of 23,143 and 2,534, respectively. With the exception of Ship Creek, catch and harvest of pink salmon at all locations showed a decline when looking at the 5-year (2011–2015) versus 10-year averages (2006–2015). Ship Creek catch estimates alternated between high and low catches year to year until 2015, when catch increased slightly from what was observed in 2014 (Table 8).

The average annual pink salmon catch and harvest at Bird Creek from 2011 to 2015 (7,044 and 780, respectively) were below the historical 10- and 20-year average catch (13,772 and 11,429, respectively) and harvest (1,352, and 1,178, respectively), and they are well below peak catches that took place in 2007 (25,861 fish), 2009 (37,299 fish), and prior to 2000 when catch estimates for Bird Creek ran as high as 20,060 pink salmon (Table 8). The 5-year (2011–2015) average catch and harvest of pink salmon on Bird Creek accounts for about 50% of the pink salmon catch and harvest in the AMA on average (calculated from Table 8). However, from 2011 to 2013, the harvest of pink salmon from Bird Creek dropped with just over 30% of the harvest for the AMA coming from this location. In 2013, pink salmon catch was highest in Ship Creek and in 2011, Bird, Ship, and “other freshwater” streams had very similar catches. In 2012 “other freshwater” streams had a higher estimated catch than Bird and Ship creeks.

The sport catch and harvest of pink salmon in “other freshwater” AMA creeks from 2011 to 2015 was 2,973 and 423, respectively (Table 8). This is a decline from the 10-year average (2006–2015) sport catch and harvest that was 5,774 and 595, respectively. In 2009, “other freshwater” catch and harvest peaked at 17,733 and 1,755, respectively. This peak was also observed in Bird and Ship creeks (Table 8). Saltwater catch and harvest of pink salmon is less than 1% of the total freshwater numbers of pink salmon caught and harvested in the AMA.

Stream surveys are not conducted specifically for pink salmon in any AMA drainage; however, in 2009, there was one conducted on Bird Creek and it was an unusually high run of pink salmon for an odd year in an Upper Cook Inlet stream (Appendix C4). No pink salmon catch or escapement of this magnitude has been observed in Bird Creek since 2009, and no additional escapement pink salmon-specific streams surveys have since been conducted.

SOCKEYE SALMON FISHERY

FISHERY DESCRIPTION

The primary Anchorage Management Area (AMA) streams that support sockeye salmon runs are Sixmile Creek and Twentymile River. Carmen Lake and its inlet tributaries are the primary sockeye salmon spawning areas in the Twentymile River drainage, and mainstem spawning has been documented (Stratton et al. 1994). Unlike Twentymile River, Sixmile Creek is presently closed to sport fishing, but fishing is allowed in the intertidal area below the high-tide mark near the creek mouth and in Lower and Upper Sixmile lakes. The intertidal site, marked with a steel cable across the stream and ADF&G markers, is growing in popularity and contributes to most of the AMA saltwater harvest. This fishery is probably even larger because some of the sockeye salmon reported in “saltwater fisheries” are probably harvested at the mouth of Sixmile Creek. Elmendorf Air Force Base (now called Joint Base Elmendorf-Richardson or JBER) personnel have operated a weir in Sixmile Creek since 1988. In 1998, this weir was moved upstream near the lake outlet. The Alaska Department of Fish and Game (ADF&G) operated this weir in 2009

and 2010 under contract with the Joint Base. Currently the weir is operated annually by JBER staff.

The most significant sockeye salmon fisheries are reported in “other freshwater” streams (probably Portage Valley streams) and from saltwater. Sockeye salmon returning to Placer River spawn in Luebner Lake. Sockeye salmon returning to Portage Creek primarily spawn in the artificially created channel in Williaw Creek where a viewing platform and information kiosk were installed and are maintained by U.S. Forest Service (USFS). Other AMA streams that support small sockeye salmon runs include Ship, and Campbell creeks, and Portage Valley streams. In Campbell Creek, most sockeye salmon spawn in North Fork and are thought to utilize beaver ponds for rearing. Campbell Creek has no natural lake system accessible to salmon, only the man-made lake near the creek mouth (Campbell Lake). An annual Chinook and coho salmon assessment via a combination of foot and float escapement surveys are performed and sockeye salmon are also enumerated in Campbell Creek. In the winter of 2006–2007 and again in 2010, the lake was drained to complete a dredging project to winterkill aquatic vegetation along the shoreline. It is unknown to what degree sockeye salmon utilize the lake for overwinter rearing, and escapements to this lake should be closely monitored. Sockeye salmon in Campbell Creek have similar run timing as Chinook salmon, so a combination of foot and float survey counts is used to estimate escapement.

In those freshwaters of the AMA open to fishing for salmon, the bag and possession limits for salmon (other than Chinook salmon) 16 inches or longer are 3 per day, 3 in possession, only 2 of which may be coho salmon (see ADF&G sport fishing regulations for exceptions; http://www.adfg.alaska.gov/index.cfm?adfg=fishregulations.sc_sportfish, accessed January 2017). In saltwater, the limits for salmon (other than Chinook salmon) are 6 per day, 6 in possession. Only 3 per day and 3 in possession may be coho salmon. Campbell Creek is closed to sockeye salmon sport fishing, and remains closed to all fishing downstream of Dimond Blvd.

FISHERY MANAGEMENT AND OBJECTIVES

There are no formal management objectives for sockeye salmon in the AMA.

STOCKING PROGRAM

ADF&G does not have a sockeye salmon stocking program in AMA waters.

FISHERY PERFORMANCE

The 2015 AMA estimated sockeye salmon sport fish catch was 342 fish of which 266 were harvested (Table 9, Figure 9). The 5-year average (2011–2015) catch and harvest was 385 and 178 sockeye salmon, respectively (Table 9). “Other freshwater” streams made up approximately 70% of the average catch and just over 50% of the average harvest of sockeye salmon in the AMA over the last 10 years, whereas approximately 20% of the average catch and 16% of the average harvest of sockeye salmon came from Twentymile River over the last 10 years.

Sockeye salmon escapement into Campbell Creek since 1996 has been 516 fish on average (calculated from Appendix C3). Survey data showed below-average escapements in 2011 and 2014 (434 and 368, respectively) but above-average escapement in 2015 (585 fish) (Appendix C3). Surveys were not conducted in 2012 and 2013 due to staff reductions and high water events (Appendix C3). No other formal sockeye salmon surveys are conducted on AMA streams.

Table 9.—Sockeye salmon sport catch (C) and harvest (H), Anchorage Management Area, 1996–2015.

Year	Bird Creek		Twentymile R.		Saltwater		Other freshwater		Area total	
	C	H	C	H	C	H	C	H	C	H
1996	467	184	233	26	185	134	1,464	540	2,349	884
1997	220	98	70	10	110	50	2,045	865	2,445	1,023
1998	574	448	84	61	595	363	847	399	2,100	1,271
1999	78	56	42	10	782	248	605	228	1,507	542
2000	678	446	42	0	59	38	403	53	1,182	537
2001	316	263	176	97	830	271	970	263	2,292	894
2002	0	0	288	95	91	77	315	158	694	330
2003	0	0	84	36	346	299	1,361	608	1,791	943
2004	0	0	291	88	328	110	393	88	1,012	286
2005	0	0	139	106	445	445	413	0	997	551
2006	0	0	22	11	11	11	297	58	330	80
2007	0	0	172	81	60	60	363	31	595	172
2008	0	0	275	0	150	136	1,294	87	1,719	223
2009	0	0	156	48	48	0	459	144	663	192
2010	0	0	180	32	209	110	229	51	618	193
2011	0	0	0	0	215	215	320	29	535	244
2012	0	0	66	33	0	0	154	31	220	64
2013	0	0	0	0	0	0	172	28	172	28
2014	0	0	155	0	20	20	481	268	656	288
2015	0	0	83	83	0	0	259	183	342	266
Average										
2011–2015	0	0	61	23	47	47	277	108	385	178
2006–2015	0	0	111	29	71	55	403	91	585	175

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.




Figure 9.–Sockeye salmon sport catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

DOLLY VARDEN–ARCTIC CHAR FISHERIES

FISHERY DESCRIPTION

Dolly Varden and Arctic char are 2 distinct fish species; however, anglers often have difficulties distinguishing between the 2 fish species. These fish are not stocked in AMA streams but Arctic char are raised at the WJHSF Hatchery and stocked in AMA lakes annually. For this section, the term “Dolly Varden” will be used but should be recognized that some of these fish called Dolly Varden could in fact be Arctic char.

Several area streams and lakes support small populations of resident Dolly Varden. Dolly Varden have been reported in Bird, Campbell, Chester, Ingram, and Ship creeks, and Placer and Twentymile rivers. On average, Campbell Creek has supported the largest catch estimates for the last 20 years (1996–2015) with 2,237 Dolly Varden caught and only 115 harvested (calculated from Table 10). In the last 10 years this has shifted and the “other streams” category has observed average catch estimates just under 1,000 Dolly Varden and average harvest estimates of 233, whereas Campbell Creek average catch and harvest estimates were 632 and 22, respectively (Table 10, Figure 10).

FISHERY MANAGEMENT AND OBJECTIVES

There are no formal management objectives for Dolly Varden in the AMA. No stream surveys are conducted to specifically count Dolly Varden.

Table 10.—Dolly Varden sport fish catch (C) and harvest (H), Anchorage Management Area, 1996–2015.

Year	Bird Creek		Campbell Creek		Eagle River		Ship Creek		Twentymile R.		Other streams		Lake total		Area total	
	C	H	C	H	C	H	C	H	C	H	C	H	C	H	C	H
1996	85	12	1,832	85	1,497	389	170	73	583	194	777	170	2,698	1,010	7,642	1,933
1997	162	14	938	0	804	378	256	32	1,154	243	836	364	1,736	559	5,886	1,590
1998	243	34	3,633	226	303	160	134	17	607	169	1,392	212	2,961	988	9,273	1,806
1999	90	0	1,693	626	814	97	44	22	190	99	331	89	1,362	1,092	4,524	2,025
2000	137	0	5,161	83	1,275	409	184	0	935	154	1,536	118	4,949	604	14,177	1,368
2001	22	0	12,760	238	87	0	648	33	2,027	189	767	151	947	290	17,258	901
2002	17	0	2,339	369	507	190	589	0	482	0	1,116	424	2,135	355	7,185	1,338
2003	70	50	2,568	228	820	0	536	10	702	153	3,257	439	2,851	1,205	10,804	2,085
2004	27	13	3,386	200	777	281	912	13	271	27	754	175	5,391	3,476	11,518	4,185
2005	39	13	4,116	35	953	0	584	0	260	81	391	142	3,634	1,087	9,977	1,358
2006	177	14	701	0	476	127	127	0	514	119	944	822	1,868	679	4,807	1,761
2007	59	0	710	15	225	115	599	0	1,177	218	970	252	2,849	667	6,589	1,267
2008	92	0	379	76	396	15	246	0	878	31	1,259	153	2,201	425	5,451	700
2009	79	33	198	0	0	0	28	0	1,174	57	1,272	384	1,797	653	4,548	1,127
2010	118	29	969	51	29	0	220	15	428	148	655	110	721	78	3,140	431
2011	19	0	504	0	376	9	1,203	16	491	198	804	139	1,882	410	5,279	772
2012	11	11	211	0	111	32	13	13	589	263	827	452	677	285	2,439	1,056
2013	13	0	1,542	35	54	19	662	35	331	35	397	0	1,772	318	4,771	442
2014	140	0	718	0	0	0	95	0	91	91	1,918	0	4,069	723	7,031	814
2015	39	0	385	41	39	0	103	0	236	0	870	19	2,112	189	3,784	249
Average																
2011–2015	44	2	672	15	116	12	415	13	348	117	963	122	2,102	385	4,661	667
2006–2015	75	9	632	22	171	32	330	8	591	116	992	233	1,995	443	4,784	862

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Note: Arctic char may be included in these estimates due to angler difficulty in distinguishing the two species.




Figure 10.—Dolly Varden sport catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Note: Arctic char may be included in these estimates due to angler difficulty in distinguishing the two species.

STOCKING PROGRAM

There is currently no Dolly Varden–Arctic char stocking program for streams in the AMA. There are some lakes in the AMA stocked with Arctic char.

FISHERY PERFORMANCE

Annual Dolly Varden catch has been declining slightly since 2006; however, on average, only around 15% of the estimated Dolly Varden catch is actually harvested in AMA streams (calculated from Table 10). Prior to 2006, Campbell Creek typically had the largest catches of Dolly Varden in the AMA streams. Since then, Campbell Creek has only had the largest catch in 2010 and 2013. The largest catch of Dolly Varden ever recorded (12,760) was in Campbell Creek in 2001 (Table 10, Figure 10).

STEELHEAD–RAINBOW TROUT FISHERIES

FISHERY DESCRIPTION

Several AMA streams support populations of resident rainbow trout. Rainbow trout have been reported in Bird, Campbell, Chester, Ingram, and Ship creeks, and Placer and Twentymile rivers. Although steelhead trout are not indigenous to the AMA, there has been angling interest from the public and unsuccessful stocking attempts were done to establish an anadromous rainbow trout population (Bosch 2010). On average, Campbell Creek has supported the largest catch estimates for over the last 20 years (1996–2015) with over 2,400 rainbow trout caught annually and less than 1% of those harvested (Table 11, Figure 11). Few steelhead trout have been caught in the AMA although stocking of these fish was historically attempted in Campbell Creek.

Since 1999, only triploid rainbow trout, which do not spawn, have been stocked into Chester and Campbell creeks. Both of these streams have populations of wild rainbow trout. However, stocking of steelhead trout from Kodiak and Anchor River origin into Campbell Creek was attempted in the 1950s and reattempted in the 1980s until it was discontinued in 1987 due to poor returns (Bosch 2010). Both creeks are urban with paved trail access along good portions of their reaches, making them popular spring and summer fisheries.

FISHERY MANAGEMENT AND OBJECTIVES

There are no formal management objectives for steelhead–rainbow trout in the AMA. ADF&G has a constitutional mandate to manage on the principle of sustained yield. Within the sustained yield principle, SF goals seek to optimize social and economic benefits and, where possible, expand opportunity to participate in diverse fisheries on these stocks. No stream surveys are conducted to specifically count rainbow trout.

FISHERY PERFORMANCE

For 2006–2015, less than 11% of the total catch of rainbow trout in the AMA came from streams (Table 11). This percentage is similar to historical catches going back to 1996 (Table 11). On average, nearly half of the rainbow trout caught in AMA streams come from Campbell Creek. In 2011, an emergency order (EO 2-RT-2-10-11) was issued to prohibit the retention of rainbow–steelhead trout in the Chester and Campbell creek drainages, including University Lake, and prohibited the use of bait in the Chester Creek drainage. This EO was issued to conserve the catchable-sized fish, and help ensure angler opportunity throughout the fishing season until the new hatchery came online in 2012. Stockings were back to near historical levels in 2012–2015 (Appendices B3–B4).

Table 11.—Rainbow trout sport fish catch and harvest, Anchorage Management Area, 1996–2015.

Year	Streams						Lakes		Area total			
	Campbell Creek		Ship Creek		Other creeks		Stream Total		Total catch	Total harvest	Catch	Harvest
	Catch	Harvest	Catch	Harvest	Catch	Harvest	Catch	Harvest				
1996	2,622	531	229	53	1,517	578	4,368	1,162	134,690	34,986	139,058	36,148
1997	2,988	215	84	84	507	221	3,579	520	167,746	36,625	171,325	37,145
1998	3,603	272	144	0	4,810	514	8,557	786	84,886	18,975	93,443	19,761
1999	2,874	711	94	47	1,362	144	4,330	902	158,680	27,874	163,010	28,776
2000	4,766	216	1,106	85	1,891	61	8,380	384	153,431	35,203	161,811	35,587
2001	14,952	369	1,094	0	2,053	332	18,099	701	91,521	20,610	109,620	21,311
2002	2,950	418	1,245	0	807	61	5,002	479	86,742	21,999	91,744	22,478
2003	3,177	257	2,359	0	3,480	483	9,016	740	53,203	17,782	62,219	18,522
2004	2,032	117	937	0	673	161	3,642	278	77,479	22,998	81,121	23,276
2005	1,455	99	1,312	0	1,034	463	3,801	562	52,067	13,314	55,868	13,876
2006	720	24	334	0	713	500	1,767	524	35,417	7,321	37,184	7,845
2007	888	11	231	0	2,325	82	3,444	93	26,167	5,212	29,611	5,305
2008	740	0	215	0	1,455	142	2,410	142	41,236	8,295	43,646	8,437
2009	310	0	105	0	314	10	729	10	31,391	5,706	32,120	5,716
2010	495	0	73	0	461	57	1,029	57	25,155	3,320	26,184	3,377
2011	920	0	239	0	2,035	40	3,194	40	12,767	2,071	15,961	2,111
2012	318	0	58	17	254	103	630	120	27,719	5,705	28,349	5,825
2013	646	140	691	82	1,034	17	2,371	239	45,864	9,008	48,235	9,247
2014	559	0	112	28	1,213	304	1,884	332	36,063	5,981	37,947	6,313
2015	1,636	82	466	156	78	17	2,180	255	63,772	15,554	65,952	15,809
Average												
2011–2015	816	44	313	57	923	96	2,052	197	37,237	7,664	39,289	7,861
2006–2015	723	26	252	28	988	127	1,964	181	34,555	6,817	36,519	6,999

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.




Figure 11.—Rainbow trout sport fish catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

LAKE FISHERIES

FISHERY DESCRIPTION

In the 1960s, ADF&G began a rainbow trout stocking program to increase sport fishing opportunities within AMA lakes; this program continues today (Appendices B3 and B4). Stocking of coho salmon into AMA lakes began in 1966 in small numbers but was discontinued by 1993 in all but Clunie Lake where 1 additional stocking took place in 2002. Chinook salmon have been stocked in AMA lakes since 1981 and in 1990 stocking of Arctic char and Arctic grayling began in select AMA lakes (Appendices B1, B5, and B6). Arctic grayling stocking was discontinued after the 2015 stocking. For all species, actual stocking numbers have varied by location and number annually (Appendices B1–B6). Recent stocked fish locations can be found in Figure 2.

The AMA stocked lakes program has increased sport fishing opportunities dramatically in the AMA. Without the current stocking program in place, harvest on wild stocks in the AMA would have to be managed with tighter restrictions due to the always increasing Anchorage population resulting in increasing number of anglers. Stocked lakes in the AMA have provided Alaska residents and nonresidents with additional sport fish opportunities, and effort on these lakes has continued to increase annually since the new WJHSF Hatchery started stocking at historical levels in 2012. An economic study prepared by Northern Economics (Northern Economics Inc.

2004) estimated the AMA stocked lakes fishery provided the Anchorage area with an annual benefit of more than \$3,000,000.

Arctic Char–Dolly Varden

Since 1990, 3 lakes in the AMA were designated as the primary stocking locations for Arctic char: Campbell Point, Clunie, and Mirror lakes. In 2002 and 2003, ADF&G increased stocking of Arctic char to include DeLong, Jewell, Sand, and Tangle lakes. In total, 7 AMA lakes were stocked in 2002 and 2003 (Appendix B5). Although these stocked Arctic char were active in cold water temperatures and added to the stocked lakes program, they did not do well in warmer waters and high mortalities were reported in shallow area lakes in the summer of 2003. In response, ADF&G evaluated the stocking program and reduced stocking of Arctic char to just Campbell Point and Sand lakes in most of the following years. In addition to Campbell Point and Sand lakes, since 2003 2 stockings have occurred in Jewel Lake and since 2011, Clunie Lake has been stocked annually with Arctic char.

ADF&G daily bag and possession limits for Arctic char–Dolly Varden in stocked waters are 5 per day, 5 in possession with no size limit. In all other waters the limits are 5 per day, 5 in possession, and only 1 may be 12 inches or longer.

Arctic Grayling

Arctic grayling have been stocked in 8 AMA lakes since 1990, but only Beach Lake was stocked almost annually from 1990 to 2015 (Appendix B6 and ADF&G hatchery records). In addition, from 2010 to 2015, Sand Lake was stocked annually with Arctic grayling. A healthy population of Arctic grayling is found in Symphony Lake but it has not been stocked since 2003. Natural reproduction from diploid grayling stocked in 2001 and 2003 is maintaining the angling opportunities in this lake. In 2010, a mark-recapture study examined the stock structure of the Symphony Lake Arctic grayling population. This project found that the abundance was higher than expected, nearing the limits of lake capacity, with a small proportion of the population greater than 12 inches in length.

Due to disease concerns, catchable Arctic grayling production ceased in 2002. Arctic grayling fingerling and subcatchable fish were released from 2003 to 2012. The first release of catchable Arctic grayling from WJHSF Hatchery occurred in 2013. The Arctic grayling stocking program was discontinued after 2015 in order to reduce operating expenses (ADF&G WJHSF Hatchery staff, personal communication).

ADF&G daily bag and possession limits for Arctic grayling in stocked lakes are 5 per day, 5 in possession. Symphony Lake has the same bag limit, however only 1 fish may be greater than 12 inches in length and sport fishing is closed in this lake from May 1 to June 30. In all other waters the bag and possession limits are 2 per day, 2 in possession.

Chinook Salmon

Chinook salmon have been stocked in 19 AMA lakes since 1990 (Appendix B1 and ADF&G hatchery records). From 2012 to 2015, only 9 AMA area lakes were stocked annually with Chinook salmon. Between 1990 and 2005, 54,000 Chinook salmon were stocked on average between AMA lakes. From 2007 to 2015, this average was lower (37,000 fish). When stocking occurred, as few as 4 lakes were stocked. In 2010 and 2011, no Chinook salmon were stocked in AMA lakes due to the hot water reduction at the hatcheries and awaiting the new hatchery to

come online. The Chinook salmon catchable production program has returned to near historical stocking levels when the first catchable fish were released from the WJHSF Hatchery in 2012 (Appendix B1).

ADF&G daily bag and possession limits for landlocked salmon were 10 per day, 10 in possession with no size restrictions through 2013. In 2014, the limit was reduced to 5 per day, 5 in possession to align with the rainbow trout bag and possession limits.

Rainbow Trout

Rainbow trout have been stocked in 35 AMA lakes since stocking began in the 1960s (ADF&G hatchery records). In 1966, six AMA lakes were stocked for the first time with rainbow trout. From 2012 to 2015, between 24 and 26 AMA lakes were stocked annually (Appendix B4). On average, 121,211 catchable rainbow trout were stocked in AMA lakes from 2012 to 2015 (calculated from Appendix B4). A total of 34,371 catchable rainbow trout were stocked between 2010 and 2011, when the hatcheries lost sufficient hot water and the new hatchery had yet to come online. The rainbow trout catchable production program returned to historical stocking levels when the first catchable fish were released from the WJHSF Hatchery in 2012.

ADF&G daily bag and possession limits for rainbow trout are 5 per day, 5 in possession, of which only 1 can be 20 inches or more in length. Anglers must immediately record rainbow trout 20 inches or more in length on the back of their sport fishing license, and for all Cook Inlet waters combined, there is a seasonal limit of 2 rainbow trout 20 inches or more in length.

FISHERY MANAGEMENT AND OBJECTIVES

The management objective for AMA lakes is to maintain 67,500 angler-days of effort annually. Stocking goals, public information programs, news releases, and community school classes are used to attain the management objective. News releases and weekly fishing reports outlining lake stocking efforts have helped to keep anglers up to date on the status of AMA lakes. Stocked AMA lakes have provided significant urban angling opportunities throughout the year and have supported on average (2011–2015) 45% of the annual AMA sport fishing effort (Table 2). Since the new hatchery came online and stocking has increased, effort in AMA lakes has also increased when compared with effort during 2005–2011. The most popular AMA lakes include Jewel and Cheney lakes in the Anchorage Bowl; Otter, Clunie, and Sixmile lakes on Joint Base Elmendorf Richardson (JBER); and Mirror Lake in Peters Creek (Tables 12 and 13).

Invasive Species

Northern pike were illegally introduced and documented in 6 AMA lakes. The presence and spread of these invasive northern pike have been identified as top concerns by area managers and regional staff. The northern pike management program began in 2003 with sampling of Lower Fire, Sand, and Cheney lakes using 30 ft and 48 ft variable mesh gillnets, baited hoop traps, fyke nets, and spears. The presence of ADF&G staff, targeting northern pike in area lakes, captured the attention of local anglers and media who reported the sampling trips and helped increase public awareness of the local northern pike problem. Educating the public and encouraging anglers to harvest northern pike is a crucial component in ADF&G's efforts to control northern pike in AMA lakes. ADF&G staff have continued to set gillnets and hoop traps in Lower Fire, Sand, Campbell, and Cheney lakes and to monitor reports of northern pike presence in other systems.

Table 12.—Angler lake sport fishing effort (lakes with names starting A–G), Anchorage Management Area, 1996–2015.

Year	Beach		Taku		Campbell Point		Cheney		Clunie		Delong		Fire (lower)		Fish		Green		Gwenn	
	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%
1996	2,928	5	1,192	2	1,798	3	4,789	9	4,465	8	3,194	6	2,967	5	1,443	3	2,313	4	2,705	5
1997	4,786	7	1,429	2	1,691	3	5,280	8	4,636	7	3,321	5	2,916	5	1,142	2	4,638	7	3,262	5
1998	2,328	5	1,731	4	329	1	2,703	6	2,921	7	2,000	5	2,442	6	1,898	4	3,371	8	1,109	3
1999	4,088	6	3,290	5	1,572	2	3,933	6	4,843	7	3,620	5	2,091	3	2,356	4	3,781	6	3,456	5
2000	3,468	5	5,494	8	1,982	3	3,935	6	5,850	8	4,649	7	2,288	3	1,974	3	4,204	6	2,756	4
2001	3,939	8	1,410	3	1,336	3	1,630	3	4,550	10	3,100	7	1,446	3	740	2	1,824	4	1,778	4
2002	5,148	13	1,399	3	1,469	4	1,616	4	1,130	3	2,215	6	2,352	6	933	2	919	2	1,046	3
2003	4,530	11	1,050	3	991	2	1,104	3	2,268	6	3,302	8	3,117	8			2,199	5	1,070	3
2004	2,313	5	2,540	5	1,680	4			6,289	13	4,499	9	1,471	3			2,192	5	1,420	3
2005	3,564	10	985	3	1,423	4	915	2	2,037	6	4,296	12	1,546	4						
2006	2,892	8	1,697	5	2,266	6	838	2	1,115	3	3,266	9	1,992	6						
2007	2,291	8	1,075	4	2,134	7	604	2	1,266	4	2,272	8	1,700	6			745	3		
2008	1,109	3			2,551	7	1,044	3	3,513	10	1,966	5	824	2			1,557	4	612	2
2009	2,455	9			2,645	9	2,307	8	1,390	5	2,375	9	629	2			883	3		
2010	784	3	728	3	2,287	9	3,485	14	1,286	5	2,353	10	1,860	8						
2011					1,109	7	1,145	7	2,265	13	1,587	9								
2012	751	3					1,771	8			2,102	9								
2013					4,320	13	5,642	18			4,130	13								
2014	1,346	3	1,854	5	2,127	5	1,330	3	2,440	6	6,228	16								
2015	1,982	6			4,043	13	1,284	4	2,350	8	3,315	11								
Average																				
2011–2015	1,360	4	1,854	5	3,085	10	2,752	8	2,395	9	3,944	12								
2006–2015	1,216	6	1,219	4	2,652	9	2,068	7	1,953	7	2,959	10	1,401	5			1,220	3	612	2

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Table 13.—Angler lake sport fishing effort (lakes with names starting H–Z and total AMA effort), Anchorage Management Area, 1996–2015.

Year	Hillberg		Jewel		Mirror		Otter		Sand		Sixmile		Symphony		Triangle		Other lakes		Total effort for all lakes
	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	Effort	%	
1996	2,130	4	4,343	8	4,292	8	7,035	13	1,885	3	3,923	7			771	1	25,852	47	54,699
1997	3,321	5	6,283	10	4,378	7	6,265	10	1,735	3	4,659	7			1,199	2	32,830	51	64,331
1998	1,557	4	5,373	12	3,874	9	4,005	9	2,061	5	2,867	7			1,022	2	22,162	50	43,905
1999	3,651	6	7,814	12	3,843	6	4,090	6	3,492	5	5,928	9			1,096	2	35,861	54	66,312
2000	3,546	5	6,173	9	4,052	6	5,048	7	4,692	7	5,101	7			1,086	2	39,448	57	69,607
2001	1,882	4	5,755	12	6,989	15	3,293	7	2,337	5	3,248	7					21,919	46	47,384
2002	898	2	6,632	16	5,384	13	2,577	6	1,715	4	1,072	3					21,790	54	40,201
2003	878	2	5,531	14	2,987	7	3,670	9	1,877	5	1,683	4					23,309	57	40,552
2004	1,185	2	7,956	17	4,103	9	2,983	6	2,768	6	1,323	3					24,568	52	47,539
2005	1,895	5	6,321	17	3,927	11	1,510	4	2,838	8	1,979	5					20,144	55	36,833
2006	761	2	4,774	13	3,965	11	1,318	4	3,451	10	1,562	4					22,169	62	35,741
2007			4,925	17	2,751	10	1,273	4	1,722	6	1,620	6					16,394	57	28,883
2008	2,901	8	3,477	10	3,071	9	2,515	7	1,628	5	1,184	3	572	2			21,180	59	35,984
2009	1,536	6	3,551	13	2,060	7			1,470	5	1,076	4	533	2			16,151	58	27,910
2010	1,431	6	2,624	11	1,017	4			764	3							14,199	59	24,152
2011			1,677	10		0	635	4	1,834	11							9,955	59	16,793
2012			2,042	9	976	4			5,354	24							14,419	65	22,265
2013			3,119	10	4,304	13			3,090	10							17,586	55	32,095
2014			12,624	32	3,879	10			2,587	7							18,091	46	39,333
2015			5,013	16	2,756	9			1,641	5							17,425	57	30,738
Average																			
2011–2015			5,700	15	2,979	7		4	2,439	11							15,495	56	28,245
2006–2015	1,657	5	4,383	14	2,499	8	1,435	5	2,232	9	1,361	4	553	2			16,757	58	29,389

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

To further eradicate northern pike in AMA lakes, Cheney Lake was treated with rotenone in October 2008 followed by continued monitoring by ADF&G throughout the winter. During May 2009, gillnet sampling failed to find northern pike in Cheney Lake, however many invertebrates species were collected and apparently continued to thrive. Cheney Lake was once again stocked from the WJHSF Hatchery with catchable-sized fish and in 2009 it was the second-most fished lake in the AMA. A grassroots effort by residents around Sand Lake helped ADF&G start northern pike removal from this lake too. In October of 2009, Sand Lake was treated with rotenone and spring sampling in 2010 failed to find any northern pike. Sand Lake was stocked with catchable fish and residents around the lake reported good rainbow trout fishing once more. AMA lakes will continue to be monitored for the presence of northern pike and removal methods will be determined on a case-by-case basis. In 2015, Otter Lake on JBER was also treated with rotenone to eradicate northern pike. Posttreatment netting efforts failed to find any northern pike in the lake. As of 2015, Lower Fire Lake is the only AMA lake where northern pike may still be present in small numbers. Efforts to monitor this lake are ongoing.

STOCKING PROGRAM

Fish used for stocking in AMA lakes have been raised in the William Jack Hernandez Sport Fish (WJHSF) Hatchery since 2012. In 2010, the Fort Richardson and Elmendorf hatcheries that raised sport fish for stocking were closed and replaced by the new WJHSF Hatchery at the old Elmendorf hatchery location. Since 2005, effort, catch, and harvest in AMA stocked lakes dropped considerably when both state hatcheries lost hot water, and thus the ability to rear fish to catchable size (Figure 12). The new WJHSF Hatchery facility has heated water and the ability to rear catchable-sized fish in 1 year. Catchable-sized fish were released into AMA lakes again in 2012. The 2016–2020 Statewide Stocking Plan including specifics for AMA lakes can be found here: <http://www.adfg.alaska.gov/static/fishing/pdfs/hatcheries/16region2.pdf>.




Figure 12.—Angler fishing effort for lakes in the Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet], 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Peak catch and harvest in AMA lakes occurred in 2000 (198,200 and 44,241, respectively) followed by a decline through 2011 (Table 14). Reductions in the number and size of fish available for release in local stocked lakes had negative effects on the popularity and productivity of local stocked lakes fisheries. For the years 2010 and 2011, all fish destined for AMA lakes had to be stocked at once, during the spring and early summer of 2010, due to the closure of Elmendorf State Fish Hatchery (Bosch 2010). In 2011, while waiting for the new hatchery to come online, the total numbers of sport fish caught and harvested in AMA lakes reached an all-time low. Total catch and harvest was 17,497 and 3,323, respectively (Table 14). In addition, in 2011, the lowest number of angler-days of effort was observed in AMA lakes (16,793; Table 13). This drop in catch, harvest, and effort can probably be attributed to the reduction of stocked fish in AMA lakes with the hatchery changeover (Appendices B1–B6). Total catch and harvest in AMA lakes has since rebounded and in 2015, total catch and harvest had increased to 70,481 and 17,721, respectively, which is much closer to what was observed historically (Table 14; Figure 13). From 2006 to 2015, angler effort in AMA lakes was on average nearly 30,000 angler-days (Table 2). From 2010 to 2012, the number of angler-days of effort in area lakes hit a low point, but appears to be increasing to prior levels. This reduction was also probably due to an emergency order (EO 2-R-2-01-11) that was issued in 2011 to reduce the bag and possession limit for steelhead–rainbow trout, Arctic char–Dolly Varden, and Arctic grayling.




Figure 13.—Lake sport fish catch and harvest, Anchorage Management Area, 2006–2015.

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

Arctic Char–Dolly Varden

The average estimated annual catch and harvest of Arctic char in AMA lakes from 2011 to 2015 was 2,102 and 385, respectively (Table 14). Harvest of Arctic char was lowest in 2010 (78 fish) while total catch was lowest in 2012 (677 fish). On average (2011–2015), approximately 78% of

the Arctic char caught in the AMA lakes sport fishery are released. However, this number increased in 2015 and just over 90% of Arctic char caught in AMA lakes were released (Table 14). Conversely in 2012, just under 60% of these fish were released. The additional harvest of Arctic char in 2012 could be attributed to fewer stocked fish available (this may also explain the additional harvest of landlocked salmon). Many of the Arctic char are probably misidentified as Dolly Varden. Only Arctic char are stocked in AMA lakes.

Arctic Grayling

The average estimated annual catch and harvest of Arctic grayling in AMA lakes from 2011 to 2015 was 1,741 and 102, respectively (Table 14). In 2011, the catch and harvest of Arctic grayling was one of the lowest of record. Few fish were stocked into Beach Lake compared to all other years and this probably contributed to the lower catch and harvest in 2011 (Appendix B6). From 2010 to 2015, all fish were stocked between Beach and Sand lakes (Appendix B6). Over 10,000 fish were stocked in 2010, but from 2013 to 2015, stocking was under 3,000 fingerlings each year. Due to budget cuts, the last year of stocking Arctic grayling was 2015 and the WJHSF Hatchery is no longer producing Arctic grayling.

Landlocked Chinook Salmon

The average estimated annual catch and harvest of Chinook salmon in AMA lakes from 2011 to 2015 was 3,318 and 1,168, respectively (Table 14). No lakes were stocked with Chinook salmon in 2010 or 2011 due to the hatchery change-over (Appendix B1). This was reflected in the catch and harvest numbers from 2010 to 2012 (Table 14). In 2012, after 2 years of reduced harvest, the first stocking of Chinook salmon did not occur until June and in some lakes, not until October. Catch and harvest increased significantly in 2013 and 2014 after multiple stockings of 11 AMA lakes. In 2014, total catch and harvest of Chinook salmon was the highest observed since 2005.

Rainbow Trout

The average estimated annual catch and harvest of rainbow trout in AMA lakes from 2011 to 2015 was 37,237 and 7,664, respectively (Table 14). Reductions in stocking during 2010 and 2011 probably contributed to the low catch and harvest in 2011 (12,767 and 2,071, respectively). From 1996 to 2005, average annual catch and harvest of rainbow trout was 106,425 and 25,093, respectively (calculated from Table 14). With increased stocking levels over the last 4 years (Appendix B4), catch and harvest of rainbow trout in AMA lakes has been increasing. In 2015, catch and harvest of rainbow trout were the largest observed since 2005 (63,772 and 15,554, respectively; Table 14). In 2012, an emergency order (EO 2-R-01-12) was issued increasing the bag limit for rainbow trout in Otter Lake. A rotenone treatment to eradicate northern pike was anticipated to occur in fall of 2012 but did not. In 2015, a similar emergency order (EO 2-RR-2-16-15) was issued increasing the bag limit for rainbow trout in Otter Lake again, until its actual scheduled rotenone treatment to eradicate northern pike in fall of 2015.

Table 14.—Lake sport fish catch (C), harvest (H), and percent released (%), Anchorage Management Area, 1996–2015.

Year	Arctic char–Dolly Varden			Arctic grayling			Northern pike			Rainbow trout			Landlocked Chinook salmon			Other (trout, other landlocked salmon)			Lake total	
	C	H	%	C	H	%	C	H	%	C	H	%	C	H	%	C	H	%	C	H
1996	2,698	1,010	63	2,103	182	91	893	183	80	134,690	34,986	74	32,682	13,280	59				173,066	49,641
1997	1,736	559	68	4,555	68	99	433	220	49	167,746	36,625	78	21,490	8,913	59				195,960	46,385
1998	2,961	988	67	1,077	59	95	551	337	39	84,886	18,975	78	25,313	12,601	50				114,788	32,960
1999	1,362	1,092	20	1,351	158	88	726	243	67	158,680	27,874	82	17,372	7,854	55	5,226	921	82	184,717	38,142
2000	4,949	604	88	122	10	92	1,685	1,274	24	153,431	35,203	77	36,723	6,952	81	1,290	198	85	198,200	44,241
2001	947	290	69	412	43	90	4,822	1,140	76	91,521	20,610	77	25,157	9,638	62	650	173	73	123,509	31,894
2002	2,135	355	83	4,463	747	83	2,151	1,190	45	86,742	21,999	75	18,352	6,093	67	199	199	0	114,042	30,583
2003	2,851	1,205	58	2,192	221	90	2,351	1,066	55	53,203	17,782	67	9,614	3,206	67	1,797	779	57	72,008	24,259
2004	5,391	3,476	36	363	90	75	1,473	663	55	77,479	22,998	70	16,790	8,468	50	593	0	100	102,089	35,695
2005	9,977	1,358	86	630	166	74	1,413	611	57	55,868	13,876	75	16,900	4,850	71	688	64	91	85,476	20,925
2006	1,817	641	65	152	55	64	1,053	364	65	35,417	7,321	79	4,295	1,595	63	253	39	85	42,987	10,015
2007	2,849	667	77	1,340	386	71	1,975	1,285	35	26,167	5,212	80	7,689	1,463	81	0	0		40,020	9,013
2008	2,201	425	81	3,409	444	87	1,267	205	84	41,236	8,295	80	5,721	1,479	74	307	246	20	54,141	11,094
2009	1,797	653	64	1,124	194	83	2,560	2,040	20	31,391	5,706	82	4,972	1,948	61	323	36	89	42,167	10,577
2010	721	78	89	599	123	79	594	507	15	25,155	3,320	87	1,864	341	82	1008	364	64	29,941	4,733
2011	1,882	410	78	367	0	100	645	616	4	12,767	2,071	84	1,782	209	88	54	17	69	17,497	3,323
2012	677	285	58	2,105	120	94	469	312	33	27,719	5,705	79	381	216	43	990	133	87	32,341	6,771
2013	1,772	318	82	2,796	124	96	2,473	2,323	6	45,864	9,008	80	3,282	897	73	1543	48	97	57,730	12,718
2014	4,069	723	82	1,982	60	97	129	100	22	36,063	5,981	83	8,339	2,903	65	246	62	75	50,828	9,829
2015	2,112	189	91	1,453	205	86	198	136	31	63,772	15,554	76	2,807	1,617	42	139	20	86	70,481	17,721
Average																				
2011–2015	2,102	385	78	1,741	102	95	783	697	20	37,237	7,664	81	3,318	1,168	62	594	56	82	45,775	10,072
2006–2015	1,990	439	77	1,533	171	86	1,136	789		34,555	6,817	81	4,113	1,267	67	486	97	74	43,813	9,579

Source: Alaska Sport Fishing Survey database [Intranet]. 1996–present. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited September 8, 2016). Available from Division of Sport Fish, Research and Technical Services.

REFERENCES CITED

- Bosch, D. 2010. Area management report for the recreational fisheries of Anchorage, 2009 and 2010. Alaska Department of Fish and Game, Fishery Management Report No. 10-53, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/Fmr10-53.pdf>
- Havens, A. C., J. B. Murray, K. J. Delaney, and K. J. Roth. 1987. Evaluation of enhancement efforts for rainbow trout, coho salmon, and Chinook salmon in southcentral Alaska, 1986. Alaska Department of Fish and Game, Fishery Data Series No. 33, Juneau. <http://www.adfg.alaska.gov/FedAidPDFs/fds-033.pdf>
- Healey, M. C. 1991. Life history of Chinook salmon (*Oncorhynchus tshawytscha*). Pages 311-393 [In] Pacific salmon life histories. Edited by C. Groot and L. Margolis. University of British Columbia Press, Vancouver, BC.
- Kari, J., and J. A. Fall. 2003. Shem Pete's Alaska, The Territory of the Upper Cook Inlet Dena'ina. University of Alaska Press, Fairbanks.
- Miller, J. 1990. Stocking records, 1990. Alaska Department of Fish and Game, Division of Fisheries Rehabilitation, Enhancement, and Development (FRED), Anchorage.
- Mills, M. J. 1989. Alaska statewide sport fisheries harvest report, 1988. Alaska Department of Fish and Game, Fishery Data Series No. 122, Juneau. <http://www.adfg.alaska.gov/FedAidPDFs/fds-122.pdf>
- Mills, M. J. 1993. Harvest, catch, and participation in Alaska sport fisheries during 1992. Alaska Department of Fish and Game, Fishery Data Series No. 93-42, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/fds93-42.pdf>
- Mills, M. J. 1994. Harvest, catch, and participation in Alaska sport fisheries during 1993. Alaska Department of Fish and Game, Fishery Data Series No. 94-28, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/fds94-28.pdf>
- Munro, A. R., and E. C. Volk. 2010. Summary of Pacific salmon escapement goals in Alaska with a review of escapements from 2001 to 2009. Alaska Department of Fish and Game, Special Publication No. 10-12, Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/SP10-12.pdf>
- Northern Economics Inc. 2004. Economic effect of ADF&G's Region II and III recreational stocking programs; Hatchery valuation analysis. Prepared for Alaska Department of Fish and Game, Northern Economics Inc, Anchorage.
- Sandercock, F. K. 1991. Life history of coho salmon (*Oncorhynchus kisutch*). Pages 397-445 [In] C. Groot and L. Margolis, editors. Pacific salmon life histories. University of British Columbia Press, Vancouver, Canada.
- Stratton, B., and P. Cyr. 1995. Annual management report for the recreational fisheries in the Anchorage area, 1994. Alaska Department of Fish and Game, Fishery Management Report No. 95-05, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/fmr95-05.pdf>
- Stratton, B., and P. Cyr. 1997. Area management report for the Anchorage area, 1995. Alaska Department of Fish and Game, Fishery Management Report No. 97-1, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/fmr97-01.pdf>
- Stratton, B., A. Hoffmann, and P. Cyr. 1994. Annual management report for the Anchorage area 1993. Alaska Department of Fish and Game, Fishery Management Report No. 94-08, Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/fmr94-08.pdf>

APPENDIX A: SPORT FISHING REGULATIONS

Appendix A1.–Sport fishing regulations for Ship Creek, 1957–2015.

Year	Sport fishing regulations for Ship Creek
1957–1959	Closed to sport fishing from April 1 through May 27. Bag limit was 10 trout daily or in possession, only two 20 inches or more in length. No salmon fishing regulations.
1960	Closed to all sport fishing.
1961–1962	Closed to salmon fishing. Closed to sport fishing from April 1 through May 27. Bag limit was 10 trout daily or in possession, only two 20 inches or more in length. Anglers were allowed up to 20 resident fish if excess were Dolly Varden.
1963	Closed to sport fishing from April 1 through May 25.
1964–1965	Closed to sport fishing from April 1 through the third Friday in May. The fishery was open to salmon fishing (except Chinook salmon) downstream of a marker 300 feet below the Chugach Power Plant Dam. Bag limit was 3 chum, sockeye, or pink salmon with an additional 3 coho salmon allowed.
1966–1967	Legal gear was defined as a single-hook-only with gap between point and shank of one-half inch or less.
1968	Closed to all fishing from January 1 through August 31. Anglers were allowed 3 salmon. Closed to Chinook salmon fishing.
1969	From September 1 through December 31 anglers were allowed 3 salmon but excess Dolly Varden was removed from the resident fish bag limit.
1970	Closed to all fishing from January 1 through August 31 except for a Chinook salmon opening from July 4 through July 19. Bag limit was 1 Chinook salmon per day and 2 per season. From September 1 through December 31, anglers were allowed 3 salmon.
1971–1972	Closed to all fishing from January 1 through August 31 except for Chinook salmon openings on June 10 through June 11 and June 17 through June 18. A Chinook salmon punch card was required and the bag limit was 1 Chinook salmon per day and 2 per season. From September 1 through December 31, anglers were allowed 3 salmon.
1973–1981	Closed to all fishing from January 1 through August 17. Closed to Chinook salmon fishing. From August 18 through December 31, anglers were allowed 3 salmon. Legal gear was single-hook-only with a gap between point and shank of one-half inch or less.
1982–1984	Same regulations as 1973–1981. In addition, rainbow trout daily bag limit was reduced to 5 fish, only one 20 inches or more in length.
1985–1986	Closed to all fishing from January 1 through July 31. Single-hook-only restriction was lifted. Anglers were allowed 3 salmon other than Chinook salmon from August 1 to December 31.

-continued-

Year	Sport fishing regulations for Ship Creek
1987–1990	The area opened to salmon fishing was downstream of a marker located 100 feet below the Chugach Power Plant Dam. In addition, the creek was open to all fishing (including Chinook salmon) on Tuesdays and Wednesdays for 5 consecutive weeks commencing the second Tuesday in June. Chinook salmon bag and possession limits were 1 and 2, respectively with no seasonal limit.
1991–1992	Chinook salmon fishing was allowed from January 1 through July 13. Daily bag and possession limits were 1 and 2, respectively with no seasonal limit. Fishing for other salmon was allowed year-round with bag and possession limits of 3 and 3, respectively. In addition, fishing for Dolly Varden, rainbow trout, and other species was allowed year-round.
1993	A seasonal limit of 5 Chinook salmon in Cook Inlet waters was added.
1997	The possession limit for Chinook salmon was reduced to 1 and a regulation went into effect that prohibited anglers from continuing to sport fish in waters open to Chinook salmon fishing after harvesting a Chinook salmon.
1999	Chinook salmon bag and possession limits were 1 per day and 1 in possession, and anglers were not allowed to fish in Ship Creek for the remainder of the day after harvesting a Chinook salmon 20 inches or longer. For salmon other than Chinook salmon, 16 inches or longer in length, the bag and possession limits were 3 fish, and all three could be coho salmon. Reeve Boulevard upstream to 300 ft upstream of Elmendorf Dam was closed to all fishing.
2001	The bag and possession limits for Chinook salmon less than 20 inches in length, and other salmon less than 16 inches in length was 10 fish. The fishery was open all year for these small salmon. In waters open to fishing for Chinook salmon 20 inches or more in length, fishing was not allowed between 11:00 PM and 6:00 AM from May 15 through July 13.
	Statewide regulation defined the bag and possession limits for Chinook salmon less than 20 inches in length (jack salmon), in fresh waters open to Chinook salmon fishing, as 10 per day and 10 in possession.
2005	In waters closed to fishing for salmon 20 inches or more, fishing was closed for salmon less than 20 inches in length.
	A coho salmon 16 inches or longer that is removed from fresh water must be retained and becomes part of the bag limit of the person who originally hooked the fish. A person may not remove a coho salmon 16 inches or longer from the water before releasing it.
2012	Sport fishing is closed from April 15 to June 14 from 100 feet upstream of the Chugach Power Plant Dam upstream to Reeves Boulevard.
	Footgear with absorbent felt or other fibrous material on the soles are prohibited while sport fishing in the fresh waters of Alaska. The regulation was adopted by the Alaska Board of Fisheries in 2010 to reduce the potential for introduction and spread of invasive organisms, including plants, into Alaska waters. Invasive organisms spread by contaminated waders and other gear can threaten resident fish stocks and important fish habitat.

Appendix A2.–Sport fishing regulations for Eagle River, 1957–2015.

Year	Sport fishing regulations for Eagle River
1957–1959	Closed to sport fishing from April 1 through May 27. Bag limit was 10 trout daily or in possession, only two 20 inches or more in length. No salmon fishing regulations.
1960	Closed to salmon fishing upstream of one-quarter mile above Glenn Highway Bridge. Bag limits were 10 salmon or trout daily, 3 could be salmon greater than 16 inches in length, and 2 could be Chinook salmon.
1961–1962	Anglers were allowed up to 20 resident fish if the excess were Dolly Varden.
1963	Closed to sport fishing from April 1 through May 25. Closed to salmon fishing upstream of one-quarter mile above Glenn Highway Bridge. Bag limit was 6 coho salmon; 3 pink, chum, or sockeye salmon; and 1 Chinook salmon. Resident fish bag limits were 10 trout daily, only 2 over 20 inches. Anglers were allowed up to 20 resident fish if the excess were Dolly Varden.
1964–1967	Closed season was from April 1 through the third Friday in May.
1968	No closed season. Bag limit was 3 salmon, 16 inches or greater in length. Closed to Chinook salmon fishing.
1969–1981	Excess Dolly Varden removed from bag limit in 1969.
1982–1986	Rainbow trout bag limit was reduced to 5 per day, only one 20 inches or greater in length in 1982. Bag limits were 10 for other resident fish.
1987–1991	South Fork Eagle River below the falls was closed to all fishing from June 1 through August 14.
1993–1996	Regulations restricted Chinook salmon fishing to a 30-day period commencing the Saturday before Memorial Day. Fishing was restricted to that portion of Eagle River upstream of Bailey Bridge on Fort Richardson to an ADF&G marker located approximately adjacent to Mile 7.4 of Eagle River Road. The area located approximately 100 yards on either side of the confluence of South Fork Eagle River was closed to fishing from June 1 through August 14. North Fork Eagle River upstream from an ADF&G marker located near its confluence with Eagle River was closed to all fishing during the Chinook salmon season. Passes were required to fish on Fort Richardson.

-continued-

Year	Sport fishing regulations for Eagle River
1999–2001	<p>In areas open for fishing Chinook salmon less than 20 inches in length and other salmon 16 inches in length or less, the fishery was open all year. Bag and possession limits for these small salmon was 10 fish.</p> <p>Statewide regulation defined the bag and possession limits as 10 per day and 10 in possession for Chinook salmon less than 20 inches in length (jack salmon) in fresh waters open to Chinook salmon fishing.</p>
2005	<p>In waters closed to fishing for salmon 20 inches or more, waters were closed to fishing for salmon less than 20 inches in length.</p>
2011	<p>A coho salmon 16 inches or longer that is removed from fresh water must be retained and becomes part of the bag limit of the person who originally hooked the fish. A person may not remove a coho salmon 16 inches or longer from the water before releasing it.</p>
2012	<p>Footgear with absorbent felt or other fibrous material on the soles are prohibited while sport fishing in the fresh waters of Alaska. The regulation was adopted by the Alaska Board of Fisheries in 2010 to reduce the potential for introduction and spread of invasive organisms, including plants, into Alaska waters. Invasive organisms spread by contaminated waders and other gear can threaten resident fish stocks and important fish habitat.</p>

Appendix A3.–Sport fishing regulations for Campbell Creek, 1957–2015.

Year	Sport fishing regulations for Campbell Creek
1957–1959	Closed to sport fishing from April 1 through May 27. Bag limit was 10 trout daily or in possession, only two 20 inches or more in length. No salmon fishing regulations.
1960	Campbell Creek was open to salmon fishing, except Chinook salmon, from August 22 through September 23. Bag limits were 10 salmon or trout daily, only 3 could be salmon greater than 16 inches in length and only 2 could be trout over 20 inches in length.
1961–1962	Anglers were allowed up to 20 resident fish if excess were Dolly Varden.
1963	Closed to sport fishing April 1 through May 25. Bag limit was 6 coho salmon; 3 pink, chum, or sockeye salmon. Resident fish bag limits were 10, only 2 over 20 inches.
1964–1967	Closed to sport fishing from April 1 through the third Friday in May. Open to salmon fishing (except Chinook salmon) from August 1 through September 30. Closed to salmon fishing above the Seward Highway. Bag limit was 6 coho, and 3 chum, sockeye, or pink salmon.
1968	Open to salmon fishing (except Chinook salmon) from August 1 through September 30. Closed to salmon fishing above the Seward Highway. Bag limit was 3 salmon 16 inches or greater in length. No closed season for resident fish.
1969–1970	Excess Dolly Varden removed from bag limit in 1969.
1971–1981	Closed to fishing above the Seward Highway and closed to salmon fishing throughout the drainage.
1982–1984	Rainbow trout bag limit was reduced to 5 per day, only one 20 inches or greater in length in 1982.
1985	Closed to all fishing above the Forks, and closed to salmon fishing below the Forks.
1986	Entire drainage was open to fishing but closed to salmon fishing.
1987–1992	Only unbaited, artificial lures could be used upstream of the Forks, and rainbow trout could not be kept.
1993–1994	Open to coho salmon fishing from July 25 through October 15, with fishing limited to that portion of Campbell Creek upstream from Dimond Boulevard to an ADF&G marker located in the vicinity of Folker Street. Bag and possession limits for coho salmon were 3 and 3, respectively. Fishing for all other salmon was closed. Campbell Lake was closed to all fishing in 1993.
1996–1998	The portion of Campbell Creek that flows through Wickersham subdivision between Lake Otis Parkway and Shelikof Street was closed to all sport fishing year-round. Coho salmon fishing was allowed from July 25 through October 1 from Dimond Boulevard upstream to C Street. Coho salmon fishing was allowed from August 5 through October 1 upstream of C Street to markers near Piper Street except for the Wickersham closed area.
1999–2001	Regulations were the same as those established by the BOF during 1996–1998.
2005	In waters closed to fishing for salmon 20 inches or more in length, waters were closed to fishing for salmon less than 20 inches in length. ADF&G created a youth-only fishery on Campbell Creek for Chinook salmon.
2011	A coho salmon 16 inches or longer that is removed from fresh water must be retained and becomes part of the bag limit of the person who originally hooked the fish. A person may not remove a coho salmon 16 inches or longer from the water before releasing it.
2012	Footgear with absorbent felt or other fibrous material on the soles are prohibited while sport fishing in the fresh waters of Alaska. The regulation was adopted by the Alaska Board of Fisheries in 2010 to reduce the potential for introduction and spread of invasive organisms, including plants, into Alaska waters. Invasive organisms spread by contaminated waders and other gear can threaten resident fish stocks and important fish habitat.

**APPENDIX B: STOCKING IN THE ANCHORAGE
MANAGEMENT AREA**

Appendix B1.—Chinook salmon stocking in the Anchorage Management Area by year (1996–2015) and site.

Year	Catchable ^a														Smolt	
	Beach	Lake												Total		Ship Creek
		Campbell Pt	Cheney	Clunie	DeLong	Green	Hillberg	Jewel	Mirror	Otter	Sand	Spring	Taku Campbell			
1996	2,989	1,588	4,880	4,023	5,020	1,558	1,587	13,929	8,191	6,776	3,929	998	1,985	1,154	58,607	228,000
1997	2,000	1,000	4,191	2,767	4,032	1,586	1,586	7,325	7,000	5,500	4,000	1,000	2,231	1,651	45,869	325,891
1998	4,533	2,036	6,364	3,000	7,291	2,016	2,062	13,865	7,275	0	6,796	1,000	3,996	1,008	61,242	204,741
1999	2,744	643	6,228	4,045	5,644	2,006	1,932	9,628	7,749	0	5,867	500	3,052	0	50,038	197,168
2000	10,709	0	^b	8,819	5,348	2,149	2,058	9,741	15,399	0	5,119	1,031	0	0	60,373	265,582
2001	4,139	3,807	^b	8,360 ^c	5,966	998	3,308	21,792	10,272	0	4,945	0	0	0	63,587	254,924
2002	3,838	2,000	^b	8,004	6,207	1,086	981	12,538	9,683	0	4,930	0	0	0	49,267	290,501
2003	4,040	1,975	^b	3,822	6,055	1,190	1,144	24,243	7,142	0	5,133	0	5,811	0	60,555	329,416
2004	4,078	2,302	^b	2,981	5,931	1,261	1,261	21,978	7,396	0	4,650	0	2,910	0	54,748	320,226
2005	3,925	3,158	^b	2,981	5,982	1,100	1,100	15,828	6,958	0	6,122	0	3,058	0	50,212	358,029
2006	0	25,723 ^d	^b	0	26,277 ^d	0	0	60,497 ^d	29,043 ^d	0	0	0	0	0	141,540	176,055
2007	0	6,500	^b	3,118	10,530	1,070	1,117	15,795	11,565	0	0	0	0	0	49,695	333,940
2008	0	3,375	^b	2,950	8,031	1,100	1,050	26,622	10,700	0	0	0	0	0	53,828	341,495
2009	0	10,190	^b	3,060	14,838	921	987	27,850	0	0	0	0	0	0	57,846	282,735
2010	0	0	^b	0	0	0	0	0	0	0	0	0	0	0	0	332,597
2011	0	0	^b	0	0	0	0	0	0	0	0	0	0	0	0	314,194
2012	1,763	3,830	1,599	13,889	8,675	2,841	2,866	9,705	710	0	1,652	0	0	0	47,530	329,082
2014	4,220	1,933	2,889	1,846	5,139	946	900	10,575	8,510	0	2,667	0	1,978	0	41,603	358,517
2015	4,123	1,980	4,676	1,944	6,992	1,109	978	11,285	9,718	0	2,442	0	2,009	0	47,256	365,246

Source: All stocking information is from ADF&G SF hatchery records.

^a “Catchable” size is about 8 inches in length.

^b Discontinued stocking beginning in 2000 due to the presence of northern pike.

^c An additional 24,498 Chinook salmon were stocked in Clunie Lake in 2001 but were undersized and considered “subcatchables.”

^d In 2006, the Fort Richardson Hatchery had an outbreak of Bacterial Kidney Disease and Chinook salmon could not be transported to the Elmendorf Hatchery. Instead, they were stocked early into 4 local lakes. They were considered “fingerling-sized” at the time of release.

Appendix B2.–Coho salmon stocking in Anchorage Management Area by year (1996–2015) and site.

Site	Smolt releases			Total
	Site			
	Bird Creek ^a	Campbell Creek	Ship Creek	
1996	147,618	75,943	227,914	451,475
1997	294,565	71,519	232,066	598,150
1998	164,211	83,317	232,765	480,293
1999	111,430	42,046	165,388	318,864
2000	97,409	63,730	260,070	421,209
2001	0	69,836	233,563	303,399
2002	0	69,836	212,639	282,475
2003	0	78,576	234,716	313,292
2004	109,949	85,790	241,006	436,745
2005	100,605	60,387	251,446	412,438
2006	104,974	78,805	252,775	436,554
2007	104,979	82,138	255,380	442,497
2008	113,035	83,421	245,490	441,946
2009	113,300	15,400	287,825	416,525
2010	157,534	50,214	252,319	460,067
2011	136,047	71,960	254,718	462,725
2012	70,004	0 ^b	243,499	313,503
2013	110,297	83,088	273,173	466,558
2014	91,443	29,028	226,576	347,047
2015	132,870	52,110	249,401	434,381

Source: All stocking information is from ADF&G SF hatchery records, which goes back to an early as 1968 for some systems.

Note: Stocking of “catchable” size fish (about 8 inches in length) was discontinued in 2003.

^a Bird Creek was not stocked from 2001 to 2003 due to construction of the parking area just north of the creek..

^b Excessive mortality occurred during early rearing in 2012. No fish were stocked in Campbell Creek to ensure adequate fish were available for Ship Creek and future broodstock needs. Excess fish beyond Ship Creek needs were stocked in Bird Creek.

Appendix B3.–Rainbow trout stocking in the Anchorage Management Area by year (1996–2005) and site.

Release type	Site	Year									
		1996	1997	1998	1999	2000	2001	2002	2003	2004	2005 ^a
Catchable ^b											
	Airstrip/Willow Pond	985	1,050	1,181	1,018	1,497	1,938	2,200	1,866	1,671	1,281
	Alder Pond	5,081	2,592	4,002	2,608	2,072	1,906	2,019	2,455	2,185	1,098
	Beach Lake	4,410	4,244	4,056	10,312	10,487	8,087	10,095	9,614	11,920	7,527
	Campbell Creek	5,104	2,686	2,462	3,030	4,563	3,909	2,291	4,264	1,560	1,697
	Campbell Pt Lake	5,114	2,906	2,172	3,027	5,652	5,533	2,561	2,456	5,829	1,442
	Cheney Lake ^c	10,254	8,946	5,795	10,963	14,473	0	0	0	0	0
	Chester Creek ^d	4,975	2,611	1,000	1,000	852	2,335	2,036	1,779	976	613
	Clunie Lake	9,244	7,662	4,152	9,346	4,669	7,804	3,932	4,613	6,027	4,895
	Delong Lake	10,246	6,207	5,684	9,904	16,589	13,190	1,231	10,182	17,205	11,363
	Dishno Lake	512	515	125	483	0	0	0	0	0	0
	Eagle River	0	0	0	0	0	0	0	0	0	0
	Edmonds Lake	985	1,017	1,195	1,009	500	1,000	1,723	1,967	1,474	943
	Fire Island L	0	0	0	0	0	0	0	0	0	0
	Fish Lake	0	1,054	1,500	2,473	1,135	300	250	532	309	370
	Goose L	0	0	0	0	0	0	0	0	0	0
	Green Lake	3,345	2,729	2,088	2,870	3,151	2,546	1,500	1,359	1,005	889
	Gwen Lake	4,993	5,299	3,929	3,969	4,807	5,153	2,073	4,994	5,001	3,002
	Hillberg Lake	3,393	3,054	2,984	4,014	4,802	1,645	1,532	1,889	1,840	1,744
	Jewel Lake	13,621	10,189	8,986	16,794	15,946	24,622	14,057	17,344	20,060	12,656
	Lake Otis	1,573	1,155	1,000	1,000	500	500	500	250	554	458
	Lower Explorer Lake	0	0	0	0	0	0	0	0	0	0
	Lower Fire Lake	5,170	3,081	3,350	2,956	3,000	3,018	2,976	2,713	2,109	1,663
	Mirror Lake	12,789	13,281	7,032	11,299	12,469	20,195	9,299	7,402	10,812	9,855
	Otter Lake	9,329	12,767	6,994	10,886	10,941	10,159	5,418	7,342	3,738	3,618
	Rabbit Lake	2,553	0	0	1,994	0	0	920	0	0	400
	Sand Lake	6,069	3,646	1,098	3,022	4,096	6,201	3,074	2,105	4,983	2,680
	Spring Lake	1,063	917	500	500	500	0	500	500	505	370
	Sundi Lake	0	0	0	0	0	0	0		0	0
	Taku Campbell Lake ^c	4,213	3,022	1,898	3,948	7,942	0	2,869	1,804	3,490	2,225

-continued-

Release type	Site	Year									
		1996	1997	1998	1999	2000	2001	2002	2003	2004	2005 ^a
Catchable ^b											
	Tangle Pond	3,004	1,247	1,181	1,733	1,000	1,713	1,031	1,021	1,607	1,075
	Thompson Lake	1,979	1,020	978	939	0	977	0	0	0	0
	Triangle Lake	1,448	989	1,000	1,007	707	0	500	500	505	218
	University–APU Lake	0	0	0	0	0	0	0	0	0	0
	Upper Six Mile Lake	3,110	3,000	2,234	4,103	5,066	2,256	2,001	2,241	1,898	1,210
	Waldon Lake	2,006	2,034	1,005	1,275	1,000	4,615	3,208	1,149	0	864
	Total	138,564	110,917	81,579	129,481	140,416	131,603	81,798	94,344	109,267	74,156
Subcatchable ^c											
	Beach Lake	0	0	0	29,844	0	0	0	0	0	0
	Campbell Pt Lake	0	0	0	0	0	0	0	0	0	0
	Cheney Lake ^c	0	0	0	0	0	0	0	0	0	0
	Chester Creek ^d	0	0	0	0	0	0	0	0	0	0
	Clunie Lake	0	0	0	46,138	0	0	0	0	0	0
	DeLong Lake	0	51,088	0	2,769	0	0	0	0	0	0
	Dishno Lake	0	0	0	0	0	0	0	0	0	0
	Eagle River	0	0	0	0	0	0	0	0	0	0
	Edmonds Lake	0	0	0	0	0	0	0	0	0	0
	Eklutna Lake	0	0	0	0	0	0	0	0	0	0
	Fish Lake	0	0	0	0	0	0	0	0	0	0
	Green Lake	0	0	0	0	0	0	0	0	0	0
	Gwen Lake	0	0	0	0	0	0	0	0	0	0
	Hillberg Lake	0	0	0	0	0	0	0	0	0	0
	Jewel Lake	0	53,919		6,772	0	0	0	0	0	0
	Lake Otis	0	0	0	0	0	0	0	0	0	0
	Lower Fire Lake	0	0	0	0	0	0	0	0	0	0
	Mirror Lake	0	0	0	38,254	0	0	0	0	0	0
	Otter Lake	0	0		49,936	0	0	0	0	0	0
	Sand Lake	0	0	0	0	0	0	0	0	0	0
	Spring Lake	0	0	0	0	0	0	0	0	0	0

-continued-

Release type	Site	Year									
		1996	1997	1998	1999	2000	2001	2002	2003	2004	2005 ^a
Subcatchable ^e											
	Taku Campbell Lake ^c	0	0	0	0	0	0	0	0	0	0
	Tangle Pond	0	0	0	0	0	0	0	0	0	0
	Thompson Lake	0	0	0	0	0	0	0	0	0	0
	Triangle Lake	0	0	0	0	0	0	0	0	0	0
	Upper Six Mile Lake	0	0	0	0	0	0	0	0	0	0
	University–APU Lake	0	0	0	0	0	0	0	0	0	0
	Waldon Lake	0	0	0	0	0	0	0	0	0	0
	Total	1,996	107,004	1,998	175,712	2,000	2,001	2,002	2,003	2,004	0

Source: All stocking information is from ADF&G SF hatchery records.

^a Hatchery switched to cold water rearing, so it took 2 years to rear a catchable-sized rainbow trout.

^b Catchable releases include catchable-sized and brood stock rainbow trout. “Catchable size” is about 8 inches in length.

^c Cheney and Taku Campbell Lakes were not stocked starting in 2001 due to illegal introduction of northern pike.

^d Includes fish stocked in University (Alaska Pacific University) Lake.

^e Subcatchable releases include fry, smolt, subcatchable-sized fish, and eyed eggs.

Appendix B4.–Rainbow trout stocking in the Anchorage Management Area by year (2006–2015) and site.

Release type	Site	Year									
		2006 ^a	2007	2008	2009	2010	2011	2012	2013	2014	2015
Catchable ^b											
	Airstrip/Willow Pond	616	500	313	200	0	0	2,017	1,534	1,640	2,308
	Alder Pond	507	0	0	0	0	0	1,889	1,444	1,493	1,766
	Beach Lake	2,802	0	0	0	0	0	18,125	14,479	10,747	12,075
	Campbell Creek	1,522	0	0	0	0	0	3,139	2,354	4,465	3,559
	Campbell Pt Lake	837	4,050	6,200	25,271	3,675	919	12,175	8,223	8,880	7,056
	Cheney Lake ^c	0	0	0	9,942	497	124	5,933	10,162	7,100	5,895
	Chester Creek ^d	326	0	0	0	0	0	935	1,000	1,000	902
	Clunie Lake	2,060	14,857	9,136	12,259	0	0	10,135	4,273	10,569	3,617
	Delong Lake	4,319	15,483	11,596	30,883	3,323	962	9,088	8,828	6,489	4,856
	Dishno Lake	0	0	0	0	0	0	0	0	0	0
	Eagle River	0	0	0	0	0	0	0	0	0	0
	Edmonds Lake	395	0	2,256	0	0	0	990	1,500	999	902
	Fire Island L	0	0	0	0	0	0	0	0	0	0
	Fish Lake	100	842	1,004	806	0	0	1,423	1,000	1,748	2,067
	Goose L	0	0	0	0	0	0	0	0	0	0
	Green Lake	408	900	1,188	1,990	100	100	3,765	2,094	2,735	2,330
	Gwen Lake	1,364	6,526	4,644	5,973	0	0	6,551	2,880	4,301	2,948
	Hillberg Lake	676	2,588	3,116	4,843	0	0	4,797	2,000	2,701	1,774
	Jewel Lake	4,999	20,397	13,089	36,946	5,970	973	25,886	13,120	15,461	10,254
	Lake Otis	275	1,856	2,648	771	0	0	1,789	1,500	1,627	1,635
	Lower Explorer Lake	0	0	0	0	0	0	0	0	0	0
	Lower Fire Lake	846	0	0	0	0	0	3,854	997	1,017	0
	Mirror Lake	4,424	19,131	7,880	0	0	0	15,332	16,058	13,758	12,907
	Otter Lake	1,827	0	0	0	0	0	0	0	0	0
	Rabbit Lake	0	0	0	0	0	0	0	0	997	0
	Sand Lake	2,098	5,400	572	0	12,791	916	5,999	11,681	6,039	5,871
	Spring Lake	180	687	621	741	0	0	500	500	500	482
	Sundi Lake	0	0	0		0	0	0	0	0	0
	Taku Campbell Lake ^c	2,674	0	0	0	0	0	5,415	5,000	4,625	4,440

-continued-

Release type	Site	Year									
		2006 ^a	2007	2008	2009	2010	2011	2012	2013	2014	2015
Catchable ^b											
	Tangle Pond	510	0	0	0	0	0	1,023	1,054	1,195	1,255
	Thompson Lake	0	0	0	0	0	0	0	0	0	0
	Triangle Lake	180	687	527	741	0	0	1,304	1,150	1,737	1,007
	University/APU Lake	0	0	0	0	0	0	0	0	0	0
	Upper Six Mile Lake	480	0	0	0	0	0	2,079	2,000	2,262	1,780
	Waldon Lake	375	2,250	2,150	0	0	0	5,788	2,083	2,206	1,969
	Total	34,800	98,161	68,948	133,375	28,366	6,005	151,943	118,927	118,305	95,670
Subcatchable ^c											
	Beach Lake	0	0	0	0	0	0	0	0	0	0
	Campbell Pt Lake	0	0	0	0	10,462	0	0	0	0	0
	Cheney Lake ^c	0	0	0	0	28,421	0	0	0	0	0
	Chester Creek ^d	0	0	0	0	0	0	0	0	0	0
	Clunie Lake	0	0	0	0	21,125	0	0	0	0	0
	Delong Lake	0	0	0	0	25,293	0	0	0	0	0
	Dishno Lake	0	0	0	0	0	0	0	0	0	0
	Eagle River	0	0	0	0	0	0	0	0	0	0
	Edmonds Lake	0	0	0	0	0	0	0	0	0	0
	Eklutna Lake	0	0	0	0	0	0	0	0	0	0
	Fish Lake	0	0	0	0	2,025	0	0	0	0	0
	Green Lake	0	0	0	0	16,755	0	0	0	0	0
	Gwen Lake	0	0	0	0	7,375	0	0	0	0	0
	Hillberg Lake	0	0	0	0	19,455	0	0	0	0	0
	Jewel Lake	0	0	0	0	24,956	0	0	0	0	0
	Lake Otis	0	0	0	0	2,970	0	0	0	0	0
	Lower Fire Lake	0	0	0	0	0	0	0	0	0	0
	Mirror Lake	0	0	0	0	0	0	0	0	0	0
	Otter Lake	0	0	0	0	0	0	0	0	0	0
	Sand Lake	0	0	0	0	36,150	180	0	0	0	0
	Spring Lake	0	0	0	0	1,015	0	0	0	0	0

-continued-

Release type	Site	Year									
		2006 ^a	2007	2008	2009	2010	2011	2012	2013	2014	2015
Subcatchable ^e											
	Taku Campbell Lake ^c	0	0	0	0	0	0	0	0	0	0
	Tangle Pond	0	0	0	0	0	0	0	0	0	0
	Thompson Lake	0	0	0	0	0	0	0	0	0	0
	Triangle Lake	0	0	0	0	2,025	0	0	0	0	0
	Upper Six Mile Lake	0	0	0	0	0	0	0	0	0	0
	University–APU Lake	0	0	0	0	0	0	0	0	0	0
	Waldon Lake	0	0	0	0	2,375	0	0	0	0	0
	Total	0	2,007	2,008	2,009	202,412	2,191	2,012	2,013	2,014	2,015

Source: All stocking information is from ADF&G SF hatchery records.

^a Hatchery switched to cold water rearing, so it took 2 years to rear a catchable-sized rainbow trout.

^b Catchable releases include catchable-sized and brood stock rainbow trout. “Catchable size” is about 8 inches in length.

^c Cheney and Taku Campbell Lakes were not stocked starting in 2001 due to illegal introduction of northern pike.

^d Includes fish stocked in University (Alaska Pacific University) Lake.

^e Subcatchable releases include fry, smolt, subcatchable-sized fish, and eyed eggs.

Appendix B5.—Arctic char stocking in the Anchorage Management Area by year (1996–2015) and lake.

Year	Lake									Total
	Campbell Point	Cheney	Clunie	Delong	Jewel	Mirror	Sand	Tangle Pond	Airstrip pond	
1996	0	0	0	0	0	0	0	0	0	0
1997	1,000	0	1,000	0	0	2,000	0	0	0	4,000
1998	852	40	2,133	0	0	3,908	0	0	0	6,933
1999	0	0	0	0	0	0	0	0	0	0
2000	1,027	0	0	0	0	2,012	0	0	0	3,039
2001	0	0	0	0	0	0	0	0	0	0
2002	2,094	0	4,387	14,820	4,000	4,845	2,522	503	0	33,171
2003	1,796	0	4,496	4,400	4,035	6,117	4,522	503	0	25,869
2004	2,096	0	0	0	0	0	2,603	0	0	4,699
2005	1,928	0	0	0	0	0	2,194	0	0	4,122
2006	2,904	0	0	0	1	0	4,332	0	0	7,237
2007	1,142	0	0	0	258	0	6,121	0	0	7,521
2008	2,102	0	0	0	0	0	3188	0	0	5,290
2009	2,017	0	0	0	0	0	0	0	0	2,017
2010	1,533	0	0	0	0	0	7,434	0	0	8,967
2011	1,301	0	769	0	0	0	2,481	0	150	4,701
2012	1,372	0	1,011	0	2,814	0	1,606	0	0	6,803
2013	2,133	0	1,100	0	0	0	2,665	0	0	5,898
2014	1,796	0	988	0	0	0	2,233	0	0	5,017
2015	1,940	0	999	0	0	0	2,170	0	0	5,109

Source: All stocking information is from ADF&G SF hatchery records.

Note: Unless otherwise noted, releases are of catchable-sized Arctic char.

Appendix B6.–Arctic grayling stocking in the Anchorage Management Area by year (1996–2015) and site.

Year	Site								Total
	APU Lake ^a	Beach Lake	Lower Fire Lake ^b	Tangle Pond ^c	Sand Lake	Walden Lake ^c	Airport–Willow Pond ^d	Symphony Lake	
1996		8,000	7,500		0			0	15,500
1997		0			0			0	0
1998		0			0			0	0
1999		1,048			0			0	1,048
2000		0			0			0	0
2001 ^c		4,749			0			2,936	7,685
2002 ^c		4,199			0			0	4,199
2003		7,081 ^d			0			4,239	11,320
2004		4,489			0			0	4,489
2005		279			0			0	279
2006		4,080			0			0	4,080
2007		5,668			0			0	5,668
2008		0			0			0	0
2009		3,192			0			0	3,192
2010		3,034			7,885			0	10,919
2011		215			6,753			0	6,968
2012		4,694			1,218			0	5,912
2013		1,000			1,477			0	2,477
2014		1,559			1,385			0	2,944
2015		1,402			1,591			0	2,993

Source: All stocking information is from ADF&G SF hatchery records.

Note: 2013 was the first year of releases that came from the WJHSF Hatchery.

Note: Unless otherwise noted, all releases are fingerlings.

^a “APU” is Alaska Pacific University. Stocking was discontinued in this Lake in 1991.

^b Stocking was discontinued in 1997.

^c Stocking was discontinued in 1995.

^d Stocking was discontinued in 1993.

APPENDIX C: ESCAPEMENT ESTIMATES

Appendix C1.—Ship Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.

Year	Chinook salmon				Coho salmon			
	Adults collected at hatchery	Used for egg take	Survey	Estimated escapement ^a	Adults collected at hatchery	Used for egg take	Survey	Estimated escapement ^a
1996	171	^b	503	674			1,013	^c
1997	108	^b	447	555			1,205	^c
1998	103	^b	360	463			1,090	^c
1999	328	^b	516	844		434	585	1,019 ^d
2000	341	^b	323	664	348	305	815	1,163
2001	258	^b	541	799	619	514	968	1,587
2002	467	336	1,492	1,959	727	378	4,225	4,952
2003	535	232	2,198	2,733	323	295	1,143	1,466
2004	468	352	1,606	2,074	953	514	1,262	2,215
2005	343	215	1,485	1,828	575	533	466	1,041
2006	458	294	1,431	1,889	757	583	1,431	2,188
2007	371	256	1,475	1,846	459	432	254	713
2008	468	252	833	1,301	820	508	891	1,711
2009	379	366	916	1,295	777	648	1,066	1,843
2010	30	30	368	398	562	446	189	751
2011	408	290	896	1,304	1,201	828	1,186	2,387
2012	569	321	227	796	1,208	534	836	2,044 ^e
2013	571	342	468	1,039	954	665	691	1,645
2014	1,048	447	423	1,471	2,239	782	2,203	4,442
2015	1,286	492	1,161	2,447	2,220	754	811	3,031
Average								
2011–2015	776	378	635	1,411	1,564	713	1,145	2,710
2006–2015	559	309	820	1,379	1,120	618	956	2,076

Note: These surveys are a combination of trap collection counts and foot surveys. Trap collection was designed to capture broodstock and not to enumerate fish. Foot surveys were designed to manage the fishery and ensure that the hatchery collected the required broodstock. After 2001, all surveys were foot only.

Note: From 1996 to 1999, the number of Chinook salmon collected and used for egg takes are unknown. The number listed was documented only as “brood.”

^a Estimated escapement is the survey counts plus the number of adults collected at the hatchery. Survey dates vary by year and final escapement of salmon into the stream could be higher or lower but can’t be verified due to varying survey conditions from year to year.

^b Total number used for egg takes is unknown.

^c Estimated escapement number would be survey number only. Accurate records for adults that made it to the hatchery are unknown.

^d Total number of adults collected is unknown. The egg take number was used. The actual escapement is probably larger than reported.

^e Survey was incomplete. Only surveyed to Reeves Blvd. Actual escapement is probably higher.

Appendix C2.–Eagle River salmon escapement estimates, Anchorage Management Area, 1996–2015.

Year	Chinook	Coho	Sockeye	Pink	Chum
1963	135	200			
1996	141 ^a				
1997	412				
1998	163 ^b				
1999	224				
2000 ^c	^b				
2001	77 ^b			19	
2002	27 ^b				
2003	167 ^b				
2004	157				
2005	122 ^d				
2006	101 ^d				
2007	117 ^d				
2008	156 ^d				
2009	152 ^d				
2010	10 ^d				
2011	7 ^d				
2012					
2013					
2014					
2015					
Average					
2011–2015	7				
2006–2015	91				

Note: Estimates are from foot surveys designed for Chinook salmon only.

^a Survey conducted after spawning occurred.

^b High water and poor visibility.

^c No survey conducted.

^d Surveys include Meadow Creek.

Appendix C3.–Campbell Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.

Year	Chinook	Coho	Sockeye	Pink	Chum
1996	369	1,612	349		
1997	1,119	1,007	294		
1998	761	2,968	646		
1999	1,035	537	435		
2000	591	3,196	109		
2001	717	2,377 ^a	163		
2002	744	7,574	1,473		
2003	745	1,799	1,857		
2004	964	713	776		
2005	1,097	1,130	654		
2006	1,052	542	589	3	1
2007	588	^b	203		
2008	439	403 ^c	42		
2009	554	766	69		
2010	290	157	249		
2011	260	^b	434		
2012	^b	^b	^b		
2013	^b	146	^b		
2014	274	3,316	368		
2015	654	1,178	585		
Average					
2011–2015	396	1,547	462		
2006–2015	514	930	317		

Note: Estimates are from foot surveys designed for Chinook, sockeye, and coho salmon.

^a Only South Fork and mainstem from Forks to Folker Street were counted.

^b No survey conducted.

^c Only North Fork, South Fork, and mainstem from Forks to Folker Street were counted.

Appendix C4.–Bird Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.

Year	Chinook	Coho	Sockeye	Pink	Chum
1984	21			420	
1996	212	169		987	214
1997	231	603		8	18
1998	131	1,446		7	5
1999	497	279		1,255	75
2000	117	703		1,873	12
2001	88	1,554	7	2,828	228
2002	48	66		1,341	633
2003	140	4	8	2,925	242
2004	307	376	4	1,902	234
2005	29	619	0	450	1
2006	^a	442			
2007	173	^a	2	^a	^a
2008	106	115		5,357	62
2009	148	278	2	32,100	179
2010	12	19		^a	
2011	112	45		^a	1
2012	^a	^a			
2013	^a	^a			
2014	^b	629			
2015	87	^a			
Average					
2011–2015	100	337			
2006–2015	106	255			

Note: Estimates are from foot surveys designed for Chinook and coho salmon.

^a No survey conducted.

^b Incomplete survey.

Appendix C5.—Rabbit Creek salmon escapement estimates, Anchorage Management Area, 1996–2015.

Year	Chinook	Coho	Sockeye	Pink	Chum
1986	10	169			
1993-1996 ^a					
1997	31				
1998	41	309		43	
1999-2000 ^a					
2001	64	697	300 ^b	7	
2002	9	1,243	0	1,004	
2003	7	348	0	33	
2004	55	1,448	0	234	
2005	73	7	71	257	
2006	39	24	275	118	
2007 ^a					
2008	15	109	4	1,562	1
2009	36	1,725	1	524	0
2010	16	7	15	0	0
2011	68		39	1	0
2012 ^a					
2013 ^a					
2014	35		14	18	
2015					
Average					
2011–2015	52		27	10	0
2006–2015	35	466	58	371	0

Note: Estimates are from foot surveys designed for Chinook and coho salmon.

^a No survey conducted.

^b Estimated from boardwalk at Potter Marsh.