

**Annual Report to the Alaska Board of Game on
Intensive Management for Caribou
with Wolf Predation Control
in Game Management Units 9B, 17B&C, and 19A&B,
the Mulchatna Caribou Herd**

**Prepared by the Division of Wildlife Conservation
February 2019**



1) **Description of IM Program¹ and Department recommendation for reporting period**

A) **This report is an annual evaluation for a predation control program authorized by the Alaska Board of Game (Board) under 5 AAC 92.111²**

B) **Month this report was submitted by the Department to the Board:**

February X (annual report) Year 2019

C) **Program name: Mulchatna Caribou Herd Predation Management Area**

D) **Existing program does not have an associated Operational Plan, it does have a detailed Intensive Management Plan in regulation (5 AAC 92.111).**

E) **Game Management Units (Units) fully or partly included in IM program area:**
Units 9B, 17B&C, and 19A&B

F) **IM objectives for caribou: population size 30,000–80,000 harvest 2,400–8,000.**

G) **Month and year the current predation control program was originally authorized by the Board:**

The plan was initially authorized in March 2011 for Units 9B and 17B&C and was modified in March 2012 to include Units 19A&B.

H) **Predation control is currently active in this IM area.**

I) **If active, month and year the current predation control program began:**

- March 1, 2012 in Regulatory Year (RY) 2011 (RY 2011 = July 1, 2011 through June 30, 2012).
- Reauthorized in March 2017 for six more years.

J) **A habitat management program funded by the Department or from other sources is currently active in this IM area (Y/N): N**

K) **Size of IM program area (square miles) and geographic description:**

39,683 sq. miles in Units 9B, 17B&C, and 19A&B.

L) **Size and geographic description of area for assessing ungulate abundance:**

Approximately 50,000 sq. miles and includes the range of the Mulchatna Caribou Herd, and extends beyond Units 9B&C, 17B&C, and 19A&B into Unit 18.

¹ For purpose and context of this report format, see *Intensive Management Protocol, section on Tools for Program Implementation and Assessment*

² [Regulatory numbers for existing IM programs formerly under 5AAC92.125 were divided into groups and given new numbers in October 2012 (see IM Plan template--Version 3, January 2013)]

³ The interim annual update may be limited only to sections that changed substantially since prior annual report [e.g., only Tables 3 and 6 in areas with a fall ungulate survey and only wolf control]

M) Size and geographic description of area for ungulate harvest reporting:

Approximately 50,000 sq. miles and includes the range of the Mulchatna Caribou Herd and extends beyond Units 9B&C, 17B&C, and 19A&B into Unit 18.

N) Size and geographic description of area for assessing predator abundance:

The area for assessing predator abundance includes all of the predation control area and is described in 'O' below.

O) Size and geographic description of predation control area:

Initially the predation control area designed in RY12 was named the Mulchatna Wolf Control Area. However this area was expanded in RY17 to include additional calving grounds and adjacent habitat used by the Mulchatna herd since RY13. Because this newly expanded area has a different starting date for SDA hunters (December 1) than the original area (February 1), we refer to these areas with different names. The initial control area was renamed, Kemuk Wolf Control Area (KWCA) while the newly added area is named Greater Mulchatna Wolf Control Area (GMWCA). Both areas combined include approximately 9,844 mi².

Kemuk Wolf Control Area:

That portion of Unit 17B south of a line between Tikchik Mountain (N 60.05, W 158.300) and Sleitat Mountain (N 60.05, W 157.067), then southeast to the Kuktuli Hills (N 59.80, W 156.300), then southwest into 17C to a point at N 59.32, W 157.066, then west to N 59.32, W 158.300, then north returning into 17B to the beginning point at Tikchik Mountain (N 60.05, W 158.300).

Greater Mulchatna Wolf Control Area:

That portion of Unit 17B east of a line between Tikchik Mountain (N 60.05, W 158.300) north to a point south of the Shotgun Hills (N 60.37, W 158.300), then east to the headwaters of Klutapak Creek (N 60.37, W 157.379), then a line northeast to a point on the Unit 17B/19B boundary (N 60.68, W 156.841) into Unit 19B northeast to a point at the junction of the S. Fork Hoholitna River and the Hoholitna River (N 60.91, W 156.243), then track east just south of the north bank of the Hoholitna River to a point at the mouth of Whitefish Lake (N 60.94, W 154.993), then a line east to a corner point at N 60.94, W 154.595, then south into Unit 17B and across the upper Mulchatna River to a point N 60.78, W 154.595, then east to N 60.77, W 154.539, south to N 60.58, W 154.539, then southwest to N 60.52, W 154.619, west to N 60.52, W 154.747 on the Unit 17B/9B boundary, then south into Unit 9B to a point N 60.42, W 154.746 and southwest crossing through a portion of Unit 9B and back into Unit 17B, crossing the Kuktuli River to the Unit 17B/9B boundary (N 59.78, W 155.566), then southwest across Unit 9B to the Unit 9B/17C boundary (N 59.33, W 155.884), then west along the drainage of Lower Klutuk Creek to the Unit 9B/17C boundary (N 59.32, W 156.988), then west to the southeast point of the Kemuk WCA boundary (N 59.32, W 157.067) then excluding the entirety of the Kemuk WCA, track northeast on the Kemuk WCA boundary to

the Kuktuli Hills (N 59.80, W 156.300), northwest to Sleitat Mountain (N 60.05, W 157.067), then west returning to the starting point at Tikchik Mountain (N 60.05, W 158.300).

P) Criteria for evaluating progress toward IM objectives:

- Fall calf-to-cow ratios
- Fall bull-to-cow ratio
- Caribou abundance

Q) Criteria for success with this program:

- Fall bull-to-cow ratio can be maintained at a minimum of 35 bulls:100 cows.
- Fall calf-to-cow ratio can be sustained above 30 calves:100 cows.
- The population can grow at a sustained rate of 5% annually.
- Caribou harvest objectives are met.

R) Department recommendation for IM program in this reporting period:

The Department recommends continuation of the predation control program during RY2018. We will continue monitoring the Mulchatna Caribou herd to determine progress towards IM objectives (details provided in Section 6).

S) IM Annual Report data and information inclusion date:

February X (annual report) Year 2019




Figure 1. Location of the Kemuk and Greater Mulchatna Wolf Control Areas in Game Management Units 9B, 17B&C, and 19B.

2) Prey data

Date(s) and method of most recent summer abundance assessment for caribou (if statistical variation available, describe method here and show result in Table 1): The last successful photocensus of post-calving aggregation was conducted on June 28–29, 2016.

Compared to IM area, was a similar trend and magnitude of difference in abundance observed in nearby non-treatment area(s) since program inception (Y/N) N/A and in the last year (Y/N) N/A? Describe comparison if necessary:

The IM area comprises a small portion of the annual range of the Mulchatna caribou herd. The annual range of the majority of caribou in the herd includes use of areas both within and outside of the IM area, but the spatial and temporal characteristics of movements within the IM area are variable. Therefore, it is difficult to quantify trends in abundance relative to treatment and non-treatment areas.

Date(s) of most recent age and sex composition survey (if statistical variation

available, describe method here and show result in Table 1): October 12–13, 2018

Compared to IM area, was a similar composition trend and magnitude of difference in composition observed in nearby non-treatment area(s) since program inception (Y/N) N/A and in the last year (Y/N) N/A? Describe comparison if necessary:

The IM area comprises a small portion of the annual range of the Mulchatna caribou herd and was initially aligned closely with the calving ground of the western segment of the population (RY2011–2013), and the summer and winter grounds of the eastern segment of the population. In recent years however (RY2014–2016), this western segment of the population calved outside the wolf control area, but close enough that they still may have benefitted from any removal of wolves. Teasing out treatment and non-treatment effects was compounded by the fact that these two areas were too close spatially to really be considered independent of one another. The composition data in Table 1 suggests the caribou in the western segment of the population were most successful in rearing calves during RY2011–2013 when they were calving within the wolf removal area. During RY2017, the wolf control area was expanded to include much of the calving grounds of the eastern segment of the Mulchatna herd in the upper Mulchatna River. Both portions of the herd experienced relatively high calf abundance in fall of 2018, likely related to a high wolf harvest across the wolf control area, (Table 1). At this point we are unable to accommodate a true experimental versus control comparison given the nearness in proximity of both calving grounds to the wolf control area.

The combined calf-to-cow ratio of 34 calves:100 cows in RY2018 was higher than RY2017. For the first time since 2014 it reached, and also exceeded the objective of 30:100 and is the highest since the control program started. The combined bull:100 cows stayed the same as in RY16, at 32 bulls:100 cows, still below the high of 39 bulls:100 cows in RY16, and below the objective of 35 bulls:100 cows.

Table 1. Caribou abundance, age and sex composition in assessment area (L) since program implementation in year 1 (not exclusively limited to inception of predation control) to reauthorization review in year 2017 in Mulchatna Caribou Herd Predation Management Area. Regulatory year is 1 July to 30 June (e.g, RY 2010 is 1 July 2010 to 30 June 2011).

Eastern Segment of the MCH

		Composition (number per 100 cows)		
Period	RY	Calves	Bulls	Total (n)
Year 0	2010	17	13	2,581
Year 1	2011	14	18	2,649
Year 2	2012	22	17	2,217
Year 3	2013	14	27	1,479
Year 4	2014	33	31	2,226

Year 5	2015	31	32	2,827
Year 6	2016	27	38	2,525
Year 7	2017	28	33	2,587
Year 8	2018	39	33	2,515

Western Segment of the MCH

		Composition (number per 100 cows)		
Period	RY	Calves	Bulls	Total (n)
Year 0	2010	23	23	2,011
Year 1	2011	28	34	1,995
Year 2	2012	38	29	2,636
Year 3	2013	23	27	1,743
Year 4	2014	27	38	2,567
Year 5	2015	27	38	2,587
Year 6	2016	18	40	2,670
Year 7	2017	18	31	2,573

All Areas Combined

		Composition (number per 100 cows)			
Period	RY	Abundance (variation)	Calves	Bulls	Total (n)
Year 0	2010	-	20	17	4,592
Year 1	2011	-	19	22	5,282 ^a
Year 2	2012	19,000-27,000 ^b	30	23	4,853
Year 3	2013	15,000-22,000 ^b	19	27	3,222
Year 4	2014	21,000-32,000	30	35	4,793
Year 5	2015	30,736-38,190	29	35	5,414
Year 6	2016	21,346-33,137	22	39	5,195
Year 7	2017	-	23	32	5,160

^a Includes caribou not assigned to the Eastern or Western Segment of the MCH.

^b Estimate of abundance based on the Rivest methodology (Rivest et al. 1998).

Describe trend in abundance or composition:

Trends in calf-to-cow ratios are variable from year to year and are still below those observed in the late 1980s–early 1990s when the herd was in a significant growth phase. Bull-to-cow ratios were on a positive trend and improved each year during RY2010–2016 but declined in RY2017. Both of these metrics are below objectives at this time. The RY2016 point estimate for abundance of 27,242 +/- 5,896 is lower than RY2015 (30,736 +/- 3,727), but higher than RY2014 (26,275 +/- 5,500). The overlap in confidence intervals between these estimates does not allow for a good understanding of the current population trend.

Table 2. Caribou harvest in assessment area (M). Methods for estimating unreported

harvest are described in Survey and Inventory reports.

Period	RY	Reported			Estimated		Total harvest	Other mortality ^a	Total
		Male	Female	Unk Sex	Unreported	Illegal			
Year 0	2010 ^b	250	220	4	Unk	Unk	470	Unk	474
Year 1	2011 ^b	242	243	9	Unk	Unk	494	Unk	494
Year 2	2012 ^b	184	173	4	Unk	Unk	361	Unk	361
Year 3	2013 ^c	70	35	1	Unk	Unk	106	Unk	106
Year 4	2014 ^c	125	52	5	Unk	Unk	182	Unk	182
Year 5	2015 ^c	159	74	2	Unk	Unk	235	Unk	235
Year 6	2016 ^c	209	119	2	Unk	Unk	330	Unk	330
Year 7	2017 ^c	250	186	4	Unk	Unk	440	Unk	440

^a Clarify (vehicle mortality, Defense of Life and Property, Mortuary, etc.).

^b Data from WinfoNet, Harvest Information, Data Download (harvest report cards).

^c Data from WinfoNet, Permitting, Hunt Statistics, General Hunt, RY, RC503.

Describe trend in harvest:

Although the reported harvest is still below objectives (2,400 – 8,000), the harvest has steadily increased from the low in RY2013. Since RY2011, the majority of harvest has been during late winter. The majority of hunters are local residents (i.e. people who live within the herd’s range, primarily residents of Unit 18). Marginal snow conditions RY2013–RY2015 prevented hunters from accessing caribou with snowmachines resulting in low harvest. Improved snow conditions in RYs 2016 and 2017 enabled hunters to access caribou by snowmachine which increased hunting success. We suspect that the actual harvest is substantially higher than the reported harvest in some years.

Describe any other harvest related trend if appropriate:

NA

3) Predator data

Date(s) and method of most recent spring abundance assessment for wolves (if statistical variation available, describe method here and list in Table 3):

See below.

Date(s) and method of most recent fall abundance assessment for wolves (if statistical variation available, describe method here and list in Table 3):

See below.

Other research or evidence of trend or abundance status in wolves:

Sightings of wolves in the IM area by local residents and department personnel continued into RY2017. Decreased numbers of public phone calls reporting wolf observations may be partially due to this year’s lack of good snow cover for tracking and spotting wolves.

In March 2017, the department initiated a study including deployment of GPS collars on wolf packs in the IM area. The objectives of the study are to map wolf pack territories,

determine seasonal pack sizes, and evaluate change in wolf density relative to the wolf removal program. During the initial capture field work, wolf tracks were common and found throughout much of the MCH WCA. Seventeen wolves were collared, comprising 5 packs and multiple lone wolves. Mean minimum observed pack size was 6 wolves during spring and 9 wolves during fall 2017. A preliminary density calculation based on 7 months of GPS data and minimum observed seasonal pack sizes resulted in spring and fall 2017 wolf densities of 2.2 and 3.0 wolves per 1000 km², respectively, in the Mulchatna and lower Nushagak River drainages. The estimated fall density of 3.0 wolves calculates to a minimum of 76 wolves comprising the packs that inhabit the MCH WCA. This estimate should be viewed cautiously, as we did not have all the known packs within the WCA collared. Further, the estimate does not include lone wolves, which are known to occur in the WCA, but that we have insufficient data to estimate their numbers.

In RY2017, favorable snow conditions and an expanded WCA boundary facilitated the highest reported wolf harvest since the wolf control program was initiated. A total of 78 wolves were reported harvested in the WCA, including 9 of 12 (75%) remaining radiocollared wolves. The density of harvested wolves alone equals 3.1 wolves per 1000 km², and in comparison to the minimum estimate of pack dwelling wolves previously mentioned, indicates a significant population reduction was obtained during RY2017. Observations during wolf capture operations in April 2018 were that both the occurrence and distribution of wolf tracks was down substantially from the previous spring, and the majority of sets of tracks encountered were of singles or pairs of wolves. During that effort we only found a total of 5 additional wolves; 1 breeding pair and 3 lone females. The 2018 mean spring pack size was 2 wolves. Three packs produced a minimum of 16 pups during the summer, and 2018 fall mean pack size was 7 wolves.

Table 3. Wolf abundance objectives and removal in the Kemuk and Greater Mulchatna Wolf Control Areas (WCAs). Removal objective is to annually remove 100 % of the wolves in the wolf control areas, so the estimated or confirmed number remaining in the control area by the May calving season each regulatory year is 0.

Subunits 9B and 17B&C, and 19B (Subunits 19A is outside of WCAs)

Period ^a	RY	Non-SDA Harvest removal from WCAs		SDA Public control removal from WCAs	Total removal ^b from WCAs	Minimum Spring abundance (variation) WCAs
		Trap	Hunt			
Year 1	2011	14	52	11	77	14
Year 2	2012	17	0	0	17	-
Year 3	2013	0	10	0	10	-
Year 4	2014	0	0	0	0	-
Year 5	2015	19	2	0	21	-
Year 6	2016	26	28	3	57	-
Year 7	2017	38	9	31	78	-

^a Each respective year of data is from the ADF&G Winfonet database.

^b Additional removal may be Defense of Life and Property, vehicle kill, etc.

4) Habitat data and nutritional condition of prey species

Where active habitat enhancement is occurring or was recommended in the Operational Plan, describe progress toward objectives:

Objective(s):

Not Applicable: There are no demonstrated methods to improve caribou habitat and no reason to believe that habitat is limiting the caribou population.

Area treated and method: N/A

Observation on treatment response: N/A

Evidence of progress toward objective(s) (choose one: Apparent Statistical): N/A

Similar trend in nearby non-treatment areas? N/A

Describe any substantial change in habitat not caused by active program: N/A

Table 4. Nutritional indicators for caribou in assessment area (L) of the Mulchatna Caribou herd Predation Management Area.

Period	RY	Pregnancy Females >2 yrs age ^a	Female Calf Weights ^b at 10.5 months in lbs. (n)
Year 0	2010	79%	124 (20)
Year 1	2011	76%	119 (13)
Year 2	2012	79%	127 (14)
Year 3	2013	90%	128 (14)
Year 4 ^c	2014	61%	133 (13)
Year 5	2015	83%	119 (23)
Year 6	2016	73%	120 (18)
Year 7	2017	80%	122 (15)

^a Pregnancy rate is based on known-aged animals from a collared sample of adult female caribou. Pregnancy status is determined in May, i.e., RY 2010 pregnancy data is collected in May 2011, based on observed characteristics of pregnancy, i.e. antler retention, udder development, and/or presence of a calf at heel.

^b Female calf weight data is collected in April of the RY, i.e., RY 2010 female calf weight data is collected in April 2011.

^c Survey delayed due to weather which affected sample size and timing of survey.

Where objectives on nutritional condition were listed in the Operational Plan, describe trend in condition indices since inception of (a) habitat enhancement or (b) enhanced harvest: N/A

Evidence of trend: N/A

Similar trend in nearby non-treatment areas? N/A

5) Costs specific to implementing Intensive Management

Table 5. Cost (\$1000 = 1.0) of agency salary based on estimate of proportional time of field level staff and cost of operations for intensive management activities (e.g., predator control or habitat enhancement beyond normal Survey and Inventory work) performed by personnel in the Department or work by other state agencies (e.g., Division of Forestry) or contractors in Mulchatna Caribou Herd Predation Management Area. Fiscal year (FY) is also 1 July to 30 June but the year is one greater than the comparable RY (e.g, FY 2010 is 1 July 2009 to 30 June 2010).

Period	FY	Predation control ^a		Other IM activities		Total IM cost	Research cost ^d
		Time ^b	Cost ^c	Time ^b	Cost ^c		
Year 1	2012	0.0	0.0	1.0	36.0	36.0	415.0
Year 2	2013	0.0	0.0	0.5	6.0	6.0	421.2
Year 3	2014	0.0	0.0	0.5	6.0	6.0	215.0
Year 4	2015	0.0	0.0	0.5	6.0	6.0	0.0
Year 5	2016	0.0	0.0	0.5	6.0	6.0	0.0
Year 6	2017	0.0	0.0	1.0	13.0	13.0	230.0
Year 7	2018	0.0	0.0	1.0	13.0	13.0	239.9

^a State or private funds only.

^b Person-months (22 days per month).

^c Salary plus operations.

^d Separate from implementing IM program but beneficial for understanding of ecological or human response to management treatment (scientific approach that is not unique to IM).

6) Department recommendations² for annual evaluation (1 February) following Year 7(RY2017) for the Mulchatna Caribou herd Predation Management Area

Has progress toward defined criteria been achieved?

Yes; The fall bull-to-cow ratio increased each year of the program until fall 2017 when it declined. It was at or above objectives during RY2014–2016 for the first time since RY2000 and remained steady at just below objective during RY2017-2018. The calf-to-cow ratios are very erratic, but did meet (RY2012, RY2014) or exceed (RY2018)

² Prior sections include primarily objective information from field surveys; Sections 6 and 7 involve professional judgment by area biologists to interpret the context of prior information for the species in the management area.

management objectives since the initiation of this IM program. The abundance estimate shows an increase of 19% during RY2012–2016 when comparing the point estimates, although the RY2016 estimate showed a decline from the previous year. However, the confidence intervals are overlapping on these estimates, so care must be taken in interpreting the results.

Has achievement of success criteria occurred?

We have mixed results in meeting the objectives of the success criteria. The bull:100 cow objective of 35:100 was met during RY2014–RY2016, but not during RY2017–2018. The calf:100 cow objective of 30:100 was met in RY2014 (30:100), narrowly missed in RY2015 (29:100), well below in RY2016–2017 (22 and 23:100 respectively) and then exceeded in RY2018 (34:100). The point estimates for the abundance estimates indicate growth in the herd during RY2012–2016, but are still below the lower bound of the population objective. Although the harvest objective of 2,400–8,000 has not been met, it has consistently increased from just under the low 4.5% of the lower objective in RY2013, to 18% in RY2018. Failure to meet the harvest objective is partially due to lack of opportunity stemming from lack of snow and frozen rivers providing access, and may also be the result of failure of hunters to report harvest.

Recommendation for IM program (choose one): Continue Modify Suspend Terminate

In fall of 2017, the WCA was expanded to 9,844 sq. mi., near the maximum 10,000 sq. mi. authorized in regulation.

Continue Same-Day Airborne Wolf Control Program in expanded WCAs.

7) Evaluation (1 February) for program renewal (following final Year 14 [RY 2024]) and Department recommendations for the Mulchatna Caribou herd Predation Management Area

Has progress toward defined criteria been achieved (describe)? See Section 6.

Has achievement of success criteria occurred (describe)? See Section 6.

Recommendation for IM program: Continue Modify Suspend Terminate

Rationale for recommendation on overall program: The program was successful in RY17 because of much improved tracking and flying conditions for SDA hunters. Increased calf:100 cow ratios following the successful RY17 season suggest the program had a significant positive impact on calf survival.

Other recommendations (if continuation is recommended, specific actions on individual practices): The increased area of the expanded WCAs increased the number of susceptible wolf packs and provided for targeted efforts in calving areas; therefore, greatly increasing the chances for a successful program. Radio-collared wolves that were subsequently removed by SDA pilots

and gunners indicate the SDA program is of value to the wolf control program.