



State of Alaska
Department of Fish and Game
Sportfish Division

Nomination Form
Anadromous Waters Catalog

MMP JB

Region Interior USGS Quad(s) COLEEN D-3 C3 C4 B4 A4 A-3

Anadromous Waters Catalog Number of Water Body 334-45-11000-2001-3271

Name of Water Body Coleen River USGS Name Local Name

Addition Deletion Correction Backup Information

For Office Use

Nomination #	<u>18-681</u>	<u>James J. Harbuck</u>	<u>10/19/2018</u>
		Fisheries Scientist	Date
Revision Year:	<u>2019</u>	<u>Rob Benker</u>	<u>10/19/18</u>
		Habitat Operations Manager	Date
Revision to:	<input checked="" type="checkbox"/> Atlas <input checked="" type="checkbox"/> Catalog	<u>JH</u>	<u>10 Sept 18</u>
		AWC Project Biologist	Date
Revision Code:	<u>A-1, B-2, B-6</u>		
		GIS Analyst	Date

OBSERVATION INFORMATION

Species	Date(s) Observed	Spawning	Rearing	Present	Anadromous
chum salmon	09/19/2017	✓			✓
<u>Entered River w/ Chum salmon spawning, ran on Chum salmon habitat</u>					

18-681

IMPORTANT: Provide all supporting documentation that this water body is important for the spawning, rearing or migration of anadromous fish, including: number of fish and life stages observed; sampling methods, sampling duration and area sampled; copies of field notes; etc. Attach a copy of a map showing location of mouth and observed upper extent of each species, as well as other information such as: specific stream reaches observed as spawning or rearing habitat; locations, types, and heights of any barriers; etc.

Comments:
We used a R44 helicopter for aerial survey of the Coleen River on September 19, 2017. The survey began at the mouth of the Coleen River. We visually observed migrating fall chum salmon starting at the mouth and up through the upper reach Lat/Long listed above. We visually observed spawning fall chum salmon starting at the lower reach Lat/Long listed above. We observed spawning fall chum salmon from the lower reach up through the upper reach listed above. We did not observe any spawning or migrating fall chum salmon above the upper reach Lat/Long listed above. The aerial survey ended at Strangle Woman Creek. We collected tissue samples from 101 spawning fall chum salmon and provided the samples to ADF&G gene conservation lab. Coordinates listed are WGS 84.
Coordinates (Lat,Long): Upper(67.79942,-142.39149) Lower(67.13314,-142.68033)

Name of Observer (please print): Brian McKenna

Signature: 209.193.24.253 (Web Nomination) Date: 09/08/2018

Agency: _____

Address: 122 1st Avenue
Fairbanks, AK 99701

This certifies that in my best professional judgment and belief the above information is evidence that this waterbody should be included in or deleted from the Anadromous Waters Catalog.

Signature of Area Biologist: _____ Date: _____ Revision 3/16

Name of Area Biologist (please print): _____

Report of Activities for ADF&G FRP SF2017-165



Teedraanjik (Salmon Fork) and Coleen Rivers, Alaska.

**Tanana Chiefs Conference, Fisheries Program
Fairbanks, Alaska
December, 2017**



Tanana Chiefs Conference

The Tanana Chiefs Conference's Fisheries Program strives to continually build educational capacity and expertise in fisheries science and management throughout the TCC region, including the Yukon and Kuskokwim River drainages. Our goals are to utilize western science and traditional knowledge to enable sustainable fisheries, and to advocate for cultural and traditional fishing and hunting rights. We endeavor to accomplish these goals by partnering with other Tribal organizations, NGO's, and State and Federal agencies to better manage, protect, and preserve our fisheries resources.

<http://www.tananachiefs.org/sustainability/fish-wildlife/>

Author:

Brian C. McKenna
Tanana Chiefs Conference
Wildlife & Parks Department
201 First Avenue, Suite 300
Fairbanks, Alaska 99701
Phone: 907-452-8251 ext. 3318
Brian.mckenna@tananachiefs.org

Overview

The purpose of this project was to collect baseline data from Chinook and fall chum salmon populations spawning in the Teedraanjik (Salmon Fork) and Coleen River watersheds. We used a combination of aerial and float surveys of the spawning grounds to document and delineate the spatial extent of spawning activity for salmon species in both rivers. Stream temperature data loggers were placed near active spawning sites, and will be collected in 2018 to describe water temperature profiles during the incubation period. Baseline data such as age, sex, and length were collected to determine population structure, and tissue samples were collected for genetic stock identification. Analyses of the baseline data collections will be completed after the conclusion of the 2018 sampling season. Nominations to the anadromous waters catalog will also be completed after completion of the 2018 field season.

Objectives

1. Conduct aerial surveys of Chinook and fall chum salmon spawning in the Teedraanjik and Coleen rivers to determine spatial extent and distribution of spawning activity.
2. Submit nominations to the State of Alaska's Anadromous Waters Catalog (AWC) if/when salmon are positively identified in areas presently not included within the AWC.
3. Determine age, sex, and length composition of adult Chinook and fall chum salmon populations in the Teedraanjik and Coleen rivers.
4. Collect axillary process tissue samples appropriate for genetic (DNA) analyses from Chinook and fall chum salmon spawning in the Teedraanjik and Coleen rivers.
5. Deploy stream temperature loggers in the Teedraanjik and Coleen rivers to monitor and characterize spawning habitat for Chinook and fall chum salmon.

Study Area:

The Teedraanjik, also known as the Salmon Fork of the Black River, is a clear water tributary of the Draanjik (Black River) and Porcupine Rivers. The Teedraanjik is a trans-boundary river originating in the North Ogilvie Mountains of Yukon Territory, Canada (Yukon Ecoregions Working Group, 2004) (Figure 1). It flows southwesterly crossing the international border just south of the Arctic Circle. West of the international boundary, the Teedraanjik crosses the Porcupine Plateau and flows westerly for roughly 74 river miles to the confluence with the Draanjik. Within Alaska, the Teedraanjik drains large portions of the Bureau of Land Management's (BLM) Eastern Interior Region, as well as the Yukon Flats National Wildlife Refuge (YFNWR).

The Coleen River is a clear water tributary of the Porcupine River located in northeastern Alaska within the Arctic National Wildlife Refuge. It originates in the Davidson Mountains and drains a portion of the southern Brooks Range flowing for 186 river miles before emptying into the Porcupine River (Figure 1).

Methods:

In the Teedraanjik watershed we used a R44 helicopter to define the spatial extent of spawning activity, collect baseline samples, and to deploy stream temperature data loggers for both Chinook and fall chum salmon. In the Coleen River watershed we used a combination of methods including a fixed wing wheeled aircraft, a R44 helicopter, and an inflatable canoe. We used the fixed wing aircraft and R44 helicopter to define the spatial extent of spawning activity in the Coleen River watershed. We also utilized the fixed wing aircraft as a transport so we could float the Coleen River to collect baseline samples and deploy stream temperature data loggers.

Rod and reel capture technique was used to capture fish for sampling. Scales were collected for aging analysis, sex was determined by visual observations, mid-eye to fork length measurements were recorded, and tissue samples were collected. All captured fish were released alive immediately after being sampled. Fresh carcasses were also sampled, and returned to the river.

Live fish, or recently deceased fish with red gills, were sampled on the spawning grounds. Collections were accomplished using rod and reel. Tissue samples were collected by clipping approximately a ½ - 1" portion of the pelvic axillary process of each individual fish according to the ADF&G Gene Conservation Laboratory's (GCL) protocol (Appendix 1). Tissue samples were stored in bulk vials partially filled with anhydrous ethyl alcohol. The bulk vials were shipped to the ADF&G GCL. Samples were shared with the USFWS and with the DFO-Canada. The samples will be genotyped by the ADF&G GCL, and will be used to improve the existing genetic baseline.

Onset Tidbit V2 stream temperature loggers were deployed in Chinook and fall chum salmon spawning locations in the Teedraanjik and Coleen watersheds. Tidbits were anchored into the substrate using duckbill anchors with stainless steel cable. They were marked with pink surveyor markers and flagging tape. GPS data was recorded for each tidbit location. The data loggers will be collected in the summer of 2018.

Chinook Salmon:

The Coleen River was surveyed via a fixed wing wheeled aircraft on July 22, 2017. Sampling occurred on the Coleen River via a float trip using an inflatable Soar Pro Pioneer canoe; July 23 and 26, 2017. A R44 helicopter was also used for one day, August 5, 2017, to collect additional samples in the Coleen River. We did not use a fixed wing aircraft, nor did we float the Teedraanjik. A R44 helicopter was used exclusively to survey and sample the Teedraanjik watershed. Teedraanjik surveys and sampling occurred between August 2 and 5, 2017. The fixed wing flight on the Coleen originated in Fairbanks. For the R44 surveys, we based operation in Circle, and made daily departures between August 2 and 5, 2017.

Fall Chum Salmon:

A R44 was used to survey the Teedraanjik watershed on September 17, 2017. The R44 helicopter was also used to provide access to Kevinjik Creek within the Teedraanjik watershed for baseline sample collections on September 17 and 18, 2017. The R44 was also used to survey the Coleen River on September 19, 2017, and to collect baseline samples on September 20, 2017. R44 operations were based in Chalkyitsik, with daily departures between September 17 and 20, 2017.

Results and Discussion

In the Teedraanjik watershed, ASL and tissue samples were collected from 47 spawning Chinook salmon and 135 spawning fall chum salmon (Figures 2 & 3). The 47 Chinook salmon tissue samples collected from the Teedraanjik in 2018 bring the total number of Chinook salmon tissue samples for the Teedraanjik up to 98 (Table 1) (Personal Communication Sara Gilk-Baumer, October 2016). A total of 12 temperature loggers were deployed in the Teedraanjik watershed; 7 were deployed on the main stem of the Teedraanjik in Chinook salmon spawning locations, and 5 were deployed in the Kevinjik Creek in fall chum salmon spawning locations (Figure 4).

In the Coleen River, ASL and tissue samples were collected from 42 spawning Chinook salmon and 101 spawning fall chum salmon (Figures 5 & 6). The 42 Chinook salmon tissue samples collected from the Coleen River in 2017 bring the total number of tissue samples for Chinook salmon in the Coleen River up to 103 (Table 2) (Personal Communication, Sara Gilk-Baumer October 2016). A total of 9 temperature loggers were deployed in the Coleen River with 3 being placed in Chinook salmon spawning locations and 6 placed in fall chum salmon spawning locations (Figure 7).

All ASL data was sent to the ADF&G division of commercial fisheries for aging analyses and will be reported on after completion of the 2018 sampling season. Tissue samples were stored in bulk vials filled with anhydrous ethyl alcohol. The bulk vials were shipped to the ADF&G GCL. Samples will be shared with the USFWS and with the DFO-Canada. The samples will be genotyped by the ADF&G GCL, and will be used to improve the existing genetic baseline. Stream temperature data loggers will be collected during the 2018 field season. Nominations for the ADF&G AWC will be submitted after the completion of the 2018 field season.

Acknowledgements

We would like to express our gratitude to the Pacific States Marine Fisheries Commission for administering the funding support for this project through the 2012 Yukon River Chinook Salmon Federal Fishery Disaster Fund. We would like to acknowledge the Chalkyitsik Village Council and the Circle Tribal Council for their support for this project. We would also like to thank Quintin Slade of InFlight Helicopters and Brian Lepping for their sampling and logistical support. We would like to thank the Canada Border Services Agency for authorizing this project to perform aerial surveys in the Yukon Territory. Finally, we also would like to thank the ADF&G division of commercial fisheries for future ASL analysis and the ADF&G GCL staff for their time in organizing sampling materials, intaking, processing, and future analysis of genetic samples.

References

Personal Communication with Sara Gilk-Baumer, October 2016.

Yukon Ecoregions Working Group, 2004. Yukon Coastal Plain. *In: Ecoregions of the Yukon Territory: Biophysical properties of Yukon landscapes*, C.A.S. Smith, J.C. Meikle and C.F. Roots (eds.), Agriculture and Agri-Food Canada, PARC Technical Bulletin No. 04-01, Summerland, British Columbia, p. 63-72.

North Ogilvie Mountains, Taiga Cordillera Ecozone, Ecoregion 168, p. 123 – 130.





Figure 1 — Map illustrating the general location of the Teedraanjik and Coleen Rivers. The Teedraanjik is denoted by the black rectangle and is a transboundary river originating in Yukon Territory and flowing into Alaska. The Coleen River is denoted by the red oval and drains a portion of the southern Brooks Range in Alaska before emptying into the Porcupine River.

Teedraanjik Chinook Salmon Tissue Sample Collections



Esri, HERE, DeLorme, MapmyIndia, © OpenStreetMap contributors, and the GIS user community

Year

-  2015
-  2016
-  2017



Tanana
Chiefs
Conference


Prepared by Brian Mokenna
Tanana Chiefs Conference
Wildlife & Parks Program
Palmdale, Alaska
December 2017

10 5 0 Miles

Coordinate System: NGS 1984 Web Mercator Auxiliary Sphere
Projection: Web Mercator Auxiliary Sphere

Figure 2 — Map illustrating the location of Chinook salmon samples collected in the Teedraanjik, 2015 - 2017.

Teedraanjik Fall Chum Salmon Sample Location



Esri, HERE, DeLorme, MapmyIndia, © OpenStreetMap contributors, and the GIS user community, Source: Esri, DigitalGlobe, GeoEye, IGN, USDA, USGS, AEC, Getmapping, Aergrid, IGN, IGP, swisstopo, and the GIS User Community

Legend

 **Nee'tinil Fall Chum Salmon**



Prepared by Brian McKenna
Tanana Chiefs Conference
Wildlife & Parks Program
Fairbanks, Alaska
December 2017



Coordinate System: NAD83 1200 Feet Universal Auxiliary Sphere
Responsible: Unknown Auxiliary Sphere

Figure 3 — Map illustrating the location of fall chum salmon samples collected in the Teedraanjik, 2017.





Figure 4 — Map illustrating the location of stream temperature data logger deployments in the Teedraanjik, 2017.

Coleen River Chinook Salmon Tissue Sample Collections



Legend

- Chinook Observations 2017
- Chinook Tissue Sample Locations 2017
- Coleen River Watershed

Prepared by Brian McKenna
 Tanana Chiefs Conference
 Wildlife & Parks Program
 Fairbanks, Alaska
 December, 2017




Tanana
 Chiefs
 Conference



Figure 5 — Map illustrating the location of spawning Chinook salmon ASL and tissue samples collections in the Coleen River, 2017.

Coleen River Fall Chum Salmon



Legend

- Fall Chum Observations 2017
- Fall Chum Tissue Sample Location 2017
- Coleen River Watershed

Prepared by Brian McKenna
Tanana Chiefs Conference
Wildlife & Parks Program
Fairbanks, Alaska
December, 2017



Tanana
Chiefs
Conference




Figure 6 — Map illustrating the location of spawning fall chum salmon ASL and tissue samples collections in the Coleen River, 2017.




Figure 7 — Map illustrating the location of stream temperature data logger deployments in the Coleen River, 2017.

Table 1 —Chinook salmon tissue samples collected from the Teedraanjik between 2015 and 2017.

Tributary (main)	Branch Tributary	Number of Samples			All years
		2015	2016	2017	
Porcupine River	Teedraanjik (Salmon Fork)	50	1	47	98
Totals		50	1	47	98

Table 2 —Chinook salmon tissue samples collected from the Coleen River between 2011 and 2017.

Tributary (main)	Branch Tributary	Number of Samples			All years
		2011	2013	2017	
Porcupine River	Coleen River	24	37	42	103
Totals		24	37	42	103

Collections in 2011 and 2013 were not conducted by the Tanana Chiefs Conference.

Appendix 1 — Non-lethal Bulk Sampling Finfish Tissues for DNA Analysis (ADF&G Gene Conservation Laboratory, Anchorage, Alaska).

Non-lethal Bulk Sampling Finfish Tissues for DNA Analysis ADF&G Gene Conservation Lab, Anchorage



I. General Information

We use axillary process samples from individual fish to determine the genetic characteristics and profile of a particular run or stock of fish. This is a non-lethal method of collecting tissue samples from adult fish for genetic analysis. The most important thing to remember in collecting samples is that **only quality tissue samples give quality results**. If sampling from carcasses: tissues need to be as "fresh" and as cold as possible and recently moribund, do not sample from fungal fins.

II. Sampling Method

Preservative used: Isopropanol/Methanol/Ethanol (EtOH) preserves tissues for later DNA extraction. Avoid extended contact with skin.

Sampling instructions are written for (N=100 fish/125ml) bulk bottle. Steps for collecting axillary process tissues:



SILLY: _____
Location: _____
Sample Date(s): ____/____/____
Sampler's name: _____
Total # fish sampled: _____
Latitude: _____
Longitude: _____
Species: _____
Comments: _____
ADF&G: Preserved in EtOH

Supplies included in sampling kit:

1. Clipper- used to cut a portion of **one** axillary process per fish
2. Sample target: 100 axillary clips/125ml bulk bottle
3. Labels on bulk sample bottles. Location, Sample date, Sampler, Total # fish sampled and comments (if any)
4. 1:125ml wide mouth bottle(s) for EtOH "refresh" step
5. Sampling instructions

- Wipe dry the axillary process "spine" prior to sampling to avoid getting excess water or fish slime into the 125ml bottle (see diagram).
- Clip off the axillary "spine" using dog nail clippers or scissors to get roughly a 1/2 - 1" **inch maximum** piece and/or about the size of a small fingernail.
- Place each tissue piece into bulk bottle (**place only one piece of axillary from each fish**).
- Repeat: **up to 100 fish /125ml bulk bottle** (into same bottle). If you don't reach this number of fish per location, that's ok. Maximum storage capacity 125ml bulk for proper preservation of axillary tissue is (N=100).
- Record on **each label**: Location, sampling date (mm/dd/yyyy), sampler's name(s), total number of fish sampled, latitude/longitude, and field notes (if any). Use pencil. This insures correct data with each collection bottle.
- If collection occurs over 4~5 day period, "refresh" EtOH at end of the collection.
- After the collection is complete and 24 hours have passed, "refresh" the axillary tissues as follows: carefully pour off 3/4 EtOH and then pour fresh EtOH into sample bottle containing axillary clips. Cap and invert bottle twice mixing EtOH and tissue.
- Freezing not required, store sample bottle in upright cool location for good tissue quality.

Return to ADF&G Anchorage lab:

ADF&G - Genetics
333 Raspberry Road
Anchorage, Alaska 99518

Lab staff: 907-267-2247
Judy Berger 907-267-2175
Freight code _____



Coleen River

Nomination to update ADF&G AWC
Fall chum salmon spawning update (2017)

TCC, CHs (67.79942, -142.391490)

AWC, Ks, CHp (67.7488, -142.4451)

Legend


-  Current AWC
-  Update to AWC

Google Earth

Image Landsat / Copernicus
© 2018 Google

50 km





-142,26704, 67,84563
Chinook salmon spawning

-142,39149, 67,79942
chum salmon spawning

extend 334-45-11000-2001-3271
w/Chinook salmon spawning,
use arc2019 for hydro

extend 334-45-11000-2001-3271
w/chum salmon spawning, replace CHP w/CHS,
use arc2019 for hydro