

Fishery Management Report No. 22-01

Annual Management Report for Shellfish Fisheries in the Bering Sea/Aleutian Islands Management Area, 2020/21

by

Ethan Nichols

Miranda Westphal

and

Janis Shaishnikoff

January 2022

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

| Weights and measures (metric) | | General | | Mathematics, statistics | |
|---|--------------------|--|-------------------------------|--|-------------------------|
| centimeter | cm | Alaska Administrative Code | | all standard mathematical signs, symbols and abbreviations | |
| deciliter | dL | | AAC | | |
| gram | g | all commonly accepted abbreviations | e.g., Mr., Mrs., AM, PM, etc. | alternate hypothesis | H _A |
| hectare | ha | | | base of natural logarithm | <i>e</i> |
| kilogram | kg | | | catch per unit effort | CPUE |
| kilometer | km | all commonly accepted professional titles | e.g., Dr., Ph.D., R.N., etc. | coefficient of variation | CV |
| liter | L | | | common test statistics | (F, t, χ^2 , etc.) |
| meter | m | at | @ | confidence interval | CI |
| milliliter | mL | compass directions: | | correlation coefficient (multiple) | R |
| millimeter | mm | east | E | correlation coefficient (simple) | r |
| Weights and measures (English) | | north | N | covariance | cov |
| cubic feet per second | ft ³ /s | south | S | degree (angular) | ° |
| foot | ft | west | W | degrees of freedom | df |
| gallon | gal | copyright | © | expected value | <i>E</i> |
| inch | in | corporate suffixes: | | greater than | > |
| mile | mi | Company | Co. | greater than or equal to | ≥ |
| nautical mile | nmi | Corporation | Corp. | harvest per unit effort | HPUE |
| ounce | oz | Incorporated | Inc. | less than | < |
| pound | lb | Limited | Ltd. | less than or equal to | ≤ |
| quart | qt | District of Columbia | D.C. | logarithm (natural) | ln |
| yard | yd | et alii (and others) | et al. | logarithm (base 10) | log |
| | | et cetera (and so forth) | etc. | logarithm (specify base) | log ₂ , etc. |
| Time and temperature | | exempli gratia | | minute (angular) | ' |
| day | d | (for example) | e.g. | not significant | NS |
| degrees Celsius | °C | Federal Information Code | FIC | null hypothesis | H ₀ |
| degrees Fahrenheit | °F | id est (that is) | i.e. | percent | % |
| degrees kelvin | K | latitude or longitude | lat or long | probability | P |
| hour | h | monetary symbols | | probability of a type I error | |
| minute | min | (U.S.) | \$, ¢ | (rejection of the null hypothesis when true) | α |
| second | s | months (tables and figures): first three letters | Jan,...,Dec | probability of a type II error | |
| Physics and chemistry | | registered trademark | ® | (acceptance of the null hypothesis when false) | β |
| all atomic symbols | | trademark | ™ | second (angular) | " |
| alternating current | AC | United States | | standard deviation | SD |
| ampere | A | (adjective) | U.S. | standard error | SE |
| calorie | cal | United States of America (noun) | USA | variance | |
| direct current | DC | U.S.C. | United States Code | population sample | Var var |
| hertz | Hz | | | | |
| horsepower | hp | | | | |
| hydrogen ion activity (negative log of) | pH | | | | |
| parts per million | ppm | U.S. state | use two-letter abbreviations | | |
| parts per thousand | ppt, ‰ | | (e.g., AK, WA) | | |
| volts | V | | | | |
| watts | W | | | | |

FISHERY MANAGEMENT REPORT NO. 22-01

**ANNUAL MANAGEMENT REPORT FOR SHELLFISH FISHERIES OF
THE BERING SEA/ALEUTIAN ISLANDS MANAGEMENT AREA, 2020/21**

by

Ethan Nichols

Alaska Department of Fish and Game, Division of Commercial Fisheries, Dutch Harbor

and

Miranda Westphal

Alaska Department of Fish and Game, Division of Commercial Fisheries, Dutch Harbor

and

Janis Shaishnikoff

Alaska Department of Fish and Game, Division of Commercial Fisheries, Dutch Harbor

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518-1565

January 2022

This report was compiled from data collection funded through Alaska Fisheries Information Network (AKFIN), Task II: Groundfish/Shellfish Fisheries Data collection and Analysis in Alaska, NOAA Cooperative Agreement NA18NMF4370300 and through the Bering Sea and Aleutian Islands Crab Rationalization program, NOAA Cooperative Agreement NA13NMF4370148.

The Fishery Management Reports series was established in 1989 by the Division of Sport Fish for the publication of an overview of management activities and goals in a specific geographic area and became a joint divisional series in 2004 with the Division of Commercial Fisheries. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.adfg.alaska.gov/sf/publications/>. This publication has undergone regional peer review.

Product names used in this publication are included for completeness and do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

*Ethan Nichols,
Alaska Department of Fish and Game, Division of Commercial Fisheries
PO Box 920587, Dutch Harbor, Alaska 99692 USA*

*Miranda Westphal,
Alaska Department of Fish and Game, Division of Commercial Fisheries
PO Box 920587, Dutch Harbor, Alaska 99692 USA*

and

*Janis Shaishnikoff
Alaska Department of Fish and Game, Division of Commercial Fisheries
PO Box 920587, Dutch Harbor, Alaska 99692 USA*

This document should be cited as follows:

E. Nichols, J. Shaishnikoff, and M. Westphal. 2022. Annual management report for shellfish fisheries of the Bering Sea/Aleutian Islands Management Area, 2020/21. Alaska Department of Fish and Game, Fishery Management Report No. 22-01, Anchorage.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375

TABLE OF CONTENTS

| | Page |
|---|------|
| LIST OF TABLES..... | iii |
| LIST OF FIGURES..... | iv |
| ABSTRACT | 1 |
| INTRODUCTION..... | 1 |
| SECTION I: BERING SEA SHELLFISH FISHERIES..... | 3 |
| BRISTOL BAY KING CRAB REGISTRATION AREA T | 3 |
| Description of Area | 3 |
| Bristol Bay Red King Crab..... | 3 |
| 2020/21 Fishery | 3 |
| Port Sampling | 3 |
| Stock Status | 4 |
| BERING SEA KING CRAB REGISTRATION AREA Q | 4 |
| Description of Area | 4 |
| Saint Matthew Island Section Blue King Crab | 5 |
| 2020/21 Fishery | 5 |
| Stock Status | 5 |
| Northern District Golden King Crab | 5 |
| 2020 Fishery | 5 |
| Fishery Management and Stock Status | 6 |
| Pribilof District Golden King Crab..... | 6 |
| 2020 Fishery | 6 |
| Port Sampling | 6 |
| Fishery Management and Stock Status | 6 |
| Bering Sea Scarlet King Crab | 7 |
| 2020 Fishery | 7 |
| Fishery Management and Stock Status | 7 |
| BERING SEA TANNER CRAB MANAGEMENT DISTRICT | 7 |
| Description of Area | 7 |
| Bering Sea Tanner Crab | 8 |
| 2020/21 Fishery East of 166°W Longitude | 8 |
| Stock Status | 8 |
| 2020/21 Fishery West of 166°W Longitude | 8 |
| Port Sampling | 9 |
| Stock Status | 9 |
| Bering Sea Snow Crab..... | 9 |
| 2020/21 Fishery | 9 |
| Port Sampling | 10 |
| Stock Status | 10 |
| Bering Sea Grooved Tanner Crab..... | 11 |
| 2020 Fishery | 11 |
| Fishery Management and Stock Status | 11 |

TABLE OF CONTENTS (Continued)

| | Page |
|--|------|
| Bering Sea Triangle Tanner Crab | 11 |
| 2020 Fishery | 11 |
| Fishery Management and Stock Status | 11 |
| NORTH PENINSULA DISTRICT DUNGENESS CRAB | 12 |
| Description of District | 12 |
| Dungeness Crab | 12 |
| 2020 Fishery | 12 |
| Fishery Management and Stock Status | 12 |
| BRISTOL BAY - BERING SEA WEATHERVANE SCALLOP REGISTRATION AREA Q | 12 |
| Description of Area | 12 |
| Bristol Bay - Bering Sea Weathervane Scallop | 12 |
| 2020/21 Fishery | 12 |
| Fishery Management and Stock Status | 12 |
| BERING SEA MISCELLANEOUS SHELLFISH SPECIES | 13 |
| Description of District | 13 |
| 2020 Fisheries | 13 |
| Octopus | 13 |
| Fishery Management and Stock Status | 13 |
| REFERENCES CITED | 14 |
| TABLES AND FIGURES | 15 |
| SECTION II: ALEUTIAN ISLANDS SHELLFISH FISHERIES | 73 |
| ALEUTIAN ISLANDS KING CRAB MANAGEMENT AREA | 73 |
| Description of Area | 73 |
| Aleutian Islands Golden King Crab | 73 |
| 2020/21 Fishery East of 174°W Longitude | 73 |
| Port Sampling | 73 |
| 2020/21 Fishery West of 174°W Longitude | 74 |
| Port Sampling | 74 |
| Fishery Management and Stock Status | 74 |
| EASTERN ALEUTIAN TANNER CRAB DISTRICT | 75 |
| Description of District | 75 |
| Tanner Crab | 75 |
| 2020 Commercial Fishery | 75 |
| Fishery Management and Stock Status | 75 |
| Subsistence | 75 |
| 2020 King and Tanner Fishery | 75 |
| ALEUTIAN DISTRICT DUNGENESS CRAB | 76 |
| Description of District | 76 |
| 2020 Fishery | 76 |
| Fishery Management and Stock Status | 76 |

TABLE OF CONTENTS (Continued)

| | Page |
|---|------|
| DUTCH HARBOR WEATHERVANE SCALLOP REGISTRATION AREA O | 76 |
| Description of Area | 76 |
| Dutch Harbor Weathervane Scallop | 76 |
| 2020/21 Fishery | 76 |
| Fishery Management and Stock Status | 76 |
| ALEUTIAN ISLANDS MISCELLANEOUS SHELLFISH | 77 |
| Description of District | 77 |
| 2020 Fisheries..... | 77 |
| Octopus..... | 77 |
| Fishery Management and Stock Status | 77 |
| REFERENCES CITED | 78 |
| TABLES AND FIGURES..... | 79 |
| SECTION III: BERING SEA/ALEUTIAN ISLANDS COMMUNITY DEVELOPMENT QUOTA AND ADAK COMMUNITY ALLOCATION SHELLFISH FISHERIES | 106 |
| Description of Area | 106 |
| Program Background..... | 106 |
| Current Fishery..... | 107 |
| TABLES..... | 109 |

LIST OF TABLES

| Table | Page |
|---|------|
| 1. Bristol Bay red king crab commercial fishery harvest data, 1966–2020/21..... | 16 |
| 2. Bristol Bay red king crab commercial fishery value and season dates, 1980–2020/21..... | 19 |
| 3. Bristol Bay red king crab commercial fishery harvest and effort by week, 2020/21..... | 21 |
| 4. Bristol Bay red king crab commercial fishery harvest and effort by statistical area, 2020/21..... | 22 |
| 5. Bristol Bay red king crab cost-recovery harvest data and charter length, 1990–2020..... | 23 |
| 6. Saint Matthew Island Section blue king commercial fishery harvest data, 1977–2020/21..... | 25 |
| 7. Saint Matthew Island Section blue king crab commercial fishery value and season dates, 1977– 2020/21..... | 27 |
| 8. Pribilof District golden king crab commercial fishery harvest data, 1981/82–2020..... | 28 |
| 9. Pribilof District golden king crab commercial fishery value and season dates, 1991–2020..... | 30 |
| 10. Pribilof District golden king crab commercial fishery harvest and effort by statistical area, 2020..... | 31 |
| 11. Bering Sea scarlet king crab commercial fishery harvest data, 1995–2020..... | 32 |
| 12. Bering Sea District Tanner crab commercial fishery harvest data, 1968–2020/21..... | 33 |
| 13. Bering Sea District Tanner crab commercial fishery value and season dates, 1974–2020/21..... | 39 |
| 14. Bering Sea District Tanner crab commercial fishery harvest and effort by week, west of 166° W long, 2020/21..... | 42 |
| 15. Bering Sea District Tanner crab commercial fishery harvest and effort by statistical area, west of 166° W long, 2020/21..... | 44 |
| 16. Bering Sea District snow crab commercial fishery harvest data, 1977/78–2020/21..... | 46 |
| 17. Bering Sea District snow crab commercial fishery value and season dates, 1980–2020/21..... | 49 |
| 18. Bering Sea District snow crab commercial fishery harvest and effort by week, 2020/21..... | 53 |
| 19. Bering Sea District snow crab commercial fishery harvest and effort by statistical area, 2020/21..... | 55 |
| 20. Bering Sea District grooved Tanner crab commercial fishery harvest data, 1992–2020..... | 57 |
| 21. Bering Sea District triangle Tanner crab commercial fishery harvest data, 1995–2020..... | 58 |
| 22. North Peninsula District commercial Dungeness crab fishery data, 1992–2020..... | 59 |

LIST OF TABLES (Continued)

| Table | Page |
|---|------|
| 23. Bristol Bay–Bering Sea weathervane scallop commercial fishery harvest data, 1987–2020/21..... | 61 |
| 24. Bering Sea commercial octopus incidental harvest in groundfish fisheries, 1995–2020. | 62 |
| 25. Aleutian Islands golden king crab commercial fishery harvest data, 1981/82–2020/21. | 80 |
| 26. Aleutian Islands golden king crab commercial fishery value and season dates, 1981/82–2020/21. | 86 |
| 27. Aleutian Islands golden king crab commercial fishery harvest by statistical area east of 174° W long, 2020/21..... | 89 |
| 28. Eastern Aleutian Islands golden king crab cost-recovery harvest data and charter length, 2013–2020. | 90 |
| 29. Eastern Aleutian District Tanner crab commercial fishery harvest data, 1973/74–2020. | 91 |
| 30. Eastern Aleutian District Tanner crab commercial fishery value and season dates, 1973/74–2020..... | 94 |
| 31. Subsistence king and Tanner crab harvest from the Eastern Aleutian Islands, west of Scotch Cap Light and east of 168°W long, 1999–2020. | 96 |
| 32. Aleutian District Dungeness crab commercial fishery data, 1974–2020..... | 97 |
| 33. Dutch Harbor weathervane scallop commercial fishery harvest data, 1982–2020/21..... | 99 |
| 34. Aleutian Islands commercial octopus incidental harvest in groundfish fisheries, 1996–2020. | 101 |
| 35. Community Development Quota and Adak Community Allocation program percent allocation by fishery to each group. | 110 |

LIST OF FIGURES

| Figure | Page |
|--|------|
| 1. Bristol Bay king crab commercial fishery Registration Area T. | 63 |
| 2. Bristol Bay red king crab commercial fishery harvest, catch per unit effort, and number of vessels, 1966–2020/21..... | 64 |
| 3. Bering Sea king crab commercial fishery Registration Area Q..... | 65 |
| 4. Saint Matthew Island section blue king crab commercial fishery harvest, catch per unit effort, and number of vessels, 1977–2020/21. | 66 |
| 5. Bering Sea District Tanner crab commercial fishery Registration Area J including subdistricts and sections..... | 67 |
| 6. Bering Sea Tanner crab commercial fishery harvest, catch per unit effort, and number of vessels, 1974/75–2020/21..... | 68 |
| 7. Bering Sea Snow crab commercial fishery harvest, catch per unit effort, and number of vessels, 1977/78–2020/21..... | 69 |
| 8. North Peninsula and Aleutian Islands Dungeness crab Districts of Registration Area J..... | 70 |
| 9. Bering Sea and Aleutian Islands weathervane scallop Registration Areas..... | 71 |
| 10. Bering Sea and Aleutian Islands Tanner crab and miscellaneous shellfish Districts of Registration Area J..... | 72 |
| 11. Aleutian Islands king crab commercial fishery Registration Area O. | 102 |
| 12. Eastern Aleutian Islands golden king crab commercial fishery harvest, catch per unit effort, and number of vessels, 1981/82–2020/21. | 103 |
| 13. Western Aleutian Islands golden king crab commercial fishery harvest, catch per unit effort, and number of vessels, 1981/82–2020/21. | 104 |
| 14. Eastern Aleutian District Tanner crab sections of Registration Area J. | 105 |

ABSTRACT

The Alaska Department of Fish and Game (ADF&G) manages commercial and subsistence shellfish fisheries in the territorial waters and Exclusive Economic Zone (EEZ) of the Bering Sea and Aleutian Islands in the northern Pacific Ocean. This report presents details on commercial and subsistence invertebrate and shellfish fisheries harvest, participation, and value in the Bering Sea and Aleutian Islands (BSAI) areas, excluding king crab fisheries north of Cape Romanzof. In 2020/21, red king crab, golden king crab, snow crab, Tanner crab, and giant Pacific octopus were taken in BSAI fisheries. Current fishery management practices, a summary of the most recent commercial fishery, and general stock status information are presented.

Keywords: Red king crab *Paralithodes camtschaticus*, golden king crab *Lithodes aequispinus*, scarlet king crab *Lithodes couesi*, snow crab *Chionoecetes opilio*, Tanner crab *C. bairdi*, Dungeness crab *Metacarcinus magister*, giant Pacific octopus *Enteroctopus dofleini*, blue king crab *P. platypus*, grooved Tanner crab *C. tanneri*, triangle Tanner crab *C. angulatus*, Community Development Quota, Crab Rationalization, Individual Fishing Quota, subsistence, guideline harvest level, Bering Sea, Aleutian Islands, North Peninsula, bycatch, confidential interviews, retained catch, species composition sample, size frequencies

INTRODUCTION

The Alaska Department of Fish and Game (ADF&G) manages commercial and subsistence invertebrate and shellfish fisheries occurring in the state waters of Alaska (0–3 nmi) and waters of the Exclusive Economic Zone (EEZ; 3–200 nmi) of the Bering Sea and Aleutian Islands. The Bering Sea (including Bristol Bay) is made up of waters north of Cape Sarichef (54°36'N lat) and west of the Maritime Boundary Agreement Line of 1990, excluding the Norton Sound Section. The Aleutian Islands area includes waters west of the longitude of Scotch Cap Light (164°44.72'W long), east of the Maritime Boundary Agreement Line of 1990, south of Cape Sarichef (58°36'N lat), to 171°W long westward, and south of a line from the latitude 55°30'N. Crab in the Bering Sea north of Cape Romanzof (61°49'N lat), including Norton Sound, are managed by ADF&G's Nome office and are not included in this report. Waters of the Bering Sea and Aleutian Islands (BSAI) historically support the largest and most valuable commercial crab fisheries in Alaska.

The BSAI is divided into registration areas for king crab management and districts for Tanner crab, Dungeness crab, and miscellaneous shellfish management. BSAI king and Tanner crab fisheries in the EEZ are managed under a federal fisheries management plan (FMP) that establishes a cooperative management structure deferring king and Tanner crab management to the State of Alaska with federal oversight. Other crab and miscellaneous shellfish fisheries that occur in territorial waters are managed solely under state jurisdiction. Since 2005, most EEZ BSAI crab fisheries are managed under the federal crab rationalization program, which resulted in consolidation of harvesting and processing sectors and substantially changed historical fishing practices.

Species commercially harvested during the 2020/21 season in the BSAI include red king crab *Paralithodes camtschaticus*, golden king crab *Lithodes aequispinus*, Tanner crab *Chionoecetes bairdi*, snow crab *Chionoecetes opilio*, and giant Pacific octopus *Enteroctopus dofleini*. Historically, waters of the BSAI have supported commercial harvests of blue king crab *P. platypus*, scarlet king crab *L. couesi*, triangle Tanner crab *C. angulatus*, Dungeness crab *Metacarcinus magister*, green sea urchins *Strongylocentrotus droebachiensis*, pandalid shrimp *Pandalidae* spp., hair crab *Erimacrus isenbeckii*, weathervane scallop *Patinopecten caurinus*, and several species of sea snails. However, fisheries for these species are currently either closed due to low abundance

or not currently commercially prosecuted. For additional background on current BSAI shellfish fisheries and information on historical fisheries not covered in this report, refer to *Annual management report for shellfish fisheries of the Bering Sea/Aleutian Islands Management Area, 2015/16* (Leon et al. 2017).

In 2020/21, 69 catcher vessels, 2 catcher-processors, and 8 shore-based processors were involved in harvesting and processing shellfish in the BSAI. Shellfish landings totaled approximately 55.9 million pounds.

SECTION I: BERING SEA SHELLFISH FISHERIES

BRISTOL BAY KING CRAB REGISTRATION AREA T

DESCRIPTION OF AREA

Bristol Bay king crab Registration Area T includes all waters north of Cape Sarichef (54°36'N lat), south of Cape Newenham (58°39'N lat), and east of 168°W long (Figure 1).

BRISTOL BAY RED KING CRAB

2020/21 Fishery

The 2020/21 Bristol Bay red king crab (BBR) fishery opened October 15 with a combined Individual Fishing Quota (IFQ) and Community Development Quota (CDQ) Total Allowable Catch (TAC) of 2,648,000 pounds (Table 1). TAC was allocated by National Marine Fisheries Service (NMFS) as 90% IFQ (2,383,200 pounds) and 10% CDQ (264,800 pounds) with all 6 of the CDQ groups participating in the harvest of the CDQ allocation. Forty-seven vessels participated in the fishery and harvested 2,646,874 pounds, of which 0.2% was deadloss (Table 1). Despite the regulatory fishing season running through January 15, 99% of the harvest occurred by mid-November, with the last delivery on January 15 (Tables 2 and 3). On average, vessels were active in the fishery for 8 days. Harvesters were paid an initial average exvessel price of \$9.11 per pound, the highest BBR exvessel price since the inception of the fishery. Total exvessel fishery value was estimated to be \$24,086,513 (Table 2).

Total effort for the 2020/21 fishery was 20,244 pot lifts. Catch per unit effort (CPUE) was 21 legal crab per pot, slightly below the post-rationalization (2005/06–2019/20) average CPUE of 26 (Table 1, Figure 2). Harvest was distributed across 11 ADF&G statistical reporting areas with the highest concentration of harvest (44%) occurring in ADF&G statistical area 625630, northeast of Amak Island (Table 4).

BBR cost recovery was conducted by ADF&G in 2020 with 124,289 pounds (21,780 legal male crab) harvested. The 18-day charter occurred from October 7 to October 24 (Table 5).

Port Sampling

During the 2020/21 BBR fishery, ADF&G personnel and onboard observers sampled red king crab from vessels at shore-based processors in Akutan, Dutch Harbor, King Cove, Kodiak, and Saint Paul Island. Biological data collected on landed red king crab consisted of carapace length, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 89 of the 95 total landings (IFQ and CDQ) during the 2020/21 BBR fishery. Average weight of sampled crab was 6.1 pounds while average carapace length was 150 mm (Table 1).

Stock Status

The 2020 NMFS trawl survey of Bristol Bay was cancelled due to COVID-19.

Based on 2019 NMFS trawl survey data (Zacher et al. 2020), estimated immature male biomass of 6.2 million pounds was below the 20-year average of 20.9 million pounds, estimated mature male biomass of 27.4 million pounds was below the 20-year average of 78.7 million pounds, estimated legal male biomass of 19.8 million pounds was below the 20-year average of 61.8 million pounds, and estimated mature female biomass of 28.9 million pounds was below the 20-year average of 71.5 million pounds.

Because there was no survey data to update the assessment model, updated data in the model included total catch and length frequencies from the 2019/20 directed fishery, bycatch in the Tanner crab fishery, and discarded catch and length frequencies from the 2019 and 2020 groundfish fisheries. The 2019 estimate of recruitment (3.8 million crab) was the lowest since 1994. Due to the cancelled 2020 survey, recruitment in the terminal year was fixed to the mean recruitment in the most recent 7 years, which gave an estimated recruitment of 18.9 million crab. The large recruitment estimate with no new survey inputs was considered highly uncertain and disregarded in the recommendation to the North Pacific Fishery Management Council (NPFMC). The assessment showed this stock has been in decline since 2009 and recruitment has been below the long-term average for the past 15 years (NPFMC 2020a, Scientific and Statistical Committee 2020).

Data from the NMFS trawl survey is incorporated into a length-based analysis (LBA) model used to establish the fishery TAC. Outputs from the 2020 LBA model showed continued declines for effective spawning biomass (ESB), legal males, females, and recruitment (in contrast to the stock assessment), although female abundance and ESB estimates were both above the threshold to allow for a fishery to occur. The regulatory harvest strategy uses a stair-step exploitation approach based on abundance. The LBA model estimated effective spawning biomass at 25.12 million pounds; therefore, an exploitation rate of 10% was applied to mature male red king crab for a 2020/21 TAC of 2.648 million pounds. This TAC was a 30% decrease from the previous year and the lowest TAC since rationalization. The harvest strategy is found in 5 AAC 34.816 *Bristol Bay red king crab harvest strategy*.

Further information on BBR stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

BERING SEA KING CRAB REGISTRATION AREA Q

DESCRIPTION OF AREA

The Bering Sea king crab Registration Area Q southern boundary is a line from 54°36'N lat, 168°W long, to 54°36'N lat, 171°W long, to 55°30'N lat, 171°W long, to 55°30'N lat, 173°30'E long. The northern boundary is the latitude of Point Hope (68°21'N lat). The eastern boundary is a line from 54°36'N lat, 168°W long, to 58°39'N lat, 168°W long, to Cape Newenham (58°39'N lat). The western boundary is the United States–Russia Maritime Boundary Line of 1990 (Figure 3). Area Q is divided into two districts: the Pribilof District, which includes waters south of Cape Newenham; and the Northern District, which includes all waters north of Cape Newenham. The Northern District is subdivided into three sections. The Saint Matthew Island

Section includes waters north of Cape Newenham and south of Cape Romanzof (61°49'N lat). The Norton Sound Section includes all waters north of Cape Romanzof and south of 66°N lat. The Kotzebue Sound Section encompasses all remaining waters of the district. The Norton Sound Section and Kotzebue Sound Section are not addressed in this report.

SAINT MATTHEW ISLAND SECTION BLUE KING CRAB

2020/21 Fishery

The 2020/21 Saint Matthew Island Section blue king crab fishery was closed due to the 2020 model-based estimate of mature male abundance being below the regulatory threshold for opening a fishery (Tables 6 and 7, Figure 4).

Stock Status

The 2020 NMFS trawl survey and ADF&G pot survey for Saint Matthew Island blue king crab were both cancelled due to COVID-19.

Based on 2019 NMFS trawl survey data (Zacher et al. 2020), estimated immature male biomass of 1.7 million pounds was below the 20-year average of 2.2 million pounds, estimated mature male biomass of 6.3 million pounds was below the 20-year average of 7.7 million pounds, estimated legal male biomass of 5.1 million pounds was at the 20-year average of 5.1 million pounds, and estimated mature female biomass of 0.86 million pounds was above the 20-year average of 0.23 million pounds. Stock assessment authors attribute increases in trawl survey biomass estimates to movement of crab out of nearshore areas and into the trawl survey zone, as opposed to actual increases in abundance. In 2018, NMFS declared the Saint Matthew Island blue king crab stock overfished and began developing a stock rebuilding plan. The stock continues to decline and is still considered overfished.

ADF&G conducted triennial pot surveys in the Saint Matthew Island Section from 1995 to 2013, with a focus on the nearshore waters with bottom topography unsuitable to trawl surveys. From 2013 to 2018, in response to a request for more detailed information from the stock assessment authors, the survey was conducted on an annual basis.

Due to the cancellation of both surveys for this stock, the only new information added to the assessment was bycatch in the groundfish fisheries. Projected estimates of mature male abundance were derived from the accepted stock assessment model and were below the threshold for opening a fishery (NPFMC 2020a, Scientific and Statistical Committee 2020).

Further information on Saint Matthew Island blue king crab stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

NORTHERN DISTRICT GOLDEN KING CRAB

2020 Fishery

One vessel harvested golden king crab in the Saint Matthew Island Section of the Northern District during 2020 under the authority of an ADF&G commissioner's permit. The permit was issued with a guideline harvest range of 15,000 to 30,000 pounds and was designed to allow for exploratory fishing and data gathering. Harvest information is confidential.

Fishery Management and Stock Status

In 2007, NMFS amended the FMP and removed Bering Sea golden king crab, which transferred sole jurisdiction for the fishery to the state (NPFMC 2007). Onboard observers have been required on most vessels that targeted deepwater crab species since 1994 and have collected information detailing the size and sex composition of the retained and nonretained golden king crab and bycatch species. This information is useful when developing management measures for deepwater crab stocks; however, there is currently no harvest strategy for this stock.

Stock biomass of golden king crab in the Northern District has been estimated using area-swept methods applied to NMFS upper continental slope trawl survey data (Hoff 2016). Although biomass estimates of golden king crab from the slope survey are available, they are highly uncertain and not currently used in fishery management. The most recent NMFS Bering Sea slope survey was conducted in 2016. Results of the 2016 NMFS Bering Sea slope survey show an estimated biomass of 0.81 million pounds (Hoff 2016).

PRIBILOF DISTRICT GOLDEN KING CRAB

2020 Fishery

The 2020 Pribilof District golden king crab (PIGKC) fishery had a guideline harvest level (GHL) of 130,000 pounds and was open January 1, 2020, through December 31, 2020. Four vessels participated in the fishery and harvested 107,679 pounds, of which 1.5% was deadloss (Table 8). Sporadic harvest occurred late January through early November. Harvest by statistical week is confidential and therefore not shown in tables. On average, vessels were active in the fishery for 48 days. Harvesters were paid an initial average exvessel price of \$5.87 per pound, the highest PIGKC exvessel price since the inception of the fishery. Total exvessel fishery value was estimated to be \$621,971 (Table 9).

Total effort for the 2020 fishery was 2,960 pot lifts. Catch per unit effort (CPUE) was 8 legal crab per pot, slightly below the 10-year (2009–2019) average CPUE of 10 (Table 8). Harvest was distributed across 6 ADF&G statistical reporting areas with the highest concentration of harvest (33%) occurring in ADF&G statistical area 705600, southwest of St. George Island (Table 10).

Port Sampling

During the 2020 PIGKC fishery, ADF&G personnel and onboard observers sampled golden king crab from vessels at shore-based processors in Dutch Harbor and Juneau. Biological data collected on landed golden king crab consisted of carapace length, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 12 of the 12 total landings during the 2020 PIGKC fishery. Average weight of sampled crab was 4.4 pounds and average carapace length was 149 mm (Table 8).

Fishery Management and Stock Status

The golden king crab fishery is managed using a GHL based on a buffered federal acceptable biological catch (ABC) that approximates long-term average harvest. In 2012, NPFMC established a total-catch overfishing level (OFL) of 200,000 pounds. The 2012 OFL used the relationship of bycatch mortality to retained catch in the Pribilof District golden king crab fishery from 2001 to 2010 (NPFMC 2011).

An ABC of 180,000 pounds was also set in 2012, by applying a 10% buffer on the Tier 5 harvest control rule OFL (NPFMC 2011). For 2015, ABC was reduced to 150,000 pounds by employing a 25% buffer on the OFL to account for uncertainty associated with limited stock condition data (NPFMC 2014). The GHL is established by reducing the ABC to account for bycatch mortality of golden king crab across all fisheries.

Stock biomass of golden king crab in the Pribilof District has been estimated using area-swept extrapolations applied to NMFS slope trawl survey data. Although biomass estimates of golden king crab from the slope survey are available, they are highly uncertain and not currently used in fishery management. The most recent NMFS Bering Sea slope survey was conducted in 2016. Results of the 2016 NMFS Bering Sea slope survey show an estimated biomass of 3.12 million pounds (Hoff 2016).

Further information on Pribilof District golden king crab stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

BERING SEA SCARLET KING CRAB

2020 Fishery

One vessel harvested scarlet king crab in the Bering Sea during 2020. Harvest was restricted to incidental harvest during the Bering Sea golden king crab fishery. Harvest information is confidential (Table 11).

Fishery Management and Stock Status

In 2007, NMFS amended the FMP and removed Bering Sea scarlet king crab, which transferred sole jurisdiction for the fishery to the state (NPFMC 2007). Onboard observers have been required on most vessels that targeted deepwater crab species since 1994 and have collected information detailing the size and sex composition of the retained and nonretained scarlet king crab and bycatch species. This information is useful when developing management measures for deepwater crab stocks; however, there is currently no harvest strategy.

Stock biomass of scarlet king crab in the Bering Sea has been estimated using area-swept methods applied to NMFS upper continental slope trawl survey data (Hoff 2016). Although biomass estimates of scarlet king crab from the slope survey are available, they are highly uncertain and not currently used in fishery management. The most recent NMFS Bering Sea slope survey was conducted in 2016. Results of the 2016 NMFS Bering Sea slope survey show an estimated biomass of 0.97 million pounds (Hoff 2016).

BERING SEA TANNER CRAB MANAGEMENT DISTRICT

DESCRIPTION OF AREA

The Bering Sea District of Tanner crab Registration Area J includes all waters north of Cape Sarichef (54°36'N lat). The district is divided into the Eastern and Western Subdistricts at 173°W long. The Eastern Subdistrict is further divided into two sections: the Norton Sound Section north of the latitude of Cape Romanzof (61°49'N lat) and east of 168°W long, and the General Section south and west of the Norton Sound Section (Figure 5).

BERING SEA TANNER CRAB

The Bering Sea Tanner crab stock is managed as two separate fisheries, east and west of 166°W long, with a separate TAC set for each fishery.

2020/21 Fishery East of 166°W Longitude

The 2020/21 Bering Sea Tanner crab fishery east of 166°W long (EBT) was closed (Tables 12 and 13, Figure 6). Model-based mature male biomass estimate east of 166°W long was below the regulatory threshold (mature male biomass must be at least 25% of the 1982–2018 average) required to open the fishery. The harvest strategy is found in 5 AAC 35.508 *Bering Sea District C. bairdi Tanner crab harvest strategy*.

Stock Status

The 2020 NMFS trawl survey of the eastern Bering Sea was cancelled due to COVID-19.

Based on 2019 NMFS trawl survey estimates east of 166°W long (Zacher et al. 2020), estimated immature male biomass of 9.7 million pounds was below the 20-year average of 21.2 million pounds, estimated mature male biomass of 14.1 million pounds was below the 20-year average of 36.3 million pounds, estimated legal male biomass of 12.2 million pounds was below the 20-year average of 28.5 million pounds, and estimated mature female biomass of 1.4 million pounds was below the 20-year average of 8.1 million pounds.

Since no survey data were available, and the fishery was closed during the 2019/20 season, new data for the 2020 assessment was limited to updating bycatch in the groundfish fishery and bycatch from other crab fisheries. Retrospective analysis shows this stock is sensitive to missing survey data when estimating terminal year recruitment, and therefore the average recruitment from 1982 through 2019 was used to approximate this year's recruitment. Projected model estimates show continued decline in industry-preferred 5-inch males, with the 2019 survey estimate the fourth lowest in the timeseries. The projected estimates of mature male abundance for EBT were below threshold to allow for a fishery opening (NPFMC 2020a, Scientific and Statistical Committee 2020). The 2019 trawl survey showed weak signs of recruitment into the mature size classes (Zacher et al. 2020).

Further information on Tanner crab stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

2020/21 Fishery West of 166°W Longitude

The 2020/21 Bering Sea Tanner crab fishery west of 166°W long (WBT) opened October 15 with a combined IFQ and CDQ TAC of 2,348,000 pounds (Table 12). TAC was allocated by NMFS as 90% IFQ (2,113,200 pounds) and 10% CDQ (234,800 pounds) with all six of the CDQ groups participating in the harvest of the CDQ allocation. Forty-one vessels participated in the fishery and harvested 1,449,543 pounds, of which 1.7% was deadloss (Table 12). Harvest occurred throughout the season, with the last delivery on April 4 (Table 13 and 14). On average, vessels were active in the fishery for 24 days. Harvesters were paid an initial average exvessel price of \$3.23 per pound. Total exvessel fishery value was estimated to be \$4,676,509 (Table 13).

Total effort for the 2020/21 fishery was 47,388 pot lifts. CPUE was 18 legal crab per pot, slightly below the post-rationalization (2005/06–2019/20) average CPUE of 20 (Table 12; Figure 6).

Harvest was distributed across 49 ADF&G statistical reporting areas with the highest concentration of harvest (37%) occurring in ADF&G statistical area 665530, southeast of Saint George Island (Table 15).

Port Sampling

During the 2020/21 WBT fishery, ADF&G personnel and onboard observers sampled Tanner crab from vessels at shore-based processors in Akutan, Dutch Harbor, King Cove, and Saint Paul Island. Biological data collected on landed Tanner crab consisted of carapace width, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 82 of the 84 total landings (IFQ and CDQ) during the 2020/21 WBT fishery. Average weight of sampled crab was 1.7 pounds and average carapace width was 134 mm (Table 12).

Stock Status

The 2020 NMFS trawl survey of the eastern Bering Sea was cancelled due to COVID-19.

Based on 2019 NMFS trawl survey estimates west of 166°W long (Zacher et al. 2020), estimated immature male biomass of 17.0 million pounds was below the 20-year average of 30.0 million pounds, estimated mature male biomass of 21.6 million pounds was below the 20-year average of 47.8 million pounds, estimated legal male biomass of 19.3 million pounds was below the 20-year average of 39.9 million pounds, and estimated mature female biomass of 9.1 million pounds was below the 20-year average of 11.4 million pounds.

Because no survey data were available and the fishery was closed during the 2019/20 season, new data for the 2020 assessment was limited to updating bycatch in the groundfish fishery and bycatch from other crab fisheries. Retrospective analysis shows that this stock is sensitive to missing survey data when estimating terminal year recruitment, and therefore the average recruitment from 1982 through 2019 was used to approximate this year's recruitment. Projected model estimates show continued decline in industry-preferred 5-inch males. The projected estimates of mature male abundance for WBT were above threshold and allowed for a fishery opening (NPFMC 2020a, Scientific and Statistical Committee 2020). Estimates from the 2019 trawl survey (Zacher et al. 2020) show strong juvenile recruitment into the WBT and could potentially be a sign of future stock recovery.

Further information on Tanner crab stock status and federal overfishing levels may be found in the *2020 Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

BERING SEA SNOW CRAB

2020/21 Fishery

The 2020/21 Bering Sea snow crab (BSS) fishery opened October 15 with a combined IFQ and CDQ TAC of 45,000,000 pounds (Table 16). TAC was allocated by NMFS as 90% IFQ (40,500,000 pounds) and 10% CDQ (4,500,000 pounds) with all six of the CDQ groups participating in the harvest of the CDQ allocation. Sixty-two vessels participated in the fishery and harvested 45,001,190 pounds, of which 1.9% was deadloss (Table 16). Harvest occurred late December through May, with the last delivery on June 2 (Table 17 and 18). On average, vessels

were active in the fishery for 90 days. Harvesters were paid an initial average exvessel price of \$3.01 per pound. Total exvessel fishery value was estimated to be \$105,928,621 (Table 17).

Total effort for the 2020/21 fishery was 171,678 pot lifts. CPUE was 218 legal crab per pot, slightly above the post-rationalization (2005/06–2019/20) average CPUE of 215 (Table 16; Figure 7). Harvest was distributed across 57 ADF&G statistical reporting areas with the highest concentration of harvest (22%) occurring in ADF&G statistical area 786030, northwest of Saint Matthew Island adjacent to the United States–Russia Maritime Boundary Line (Table 19).

Port Sampling

During the 2020/21 Bering Sea snow crab fishery, ADF&G personnel and onboard observers sampled snow crab from vessels at shore-based processors in Akutan, Dutch Harbor, King Cove, and Saint Paul. Biological data collected on landed snow crab consisted of carapace width, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 389 of the 407 total landings (IFQ and CDQ) during the 2020/21 Bering Sea snow crab fishery. Average weight of sampled crab was 1.2 pounds and average carapace width was 107 mm (Table 16).

Stock Status

The 2020 NMFS trawl survey of the eastern Bering Sea was cancelled due to COVID-19.

Based on 2019 NMFS trawl survey estimates (Zacher et al. 2020), estimated immature male biomass of 626.5 million pounds was above the 20-year average of 297.8 million pounds, estimated mature male biomass of 120.3 million pounds was below the 20-year average of 148.4 million pounds, estimated legal male biomass of 387.8 million pounds was above the 20-year average of 234.8 million pounds, and estimated mature female biomass of 235.5 million pounds was above the 20-year average of 179.1 million pounds.

The snow crab stock assessment, in particular, is highly sensitive to loss of terminal-year survey data, in contrast to the other major crab stocks. Because the survey was cancelled, updated data in the 2020 assessment included retained and total catch and length frequencies from the 2019/20 directed fishery, and discard catch and length frequencies from the 2019/20 groundfish fisheries. The crab plan team initially recommended a new model for 2020 that was based on a generalized model for Alaskan crab stocks (GMACS) framework. The new GMACS framework resulted in estimated stock abundance and an OFL/ABC that were substantially higher than in previous years. The GMACS model was rejected by the NPFMC Scientific and Statistical Committee and the assessment reverted to using mythology based on the accepted 2019 model. Results from the 2019 model using available 2020 data were much lower than GMACs but still high compared to previous seasons (NPFMC 2020a, Scientific and Statistical Committee 2020).

All model estimates showed a large increase in abundance of industry-preferred 4-inch males, but high uncertainty associated with the magnitude of model estimates was notable given lack of survey 2020 data and the observed decline in survey abundance between the 2018 and 2019 eastern Bering Sea trawl surveys. However, given projected high recruitment estimates in both stock assessment models, managers concluded a surplus of industry-preferred 4-inch males was available to the fishery despite not having current survey data to inform the decision. Additionally, managers assumed 75% old-shell selectivity for TAC computation with selectivity scaled to reflect

anticipated old-shell selectivity in the 2020/21 fishery given the likely small proportion of old-shell 4-inch males in the population. The TAC for 2020/21 was set by calculating an average of the three-year trend of model survey observed abundance, resulting in a TAC of 45 million pounds.

Further information on snow crab stock status and federal overfishing levels may be found in the *2020 Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

BERING SEA GROOVED TANNER CRAB

2020 Fishery

No vessels harvested grooved Tanner crab in the Bering Sea during 2020 (Table 20).

Fishery Management and Stock Status

In 2007, NMFS amended the FMP and removed Bering Sea grooved Tanner crab, which transferred sole jurisdiction for the fishery to the state (NPFMC 2007). Onboard observers have been required on most vessels that targeted deepwater crab species since 1994 and have collected information detailing the size and sex composition of the retained and nonretained grooved Tanner crab and bycatch species. This information is useful when developing management measures for deepwater crab stocks; however, there is currently no harvest strategy.

Stock biomass of grooved Tanner crab in the Bering Sea has been estimated using area-swept methods applied to NMFS upper continental slope trawl survey data (Hoff 2016). Although biomass estimates of grooved Tanner crab from the slope survey are available, they are highly uncertain and not currently used in fishery management. The most recent NMFS Bering Sea slope survey was conducted in 2016. Results of the 2016 NMFS Bering Sea slope survey show an estimated biomass of 3.81 million pounds in the eastern Bering Sea (Hoff 2016).

BERING SEA TRIANGLE TANNER CRAB

2020 Fishery

No vessels harvested triangle Tanner crab in the Bering Sea during 2020 (Table 21).

Fishery Management and Stock Status

In 2007, NMFS amended the FMP and removed Bering Sea triangle Tanner crab, which transferred sole jurisdiction for the fishery to the state (NPFMC 2007).

Triangle Tanner crab are currently managed as bycatch only to other nonrationalized fisheries. Stock biomass of triangle Tanner crab in the Bering Sea has been estimated using area-swept methods applied to NMFS upper continental slope trawl survey data (Hoff 2016). Although biomass estimates of triangle Tanner crab from the slope survey are available, they are highly uncertain and not currently used in fishery management. The most recent NMFS Bering Sea slope survey was conducted in 2016. Results of the 2016 NMFS Bering Sea slope survey show an estimated biomass of 13.5 million pounds in the eastern Bering Sea (Hoff 2016).

NORTH PENINSULA DISTRICT DUNGENESS CRAB

DESCRIPTION OF DISTRICT

The North Peninsula District for Dungeness crab includes all waters of Registration Area J north of the latitude of Cape Sarichef at 54°36'N lat (Figure 8).

DUNGENESS CRAB

2020 Fishery

The North Peninsula Dungeness crab fishery opened May 1, 2020. Two vessels participated in the fishery; harvest information is confidential (Table 22).

Fishery Management and Stock Status

The North Peninsula Dungeness crab fishery is managed using 3-S management (size, sex, and season). Only male Dungeness crab 6.5 inches (165 mm) or greater in carapace width may be retained from May 1 to October 18. No stock assessment is available and limited biological and fishery data have been collected through dockside sampling.

BRISTOL BAY - BERING SEA WEATHERVANE SCALLOP REGISTRATION AREA Q

DESCRIPTION OF AREA

Bristol Bay–Bering Sea weathervane scallop Registration Area Q is a combination of the Bristol Bay king crab Registration Area T and Bering Sea king crab Registration Area Q (Figure 9).

BRISTOL BAY - BERING SEA WEATHERVANE SCALLOP

2020/21 Fishery

The 2020/21 Bristol Bay–Bering Sea weathervane scallop fishery opened July 1, 2020, with a GHF of 7,500 pounds of shucked scallop meat. No vessels participated in the fishery (Table 23) and it closed by regulation on February 15, 2021.

Fishery Management and Stock Status

Fishing effort in Bristol Bay–Bering Sea weathervane scallop Registration Area Q primarily occurs north–northwest of Unimak Island. The fishery CPUE of 20 pounds of shucked scallop meat per dredge hour in the 2019/20 season was a time-series low and a continuation of overall decline in fishery performance since the 2014/15 season. Area Q contributed substantially to statewide scallop harvest until the 2014/15 season when fishery performance declined rapidly and harvesters reported weak meat condition. Tissue samples collected by fishery observers have shown a high prevalence of an apicomplexan parasite in the Bering Sea beds (NPFMC 2017), similar to the apicomplexan seen in Icelandic scallop beds, causing the weak meat condition. Infection in the scallop beds is thought to be the primary driver of poor fishery performance, although other environmental or recruitment factors may contribute. Absent survey data, minimal GHFs have been set since 2015 to monitor bed condition and maintain the fishery data time series.

Area Q has never been surveyed with dredge gear as part of the Statewide Weathervane Scallop Dredge survey. The NMFS annual trawl survey overlaps with the Area Q fishing grounds, but

scallop shell height composition data is not collected. Therefore, the stock is assessed annually using available fishery and observer data. The harvest strategy is found in 5 AAC 38.076 *Alaska Scallop Fishery Management Plan* and 5 AAC 38.078 *State-Waters Weathervane Scallop Management Plan*.

Further information on weathervane scallop stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the Scallop Fishery off Alaska* (NPFMC 2020b).

BERING SEA MISCELLANEOUS SHELLFISH SPECIES

DESCRIPTION OF DISTRICT

The Bering Sea District of Registration Area J for miscellaneous shellfish includes all waters north of the latitude of Cape Sarichef at 54°36'N lat and east of the United States–Russia Maritime Boundary Line of 1990 (Figure 10).

2020 FISHERIES

Octopus

In 2020, harvest from state waters was 331 pounds from 8 vessels and 12 landings. Harvest from state and federal waters combined was 42,438 pounds from 120 vessels and 168 landings. All reported harvest was incidental to groundfish fisheries. Average exvessel price based on landed weight of octopus in 2020 was \$0.19 per pound (Table 24).

Fishery Management and Stock Status

Octopus are considered a shellfish species under State of Alaska regulation. Limited directed fishing within state waters may occur under the authority of a commissioner's permit; however, octopus are primarily retained as bycatch during state and federal groundfish fisheries. Currently, vessels may retain incidentally caught octopus up to 20% of the weight of the target groundfish species or halibut onboard. Most octopus are retained as bycatch in Pacific cod pot gear fisheries.

Incidental harvest of octopus in the Bering Sea is dominated by giant Pacific octopus *Enteroctopus dofleini*, although at least nine other species of octopus are known to occur in the Bering Sea. The 2020 NMFS trawl survey of the eastern Bering Sea was cancelled due to COVID-19. Results from the 2019 NMFS Eastern Bering Sea shelf trawl survey estimate octopus (all species) biomass at 13.1 million pounds with 94% of the estimate composed of *E. dofleini*. The 2019 octopus (all species) biomass estimate is above the 5-year average (2014–2018) of 10.5 million pounds. Biomass estimates for octopus in the Bering Sea are highly variable and do not necessarily reflect the sizes of octopus caught in fisheries (Ormseth et al. 2020). General knowledge of the stock is limited and there is currently no reliable estimate of octopus biomass (Ormseth et al. 2020).

Further information on octopus stock status and federal overfishing levels may be found in the 2020 *Assessment of the Octopus Stock Complex in the Bering Sea and Aleutian Islands* (Ormseth et al. 2020).

REFERENCES CITED

- Ormseth, O. A., M. E. Conners, K. Aydin, and C. Conrath. 2020. Assessment of the Octopus Stock Complex in the Bering Sea and Aleutian Islands. [*In*] Stock Assessment and Fishery Evaluation Report of the Bering Sea/Aleutian Islands Regions. North Pacific Fishery Management Council, Anchorage.
- Hoff, G. R. 2016. Results of the 2016 eastern Bering Sea upper continental slope survey of groundfish and invertebrate resources. U.S. Department of Commerce, NOAA Technical Memo NMFS-AFSC-339.
- Leon, J. M., J. Shaishnikoff, E. Nichols, and M. Westphal. 2017. Annual management report for shellfish fisheries of the Bering Sea/Aleutian Islands Management Area, 2015/16. Alaska Department of Fish and Game, Fishery Management Report No. 17-10, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2007. Fisheries management plan for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2011. Stock assessment and fishery evaluation report for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions: 2011 crab SAFE. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2014. Stock assessment and fishery evaluation report for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions: 2014 crab SAFE. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2017. Stock assessment and fishery evaluation report for the weathervane scallop fishery off Alaska: 2017 scallop SAFE. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2020a. Stock assessment and fishery evaluation report for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions: 2020 crab SAFE. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2020b. Stock assessment and fishery evaluation report for the weathervane scallop fishery off Alaska: 2020 scallop SAFE. North Pacific Fishery Management Council, Anchorage.
- Scientific and Statistical Committee final report to the North Pacific Fishery Management Council. 2020, September 28 – October 2. Virtual Meeting. Retrieved from <https://meetings.npfmc.org>.
- Zacher, L. S., J. I. Richar, and R. J. Foy. 2020. The 2019 eastern and northern Bering Sea continental shelf bottom trawl surveys: Results for commercial crab species. U.S. Department of Commerce, NOAA Technical Memo NMFS-AFSC-400.

TABLES AND FIGURES

Table 1.—Bristol Bay red king crab commercial fishery harvest data, 1966–2020/21.

| Season | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|--------|----------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 1966 | — | 997,321 | ND | 9 | 15 | 140,554 | 2,720 | 52 | ND | ND |
| 1967 | — | 3,102,443 | ND | 20 | 61 | 397,307 | 10,621 | 37 | ND | ND |
| 1968 | — | 8,686,546 | ND | 59 | 261 | 1,278,592 | 47,496 | 27 | ND | ND |
| 1969 | — | 10,403,283 | ND | 65 | 377 | 1,749,022 | 98,426 | 18 | ND | ND |
| 1970 | — | 8,559,178 | ND | 51 | 309 | 1,682,591 | 96,658 | 17 | ND | ND |
| 1971 | — | 12,955,776 | ND | 52 | 394 | 2,404,681 | 118,522 | 20 | ND | ND |
| 1972 | — | 21,744,924 | ND | 64 | 611 | 3,994,356 | 205,045 | 19 | ND | ND |
| 1973 | — | 26,913,636 | ND | 67 | 441 | 4,825,963 | 194,095 | 25 | 5.6 | ND |
| 1974 | — | 42,266,274 | ND | 104 | 605 | 7,710,317 | 212,915 | 36 | 5.5 | ND |
| 1975 | — | 51,326,259 | 1,639,483 | 102 | 592 | 8,745,294 | 205,096 | 43 | 5.7 | ND |
| 1976 | — | 63,919,728 | 875,327 | 141 | 984 | 10,603,367 | 321,010 | 33 | 6.0 | 148 |
| 1977 | — | 69,967,868 | 730,279 | 130 | 1,020 | 11,733,101 | 451,273 | 26 | 5.9 | 148 |
| 1978 | — | 87,618,320 | 1,273,037 | 162 | 926 | 14,745,709 | 406,165 | 36 | 5.9 | 147 |
| 1979 | — | 107,828,057 | 3,555,891 | 236 | 889 | 16,808,605 | 315,226 | 53 | 6.4 | 152 |
| 1980 | 70–120 million | 129,948,463 | 1,858,668 | 236 | 1,251 | 20,845,350 | 267,292 | 37 | 6.2 | 151 |
| 1981 | 10–100 million | 33,372,832 | 706,489 | 177 | 1,013 | 5,273,530 | 536,646 | 10 | 6.3 | 151 |
| 1982 | 10–20 million ^g | 2,990,082 | 95,834 | 89 | 253 | 538,925 | 140,492 | 4 | 5.5 | 145 |
| 1983 | | | | No Commercial Fishery | | | | | | |
| 1984 | 2.5–6 million | 4,083,612 | 35,101 | 89 | 133 | 793,046 | 107,406 | 7 | 5.2 | 142 |
| 1985 | 3–5 million | 4,090,305 | 6,436 | 125 | 130 | 780,791 | 84,443 | 9 | 5.2 | 142 |
| 1986 | 6–13 million | 11,306,084 | 284,126 | 157 | 229 | 2,083,496 | 175,753 | 12 | 5.4 | 142 |
| 1987 | 8.5–17.7 million | 12,289,067 | 120,388 | 230 | 311 | 2,122,341 | 220,971 | 10 | 5.8 | 145 |
| 1988 | 7,500,000 | 7,361,026 | 23,537 | 200 | 201 | 1,231,731 | 146,179 | 8 | 6.0 | 147 |

-continued-

Table 1.—Page 2 of 3.

| Season | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|----------------------|------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 1989 | 16,500,000 | 10,156,849 | 81,334 | 207 | 280 | 1,667,405 | 205,528 | 8 | 6.1 | 148 |
| 1990 | 17,100,000 | 20,443,043 | 141,067 | 241 | 333 | 3,135,382 | 262,761 | 12 | 6.5 | 152 |
| 1991 | 18,000,000 | 17,177,894 | 119,670 | 300 | 325 | 2,630,446 | 227,555 | 12 | 6.5 | 152 |
| 1992 | 10,300,000 | 8,070,129 | 9,000 | 280 | 289 | 1,201,129 | 206,815 | 6 | 6.7 | 153 |
| 1993 | 16,800,000 | 14,587,704 | 134,314 | 292 | 360 | 2,254,989 | 254,389 | 9 | 6.5 | 152 |
| 1994–1995 | | | | No Commercial Fishery | | | | | | |
| 1996 | 5,000,000 | 8,523,114 | 26,084 | 197 | 200 | 1,266,048 | 77,081 | 16 | 6.7 | 153 |
| 1997 | 7,000,000 | 8,911,387 | 32,012 | 257 | 269 | 1,340,591 | 91,085 | 15 | 6.6 | 152 |
| 1998 ^h | 16,400,000 | 15,003,359 | 87,978 | 275 | 301 | 2,241,489 | 145,689 | 15 | 6.7 | 152 |
| 1999 | 10,660,000 | 11,835,930 | 44,807 | 258 | 284 | 1,927,105 | 151,212 | 13 | 6.1 | 148 |
| 2000 ⁱ | 8,350,000 | 8,240,644 | 37,230 | 244 | 270 | 1,272,382 | 104,056 | 12 | 6.5 | 151 |
| 2001 ⁱ | 7,150,000 | 8,523,495 | 59,973 | 230 | 257 | 1,305,396 | 66,947 | 19 | 6.5 | 151 |
| 2002 ⁱ | 9,270,489 | 9,666,847 | 35,122 | 242 | 272 | 1,498,574 | 72,514 | 21 | 6.5 | 151 |
| 2003 ⁱ | 15,713,000 | 15,728,256 | 230,467 | 250 | 296 | 2,524,506 | 134,515 | 19 | 6.2 | 149 |
| 2004 ^{h,i} | 15,424,000 | 15,447,030 | 163,750 | 251 | 294 | 2,272,184 | 97,621 | 23 | 6.8 | 154 |
| 2005/06 ^j | 18,329,000 | 18,309,335 | 86,288 | 89 | 296 | 2,732,563 | 114,944 | 24 | 6.7 | 152 |
| 2006/07 | 15,527,000 | 15,616,816 | 118,227 | 81 | 213 | 2,455,618 | 72,106 | 34 | 6.4 | 151 |
| 2007/08 | 20,383,000 | 20,366,065 | 140,384 | 74 | 281 | 3,139,336 | 113,214 | 28 | 6.5 | 151 |
| 2008/09 | 20,364,000 | 20,329,402 | 173,163 | 78 | 289 | 3,066,286 | 139,937 | 22 | 6.6 | 153 |
| 2009/10 | 16,009,000 | 15,932,654 | 122,207 | 70 | 233 | 2,537,221 | 118,521 | 21 | 6.3 | 150 |
| 2010/11 | 14,839,000 | 14,833,829 | 106,874 | 65 | 254 | 2,398,490 | 131,627 | 18 | 6.2 | 150 |
| 2011/12 | 7,834,000 | 7,833,594 | 32,068 | 62 | 161 | 1,279,054 | 45,166 | 28 | 6.1 | 149 |
| 2012/13 | 7,853,000 | 7,849,835 | 30,050 | 64 | 141 | 1,157,364 | 38,159 | 30 | 6.8 | 154 |

-continued-

Table 1.—Page 3 of 3.

| Season | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|------------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 2013/14 | 8,600,000 | 8,600,476 | 62,749 | 63 | 156 | 1,242,705 | 45,927 | 27 | 6.9 | 155 |
| 2014/15 | 9,986,000 | 9,987,008 | 101,241 | 63 | 159 | 1,498,538 | 58,702 | 26 | 6.7 | 152 |
| 2015/16 | 9,974,000 | 9,969,964 | 182,833 | 64 | 152 | 1,497,783 | 48,008 | 31 | 6.7 | 153 |
| 2016/17 | 8,469,000 | 8,466,701 | 41,120 | 63 | 148 | 1,253,967 | 33,126 | 38 | 6.8 | 154 |
| 2017/18 | 6,601,000 | 6,600,922 | 24,880 | 61 | 142 | 964,593 | 48,242 | 20 | 6.8 | 156 |
| 2018/19 | 4,308,000 | 4,307,946 | 27,575 | 55 | 121 | 606,421 | 30,722 | 20 | 7.1 | 156 |
| 2019/20 | 3,797,000 | 3,791,569 | 8,874 | 56 | 116 | 531,329 | 34,458 | 15 | 7.1 | 156 |
| 2020/21 | 2,648,000 | 2,646,874 | 3,905 | 47 | 95 | 433,482 | 20,244 | 21 | 6.1 | 150 |

Note: En dashes indicate harvest limits were not set for this fishery prior to 1980; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Guideline harvest level (GHL) began in 1980; total allowable catch (TAC) began in 2005/06.

^c Deadloss included.

^d Number of legal crab per pot lift.

^e Retained catch.

^f Carapace length in millimeters.

^g Inseason revision to 4.7 million pounds.

^h Total GHL announced prior to general fishery opening; CDQ GHL adjusted based on general fishery harvest.

ⁱ Includes American Fisheries Act (AFA) harvest data.

^j Crab rationalization begins.

Table 2.—Bristol Bay red king crab commercial fishery value and season dates, 1980–2020/21.

| Season | Value | | Season length | | |
|----------------------|-----------------------|---------------|---------------|-----------------------|------|
| | Exvessel ^a | Total | Opened | Closed | Days |
| 1980 | \$0.90 | \$115,280,816 | 09/10/80 | 10/20/80 | 41 |
| 1981 | \$2.03 | \$66,312,676 | 09/10/81 | 12/15/81 | 97 |
| 1982 | \$3.20 | \$9,261,594 | 09/10/82 | 10/10/82 | 31 |
| 1983 | No Commercial Fishery | | | | |
| 1984 | \$2.72 | \$11,011,950 | 10/01/84 | 10/16/84 | 16 |
| 1985 | \$2.90 | \$11,843,220 | 09/25/85 | 10/02/85 | 8 |
| 1986 | \$4.05 | \$44,638,930 | 09/25/86 | 10/07/86 | 13 |
| 1987 | \$3.95 | \$48,070,777 | 09/25/87 | 10/06/87 | 12 |
| 1988 | \$4.85 | \$35,610,150 | 09/25/88 | 10/02/88 | 8 |
| 1989 | \$4.68 | \$47,202,150 | 09/25/89 | 10/06/89 | 12 |
| 1990 | \$5.03 | \$102,198,044 | 11/01/90 | 11/13/90 | 13 |
| 1991 | \$3.00 | \$51,174,672 | 11/01/91 | 11/08/91 | 8 |
| 1992 | \$5.13 | \$41,323,471 | 11/01/92 | 11/08/92 | 8 |
| 1993 | \$4.48 | \$64,750,431 | 11/01/93 | 11/10/93 | 10 |
| 1994–1995 | No Commercial Fishery | | | | |
| 1996 | \$4.01 | \$34,104,724 | 11/01/96 | 11/05/96 | 5 |
| 1997 | \$3.27 | \$29,033,365 | 11/01/97 | 11/05/97 | 5 |
| 1998 | \$2.60 | \$38,852,795 | 11/01/98 | 11/6/98 ^b | 6 |
| 1999 | \$6.23 | \$73,473,949 | 10/15/99 | 10/20/99 ^b | 6 |
| 2000 | \$4.82 | \$39,542,935 | 10/16/00 | 10/20/00 ^b | 5 |
| 2001 | \$4.95 | \$41,934,728 | 10/15/01 | 10/18/01 ^b | 4 |
| 2002 | \$6.13 | \$59,016,000 | 10/15/02 | 10/18/02 ^b | 4 |
| 2003 | \$5.05 | \$78,253,116 | 10/15/03 | 10/20/03 ^b | 6 |
| 2004 | \$4.64 | \$70,936,658 | 10/15/04 | 10/18/04 ^b | 4 |
| 2005/06 ^c | \$3.85 | \$70,086,741 | 10/15/05 | 01/15/06 | 93 |
| 2006/07 | \$3.37 | \$52,277,156 | 10/15/06 | 01/15/07 | 93 |
| 2007/08 | \$4.16 | \$84,211,504 | 10/15/07 | 01/15/08 | 93 |
| 2008/09 | \$4.97 | \$100,222,813 | 10/15/08 | 01/15/09 | 93 |
| 2009/10 | \$4.44 | \$70,172,988 | 10/15/09 | 01/15/10 | 93 |
| 2010/11 | \$6.31 | \$92,924,994 | 10/15/10 | 01/15/11 | 93 |
| 2011/12 | \$8.91 | \$69,479,174 | 10/15/11 | 01/15/12 | 93 |
| 2012/13 | \$7.28 | \$56,914,521 | 10/15/12 | 01/15/13 | 93 |
| 2013/14 | \$6.41 | \$54,763,067 | 10/15/13 | 01/15/14 | 93 |
| 2014/15 | \$6.05 | \$59,849,844 | 10/15/14 | 01/15/15 | 93 |

-continued-

Table 2.–Page 2 of 2.

| Season | Value | | Season length | | |
|---------|-----------------------|--------------|---------------|----------|------|
| | Exvessel ^a | Total | Opened | Closed | Days |
| 2015/16 | \$7.02 | \$68,754,179 | 10/15/15 | 01/15/16 | 93 |
| 2016/17 | \$9.06 | \$76,311,556 | 10/15/16 | 01/15/17 | 93 |
| 2017/18 | \$8.33 | \$54,792,098 | 10/15/17 | 01/15/18 | 93 |
| 2018/19 | \$8.45 | \$36,176,631 | 10/15/18 | 01/15/19 | 93 |
| 2019/20 | \$9.04 | \$34,208,234 | 10/15/19 | 01/15/20 | 93 |
| 2020/21 | \$9.11 | \$24,086,513 | 10/15/20 | 01/15/21 | 93 |

^a Average price per pound.

^b CDQ fishery opened after general fishery.

^c Crab rationalization begins.

Table 3.—Bristol Bay red king crab commercial fishery harvest and effort by week, 2020/21.

| Week ending | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| Oct-17 | 2,025,027 | 3,528 | 41 | 63 | 332,641 | 15,811 | 21 | 6.1 |
| Oct-24 | 246,559 | 67 | 7 | 13 | 40,500 | 1,380 | 29 | 6.1 |
| Oct-31 | 294,378 | 310 | 7 | 12 | 48,458 | 1,817 | 27 | 6.1 |
| Nov-7 | 116,488 | 192 | 3 | 8 | 16,331 | 667 | 24 | 7.1 |
| Nov-14 | CF | CF | 2 | 3 | CF | CF | CF | CF |
| ... ^e | — | — | — | — | — | — | — | — |
| Jan-9 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Jan-16 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Total | 2,646,874 | 3,905 | 47 | 95 | 433,482 | 20,244 | 21 | 6.1 |

Note: CF indicates confidential data

^a In pounds.

^b Deadloss included.

^c Number of retained crab per pot lift.

^d Retained catch.

^e Consecutive weeks with no harvest.

Table 4.—Bristol Bay red king crab commercial fishery harvest and effort by statistical area, 2020/21.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 615601 | 69,334 | 113 | 6 | 16 | 11,615 | 367 | 32 | 6.0 |
| 615630 | 449,179 | 1,047 | 19 | 37 | 75,202 | 2,678 | 28 | 6.0 |
| 615700 | 38,974 | 29 | 5 | 7 | 6,540 | 219 | 30 | 6.0 |
| 625600 | 326,005 | 278 | 16 | 40 | 54,783 | 3,010 | 18 | 6.0 |
| 625630 | 1,162,019 | 1,760 | 34 | 67 | 190,567 | 8,145 | 23 | 6.1 |
| 635600 | 71,238 | 59 | 13 | 19 | 11,696 | 1,072 | 11 | 6.1 |
| 635630 | 325,705 | 384 | 19 | 33 | 51,688 | 2,629 | 20 | 6.3 |
| 645600 | 22,676 | 14 | 13 | 18 | 3,514 | 657 | 5 | 6.5 |
| 645630 | 179,997 | 218 | 14 | 25 | 27,583 | 1,448 | 19 | 6.5 |
| Other ^e | 1,747 | 4 | 2 | 2 | 294 | 19 | 15 | 5.9 |
| Total | 2,646,874 | 3,905 | 47 | 95 | 433,482 | 20,244 | 21 | 6.1 |

^a In pounds.^b Deadloss included.^c Number of legal crab per pot lift.^d Retained catch.^e Combination of statistical areas (2) in which landings were made by fewer than three vessels.

Table 5.—Bristol Bay red king crab cost-recovery harvest data and charter length, 1990–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Charter Length | |
|---------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------------------|-------------------|-----------------------|---------------------------------------|------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Dates | Days |
| 1990 ^e | 80,701 | 24,540 | 1 | 3 | 9,567 | 870 | 16 | 5.9 | 08/07–09/07 | 30 |
| 1991 ^e | 205,851 | 12,817 | 1 | 2 | 30,351 | 518 | 62 | 6.4 | 09/02–10/07 | 35 |
| 1992 ^e | 74,089 | 3,000 | 1 | 1 | 11,213 | 670 | 17 | 6.3 | 10/08–10/23 | 15 |
| 1993 ^e | 53,200 | 800 | 1 | 1 | 8,384 | 464 | 18 | 6.3 | 08/20–09/20 | 31 |
| 1994 ^e | 93,336 | 4,500 | 1 | 1 | 14,806 | 732 | 21 | 6.0 | 09/25–10/25 | 30 |
| 1995 ^e | 80,158 | 2,339 | 1 | 2 | 14,123 | 564 | 26 | 5.5 | 08/01–08/31 | 31 |
| 1996 ^e | 107,955 | 1,918 | 1 | 3 | 15,390 | 355 | 44 | 6.9 | 08/01–08/31 | 31 |
| 1997 ^e | 154,739 | 18,040 | 1 | 4 | 21,698 | 658 | 37 | 6.3 | 07/25–08/21 | 28 |
| 1998 ^e | 188,176 | 32,564 | 1 | 2 | 22,230 | 738 | 36 | 7.0 | 08/01–08/28 | 28 |
| 1999 ^f | 185,944 | 410 | 2 | 4 | 29,368 | 1,239 | 24 | 6.3 | 09/25–10/11, 10/25–11/10 | 34 |
| 2000 ^e | 86,218 | 347 | 1 | 2 | 14,196 | 702 | 20 | 6.1 | 09/20–10/04 | 15 |
| 2001 ^f | 120,435 | 138 | 2 | 3 | 17,605 | 597 | 29 | 6.8 | 09/22–10/10, 10/23–11/08 | 36 |
| 2002 ^f | 96,221 | 181 | 2 | 2 | 14,528 | 277 | 52 | 6.6 | 09/23–10/09, 10/17–10/27 | 27 |
| 2003 ^{e,g} | 33,817 | 143 | 1 | 1 | 5,327 | 584 | 9 | 6.4 | 09/01–10/04 | 34 |
| 2004 ^f | 201,579 | 638 | 2 | 3 | 29,733 | 1,286 | 23 | 6.8 | 10/21–10/25, 10/23–10/31, 10/27–11/01 | 20 |
| 2005 ^f | 208,828 | 1,500 | 1 | 4 | 30,585 | 1,376 | 22 | 6.8 | 11/12–12/02 | 19 |
| 2006 ^f | 303,867 | 3,313 | 1 | 4 | 47,215 | 1,067 | 44 | 6.4 | 09/23–10/23 | 31 |
| 2007 ^f | 145,619 | 469 | 1 | 4 | 22,951 | 734 | 31 | 6.3 | 10/02–10/23 | 22 |
| 2008 | | | | | | No Cost-Recovery Effort | | | | |
| 2009 ^f | 100,400 | 463 | 1 | 3 | 15,726 | 646 | 24 | 6.4 | 09/25–10/12 | 18 |
| 2010 ^f | 72,787 | 69 | 1 | 3 | 11,462 | 556 | 21 | 6.4 | 09/27–10/20 | 25 |
| 2011 ^f | 118,690 | 199 | 1 | 3 | 18,963 | 618 | 31 | 6.3 | 09/30–10/21 | 21 |
| 2012 ^f | 134,712 | 286 | 3 | 7 | 18,388 | 726 | 25 | 7.3 | 10/09–10/22 | 14 |

-continued-

Table 5.–Page 2 of 2.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Charter Length | |
|-------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|---------------------------------------|------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Dates | Days |
| 2013 ^f | 198,158 | 1,332 | 1 | 3 | 29,568 | 662 | 45 | 6.7 | 10/01–10/08, 10/12–10/17 | 14 |
| 2014 ^f | 190,269 | 940 | 1 | 3 | 27,044 | 665 | 41 | 7.0 | 10/02–10/21 | 20 |
| 2015 ^f | 201,471 | 1,143 | 1 | 3 | 29,191 | 755 | 39 | 6.9 | 10/01–10/11, 10/13–10/22 | 21 |
| 2016 ^f | 183,948 | 1,107 | 1 | 3 | 27,227 | 482 | 56 | 6.8 | 10/1–10/10, 10/12–10/19 | 18 |
| 2017 ^f | 219,509 | 3,562 | 1 | 4 | 32,621 | 927 | 35 | 6.7 | 10/01–10/09, 10/11–10/23, 11/02–11/08 | 29 |
| 2018 ^f | 159,630 | 2,720 | 1 | 4 | 23,486 | 1,253 | 19 | 6.8 | 10/02–10/10, 10/11–10/16, 10/19–10/22 | 19 |
| 2019 ^f | 122,372 | 1,096 | 1 | 2 | 17,187 | 575 | 30 | 7.1 | 10/01–10/13 | 13 |
| 2020 ^f | 124,289 | 789 | 1 | 3 | 21,780 | 1,102 | 20 | 5.7 | 10/07–10/24 | 18 |

^a In pounds.^b Deadloss included.^c Number of legal crab per pot lift.^d Retained catch.^e Bering Sea and Aleutian Islands shellfish research program cost recovery.^f Bering Sea and Aleutian Islands shellfish research and observer program cost recovery.^g Includes 1,222 pounds harvested in the Pribilof District.

Table 6.—Saint Matthew Island Section blue king commercial fishery harvest data, 1977–2020/21.

| Season | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------------------------|------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 1977 | — | 1,202,066 | 129,148 | 10 | 24 | 281,665 | 17,370 | 16 | 4.3 | 130 |
| 1978 | — | 1,984,251 | 116,037 | 22 | 70 | 436,126 | 43,754 | 9 | 4.5 | 132 |
| 1979 | — | 210,819 | 56,147 | 18 | 25 | 52,966 | 9,877 | 5 | 4.0 | 129 |
| 1980 | — | CF | CF | 2 | 2 | CF | CF | CF | CF | CF |
| 1981 | — | 4,627,761 | 53,355 | 31 | 119 | 1,045,619 | 58,550 | 18 | 4.4 | ND |
| 1982 | — | 8,844,789 | 142,973 | 96 | 269 | 1,935,886 | 165,618 | 12 | 4.6 | 135 |
| 1983 | 8,000,000 | 9,454,323 | 828,994 | 164 | 235 | 1,931,990 | 133,944 | 14 | 4.8 | 137 |
| 1984 | 2–4 million | 3,764,592 | 31,983 | 90 | 169 | 841,017 | 73,320 | 11 | 4.5 | 135 |
| 1985 | 0.9–1.9 million | 2,175,087 | 2,613 | 74 | 95 | 436,021 | 46,988 | 9 | 5.0 | 139 |
| 1986 | 0.2–0.5 million | 1,003,162 | 32,560 | 38 | 43 | 219,548 | 22,073 | 10 | 4.6 | 134 |
| 1987 | 0.6–1.3 million | 1,039,779 | 600 | 60 | 61 | 227,447 | 28,230 | 8 | 4.6 | 134 |
| 1988 | 0.7–1.5 million | 1,236,462 | 7,760 | 45 | 45 | 280,401 | 21,678 | 13 | 4.4 | 133 |
| 1989 | 1,700,000 | 1,166,258 | 3,754 | 69 | 69 | 247,641 | 30,803 | 8 | 4.7 | 135 |
| 1990 | 1,900,000 | 1,725,349 | 17,416 | 31 | 38 | 391,405 | 26,264 | 15 | 4.4 | 134 |
| 1991 | 3,200,000 | 3,372,066 | 216,459 | 68 | 69 | 726,519 | 37,104 | 20 | 4.6 | 134 |
| 1992 | 3,100,000 | 2,475,916 | 1,836 | 174 | 179 | 545,222 | 56,630 | 10 | 4.5 | 134 |
| 1993 | 4,400,000 | 3,003,089 | 3,168 | 92 | 136 | 630,353 | 58,647 | 11 | 4.8 | 135 |
| 1994 | 3,000,000 | 3,764,262 | 46,699 | 87 | 133 | 827,015 | 60,860 | 14 | 4.6 | 133 |
| 1995 | 2,400,000 | 3,166,093 | 91,041 | 90 | 111 | 666,905 | 48,560 | 14 | 4.7 | 135 |
| 1996 | 4,300,000 | 3,078,959 | 36,892 | 122 | 188 | 660,665 | 91,085 | 7 | 4.7 | 135 |
| 1997 | 5,000,000 | 4,649,660 | 209,490 | 117 | 166 | 939,822 | 81,117 | 12 | 4.9 | 140 |
| 1998 | 4,099,512 | 2,968,573 | 15,554 | 132 | 266 | 635,370 | 91,826 | 7 | 4.7 | 134 |
| 1999–2008/09 ^g | | | | No Commercial Fishery | | | | | | |

-continued-

Table 6.–Page 2 of 2.

| Season | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|-----------------|------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 2009/10 | 1,167,000 | 460,859 | 10,484 | 7 | 30 | 103,376 | 10,697 | 10 | 4.5 | 130 |
| 2010/11 | 1,600,000 | 1,263,982 | 10,206 | 11 | 70 | 298,668 | 29,346 | 10 | 4.2 | 123 |
| 2011/12 | 2,359,000 | 1,881,322 | 26,588 | 18 | 90 | 437,862 | 48,554 | 9 | 4.3 | 126 |
| 2012/13 | 1,630,000 | 1,616,054 | 21,052 | 17 | 92 | 379,386 | 37,065 | 10 | 4.3 | 130 |
| 2013/14 | | | | No Commercial Fishery | | | | | | |
| 2014/15 | 655,000 | 308,582 | 5,552 | 4 | 26 | 69,109 | 10,133 | 7 | 4.5 | 132 |
| 2015/16 | 411,000 | 106,449 | 1,439 | 3 | 14 | 24,407 | 5,475 | 4 | 4.4 | 132 |
| 2016/17–2020/21 | | | | No Commercial Fishery | | | | | | |

Note: En dashes indicate harvest limits were not set for this fishery prior to 1983; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Guideline harvest level (GHL) began in 1983; total allowable catch (TAC) began in 2005/06.

^c Deadloss included.

^d Number of legal crab per pot lift.

^e Retained catch.

^f Carapace length in millimeters.

^g Crab rationalization begins.

Table 7.—Saint Matthew Island Section blue king crab commercial fishery value and season dates, 1977–2020/21.

| Season | Value | | Season length | | |
|---------------------------|-----------------------|--------------|---------------|----------|------|
| | Exvessel ^a | Total | Opened | Closed | Days |
| 1977 | \$1.00 | \$1,072,918 | 06/07/77 | 08/16/77 | 71 |
| 1978 | \$0.95 | \$1,774,803 | 07/15/78 | 09/03/78 | 51 |
| 1979 | \$0.70 | \$108,270 | 07/15/79 | 08/24/79 | 41 |
| 1980 | CF | CF | 07/15/80 | 09/03/80 | 51 |
| 1981 | \$0.90 | \$4,116,965 | 07/15/81 | 08/21/81 | 38 |
| 1982 | \$2.00 | \$17,403,632 | 08/01/82 | 08/16/82 | 16 |
| 1983 ^b | \$3.00 | \$25,875,987 | 08/23/83 | 09/06/83 | 15 |
| 1984 | \$1.50 | \$5,598,914 | 09/01/84 | 09/08/84 | 8 |
| 1985 | \$1.60 | \$3,475,958 | 09/01/85 | 09/06/85 | 6 |
| 1986 | \$2.96 | \$2,870,927 | 09/01/86 | 09/06/86 | 6 |
| 1987 | \$2.73 | \$2,833,149 | 09/01/87 | 09/05/87 | 5 |
| 1988 | \$3.10 | \$3,811,700 | 09/01/88 | 09/05/88 | 5 |
| 1989 | \$2.90 | \$3,371,233 | 09/01/89 | 09/04/89 | 4 |
| 1990 | \$3.42 | \$5,844,906 | 09/01/90 | 09/07/90 | 7 |
| 1991 | \$2.80 | \$8,835,700 | 09/16/91 | 09/20/91 | 5 |
| 1992 | \$2.92 | \$7,228,333 | 09/04/92 | 09/07/92 | 4 |
| 1993 | \$3.05 | \$9,159,080 | 09/15/93 | 09/21/93 | 7 |
| 1994 | \$4.13 | \$15,338,950 | 09/15/94 | 09/22/94 | 8 |
| 1995 | \$2.32 | \$7,123,102 | 09/15/95 | 09/22/95 | 8 |
| 1996 | \$2.21 | \$6,712,037 | 09/15/96 | 09/16/96 | 2 |
| 1997 | \$2.23 | \$9,900,019 | 09/15/97 | 09/22/97 | 8 |
| 1998 | \$1.87 | \$5,523,187 | 09/15/98 | 09/26/98 | 12 |
| 1999–2008/09 ^b | No Commercial Fishery | | | | |
| 2009/10 | \$2.19 | \$985,204 | 10/15/09 | 02/01/10 | 110 |
| 2010/11 | \$4.12 | \$5,160,670 | 10/15/10 | 02/01/11 | 110 |
| 2011/12 | \$4.33 | \$8,034,387 | 10/15/11 | 02/01/12 | 110 |
| 2012/13 | \$3.77 | \$6,008,470 | 10/15/12 | 02/01/13 | 110 |
| 2013/14 | No Commercial Fishery | | | | |
| 2014/15 | \$3.38 | \$1,025,161 | 10/15/14 | 02/01/15 | 110 |
| 2015/16 | \$4.03 | \$423,508 | 10/15/15 | 02/01/16 | 110 |
| 2016/17–2020/21 | No Commercial Fishery | | | | |

Note: CF indicates confidential data

^a Average price per pound.

^b Crab rationalization begins.

Table 8.—Pribilof District golden king crab commercial fishery harvest data, 1981/82–2020.

| Season | GHL ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|-----------|--------------------|------------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 1981/82 | — | CF | CF | 2 | 3 | CF | CF | CF | CF | CF |
| 1982/83 | — | 69,970 | 570 | 10 | 19 | 15,330 | 5,252 | 3 | 4.6 | 151 |
| 1983/84 | — | 856,475 | 20,041 | 50 | 115 | 253,162 | 26,035 | 10 | 3.4 | 127 |
| 1984 | — | No Commercial Fishing Effort | | | | | | | | |
| 1985 | — | CF | CF | 1 | 1 | CF | CF | CF | CF | CF |
| 1986 | — | No Commercial Fishing Effort | | | | | | | | |
| 1987 | — | CF | CF | 2 | 2 | CF | CF | CF | CF | CF |
| 1988 | — | CF | CF | 1 | 2 | CF | CF | CF | CF | CF |
| 1989 | — | CF | CF | 2 | 4 | CF | CF | CF | CF | CF |
| 1990–1992 | — | No Commercial Fishing Effort | | | | | | | | |
| 1993 | — | 67,458 | | 5 | 15 | 17,643 | 15,395 | 1 | 3.8 | ND |
| 1994 | — | 88,985 | 730 | 3 | 5 | 21,477 | 1,845 | 12 | 4.1 | ND |
| 1995 | — | 341,908 | 2950 | 7 | 23 | 82,489 | 9,551 | 9 | 4.1 | ND |
| 1996 | — | 329,009 | 12,409 | 6 | 32 | 91,947 | 9,952 | 9 | 3.6 | ND |
| 1997 | — | 179,249 | 5,554 | 7 | 23 | 43,305 | 4,673 | 9 | 4.1 | ND |
| 1998 | — | 35,722 | 474 | 3 | 9 | 9,205 | 1,530 | 6 | 3.9 | ND |
| 1999 | 200,000 | 177,108 | 319 | 3 | 9 | 44,098 | 2,995 | 15 | 4.0 | ND |
| 2000 | 150,000 | 127,217 | 4,599 | 7 | 19 | 29,145 | 5,450 | 5 | 4.4 | ND |
| 2001 | 150,000 | 145,876 | 8,227 | 6 | 14 | 33,723 | 4,262 | 8 | 4.3 | 143 |
| 2002 | 150,000 | 150,434 | 8,984 | 8 | 20 | 34,860 | 5,279 | 7 | 4.3 | 144 |
| 2003 | 150,000 | CF | CF | 3 | 6 | CF | CF | CF | CF | CF |
| 2004 | 150,000 | CF | CF | 5 | 8 | CF | CF | CF | CF | CF |
| 2005 | 150,000 | CF | CF | 4 | 8 | CF | CF | CF | CF | CF |

-continued-

Table 8.—Page 2 of 2.

| Season | GHL ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|-------------------|--------------------|------------------------|-----------------------|------------------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} | Length ^{e,f} |
| 2006–2009 | 150,000 | | | No Commercial Fishing Effort | | | | | | |
| 2010 | 150,000 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF |
| 2011 | 150,000 | CF | CF | 2 | 4 | CF | CF | CF | CF | CF |
| 2012 | 150,000 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF |
| 2013 | 150,000 | CF | CF | 1 | 2 | CF | CF | CF | CF | CF |
| 2014 | 150,000 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF |
| 2015–2016 | 130,000 | | | No Commercial Fishing Effort | | | | | | |
| 2017 ^g | 130,000 | CF | CF | 2 | 6 | CF | CF | CF | CF | CF |
| 2018 | 130,000 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF |
| 2019 | 130,000 | CF | CF | 2 | 5 | CF | CF | CF | CF | CF |
| 2020 | 130,000 | 107,679 | 1,638 | 4 | 12 | 24,301 | 2,960 | 8 | 4.4 | 149 |

Note: En dashes indicate harvest limits were not set for this fishery prior to 1999; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Guideline harvest level (GHL) began in 1999.

^c Deadloss included.

^d Number of legal crab per pot lift.

^e Retained catch.

^f Carapace length in millimeters.

^g Includes harvest from one vessel that was incorrectly landed using a T91Q CFEC permit card.

Table 9.—Pribilof District golden king crab commercial fishery value and season dates, 1991–2020.

| Season | Value | | Season length | | |
|--------|------------------------------|-----------|---------------|----------|------|
| | Exvessel ^a | Total | Opened | Closed | Days |
| 1993 | \$2.58 | \$173,975 | 01/01/93 | 12/31/93 | 365 |
| 1994 | \$3.97 | \$350,375 | 01/01/94 | 12/31/94 | 365 |
| 1995 | \$2.82 | \$955,522 | 01/01/95 | 12/31/95 | 365 |
| 1996 | \$2.13 | \$674,660 | 01/01/96 | 12/31/96 | 365 |
| 1997 | \$2.25 | \$391,339 | 01/01/97 | 12/31/97 | 365 |
| 1998 | \$1.96 | \$68,930 | 01/01/98 | 12/31/98 | 365 |
| 1999 | \$2.39 | \$421,674 | 01/01/99 | 06/10/99 | 161 |
| 2000 | \$3.22 | \$395,325 | 01/01/00 | 12/31/00 | 365 |
| 2001 | \$3.15 | \$433,922 | 01/01/01 | 04/15/01 | 105 |
| 2002 | \$3.09 | \$436,584 | 01/01/02 | 05/14/02 | 134 |
| 2003 | CF | CF | 01/01/03 | 05/01/03 | 121 |
| 2004 | CF | CF | 01/01/04 | 03/12/04 | 72 |
| 2005 | CF | CF | 01/01/05 | 12/31/05 | 365 |
| 2006 | No Commercial Fishing Effort | | 01/01/06 | 12/31/06 | 365 |
| 2007 | No Commercial Fishing Effort | | 01/01/07 | 12/31/07 | 365 |
| 2008 | No Commercial Fishing Effort | | 01/01/08 | 12/31/08 | 365 |
| 2009 | No Commercial Fishing Effort | | 01/01/09 | 12/31/09 | 365 |
| 2010 | CF | CF | 01/01/10 | 12/31/10 | 365 |
| 2011 | CF | CF | 01/01/11 | 12/31/11 | 365 |
| 2012 | CF | CF | 01/01/12 | 12/31/12 | 365 |
| 2013 | CF | CF | 01/01/13 | 12/31/13 | 365 |
| 2014 | CF | CF | 01/01/14 | 12/31/14 | 365 |
| 2015 | No Commercial Fishing Effort | | 01/01/15 | 12/31/15 | 365 |
| 2016 | No Commercial Fishing Effort | | 01/01/16 | 12/31/16 | 365 |
| 2017 | CF | CF | 01/01/17 | 06/12/17 | 163 |
| 2018 | CF | CF | 01/01/18 | 12/31/18 | 365 |
| 2019 | CF | CF | 01/01/19 | 05/31/19 | 151 |
| 2020 | \$5.87 | \$621,971 | 01/01/20 | 12/31/20 | 365 |

Note: CF indicates confidential data.

^a Average price per pound.

Table 10.—Pribilof District golden king crab commercial fishery harvest and effort by statistical area, 2020.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 695530 | 13,887 | 183 | 3 | 5 | 3,122 | 409 | 8 | 4.5 |
| 695600 | 27,686 | 513 | 4 | 10 | 5,976 | 1,271 | 5 | 4.6 |
| 705530 | 24,591 | 259 | 3 | 6 | 5,696 | 415 | 14 | 4.3 |
| 705600 | 35,862 | 553 | 3 | 7 | 8,220 | 741 | 11 | 4.4 |
| Other ^e | 5,654 | 1,287 | 2 | 2 | 1,287 | 124 | 10 | 4.4 |
| Total | 107,679 | 1,638 | 4 | 12 | 24,301 | 2,960 | 8 | 4.4 |

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Combination of statistical areas (2) in which landings were made by fewer than three vessels.

Table 11.—Bering Sea scarlet king crab commercial fishery harvest data, 1995–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|---------------------|------------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|----------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 1995 | 26,684 | 465 | 4 | 25 | 11,048 | 24,551 | <1 | 2.4 | \$2.45 | \$64,237 |
| 1996 | CF | CF | 2 | 7 | CF | CF | CF | CF | CF | CF |
| 1997–2001 | No Commercial Fishing Effort | | | | | | | | | |
| 2002 | CF | CF | 1 | 4 | CF | CF | CF | CF | CF | CF |
| 2003 ^f | CF | CF | 3 | 11 | CF | CF | CF | CF | CF | CF |
| 2004 ^f | CF | CF | 2 | 4 | CF | CF | CF | CF | CF | CF |
| 2005 ^f | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2006–2016 | No Commercial Fishing Effort | | | | | | | | | |
| 2017 ^{f,g} | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2018–2019 | No Commercial Fishing Effort | | | | | | | | | |
| 2020 ^f | CF | CF | 1 | 2 | CF | CF | CF | CF | CF | CF |

Note: CF indicates confidential data. Harvest is bycatch only; therefore, harvest limits are not set for this stock.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Average price per pound.

^f Restricted to incidental harvest during Bering Sea golden king crab and grooved Tanner crab fisheries.

^g Includes harvest from one vessel that was incorrectly landed using a T91Q CFEC permit card.

Table 12.—Bering Sea District Tanner crab commercial fishery harvest data, 1968–2020/21.

| Season | Location ^a | GHL/TAC ^{b,c} | Harvest ^{b,d} | Deadloss ^b | Number of | | | | Average | | |
|---------|-----------------------|------------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | | Vessels | Landings | Crab ^d | Pots lifted | CPUE ^e | Weight ^{b,f} | Width ^{f,g} |
| 1968 | | — | 17,900 | ND | ND | 7 | 6,400 | 1,400 | 5 | 2.8 | ND |
| 1969 | | — | 1,008,900 | ND | ND | 131 | 353,300 | 29,800 | 12 | 2.9 | ND |
| 1970 | | — | 1,014,700 | ND | ND | 66 | 482,300 | 16,400 | 29 | 2.9 | ND |
| 1971 | | — | 166,100 | ND | ND | 22 | 61,300 | 7,300 | 8 | 2.7 | ND |
| 1972 | | — | 107,761 | ND | ND | 14 | 42,061 | 4,260 | 10 | 2.6 | ND |
| 1973 | | — | 231,668 | ND | ND | 44 | 93,595 | 15,730 | 6 | 2.5 | ND |
| 1974 | | — | 5,044,197 | ND | ND | 69 | 2,531,825 | 22,014 | 115 | 2.0 | ND |
| 1974/75 | Southeastern | — | 6,504,984 | ND | ND | 72 | 2,526,687 | 32,275 | 78 | 2.6 | ND |
| | Pribilofs | — | 523,394 | ND | ND | 8 | 247,083 | 3,923 | 63 | 2.1 | ND |
| | TOTAL | — | 7,028,378 | ND | 28 | 80 | 2,773,770 | 36,198 | 77 | 2.5 | ND |
| 1975/76 | Southeastern | — | 16,643,194 | ND | ND | 230 | 6,682,232 | 106,445 | 63 | 2.5 | ND |
| | Pribilofs | — | 5,714,913 | ND | ND | 74 | 2,273,804 | 34,761 | 65 | 2.5 | ND |
| | TOTAL | — | 22,358,107 | ND | 66 | 304 | 8,956,036 | 141,206 | 63 | 2.5 | ND |
| 1976/77 | Southeastern | — | 41,007,736 | ND | ND | 437 | 16,089,057 | 233,667 | 69 | 2.6 | ND |
| | Pribilofs | — | 10,447,485 | ND | ND | 104 | 4,162,451 | 63,804 | 65 | 2.5 | ND |
| | TOTAL | — | 51,455,221 | ND | 83 | 541 | 20,251,508 | 297,471 | 68 | 2.5 | ND |
| 1977/78 | Southeastern | — | 53,278,012 | ND | ND | 706 | 21,055,527 | 408,437 | 52 | 2.5 | ND |
| | Pribilofs | — | 13,152,843 | ND | ND | 155 | 5,210,170 | 107,913 | 48 | 2.5 | ND |
| | TOTAL | — | 66,430,855 | ND | 120 | 861 | 26,265,697 | 516,350 | 51 | 2.5 | ND |
| 1978/79 | Southeastern | — | 39,694,205 | 75,400 | ND | 758 | 15,601,891 | 356,594 | 44 | 2.5 | ND |
| | Pribilofs | — | 2,852,969 | 600 | ND | 59 | 1,124,627 | 46,103 | 24 | 2.5 | ND |
| | TOTAL | — | 42,547,174 | 76,000 | 144 | 817 | 16,726,518 | 402,697 | 42 | 2.5 | 153 |

-continued-

Table 12.—Page 2 of 6.

| Season | Location ^a | GHL/TAC ^{b,c} | Harvest ^{b,d} | Deadloss ^b | Number of | | | | Average | | |
|-----------|-----------------------|------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | | Vessels | Landings | Crab ^d | Pots lifted | CPUE ^e | Weight ^{b,f} | Width ^{f,g} |
| 1979/80 | Southeastern | — | 35,724,003 | 56,446 | ND | 789 | 14,329,889 | 476,410 | 30 | 2.5 | ND |
| | Pribilofs | — | 890,312 | 0 | ND | 15 | 355,722 | 12,024 | 30 | 2.5 | ND |
| | TOTAL | 28–36 million | 36,614,315 | 56,446 | 152 | 804 | 14,685,611 | 488,434 | 30 | 2.5 | 151 |
| 1981 | Southeastern | — | 26,684,956 | 97,398 | ND | 674 | 10,532,007 | 496,751 | 21 | 2.5 | ND |
| | Pribilofs | — | 2,945,536 | 4,196 | ND | 87 | 1,313,951 | 62,875 | 21 | 2.2 | ND |
| | TOTAL | 28–36 million | 29,630,492 | 101,594 | 165 | 761 | 11,845,958 | 559,626 | 21 | 2.5 | 149 |
| 1982 | Southeastern | — | 8,812,302 | 69,829 | ND | 539 | 3,825,433 | 322,634 | 12 | 2.3 | ND |
| | Pribilofs | — | 2,196,477 | 68,330 | ND | 252 | 1,005,547 | 167,465 | 6 | 2.2 | ND |
| | TOTAL | 12–16 million | 11,008,779 | 138,159 | 125 | 791 | 4,830,980 | 490,099 | 10 | 2.3 | 149 |
| 1983 | Northern | — | 48,454 | 167 | ND | 10 | 29,578 | 5,950 | 5 | 1.6 | ND |
| | Southeastern | — | 4,633,354 | 52,879 | ND | 287 | 1,984,673 | 192,538 | 10 | 2.3 | ND |
| | Pribilofs | — | 592,073 | 6,983 | ND | 151 | 272,505 | 83,528 | 3 | 2.2 | ND |
| | TOTAL | 5,600,000 | 5,273,881 | 60,029 | 108 | 448 | 2,286,756 | 282,016 | 8 | 2.3 | 149 |
| 1984 | Southeastern | — | 1,099,142 | 4,688 | ND | 91 | 470,181 | 44,546 | 11 | 2.3 | ND |
| | Pribilofs | — | 109,081 | 337 | ND | 43 | 46,759 | 16,811 | 3 | 2.3 | ND |
| | TOTAL | 7,100,000 | 1,208,223 | 5,025 | 41 | 134 | 516,940 | 61,357 | 8 | 2.3 | 147 |
| 1985 | Southeastern | — | 3,023,193 | 14,096 | 38 | 143 | 1,266,567 | 85,926 | 13 | 2.4 | ND |
| | Pribilofs | — | 13,742 | 0 | 15 | 23 | 5,934 | 8,606 | 1 | 2.3 | ND |
| | TOTAL | 3,000,000 | 3,036,935 | 14,096 | 44 | 166 | 1,272,501 | 94,532 | 12 | 2.4 | 150 |
| 1986–1987 | | | | | No Commercial Fishery | | | | | | |

-continued-

Table 12.—Page 3 of 6.

| Season | Location ^a | GHL/TAC ^{b,c} | Harvest ^{b,d} | Deadloss ^b | Number of | | | | Average | | |
|---------|-----------------------|------------------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | | Vessels | Landings | Crab ^d | Pots lifted | CPUE ^e | Weight ^{b,f} | Width ^{f,g} |
| 1988 | Eastern | — | 2,294,997 | 10,724 | 102 | 249 | 957,318 | 114,384 | 8 | 2.4 | ND |
| | Western | No Commercial Fishing Effort | | | | | | | | | |
| | TOTAL | 5,600,000 | 2,294,997 | 10,724 | 102 | 249 | 957,318 | 114,384 | 8 | 2.4 | 144 |
| 1989 | Eastern | — | 6,982,865 | 34,664 | 108 | 358 | 2,894,480 | 183,692 | 16 | 2.4 | ND |
| | Western | No Commercial Fishing Effort | | | | | | | | | |
| | TOTAL | 13,500,000 | 6,982,865 | 34,664 | 108 | 358 | 2,894,480 | 183,692 | 16 | 2.4 | 149 |
| 1990 | Eastern | — | 24,529,592 | 86,603 | 184 | 1,016 | 10,701,363 | 695,667 | 15 | 2.3 | ND |
| | Western | — | 17,956 | 0 | 15 | 18 | 7,975 | 9,548 | 1 | 2.3 | ND |
| | TOTAL | 29,500,000 | 24,547,548 | 86,603 | 184 | 1,034 | 10,709,338 | 705,215 | 15 | 2.3 | 148 |
| 1990/91 | Eastern | 42,800,000 | 40,081,555 | 210,769 | 248 | 1,756 | 16,608,625 | 883,391 | 19 | 2.4 | 150 |
| 1991/92 | Eastern | — | 31,742,117 | 279,741 | 284 | 2,316 | 12,901,364 | 1,209,180 | 11 | 2.5 | ND |
| | Western | — | 52,265 | 0 | 19 | 33 | 22,738 | 15,779 | 1 | 2.3 | ND |
| | TOTAL | 32,800,000 | 31,794,382 | 279,741 | 284 | 2,349 | 12,924,102 | 1,224,959 | 11 | 2.5 | 150 |
| 1992/93 | Eastern | — | 34,821,009 | 346,505 | 293 | 2,010 | 15,074,069 | 1,151,849 | 13 | 2.3 | ND |
| | Western | — | 309,823 | 3,000 | 70 | 96 | 191,796 | 50,051 | 4 | 1.6 | ND |
| | TOTAL | 38,100,000 ^h | 35,130,832 | 349,505 | 294 | 2,106 | 15,265,865 | 1,201,900 | 13 | 2.3 | 148 |
| 1993/94 | East of 168°W | 10,700,000 ⁱ | 4,134,529 | 119,715 | 285 | 350 | 1,699,750 | 250,826 | 7 | 2.4 | ND |
| | 163°W to 173°W | 9,100,000 ^{i,k} | 12,776,371 | 155,557 | 261 | 515 | 5,539,068 | 325,963 | 17 | 2.3 | ND |
| | TOTAL | 19,800,000 | 16,910,900 | 275,272 | 296 | 865 | 7,238,818 | 576,789 | 13 | 2.3 | 151 |
| 1994 | 163°W to 173°W | 7,500,000 | 7,766,886 | 132,780 | 183 | 349 | 3,351,639 | 249,536 | 13 | 2.3 | 150 |
| 1995 | 163°W to 173°W | 5,500,000 | 4,233,061 | 44,523 | 196 | 256 | 1,877,303 | 247,853 | 8 | 2.3 | 149 |

-continued-

Table 12.–Page 4 of 6.

| Season | Location ^a | GHL/TAC ^{b,c} | Harvest ^{b,d} | Deadloss ^b | Number of | | | | Average | | |
|----------------------|-----------------------|---------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | | Vessels | Landings | Crab ^d | Pots lifted | CPUE ^e | Weight ^{b,f} | Width ^{f,g} |
| 1996 | East of 168°W | 2,200,000 ^{g(i)} | 811,301 | 6,149 | 135 | 152 | 341,039 | 73,522 | 5 | 2.4 | ND |
| | 163°W to 173°W | 6,200,000 ^{h(i)} | 994,776 | 8,464 | 192 | 195 | 393,257 | 75,753 | 5 | 2.5 | ND |
| | TOTAL ^l | 8,400,000 | 1,806,077 | 14,613 | 196 | 347 | 734,296 | 149,275 | 5 | 2.5 | 152 |
| 1997–2004 | | | | | No Commercial Fishery | | | | | | |
| 2005/06 ^m | East of 166°W | | | | No Commercial Fishery | | | | | | |
| | West of 166°W | 1,620,000 | 952,887 | 15,174 | 43 | 103 | 443,977 | 32,389 | 14 | 2.1 | 145 |
| | TOTAL ^l | 1,620,000 | 952,887 | 15,174 | 43 | 103 | 443,977 | 32,389 | 14 | 2.1 | 145 |
| 2006/07 | East of 166°W | 1,875,000 | 1,401,743 | 9,256 | 37 | 63 | 585,480 | 29,129 | 20 | 2.4 | 151 |
| | West of 166°W | 1,094,000 | 720,846 | 19,696 | 39 | 78 | 340,623 | 28,140 | 12 | 2.1 | 146 |
| | TOTAL ^l | 2,969,000 | 2,122,589 | 28,952 | 52 | 141 | 926,103 | 57,269 | 16 | 2.3 | 150 |
| 2007/08 | East of 166°W | 3,445,000 | 1,582,858 | 16,117 | 20 | 65 | 685,491 | 33,515 | 20 | 2.3 | 148 |
| | West of 166°W | 2,176,000 | 523,796 | 4,676 | 34 | 61 | 241,673 | 21,938 | 11 | 2.2 | 146 |
| | TOTAL ^l | 5,621,000 | 2,106,654 | 20,793 | 41 | 126 | 927,164 | 55,453 | 17 | 2.3 | 148 |
| 2008/09 | East of 166°W | 2,763,000 | 1,830,031 | 13,543 | 22 | 66 | 778,898 | 36,698 | 21 | 2.3 | 150 |
| | West of 166°W | 1,537,000 | 109,552 | 3,737 | 42 | 101 | 51,471 | 30,175 | 2 | 2.1 | 147 |
| | TOTAL ^l | 4,300,000 | 1,939,583 | 17,280 | 50 | 167 | 830,369 | 66,873 | 12 | 2.3 | 149 |
| 2009/10 | East of 166°W | 1,350,000 | 1,324,578 | 8,376 | 17 | 51 | 483,419 | 16,770 | 29 | 2.7 | 157 |
| | West of 166°W | | | | No Commercial Fishery | | | | | | |
| | TOTAL ^l | 1,350,000 | 1,328,356 | 12,154 | 41 | 113 | 485,963 | 42,006 | 12 | 2.7 | 157 |
| 2010/11–2012/13 | | | | | No Commercial Fishery | | | | | | |

-continued-

Table 12.—Page 5 of 6.

| Season | Location ^a | GHL/TAC ^{b,c} | Harvest ^{b,d} | Deadloss ^b | Number of | | | | Average | | |
|---------|-----------------------|------------------------|------------------------|-----------------------|-----------------------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | | Vessels | Landings | Crab ^d | Pots lifted | CPUE ^e | Weight ^{b,f} | Width ^{f,g} |
| 2013/14 | East of 166°W | 1,463,000 | 1,456,357 | 6,254 | 30 | 74 | 710,043 | 26,468 | 27 | 2.1 | 145 |
| | West of 166°W | 1,645,000 | 1,330,488 | 22,546 | 64 | 261 | 735,725 | 131,524 | 6 | 1.8 | 138 |
| | TOTAL ¹ | 3,108,000 | 2,786,845 | 28,800 | 66 | 335 | 1,445,768 | 157,992 | 9 | 1.9 | 142 |
| 2014/15 | East of 166°W | 8,480,000 | 8,450,485 | 59,788 | 42 | 143 | 4,381,890 | 87,875 | 50 | 1.9 | 141 |
| | West of 166°W | 6,625,000 | 5,253,942 | 96,921 | 58 | 237 | 3,140,954 | 142,820 | 22 | 1.7 | 134 |
| | TOTAL ¹ | 15,105,000 | 13,704,427 | 156,709 | 64 | 380 | 7,522,844 | 230,695 | 33 | 1.8 | 138 |
| 2015/16 | East of 166°W | 11,272,000 | 11,263,562 | 120,187 | 49 | 202 | 6,000,262 | 139,171 | 43 | 1.9 | 141 |
| | West of 166°W | 8,396,000 | 8,378,816 | 52,546 | 62 | 247 | 4,856,156 | 145,638 | 33 | 1.7 | 135 |
| | TOTAL ¹ | 19,668,000 | 19,642,378 | 172,733 | 70 | 449 | 10,856,418 | 284,809 | 38 | 1.8 | 138 |
| 2016/17 | | | | | No Commercial Fishery | | | | | | |
| 2017/18 | East of 166°W | | | | No Commercial Fishery | | | | | | |
| | West of 166°W | 2,500,200 | 2,496,734 | 16,212 | 34 | 91 | 1,340,230 | 29,903 | 45 | 1.9 | 139 |
| | TOTAL ¹ | 2,500,200 | 2,497,033 | 16,249 | 40 | 107 | 1,340,394 | 33,738 | 40 | 1.9 | 139 |
| 2018/19 | East of 166°W | | | | No Commercial Fishery | | | | | | |
| | West of 166°W | 2,439,000 | 2,441,201 | 40,450 | 36 | 101 | 1,380,990 | 41,922 | 33 | 1.8 | 137 |
| | TOTAL ¹ | 2,439,000 | 2,441,227 | 40,476 | 37 | 111 | 1,381,008 | 45,984 | 30 | 1.8 | 137 |
| 2019/20 | | | | | No Commercial Fishery | | | | | | |
| 2020/21 | East of 166°W | | | | No Commercial Fishery | | | | | | |
| | West of 166°W | 2,348,000 | 1,449,543 | 25,334 | 41 | 92 | 873,650 | 47,388 | 18 | 1.7 | 134 |
| | TOTAL ¹ | 2,348,000 | 1,449,545 | 25,336 | 41 | 93 | 873,651 | 48,028 | 18 | 1.7 | 134 |

-continued-

Table 12.–Page 6 of 6.

Note: En dashes indicate harvest limits were not set for these fisheries prior to 1979/80 and area-specific GHGs were not set prior to 1993/94; ND indicates these data were not collected or cannot be derived.

^a From 1974/75 through 1984/85, Bering Sea Tanner crab subdistricts were Southeastern, Pribilof, and Northern (includes the Norton Sound and General Sections). From 1987/88 through 1992/93, harvest subdistricts were divided east and west of 173°W long. From 1993/94 through 1996/97, fishery east of 168°W long is concurrent with the Bristol Bay red king crab fishery and the fishery from 163°W long to 173°W long is a directed Tanner crab fishery. From 2005/06 to current, the fishery is divided east and west of 166°W long, and harvest east of 163°W long is only allowed as incidental catch during the Bristol Bay red king crab fishery.

^b In pounds.

^c Guideline harvest level (GHL) and total allowable catch (TAC) began in 2005/06.

^d Deadloss included.

^e Number of legal crab per pot lift.

^f Retained catch.

^g Carapace width in millimeters.

^h Initial GHL announcement was 39.2 million pounds. GHL was later adjusted to 38.1 million pounds.

ⁱ GHL for waters east of 163°W long.

^j GHL for waters west of 163°W long.

^k Harvest concurrent with the Bristol Bay red king crab fishery was estimated to be well below the GHL and minimal harvest was thought to have occurred west of 163°W long; therefore, the GHL was adjusted to 16.1 million pounds for the directed Tanner crab fishery from 163° to 173°W long.

^l Bering Sea District totals include limited tanner crab harvest incidental to the Bering Sea Snow and Bristol Bay Red King crab fisheries. Includes incidental harvest, participation, and effort.

^m Crab rationalization begins.

Table 13.—Bering Sea District Tanner crab commercial fishery value and season dates, 1974–2020/21.

| Season | Value | | Season length | | | Location/Comments |
|-----------|-----------------------|--------------|-----------------------|----------|------|--------------------------|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 1974/75 | \$0.20 | \$1,405,676 | 07/29/74 | 06/15/75 | 322 | Southeastern Subdistrict |
| | | | | 06/15/75 | 322 | Pribilof Subdistrict |
| 1975/76 | \$0.19 | \$4,248,040 | 08/01/75 | 07/15/76 | 350 | Southeastern Subdistrict |
| | | | | 07/15/76 | 350 | Pribilof Subdistrict |
| 1976/77 | \$0.30 | \$15,436,566 | 08/01/76 | 07/15/77 | 349 | Southeastern Subdistrict |
| | | | | 07/15/77 | 349 | Pribilof Subdistrict |
| 1977/78 | \$0.38 | \$25,243,725 | 09/15/77 | 06/15/78 | 274 | Southeastern Subdistrict |
| | | | | 06/15/78 | 274 | Pribilof Subdistrict |
| 1978/79 | \$0.52 | \$22,085,010 | 11/01/78 | 05/24/79 | 205 | Southeastern Subdistrict |
| | | | | 05/24/79 | 205 | Pribilof Subdistrict |
| 1979/80 | \$0.52 | \$19,010,092 | 11/01/79 | 05/11/80 | 193 | Southeastern Subdistrict |
| | | | | 05/11/80 | 193 | Pribilof Subdistrict |
| 1981 | \$0.58 | \$17,126,761 | 01/15/81 | 05/07/81 | 113 | Southeastern Subdistrict |
| | | | | 05/07/81 | 113 | Pribilof Subdistrict |
| 1982 | \$1.06 | \$11,522,857 | 02/15/82 | 06/15/82 | 121 | Southeastern Subdistrict |
| | | | | 06/15/82 | 121 | Pribilof Subdistrict |
| 1983 | \$1.20 | \$6,256,622 | 02/15/83 | 06/15/83 | 121 | Northern Subdistrict |
| | | | | 05/22/83 | 97 | Southeastern Subdistrict |
| | | | | 05/22/83 | 97 | Pribilof Subdistrict |
| 1984 | \$0.95 | \$1,143,038 | 02/15/84 | 06/15/84 | 122 | Southeastern Subdistrict |
| | | | | 06/15/84 | 122 | Pribilof Subdistrict |
| 1985 | \$1.40 | \$4,231,975 | 01/15/85 | 06/15/85 | 152 | Southeastern Subdistrict |
| | | | | 06/15/85 | 152 | Pribilof Subdistrict |
| 1986–1987 | | | No Commercial Fishery | | | |
| 1988 | \$2.20 | \$5,029,284 | 01/15/88 | | | Eastern Subdistrict |
| | | | | 04/20/88 | 97 | East of 165°W long |
| | | | | 03/29/88 | 75 | West of 165°W long |
| | | | | 01/15/88 | 75 | Western Subdistrict |
| | | | | | | |
| 1989 | \$2.99 | \$20,772,065 | 01/15/89 | | | Eastern Subdistrict |
| | | | | 03/26/89 | 71 | East of 165°W long |
| | | | | 05/07/89 | 113 | West of 165°W long |
| | | | | 01/15/89 | 113 | Western Subdistrict |

-continued-

Table 13.—Page 2 of 3.

| Season | Value | | Season length | | | Location/Comments |
|----------------------|-----------------------|--------------|-----------------------|----------|------|---------------------------------|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 1990 | \$1.97 | \$48,253,834 | 01/15/90 | | | Eastern Subdistrict |
| | | | | 04/09/90 | 85 | East of 165°W long |
| | | | | 04/24/90 | 100 | West of 165°W long |
| | | | 01/15/90 | 04/24/90 | 100 | Western Subdistrict |
| 1990/91 | \$1.06 | \$42,323,199 | | | | Eastern Subdistrict |
| | | | 11/20/90 | 03/25/91 | 126 | East of 166°W long |
| | | | 11/20/90 | 03/25/91 | 126 | West of 166°W long |
| 1991/92 | \$1.84 | \$58,127,840 | | | | Eastern Subdistrict |
| | | | 11/15/91 | 03/31/92 | 138 | East of 166°W long |
| | | | 11/15/91 | 03/31/92 | 138 | West of 166°W long |
| 1992/93 | \$1.96 | \$68,296,684 | 11/15/92 | | | Eastern Subdistrict |
| | | | | 03/31/93 | 137 | East of 166°W long |
| | | | | 03/31/93 | 137 | West of 166°W long |
| | | | 11/15/92 | 03/31/93 | 137 | Western Subdistrict |
| 1993/94 | \$2.33 | \$38,832,369 | 11/01/93 | 11/10/93 | 10 | East of 168°W long ^b |
| | | | 11/20/93 | 01/01/94 | 43 | 163° to 173°W long |
| 1994 | \$4.13 | \$31,543,340 | 11/01/94 | 11/21/94 | 21 | 163° to 173°W long |
| 1995 | \$4.14 | \$17,348,234 | 11/01/95 | 11/16/95 | 16 | 163° to 173°W long |
| 1996 | \$2.50 | \$4,471,234 | 11/01/96 | 11/05/96 | 5 | East of 168°W long ^b |
| | | | 11/15/96 | 11/27/96 | 13 | 163° to 173°W long |
| 1997–2004 | | | No Commercial Fishery | | | |
| 2005/06 ^c | | | No Commercial Fishery | | | 163° to 166°W long ^d |
| | \$1.19 | \$1,118,129 | 10/15/05 | 03/31/06 | 168 | West of 166°W long |
| 2006/07 | \$1.31 | \$2,742,429 | 10/15/06 | 03/31/07 | 168 | 163° to 166°W long ^d |
| | | | | 03/31/07 | 168 | West of 166°W long |
| 2007/08 | \$1.52 | \$3,171,664 | 10/15/07 | 03/31/08 | 169 | 163° to 166°W long ^d |
| | | | | 03/31/08 | 169 | West of 166°W long |
| 2008/09 | \$1.49 | \$2,862,365 | 10/15/08 | 03/31/09 | 168 | 163° to 166°W long ^d |
| | | | | 03/31/09 | 168 | West of 166°W long |
| 2009/10 | \$1.64 | \$2,155,355 | 10/15/09 | 03/31/10 | 168 | 163° to 166°W long ^d |
| | | | No Commercial Fishery | | | West of 166°W long |
| 2010/11–2012/13 | | | No Commercial Fishery | | | |
| 2013/14 | \$2.29 | \$6,309,202 | 10/15/13 | 03/31/14 | 168 | 163° to 166°W long ^d |
| | | | | 03/31/14 | 168 | West of 166°W long |

-continued-

Table 13.—Page 3 of 3.

| Season | Value | | Season length | | | Location/Comments |
|---------|-----------------------|--------------|-----------------------|----------|------|---------------------------------|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 2014/15 | \$2.15 | \$29,073,491 | 10/15/14 | 03/31/15 | 168 | 163° to 166°W long ^d |
| | | | | 03/31/15 | 168 | West of 166°W long |
| 2015/16 | \$2.18 | \$42,432,982 | 10/15/15 | 03/31/16 | 169 | 163° to 166°W long ^d |
| | | | | 03/31/16 | 169 | West of 166°W long |
| 2016/17 | | | No Commercial Fishery | | | |
| 2017/18 | | | No Commercial Fishery | | | 163° to 166°W long ^d |
| | \$3.27 | \$8,102,596 | 10/15/17 | 03/31/18 | 168 | West of 166°W long |
| 2018/19 | | | No Commercial Fishery | | | 163° to 166°W long ^d |
| | \$3.30 | \$7,913,588 | 10/15/18 | 03/31/19 | 168 | West of 166°W long |
| 2019/20 | | | No Commercial Fishery | | | |
| 2020/21 | | | No Commercial Fishery | | | 163° to 166°W long ^d |
| | \$3.23 | \$4,676,509 | 10/15/20 | 03/31/21 | 168 | West of 166°W long |

^a Average price per pound.

^b Concurrent with Bristol Bay red king crab fishery.

^c Crab rationalization begins.

^d Directed fishery open between 163° and 166°W long. Incidental harvest allowed in entire area east of 166°W long during Bristol Bay red king crab fishery; however, no incidental harvest allowed when the directed fishery is closed.

Table 14.—Bering Sea District Tanner crab commercial fishery harvest and effort by week, west of 166° W long, 2020/21.

| Week ending | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| Oct-24 | CF | CF | 2 | 2 | CF | CF | CF | CF |
| Oct-31 | 340,735 | 3,152 | 17 | 19 | 198,494 | 11,476 | 17 | 1.7 |
| Nov-7 | 47,776 | 188 | 7 | 7 | 27,245 | 2,090 | 13 | 1.8 |
| Nov-14 | 117,264 | 535 | 3 | 4 | 68,615 | 2,094 | 33 | 1.7 |
| Nov-21 | CF | CF | 2 | 3 | CF | CF | CF | CF |
| ... ^c | | | | | | | | |
| Jan-9 | 38,344 | 338 | 3 | 3 | 22,040 | 2,532 | 9 | 2 |
| ... ^c | | | | | | | | |
| Jan-23 | 49,425 | 1,627 | 4 | 4 | 28,532 | 2,447 | 12 | 2 |
| Jan-30 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Feb-6 | 4,962 | 18 | 3 | 3 | 2,903 | 2,304 | 1 | 2 |
| Feb-13 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Feb-20 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Feb-27 | 225,105 | 1,918 | 9 | 10 | 135,738 | 5,225 | 26 | 1.7 |
| Mar-6 | 203,966 | 7,146 | 3 | 5 | 124,551 | 2,896 | 43 | 1.6 |
| Mar-13 | 126,241 | 6,483 | 9 | 10 | 87,540 | 6,274 | 14 | 1.4 |
| Mar-20 | CF | CF | 2 | 3 | CF | CF | CF | CF |
| Mar-27 | 46,877 | 1,396 | 4 | 5 | 30,924 | 2,254 | 14 | 1.5 |
| Apr-3 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| ... ^c | | | | | | | | |
| Apr-17 | CF | CF | 2 | 2 | CF | CF | CF | CF |
| Total | 1,449,543 | 25,334 | 41 | 84 | 873,650 | 47,388 | 18 | 1.7 |

-continued-

Table 14.—Page 2 of 2.

Note: CF indicates confidential data.

^a In pounds.

^b Deadloss included.

^c Number of retained crab per pot lift.

^d Retained catch.

^e Consecutive weeks with no harvest.

Table 15.—Bering Sea District Tanner crab commercial fishery harvest and effort by statistical area, west of 166° W long, 2020/21.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|---------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 665430 | 52,527 | 995 | 4 | 6 | 31,881 | 1,037 | 31 | 1.6 |
| 665500 | 296,192 | 4,365 | 13 | 25 | 178,746 | 5,042 | 35 | 1.7 |
| 665530 | 540,028 | 14,376 | 16 | 30 | 336,249 | 8,861 | 38 | 1.6 |
| 665600 | 17,772 | 189 | 10 | 11 | 10,562 | 1,067 | 10 | 1.7 |
| 665630 | 1,710 | 34 | 3 | 4 | 1,011 | 177 | 6 | 1.7 |
| 675530 | 1,482 | 76 | 3 | 5 | 939 | 85 | 11 | 1.6 |
| 675600 | 4,925 | 51 | 10 | 15 | 2,904 | 687 | 4 | 1.7 |
| 675630 | 1,777 | 26 | 8 | 11 | 1,062 | 391 | 3 | 1.7 |
| 675700 | 148 | 1 | 3 | 3 | 85 | 40 | 2 | 1.7 |
| 685530 | 3,843 | 16 | 3 | 4 | 2,189 | 250 | 9 | 1.8 |
| 685600 | 73,934 | 240 | 11 | 17 | 42,981 | 2,189 | 20 | 1.7 |
| 685630 | 65,048 | 459 | 13 | 18 | 37,809 | 3,416 | 11 | 1.7 |
| 695600 | 22,021 | 125 | 4 | 7 | 12,918 | 514 | 25 | 1.7 |
| 705600 | 21,465 | 745 | 4 | 4 | 12,595 | 781 | 16 | 1.7 |
| 705630 | 92,797 | 1,406 | 9 | 9 | 53,853 | 2,529 | 21 | 1.7 |
| 715630 | 3,457 | 24 | 6 | 8 | 2,002 | 204 | 10 | 1.7 |
| 715700 | 137,649 | 773 | 13 | 18 | 80,091 | 4,346 | 18 | 1.7 |
| 715730 | 107,074 | 906 | 10 | 14 | 62,426 | 3,422 | 18 | 1.7 |
| 715800 | 425 | 336 | 4 | 4 | 245 | 139 | 2 | 1.7 |
| 725700 | 11 | 11 | 7 | 7 | 8 | 637 | <1 | 1.4 |
| 725730 | 7 | 7 | 4 | 4 | 5 | 1,031 | <1 | 1.4 |
| 735730 | 0 | 0 | 3 | 3 | 0 | 159 | 0 | 0 |
| 735900 | 1 | 1 | 3 | 3 | 1 | 72 | <1 | 1.0 |

-continued-

Table 15.–Page 2 of 2.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 745900 | 13 | 13 | 4 | 4 | 9 | 920 | <1 | 1.5 |
| 755930 | 8 | 8 | 4 | 5 | 5 | 741 | <1 | 1.7 |
| 765930 | 3,620 | 21 | 5 | 7 | 2,117 | 1,343 | 2 | 1.7 |
| 766000 | 417 | 9 | 5 | 5 | 246 | 825 | <1 | 1.7 |
| 775930 | 8 | 8 | 3 | 3 | 5 | 413 | <1 | 1.6 |
| 786000 | 19 | 19 | 5 | 6 | 12 | 1,158 | <1 | 1.6 |
| 786030 | 66 | 66 | 6 | 7 | 44 | 2,610 | <1 | 1.5 |
| Other ^e | 1,100 | 29 | 13 | 28 | 651 | 2,254 | <1 | 1.7 |
| Total | 1,449,543 | 25,334 | 41 | 84 | 873,651 | 47,340 | 18 | 1.7 |

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Combination of statistical areas (19) in which landings were made by fewer than three vessels.

Table 16.—Bering Sea District snow crab commercial fishery harvest data, 1977/78–2020/21.

| Season | GHL/TAC ^{a,b} | Harvest ^{b,c} | Deadloss ^b | Number of | | | | Average | | |
|---------|------------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{b,e} | Width ^{e,f} |
| 1977/78 | — | 1,716,124 | ND | 15 | 38 | 1,267,546 | 13,247 | 96 | 1.4 | ND |
| 1978/79 | — | 32,187,039 | 759,137 | 102 | 490 | 22,118,498 | 190,746 | 116 | 1.5 | 113 |
| 1979/80 | — | 39,572,668 | 228,345 | 134 | 597 | 25,286,777 | 255,102 | 99 | 1.6 | 118 |
| 1981 | 39.5–91 million | 52,750,034 | 2,269,979 | 153 | 867 | 34,415,322 | 435,742 | 79 | 1.5 | 117 |
| 1982 | 16–22 million | 29,355,374 | 1,092,655 | 122 | 803 | 24,089,562 | 469,091 | 51 | 1.2 | 109 |
| 1983 | 15,800,000 | 26,128,410 | 1,324,466 | 109 | 461 | 23,853,647 | 287,127 | 83 | 1.1 | ND |
| 1984 | 49,000,000 | 26,813,074 | 798,795 | 52 | 367 | 24,009,935 | 173,591 | 138 | 1.1 | 105 |
| 1985 | 98,000,000 | 66,010,484 | 1,064,184 | 75 | 718 | 52,908,591 | 372,045 | 142 | 1.2 | 108 |
| 1986 | 57,000,000 | 97,684,139 | 1,378,533 | 87 | 990 | 76,319,307 | 542,346 | 141 | 1.3 | 110 |
| 1987 | 56,400,000 | 101,903,388 | 978,449 | 103 | 1,038 | 81,307,659 | 616,113 | 132 | 1.3 | 109 |
| 1988 | 110,700,000 | 134,060,185 | 3,242,478 | 171 | 1,285 | 105,933,542 | 747,395 | 142 | 1.3 | 110 |
| 1989 | 132,000,000 | 148,306,262 | 1,940,482 | 169 | 1,300 | 112,704,215 | 665,242 | 169 | 1.3 | 111 |
| 1990 | 139,800,000 | 161,656,405 | 1,798,664 | 189 | 1,563 | 128,859,645 | 912,718 | 141 | 1.3 | 109 |
| 1991 | 315,000,000 | 328,648,169 | 3,464,936 | 219 | 2,788 | 265,124,637 | 1,394,897 | 190 | 1.2 | 110 |
| 1992 | 333,000,000 | 315,302,034 | 2,329,852 | 248 | 2,763 | 227,376,582 | 1,281,796 | 177 | 1.4 | 112 |
| 1993 | 207,200,000 | 230,754,145 | 1,577,102 | 254 | 1,835 | 169,531,168 | 972,118 | 174 | 1.4 | 112 |
| 1994 | 105,800,000 | 149,792,718 | 1,799,763 | 272 | 1,293 | 114,810,186 | 716,524 | 160 | 1.3 | 110 |
| 1995 | 55,700,000 | 75,309,187 | 1,291,135 | 253 | 870 | 60,591,399 | 507,603 | 119 | 1.2 | 109 |
| 1996 | 50,700,000 | 65,696,173 | 1,335,372 | 234 | 771 | 52,892,320 | 520,671 | 102 | 1.2 | 108 |
| 1997 | 117,000,000 | 119,589,339 | 2,351,555 | 226 | 1,127 | 100,013,816 | 754,140 | 133 | 1.2 | 107 |
| 1998 | 234,100,000 | 252,339,284 | 3,037,499 | 230 | 1,853 | 193,618,550 | 930,794 | 208 | 1.3 | 111 |
| 1999 | 196,000,000 | 194,363,869 | 1,926,497 | 241 | 1,734 | 151,183,798 | 945,533 | 160 | 1.3 | 110 |
| 2000 | 28,500,000 | 33,291,344 | 353,125 | 231 | 315 | 25,081,681 | 182,634 | 137 | 1.3 | 111 |

-continued-

Table 16.–Page 2 of 3.

| Season | GHL/TAC ^{a,b} | Harvest ^{b,c} | Deadloss ^b | Number of | | | | Average | | |
|----------------------|------------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|----------------------|
| | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{b,e} | Width ^{e,f} |
| 2001 | 27,300,000 | 25,256,384 | 452,781 | 207 | 322 | 18,612,605 | 191,200 | 97 | 1.4 | 111 |
| 2002 ^g | 30,820,000 | 32,633,210 | 658,456 | 191 | 436 | 25,155,221 | 326,977 | 77 | 1.3 | 110 |
| 2003 ^h | 25,610,000 | 28,316,923 | 680,787 | 190 | 285 | 23,252,904 | 153,862 | 151 | 1.2 | 107 |
| 2004 | 20,831,000 | 23,942,373 | 248,576 | 189 | 265 | 18,669,591 | 123,709 | 151 | 1.3 | 110 |
| 2005 | 20,932,000 | 24,892,128 | 235,479 | 168 | 219 | 17,985,745 | 73,208 | 246 | 1.4 | 114 |
| 2005/06 ⁱ | 37,184,000 | 36,973,890 | 357,200 | 78 | 350 | 24,551,986 | 121,029 | 203 | 1.5 | 117 |
| 2006/07 | 36,566,000 | 36,355,649 | 413,743 | 69 | 307 | 29,620,685 | 89,419 | 331 | 1.2 | 109 |
| 2007/08 | 63,034,000 | 63,028,036 | 551,429 | 78 | 513 | 50,327,591 | 144,110 | 349 | 1.3 | 109 |
| 2008/09 | 58,550,000 | 58,547,849 | 434,622 | 77 | 487 | 45,945,092 | 163,537 | 281 | 1.3 | 110 |
| 2009/10 | 48,017,000 | 48,014,089 | 536,688 | 69 | 354 | 35,289,022 | 137,292 | 257 | 1.4 | 113 |
| 2010/11 | 54,281,000 | 54,263,200 | 352,388 | 68 | 386 | 37,758,496 | 147,478 | 256 | 1.4 | 115 |
| 2011/12 | 88,894,000 | 88,830,652 | 637,432 | 72 | 724 | 60,555,105 | 270,602 | 224 | 1.5 | 115 |
| 2012/13 | 66,350,000 | 66,254,528 | 465,522 | 70 | 505 | 47,455,883 | 225,627 | 210 | 1.4 | 113 |
| 2013/14 | 53,983,000 | 53,983,286 | 405,129 | 70 | 450 | 41,926,542 | 231,614 | 181 | 1.3 | 110 |
| 2014/15 | 67,950,000 | 67,941,587 | 596,641 | 71 | 543 | 55,029,818 | 286,920 | 192 | 1.2 | 110 |
| 2015/16 | 40,611,000 | 40,611,446 | 379,167 | 74 | 390 | 29,614,529 | 217,054 | 136 | 1.4 | 113 |
| 2016/17 | 21,570,000 | 21,570,915 | 250,474 | 63 | 266 | 16,412,386 | 118,548 | 138 | 1.3 | 112 |
| 2017/18 | 18,961,000 | 18,963,473 | 172,569 | 63 | 261 | 15,695,007 | 118,034 | 133 | 1.2 | 108 |
| 2018/19 | 27,581,000 | 27,578,244 | 268,204 | 61 | 313 | 22,470,886 | 127,432 | 176 | 1.2 | 109 |
| 2019/20 | 34,019,000 | 34,024,553 | 417,075 | 59 | 373 | 28,626,114 | 188,958 | 151 | 1.2 | 108 |
| 2020/21 | 45,000,000 | 45,001,190 | 863,626 | 62 | 407 | 37,492,237 | 171,678 | 218 | 1.2 | 107 |

-continued-

Table 16.—Page 3 of 3.

Note: En dashes indicate harvest limits were not set for these fisheries prior to 1981; ND indicates these data were not collected or cannot be derived.

^a Guideline harvest level (GHL) and total allowable catch (TAC) beginning in 2005/06.

^b In pounds.

^c Deadloss included.

^d Number of legal crab per pot lift.

^e Retained catch.

^f Carapace width in millimeters

^g Total harvest includes 30,919 pounds taken from an unidentified statistical area.

^h Includes 181,457 pounds illegally taken in Russian waters.

ⁱ Crab rationalization begins.

Table 17.—Bering Sea District snow crab commercial fishery value and season dates, 1980–2020/21.

| Season | Value | | Season length | | | Location/Comments |
|---------|-----------------------|--------------|---------------|----------|------|---|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 1979/80 | \$0.21 | \$8,262,308 | 11/01/79 | 08/15/80 | 289 | Bering Sea District state closure |
| | | | | 09/03/80 | 308 | Bering Sea District federal closure |
| 1981 | \$0.27 | \$13,629,615 | 01/15/81 | 09/01/81 | 230 | Bering Sea District closure |
| 1982 | \$0.83 | \$23,458,057 | 02/15/82 | 08/01/82 | 168 | Bering Sea District closure |
| 1983 | \$0.38 | \$9,425,499 | 02/15/83 | 05/22/83 | 97 | Bering Sea District closure south of 57°30'N lat |
| | | | | 08/01/83 | 168 | Bering Sea District closure north of 57°30'N lat |
| 1984 | \$0.30 | \$7,804,284 | 02/15/84 | 08/01/84 | 169 | Bering Sea District closure south of 58°N lat |
| | | | | 08/22/84 | 190 | Bering Sea District closure north of 58°N lat to allow for an orderly start to king crab season |
| 1985 | \$0.30 | \$19,483,890 | 01/15/85 | 05/08/85 | 114 | Pribilof Subdistrict closure south of 58°N lat |
| | | | | 08/01/85 | 199 | Bering Sea District closure south of 58°39'N lat |
| | | | | 08/22/85 | 220 | Northern Subdistrict closure to allow for an orderly start to king crab season |
| 1986 | \$0.55 | \$52,721,553 | 01/15/86 | 04/21/86 | 97 | Southeastern Subdistrict closure west of 164°W long |
| | | | | 06/01/86 | 138 | Pribilof Subdistrict closure |
| | | | | 08/01/86 | 199 | Northern Subdistrict closure east of 175°W long |
| | | | | 08/24/86 | 222 | Northern Subdistrict closure west of 175°W long |
| 1987 | \$0.77 | \$77,815,458 | 01/15/87 | 04/12/87 | 88 | Southeastern Subdistrict west of 164°W long and Pribilof Subdistrict closure |
| | | | | 06/01/87 | 138 | Northern Subdistrict south of 60°30'N lat and east of 178°W long closure |
| | | | | 06/22/87 | 159 | Northern Subdistrict north of 60°30'N lat and west of 178°W long closure |

-continued-

Table 17.—Page 2 of 4.

| Season | Value | | Season length | | | Location/Comments |
|--------|-----------------------|---------------|---------------|----------|------|--|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 1988 | \$0.83 | \$108,645,549 | 01/15/88 | 03/29/88 | 75 | Bering Sea District closure (Western Subdistrict to assist in an orderly closure) |
| | | | 05/15/88 | 06/30/88 | 47 | Western Subdistrict reopen and closure |
| 1989 | \$0.77 | \$112,144,331 | 01/15/89 | 03/26/89 | 71 | Eastern Subdistrict closure |
| | | | | 05/07/89 | 113 | Western Subdistrict closure |
| 1990 | \$0.77 | \$122,459,171 | 01/15/90 | 04/09/90 | 85 | Eastern Subdistrict east of 165°W long closure |
| | | | | 04/24/90 | 100 | Eastern Subdistrict west of 165°W long closure |
| | | | | 06/12/90 | 149 | Western Subdistrict closure |
| 1991 | \$0.49 | \$158,683,336 | 01/15/91 | 05/05/91 | 111 | Eastern Subdistrict closure |
| | | | | 06/23/91 | 160 | Western Subdistrict closure |
| 1992 | \$0.59 | \$184,951,171 | 01/15/92 | 04/22/92 | 99 | Bering Sea District closure |
| 1993 | \$0.93 | \$212,377,174 | 01/15/93 | 03/15/93 | 60 | Bering Sea District closure |
| 1994 | \$1.58 | \$233,301,131 | 01/15/94 | 03/01/94 | 46 | Bering Sea District closure |
| 1995 | \$2.44 | \$180,489,720 | 01/15/95 | 02/17/95 | 34 | Bering Sea District closure |
| 1996 | \$1.31 | \$84,200,277 | 01/15/96 | 02/29/96 | 46 | Bering Sea District closure |
| 1997 | \$0.71 | \$83,787,925 | 01/15/97 | 03/21/97 | 66 | Bering Sea District closure |
| 1998 | \$0.55 | \$137,642,743 | 01/15/98 | 03/20/98 | 65 | Bering Sea District closure |
| 1999 | \$0.88 | \$169,366,879 | 01/15/99 | 03/22/99 | 67 | Bering Sea District closure |
| 2000 | \$1.81 | \$59,659,349 | 04/01/00 | 04/08/00 | 8 | Bering Sea District closure |
| 2001 | \$1.49 | \$36,833,621 | 01/15/01 | 02/14/01 | 31 | Bering Sea District closure |
| 2002 | \$1.84 | \$58,674,555 | 01/15/02 | 02/08/02 | 25 | Bering Sea District closure |
| 2003 | \$1.75 | \$48,496,278 | 01/15/03 | 01/25/03 | 11 | Bering Sea District closure |
| 2004 | \$2.04 | \$48,359,092 | 01/15/04 | 01/23/04 | 9 | Bering Sea District closure |

-continued-

Table 17.—Page 3 of 4.

| Season | Value | | Season length | | | Location/Comments |
|----------------------|-----------------------|---------------|---------------|----------|------|---|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 2005 | \$1.79 | \$44,144,504 | 01/15/05 | 01/20/05 | 6 | Bering Sea District closure |
| 2005/06 ^b | \$1.51 | \$55,291,202 | 10/15/05 | 05/15/06 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/06 | 229 | Western Subdistrict closure |
| 2006/07 | \$1.37 | \$49,111,061 | 10/15/06 | 05/15/07 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/07 | 229 | Western Subdistrict closure |
| 2007/08 | \$1.63 | \$102,072,731 | 10/15/07 | 05/15/08 | 214 | Eastern Subdistrict closure |
| | | | | 05/31/08 | 230 | Western Subdistrict closure |
| 2008/09 | \$1.37 | \$79,464,730 | 10/15/08 | 05/15/09 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/09 | 229 | Western Subdistrict closure |
| 2009/10 | \$1.13 | \$53,645,621 | 10/15/09 | 05/15/10 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/10 | 229 | Western Subdistrict closure |
| 2010/11 | \$2.14 | \$115,523,133 | 10/15/10 | 05/15/11 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/11 | 229 | Western Subdistrict closure |
| 2011/12 | \$1.89 | \$166,973,717 | 10/15/11 | 05/31/12 | 230 | Eastern Subdistrict east of 171°W long closure |
| | | | | 06/15/12 | 245 | Western and Eastern Subdistricts west of 171°W long closure |
| 2012/13 | \$2.02 | \$133,088,320 | 10/15/12 | 05/15/13 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/13 | 229 | Western Subdistrict closure |
| 2013/14 | \$2.15 | \$115,438,494 | 10/15/13 | 05/15/14 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/14 | 229 | Western Subdistrict closure |
| 2014/15 | \$1.67 | \$112,275,497 | 10/15/14 | 05/15/15 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/15 | 229 | Western Subdistrict closure |
| 2015/16 | \$2.01 | \$80,936,867 | 10/15/15 | 05/15/16 | 214 | Eastern Subdistrict closure |
| | | | | 05/31/16 | 230 | Western Subdistrict closure |

-continued-

Table 17.—Page 4 of 4.

| Season | Value | | Season length | | | Location/Comments |
|---------|-----------------------|---------------|---------------|----------|------|-----------------------------|
| | Exvessel ^a | Total | Opened | Closed | Days | |
| 2016/17 | \$2.72 | \$58,026,393 | 10/15/16 | 05/15/17 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/17 | 229 | Western Subdistrict closure |
| 2017/18 | \$3.00 | \$56,464,897 | 10/15/17 | 05/15/18 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/18 | 229 | Western Subdistrict closure |
| 2018/19 | \$3.00 | \$82,036,383 | 10/15/18 | 05/15/19 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/19 | 229 | Western Subdistrict closure |
| 2019/20 | \$3.15 | \$105,928,621 | 10/15/19 | 05/15/20 | 214 | Eastern Subdistrict closure |
| | | | | 05/31/20 | 230 | Western Subdistrict closure |
| 2020/21 | \$3.01 | \$132,857,821 | 10/15/20 | 05/15/21 | 213 | Eastern Subdistrict closure |
| | | | | 05/31/21 | 229 | Western Subdistrict closure |

^a Average price per pound.^b Crab rationalization begins.

Table 18.—Bering Sea District snow crab commercial fishery harvest and effort by week, 2020/21.

| Week ending | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| Oct-17 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| ... ^e | | | | | | | | |
| Dec-26 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Jan-2 | 716,595 | 9,427 | 4 | 5 | 577,939 | 2,754 | 210 | 1.2 |
| Jan-9 | 2,908,755 | 46,403 | 15 | 20 | 2,404,260 | 11,824 | 203 | 1.2 |
| Jan-16 | 1,562,546 | 17,887 | 9 | 12 | 1,298,990 | 5,264 | 247 | 1.2 |
| Jan-23 | 3,227,069 | 60,258 | 20 | 26 | 2,730,900 | 13,959 | 196 | 1.2 |
| Jan-30 | 2,888,591 | 66,969 | 16 | 20 | 2,382,376 | 9,562 | 249 | 1.2 |
| Feb-6 | 3,507,942 | 96,814 | 30 | 34 | 2,942,077 | 15,837 | 186 | 1.2 |
| Feb-13 | 2,975,544 | 48,148 | 19 | 30 | 2,536,560 | 12,457 | 204 | 1.2 |
| Feb-20 | 2,607,643 | 61,989 | 28 | 33 | 2,168,857 | 14,885 | 146 | 1.2 |
| Feb-27 | 1,739,996 | 34,928 | 18 | 21 | 1,483,382 | 8,039 | 185 | 1.2 |
| Mar-6 | 1,416,901 | 15,191 | 7 | 12 | 1,169,302 | 4,597 | 254 | 1.2 |
| Mar-13 | 6,252,555 | 113,714 | 38 | 50 | 5,235,883 | 20,483 | 256 | 1.2 |
| Mar-20 | 1,683,710 | 22,278 | 11 | 13 | 1,382,155 | 5,476 | 252 | 1.2 |
| Mar-27 | 4,194,002 | 55,002 | 24 | 36 | 3,475,995 | 10,903 | 319 | 1.2 |
| Apr-3 | 1,973,843 | 27,353 | 16 | 21 | 1,635,159 | 6,155 | 266 | 1.2 |
| Apr-10 | 2,638,155 | 47,507 | 19 | 27 | 2,191,503 | 10,526 | 208 | 1.2 |
| Apr-17 | 1,401,698 | 15,609 | 9 | 15 | 1,148,935 | 4,471 | 257 | 1.2 |
| Apr-24 | 1,042,251 | 50,270 | 8 | 9 | 880,084 | 4,268 | 206 | 1.2 |
| May-1 | 1,053,713 | 41,890 | 6 | 13 | 870,657 | 4,501 | 193 | 1.2 |
| May-8 | 455,141 | 22,471 | 3 | 3 | 373,811 | 2,609 | 143 | 1.2 |
| May-15 | CF | CF | 1 | 2 | CF | CF | CF | CF |

-continued-

Table 18.—Page 2 of 2.

| Week ending | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|-------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| May-22 | CF | CF | 2 | 2 | CF | CF | CF | CF |
| May-29 | CF | CF | 1 | 1 | CF | CF | CF | CF |
| Total | 45,001,190 | 863,626 | 62 | 407 | 37,492,237 | 171,678 | 218 | 1.2 |

Note: CF indicates confidential data.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Consecutive weeks with no harvest.

Table 19.—Bering Sea District snow crab commercial fishery harvest and effort by statistical area, 2020/21.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|---------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 665500 | 64,719 | 437 | 4 | 5 | 51,386 | 520 | 99 | 1.3 |
| 665530 | 61,556 | 837 | 4 | 5 | 51,752 | 272 | 190 | 1.2 |
| 675530 | 193 | 7 | 3 | 4 | 165 | 56 | 3 | 1.2 |
| 675600 | 1,734 | 62 | 3 | 4 | 1,441 | 121 | 12 | 1.2 |
| 715630 | 598,817 | 6,964 | 20 | 42 | 536,163 | 4,761 | 113 | 1.1 |
| 715700 | 143,639 | 2,545 | 18 | 33 | 129,027 | 1,268 | 102 | 1.1 |
| 715730 | 302,447 | 5,549 | 13 | 32 | 274,896 | 2,277 | 121 | 1.1 |
| 715800 | 20,249 | 466 | 6 | 9 | 17,859 | 176 | 101 | 1.1 |
| 725630 | 573,464 | 7,221 | 15 | 45 | 513,176 | 4,125 | 124 | 1.1 |
| 725700 | 652,519 | 11,534 | 32 | 72 | 578,274 | 5,898 | 98 | 1.1 |
| 725730 | 891,386 | 12,896 | 33 | 77 | 781,984 | 8,499 | 92 | 1.1 |
| 725800 | 3,830 | 57 | 4 | 7 | 3,532 | 135 | 26 | 1.1 |
| 735700 | 6,159 | 110 | 7 | 9 | 5,289 | 306 | 17 | 1.2 |
| 735730 | 59,098 | 731 | 18 | 33 | 48,545 | 975 | 50 | 1.2 |
| 735800 | 13,541 | 332 | 16 | 26 | 11,534 | 667 | 17 | 1.2 |
| 735830 | 58,150 | 658 | 13 | 29 | 48,590 | 798 | 61 | 1.2 |
| 735900 | 158,231 | 1,694 | 7 | 16 | 132,287 | 724 | 183 | 1.2 |
| 745800 | 45 | 1 | 8 | 11 | 41 | 182 | <1 | 1.1 |
| 745830 | 55,846 | 1,124 | 17 | 30 | 46,387 | 718 | 65 | 1.2 |
| 745900 | 1,160,642 | 17,675 | 14 | 36 | 995,001 | 5,917 | 168 | 1.2 |
| 745930 | 214,152 | 5,096 | 18 | 38 | 169,547 | 1,332 | 127 | 1.3 |
| 755830 | 405 | 48 | 5 | 9 | 350 | 135 | 3 | 1.2 |

-continued-

Table 19.—Page 2 of 2.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|---------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 755900 | 116,282 | 2,030 | 13 | 19 | 95,852 | 738 | 130 | 1.2 |
| 755930 | 6,186,967 | 144,586 | 39 | 113 | 4,990,042 | 24,694 | 202 | 1.2 |
| 756000 | 353,636 | 9,206 | 24 | 46 | 292,396 | 1,526 | 192 | 1.2 |
| 765830 | 606 | 71 | 3 | 5 | 523 | 130 | 4 | 1.2 |
| 765900 | 363,213 | 19,416 | 19 | 34 | 303,339 | 2,229 | 136 | 1.2 |
| 765930 | 2,425,276 | 58,297 | 44 | 103 | 2,019,822 | 10,712 | 189 | 1.2 |
| 766000 | 8,558,593 | 163,648 | 50 | 130 | 7,116,869 | 29,656 | 240 | 1.2 |
| 766030 | 144,377 | 3,277 | 13 | 18 | 121,304 | 840 | 144 | 1.2 |
| 775900 | 1,852,736 | 20,423 | 18 | 37 | 1,513,590 | 6,918 | 219 | 1.2 |
| 775930 | 3,411,308 | 61,767 | 29 | 65 | 2,778,620 | 10,431 | 266 | 1.2 |
| 776000 | 2,326,345 | 37,785 | 28 | 56 | 1,951,126 | 8,765 | 223 | 1.2 |
| 776030 | 1,227,552 | 49,581 | 19 | 39 | 1,027,708 | 3,511 | 293 | 1.2 |
| 785900 | 0 | 0 | 3 | 3 | 0 | 88 | 0 | - |
| 785930 | 14,912 | 160 | 5 | 6 | 12,152 | 233 | 52 | 1.2 |
| 786000 | 3,070,981 | 50,135 | 27 | 68 | 2,517,274 | 7,611 | 331 | 1.2 |
| 786030 | 9,851,465 | 166,307 | 36 | 103 | 8,306,849 | 22,632 | 367 | 1.2 |
| Other ^e | 56,116 | 895 | 12 | 33 | 47,545 | 1,102 | 43 | 1.2 |
| Total | 45,001,190 | 863,626 | 62 | 407 | 37,492,237 | 171,678 | 218 | 1.2 |

^a In pounds.^b Deadloss included.^c Number of legal crab per pot lift.^d Retained catch.^e Combination of statistical areas (19) in which landings were made by fewer than three vessels.

Table 20.—Bering Sea District grooved Tanner crab commercial fishery harvest data, 1992–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|-------------------|------------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|-------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 1992 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 1993 | 658,796 | 71,000 | 6 | 18 | 342,095 | 35,650 | 9 | 1.9 | \$0.92 | \$540,772 |
| 1994 | 322,444 | 30,585 | 4 | 12 | 165,365 | 13,739 | 11 | 2.0 | \$2.65 | \$773,426 |
| 1995 | 984,648 | 67,329 | 8 | 51 | 461,401 | 59,028 | 7 | 2.1 | \$2.09 | \$1,917,197 |
| 1996 | 95,795 | 11,120 | 3 | 11 | 46,338 | 10,802 | 4 | 2.1 | \$1.12 | \$94,836 |
| 1997–1999 | No Commercial Fishing Effort | | | | | | | | | |
| 2000 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2001 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2002 | No Commercial Fishing Effort | | | | | | | | | |
| 2003 | CF | CF | 1 | 7 | CF | CF | CF | CF | CF | CF |
| 2004 | CF | CF | 2 | 10 | CF | CF | CF | CF | CF | CF |
| 2005 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2006–2016 | No Commercial Fishing Effort | | | | | | | | | |
| 2017 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2018 | No Commercial Fishing Effort | | | | | | | | | |
| 2019 ^f | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2020 | No Commercial Fishing Effort | | | | | | | | | |

Note: CF indicates confidential data.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Average price per pound.

^f Restricted to incidental harvest during Bering Sea golden king crab fishery.

Table 21.—Bering Sea District triangle Tanner crab commercial fishery harvest data, 1995–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|-------------------|------------------------|-----------------------|-----------|----------|------------------------------|-------------|-------------------|-----------------------|-----------------------|----------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 1995 | 49,007 | 14,147 | 4 | 26 | 41,914 | 22,180 | 2 | 1.2 | \$1.44 | \$50,198 |
| 1996 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 1997–1999 | | | | | No Commercial Fishing Effort | | | | | |
| 2000 ^f | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2001 ^f | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2002 | | | | | No Commercial Fishing Effort | | | | | |
| 2003 ^f | CF | CF | 1 | 7 | CF | CF | CF | CF | CF | CF |
| 2004 ^f | CF | CF | 3 | 13 | CF | CF | CF | CF | CF | CF |
| 2005–2016 | | | | | No Commercial Fishing Effort | | | | | |
| 2017 ^f | CF | CF | 1 | 2 | CF | CF | CF | CF | CF | CF |
| 2018–2020 | | | | | No Commercial Fishing Effort | | | | | |

Note: CF indicates confidential data. Harvest is bycatch only; therefore, harvest limits are not set for this stock.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Average price per pound.

^f Restricted to incidental harvest during Bering Sea golden king crab and grooved Tanner crab fisheries.

Table 22.—North Peninsula District commercial Dungeness crab fishery data, 1992–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|--------|------------------------|-----------------------|-----------|----------|------------------------------|-------------|-------------------|-----------------------|-----------------------|-------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 1992 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 1993 | CF | CF | 2 | 8 | CF | CF | CF | CF | CF | CF |
| 1994 | CF | CF | 2 | 13 | CF | CF | CF | CF | CF | CF |
| 1995 | 134,407 | 367 | 6 | 18 | 63,732 | 34,499 | 4 | 2.1 | \$1.32 | \$176,933 |
| 1996 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 1997 | CF | CF | 1 | 2 | CF | CF | CF | CF | CF | CF |
| 1998 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 1999 | | | | | No Commercial Fishing Effort | | | | | |
| 2000 | CF | CF | 1 | 5 | CF | CF | CF | CF | CF | CF |
| 2001 | | | | | No Commercial Fishing Effort | | | | | |
| 2002 | CF | CF | 2 | 2 | CF | CF | CF | CF | CF | CF |
| 2003 | | | | | No Commercial Fishing Effort | | | | | |
| 2004 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2005 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 2006 | | | | | No Commercial Fishing Effort | | | | | |
| 2007 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 2008 | | | | | No Commercial Fishing Effort | | | | | |
| 2009 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 2010 | 795,392 | 10,414 | 5 | 24 | 391,849 | 60,985 | 6 | 2.0 | \$1.73 | \$1,358,012 |
| 2011 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 2012 | CF | CF | 1 | 2 | CF | CF | CF | CF | CF | CF |
| 2013 | | | | | No Commercial Fishing Effort | | | | | |
| 2014 | CF | CF | 1 | 4 | CF | CF | CF | CF | CF | CF |

-continued-

Table 22.—Page 2 of 2.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|-----------|------------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|-------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 2015 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 2016 | CF | CF | 1 | 6 | CF | CF | CF | CF | CF | CF |
| 2017 | CF | CF | 1 | 2 | CF | CF | CF | CF | CF | CF |
| 2018–2019 | No Commercial Fishing Effort | | | | | | | | | |
| 2020 | CF | CF | 2 | 19 | CF | CF | CF | CF | CF | CF |

Note: CF indicates confidential data; harvest limits are not set for this stock.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Average price per pound.

Table 23.—Bristol Bay - Bering Sea (Area Q) weathervane scallop commercial fishery harvest data, 1987–2020/21.

| Season | GHL ^{a,b} | Harvest ^a | Number of | | Meat weight CPUE ^c |
|---------|--------------------|----------------------|------------------------------|--------------|----------------------------------|
| | | | Vessels | Dredge hours | |
| 1987 | — | CF | 2 | CF | CF |
| 1988 | — | | No Commercial Fishing Effort | | |
| 1989 | — | | No Commercial Fishing Effort | | |
| 1990 | — | | No Commercial Fishing Effort | | |
| 1991 | — | CF | 2 | CF | CF |
| 1992 | — | | No Commercial Fishing Effort | | |
| 1993/94 | — | 605,953 | 10 | ND | ND |
| 1994/95 | — | 505,439 | 9 | 11,113 | 45 |
| 1995/96 | | | No Commercial Fishery | | |
| 1996/97 | 600,000 | 150,295 | 1 | 2,313 | 65 |
| 1997/98 | 600,000 | 97,002 | 2 | 2,246 | 43 |
| 1998/99 | 400,000 | 96,795 | 4 | 2,319 | 42 |
| 1999/00 | 400,000 | 164,929 | 2 | 3,294 | 50 |
| 2000/01 | 200,000 | 205,520 | 3 | 3,355 | 61 |
| 2001/02 | 200,000 | 140,871 | 3 | 3,072 | 46 |
| 2002/03 | 105,000 | 92,240 | 2 | 2,038 | 45 |
| 2003/04 | 105,000 | 42,590 | 2 | 1,020 | 42 |
| 2004/05 | 50,000 | 10,050 | 1 | 275 | 37 |
| 2005/06 | 50,000 | 23,220 | 1 | 602 | 39 |
| 2006/07 | 50,000 | 48,246 | 1 | 1,138 | 42 |
| 2007/08 | 50,000 | 49,995 | 2 | 1,084 | 46 |
| 2008/09 | 50,000 | 49,995 | 1 | 960 | 52 |
| 2009/10 | 50,000 | 48,921 | 1 | 1,275 | 38 |
| 2010/11 | 50,000 | 50,100 | 2 | 972 | 52 |
| 2011/12 | 50,000 | 50,275 | 2 | 984 | 51 |
| 2012/13 | 50,000 | 50,045 | 1 | 943 | 53 |
| 2013/14 | 50,000 | 49,989 | 2 | 1,086 | 46 |
| 2014/15 | 50,000 | 12,445 | 2 | 525 | 24 |
| 2015/16 | 7,500 | 7,500 | 1 | 307 | 24 |
| 2016/17 | 7,500 | 7,575 | 1 | 275 | 28 |
| 2017/18 | 7,500 | 7,535 | 1 | 316 | 24 |
| 2018/19 | 7,500 | 7,540 | 1 | 357 | 21 |
| 2019/20 | 7,500 | 7,130 | 1 | 365 | 20 |
| 2020/21 | 7,500 | | No Commercial Fishing Effort | | |

Note: En dashes indicate harvest limits were not set for this fishery prior to 1996/97; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds of shucked scallop meat.

^b Guideline harvest level (GHL).

^c Pounds of shucked scallop meat per dredge hour.

Table 24.—Bering Sea commercial octopus incidental harvest in groundfish fisheries, 1995–2020.

| Season | State waters | | | State and federal waters | | | | |
|--------|--------------|----------|---------------------------|--------------------------|----------|---------------------------|-----------------|-----------------------------|
| | Vessels | Landings | Whole pounds ^a | Vessels | Landings | Whole pounds ^a | At-sea discards | Exvessel value ^b |
| 1995 | 5 | 12 | 2,252 | 49 | 135 | 17,554 | 5,587 | \$0.14 |
| 1996 | 6 | 10 | 1,195 | 63 | 191 | 26,343 | 21,144 | \$0.33 |
| 1997 | 3 | 3 | 59 | 44 | 92 | 12,202 | 5,205 | \$0.20 |
| 1998 | 4 | 8 | 673 | 47 | 81 | 8,204 | 5,624 | \$0.03 |
| 1999 | 2 | 2 | CF | 22 | 56 | 7,002 | 6,593 | ND |
| 2000 | 4 | 6 | 551 | 78 | 272 | 39,915 | 23,611 | \$0.03 |
| 2001 | 2 | 3 | CF | 62 | 158 | 49,641 | 41,215 | \$0.03 |
| 2002 | 2 | 2 | CF | 68 | 187 | 56,078 | 16,628 | \$0.05 |
| 2003 | 4 | 7 | 4,064 | 80 | 237 | 122,443 | 27,780 | \$0.63 |
| 2004 | 4 | 6 | 4,615 | 92 | 279 | 88,534 | 25,527 | \$0.39 |
| 2005 | 5 | 19 | 4,033 | 80 | 271 | 156,381 | 12,583 | \$0.65 |
| 2006 | 6 | 8 | 1,004 | 88 | 304 | 93,624 | 5,310 | \$0.63 |
| 2007 | 4 | 6 | 1,946 | 110 | 375 | 102,128 | 37,436 | \$0.45 |
| 2008 | 5 | 7 | 7,177 | 82 | 252 | 66,742 | 14,071 | \$0.47 |
| 2009 | 1 | 1 | CF | 67 | 144 | 20,107 | 7,858 | \$0.30 |
| 2010 | 0 | 0 | 0 | 81 | 201 | 67,187 | 35,477 | \$0.24 |
| 2011 | 2 | 3 | CF | 124 | 470 | 193,222 | 158,042 | \$0.21 |
| 2012 | 0 | 0 | 0 | 104 | 312 | 48,938 | 36,780 | \$0.34 |
| 2013 | 2 | 2 | CF | 100 | 310 | 59,351 | 40,453 | ND |
| 2014 | 0 | 0 | 0 | 121 | 403 | 60,445 | 29,697 | \$0.05 |
| 2015 | 3 | 5 | 90 | 131 | 519 | 81,685 | 49,719 | \$0.14 |
| 2016 | 1 | 2 | CF | 103 | 465 | 164,200 | 85,174 | \$0.05 |
| 2017 | 4 | 7 | 1,150 | 119 | 427 | 106,883 | 62,250 | \$0.35 |
| 2018 | 16 | 81 | 7,927 | 131 | 455 | 105,682 | 53,158 | \$0.35 |
| 2019 | 17 | 84 | 6,782 | 122 | 485 | 78,188 | 31,612 | \$0.62 |
| 2020 | 8 | 12 | 331 | 120 | 168 | 42,438 | 23,656 | \$0.19 |

Note: CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a Includes discards.

^b Average price per pound, based on landed weight.




Figure 1.—Bristol Bay king crab commercial fishery Registration Area T.




Figure 2.—Bristol Bay red king crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1966–2020/21.




Figure 3.—Bering Sea king crab commercial fishery Registration Area Q.




Figure 4.—Saint Matthew Island section blue king crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1977–2020/21.




Figure 5.—Bering Sea District Tanner crab commercial fishery Registration Area J including subdistricts and sections.




Figure 6.—Bering Sea Tanner crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1974/75–2020/21.




Figure 7.—Bering Sea Snow crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1977/78–2020/21.




Figure 8.—North Peninsula and Aleutian Islands Dungeness crab Districts of Registration Area J.




Figure 9.—Bering Sea and Aleutian Islands weathervane scallop Registration Areas.




Figure 10.—Bering Sea and Aleutian Islands Tanner crab and miscellaneous shellfish Districts of Registration Area J.

SECTION II: ALEUTIAN ISLANDS SHELLFISH FISHERIES

ALEUTIAN ISLANDS KING CRAB MANAGEMENT AREA

DESCRIPTION OF AREA

The Aleutian Islands king crab Registration Area O eastern boundary is the longitude of Scotch Cap Light (164°44.72'W long); the northern boundary is a line from Cape Sarichef (54°36'N lat) to 171°W long, north to 55°30'N lat; and the western boundary the United States–Russia Maritime Boundary Line of 1990 (Figure 11).

ALEUTIAN ISLANDS GOLDEN KING CRAB

The Aleutian Islands golden king crab (AIGKC) stock is managed as two separate fisheries, east and west of 174°W long, with a separate TAC set for each fishery.

2020/21 Fishery East of 174°W Longitude

The 2020/21 Aleutian Islands golden king crab fishery east of 174°W long (EAG) opened August 1 with a combined IFQ and CDQ TAC of 3,650,000 pounds (Table 25). TAC was allocated by NMFS as 90% IFQ (3,285,000 pounds) and 10% CDQ (365,000 pounds) with five of the six CDQ groups participating in the harvest of the CDQ allocation. Three vessels participated in the fishery and harvested 3,650,255 pounds, of which 0.9% was deadloss (Table 25). Despite the regulatory fishing season running through April 30, harvest occurred August through mid-February, with the last delivery on February 21 (Table 26). Harvest by statistical week is confidential and therefore not shown in tables. On average, vessels were active in the fishery for 114 days. Harvesters were paid an initial average price of \$4.56 per pound. Total exvessel fishery value was estimated to be \$16,492,203 (Table 26).

Total effort for the 2020/21 fishery was 28,833 pot lifts. CPUE was 30 legal crab per pot, slightly below the post-rationalization (2005/06–2019/20) average CPUE of 32 (Table 25, Figure 12). Harvest was distributed across 22 ADF&G statistical reporting areas with the highest concentration of harvest (20%) occurring in ADF&G statistical area 705232, around the Islands of Four Mountains (Table 27).

EAG cost recovery was conducted by ADF&G in 2020 with 170,863 pounds (38,833 legal male crab) harvested. The 14-day charter occurred from September 6 to September 8, and September 10 to September 20 (Table 28).

Port Sampling

During the 2020/21 EAG fishery, ADF&G personnel and onboard observers sampled golden king crab from vessels at shore-based processors in Akutan, Dutch Harbor, and King Cove. Biological data collected on landed golden king crab consisted of carapace length, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting

areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 47 of the 47 total landings (IFQ and CDQ) during the 2020/21 EAG fishery. Average weight of sampled crab was 4.2 pounds while average carapace length was 148 mm (Table 25).

2020/21 Fishery West of 174°W Longitude

The 2020/21 Aleutian Islands golden king crab fishery west of 174°W long (WAG) opened August 1 with a combined IFQ and CDQ TAC of 2,960,000 pounds (Table 25). TAC was allocated by NMFS as 90% IFQ (2,664,000 pounds) and 10% Adak Community Allocation (ACA; 296,000 pounds). Three vessels participated in the fishery and harvested 2,792,835 pounds, of which 2.2% was deadloss (Table 25). Harvest occurred throughout the season, with the last delivery on May 18 (Table 26). The regulatory closure date of April 30 was extended by ADF&G Emergency Order to May 13 at the request of industry to allow additional harvesting opportunity given harvesting inefficiencies due to COVID-19. Golden king crab mating and molting are asynchronous. Thus, the current regulatory season dates are not based on protecting crab during critical reproductive periods; rather, they largely reflect historical fishery and stock assessment timing. Harvest by statistical week is confidential and therefore not shown in tables. On average, vessels were active in the fishery for 264 days. Harvesters were paid an initial average price of \$4.51 per pound. Total exvessel fishery value was estimated to be \$12,311,834 (Table 26).

Total effort for the 2020/21 fishery was 46,701 pot lifts. CPUE was 15 legal crab per pot, below the post-rationalization (2005/06–2018/19) average CPUE of 19 (Table 25, Figure 13). Harvest was distributed across 64 ADF&G statistical reporting areas; harvest by statistical area is confidential and therefore not shown in tables.

Port Sampling

During the 2020/21 WAG fishery, ADF&G personnel and onboard observers sampled golden king crab from vessels at shore-based processors in Dutch Harbor and King Cove. Biological data collected on landed golden king crab consisted of carapace length, shell condition, and average weight. Confidential interviews, supplemented by daily fishing log records, were conducted with vessel operators to acquire detailed information regarding statistical reporting areas fished, effort, and fishery performance. Data were collected by ADF&G port samplers and onboard observers from 35 of the 35 total landings (IFQ and CDQ) during the 2020/21 WAG fishery. Average weight of sampled crab was 4.1 pounds and average carapace length was 150 mm (Table 25).

Fishery Management and Stock Status

The AIGKC fishery has been managed since 1996/97 under a constant-catch harvest strategy with TAC fixed in regulation both east and west of 174°W long (5 AAC 34.612). Only a small portion of the area in which golden king crab are commercially harvested has been surveyed historically. In 2014, methods testing for an industry cooperative in-fishery survey began with ADF&G personnel deployed on eastern Aleutian Islands golden king crab fishing vessels. The annual in-fishery survey is on-going with the goal of having a time series of these data to inform the current stock assessment model.

After adoption of a new stock assessment modelling framework in 2016, the Board of Fisheries modified the harvest strategy to give the department flexibility to modify the regulatory harvest levels based on the best scientific information available. The current stock assessment model had been in development for some time when it was adopted by the NPFMC and first used to set OFL

and ABC for the 2017/18 fishery, but a harvest strategy had not yet been developed to utilize management output quantities for TAC setting. Department scientists worked with industry to develop a new harvest strategy using model outputs for TAC setting after the stock assessment model was adopted. The current harvest strategy employing stock assessment model outputs has been used to set TACs since the 2019/20 fishery.

Further information on AIGKC stock status and federal overfishing levels may be found in the *2020 Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands Regions* (NPFMC 2020a).

EASTERN ALEUTIAN TANNER CRAB DISTRICT

DESCRIPTION OF DISTRICT

The Eastern Aleutian District (EAD) for Tanner crab encompasses all waters of Registration Area J between the longitude of Scotch Cap Light (164°44'W long), west to 172°W long, and south of the latitude of Cape Sarichef (54°36'N lat; Figure 14).

TANNER CRAB

2020 Commercial Fishery

The 2020 EAD commercial Tanner crab fishery was closed. The minimum mature male abundance thresholds necessary for a commercial fishery were not met in the Unalaska/Kalekta Bay or Akutan Bay Sections. The Makushin/Skan Bay Section exceeded the minimum mature male abundance threshold, but the calculated GHL was well below the minimum GHL of 35,000 pounds required to open the fishery (Tables 29 and 30).

Fishery Management and Stock Status

The EAD Tanner crab fishery has a total of 300 pots allowed in the fishery with no more than 50 pots per vessel; pots may be operated only from 8:00 AM until 5:59 PM. Vessel length is restricted to 58 feet or less in the Unalaska Section. In the remainder of the EAD, vessel length is restricted to 58 feet or less when the GHL for Tanner crab is 1,000,000 pounds or less. The EAD Tanner crab fishery is an open access fishery. The harvest strategy is found in 5 AAC 35.509 *Eastern Aleutian District Tanner crab harvest strategy*.

The 2020 ADF&G survey of the Eastern Aleutian District was cancelled due to COVID-19. In August 2019, Akutan Bay, Unalaska/Kalekta Bay, and Makushin/Skan Bay were surveyed with trawl gear using the ADF&G R/V *Resolution* (Spalinger and Knutson 2020). Total estimated abundance was 5.8 million crab. Legal male abundance was estimated at 0.019 million crab, well below the trawl survey time series average of 0.261 million crab (1990–2018).

SUBSISTENCE

2020 King and Tanner Fishery

In 2020, ADF&G issued 156 subsistence permits, of which 56, or 36%, were returned. The returned permits account for a reported harvest of 119 Tanner crab and 9 red king crab (Table 31). Harvest of Tanner crab ranged from 0 to 31 crab per permit holder and harvest of red king crab ranged from 0 to 7 crab per permit holder. Subsistence effort was primarily focused in Unalaska Bay. King and Tanner crab harvested in the EAD were taken with pot gear.

ALEUTIAN DISTRICT DUNGENESS CRAB

DESCRIPTION OF DISTRICT

The Aleutian District for Dungeness crab includes all waters of Registration Area J west of the longitude of Scotch Cap Light (164°44'W long), south of the latitude of Cape Sarichef (54°35.89'N lat), and east of the United States–Russia Maritime Boundary Line of 1990 (Figure 8).

2020 FISHERY

The Aleutian District Dungeness crab fishery opened May 1, 2020. No vessels registered to fish for Dungeness crab in the Aleutian Islands during 2020 (Table 32).

Fishery Management and Stock Status

The Aleutian Islands Dungeness crab fishery is managed using 3-S management (size, sex, and season). Only male Dungeness crab 6.5 inches (165 mm) or greater in carapace width may be retained in the Aleutian District from May 1 to October 31. No stock assessment is available and limited biological and fishery data have been collected through dockside sampling. Stock status of Dungeness crab in the Aleutian Islands is unknown, but the resource is believed to be limited by availability of suitable habitat.

DUTCH HARBOR WEATHERVANE SCALLOP REGISTRATION AREA O

DESCRIPTION OF AREA

The Dutch Harbor weathervane scallop Registration Area O eastern boundary is the longitude of Scotch Cap Light (164°44.72'W long), the northern boundary is the latitude of Cape Sarichef (54°36'N lat), and the western boundary 171°W long (Figure 9).

DUTCH HARBOR WEATHERVANE SCALLOP

2020/21 Fishery

The 2020/21 Dutch Harbor weathervane scallop fishery opened July 1, 2020, with a GHL of 5,000 pounds of shucked scallop meat on the Bering Sea side of Registration Area O; the Pacific Ocean side was closed. No vessels participated in the fishery (Table 33) and it closed by regulation on February 15, 2021.

Fishery Management and Stock Status

The Dutch Harbor weathervane scallop Registration Area O is managed by individual scallop beds, one on the Bering Sea side of Umnak Island and the other on the Pacific Ocean side of Unalaska Island. CPUE of 20 pounds of shucked scallop meat per dredge hour from the Bering Sea side in the 2019/20 season was a slight increase from the historically low CPUEs from the previous two seasons but remains low relative to time series high CPUE of 93 in the 2013/14 season. The Pacific Ocean side was closed in 2018/19 for a three-year period due to concerns about poor fishery performance. The Pacific side reopened for the 2020/21 season.

Area O has never been surveyed with dredge gear as part of the Statewide Weathervane Scallop Dredge survey, and the Westward Region Large-Mesh Bottom Trawl Survey does not generally overlap with scallop fishing grounds in the area. Therefore, the stock is assessed annually using

available fishery and observer data. The harvest strategy is found in 5 AAC 38.076 *Alaska Scallop Fishery Management Plan* and 5 AAC 38.078 *State-Waters Weathervane Scallop Management Plan*.

Further information on weathervane scallop stock status and federal overfishing levels may be found in the 2020 *Stock Assessment and Fishery Evaluation Report for the Scallop Fishery off Alaska* (NPFMC 2020b).

ALEUTIAN ISLANDS MISCELLANEOUS SHELLFISH

DESCRIPTION OF DISTRICT

The Aleutian Islands District of miscellaneous shellfish Registration Area J includes all waters south of the latitude of Cape Sarichef (54°36'N lat), west of the longitude of Scotch Cap Light (164°44'W long), and east of the United States–Russia Maritime Boundary Line of 1990 (Figure 10).

2020 FISHERIES

Octopus

In 2020, harvest from state waters was 228,716 pounds from 44 vessels and 253 landings. Harvest from state and federal waters combined was 334,728 pounds from 105 vessels and 465 landings. All reported harvest was incidental to groundfish fisheries. Average exvessel value based on landed weight of octopus in 2020 was \$0.37 per pound (Table 34).

Fishery Management and Stock Status

Octopus are considered a shellfish species under State of Alaska regulation. Limited directed fishing within state waters may occur under the authority of a commissioner's permit; however, octopus are primarily retained as bycatch during state and federal groundfish fisheries. Currently, vessels may retain incidentally caught octopus up to 20% of the weight of the target groundfish species or halibut onboard. Most octopus are retained as bycatch in Pacific cod pot gear fisheries.

Incidental harvest of octopus in the Aleutian Islands is dominated by giant Pacific octopus, *Enteroctopus dofleini*, although at least six other species of octopus are known to occur in the Aleutian Islands. The 2020 NMFS Aleutian Islands trawl survey was cancelled due to COVID-19. Results from the 2018 NMFS Aleutian Islands trawl survey estimate octopus (all species) biomass at 5.0 million pounds with 98% of the estimate composed of *E. dofleini*. Biomass estimates for octopus in the Aleutian Islands are highly variable and do not necessarily reflect the sizes of octopus caught in fisheries (Ormseth et al. 2020). General knowledge of the stock is limited and there is currently no reliable estimate of octopus biomass (Ormseth et al. 2020).

Further information on octopus stock status and federal overfishing levels may be found in the 2020 *Assessment of the Octopus Stock Complex in the Bering Sea and Aleutian Islands* (Ormseth et al. 2020).

REFERENCES CITED

- Ormseth, O. A., Conners, M. E., K. Aydin, and C. Conrath. 2020. Assessment of the octopus stock complex in the Bering Sea and Aleutian Islands. [In] Stock assessment and fishery evaluation report of the Bering Sea/Aleutian Islands regions. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2020a. Stock assessment and fishery evaluation report for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions: 2020 crab SAFE. North Pacific Fishery Management Council, Anchorage.
- NPFMC (North Pacific Fishery Management Council). 2020b. Stock assessment and fishery evaluation report for the weathervane scallop fishery off Alaska: 2020 scallop SAFE. North Pacific Fishery Management Council, Anchorage.
- Spalinger, K., and M. Knutson. 2020. Large-mesh bottom trawl survey of crab and groundfish: Kodiak, Chignik, South Peninsula, and Eastern Aleutian management districts, 2019. Alaska Department of Fish and Game, Fishery Management Report No. 20-16, Anchorage.

TABLES AND FIGURES

Table 25.—Aleutian Islands golden king crab commercial fishery harvest data, 1981/82–2020/21.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^c | Pots lifted | CPUE ^e | Weight ^{a,f} | Length ^{f,g} |
| 1981/82 | East of 172°W | — | 115,715 | 8,752 | 6 | 16 | 22,666 | 2,906 | 8 | 5.1 | 158 |
| | West of 172°W | — | 1,194,046 | 22,064 | 14 | 76 | 217,700 | 24,627 | 9 | 5.5 | 160 |
| | TOTAL | — | 1,309,761 | 30,816 | ND | 92 | 240,366 | 27,533 | 9 | 5.4 | ND |
| 1982/83 | East of 172°W | — | 1,184,971 | 47,479 | 49 | 136 | 227,471 | 29,369 | 8 | 5.2 | 158 |
| | West of 172°W | — | 8,006,274 | 220,743 | 99 | 501 | 1,509,001 | 150,103 | 10 | 5.3 | 158 |
| | TOTAL | — | 9,191,245 | 268,222 | ND | 637 | 1,736,472 | 179,472 | 10 | 5.3 | ND |
| 1983/84 | East of 172°W | — | 1,810,973 | 45,268 | 47 | 132 | 328,353 | 29,595 | 11 | 5.5 | ND |
| | West of 172°W | — | 8,128,029 | 171,021 | 157 | 1,002 | 1,534,909 | 226,798 | 7 | 5.3 | ND |
| | TOTAL | — | 9,939,002 | 216,289 | ND | 1,134 | 1,863,262 | 256,393 | 7 | 5.3 | ND |
| 1984/85 | East of 171°W | — | 1,521,142 | 70,362 | 13 | 67 | 327,440 | 24,044 | 14 | 4.6 | 161 |
| | West of 171°W | — | 4,141,052 | 130,377 | 38 | 107 | 863,367 | 77,049 | 11 | 4.8 | 157 |
| | TOTAL | — | 5,662,194 | 200,739 | ND | 174 | 1,190,807 | 101,093 | 12 | 4.8 | ND |
| 1985/86 | East of 171°W | — | 1,733,878 | 25,223 | 11 | 59 | 364,097 | 31,322 | 12 | 4.8 | 156 |
| | West of 171°W | — | 11,024,759 | 197,753 | 53 | 386 | 2,452,216 | 205,279 | 12 | 4.5 | 151 |
| | TOTAL | — | 12,758,637 | 222,976 | ND | 445 | 2,816,313 | 236,601 | 12 | 4.5 | ND |
| 1986/87 | East of 171°W | — | 1,869,180 | 9,510 | 17 | 71 | 400,389 | 37,585 | 11 | 4.7 | ND |
| | West of 171°W | — | 12,869,564 | 276,741 | 62 | 528 | 2,940,238 | 395,435 | 7 | 4.4 | 150 |
| | TOTAL | — | 14,738,744 | 286,251 | ND | 599 | 3,340,627 | 433,020 | 8 | 4.4 | ND |
| 1987/88 | East of 171°W | — | 1,388,983 | 25,060 | 23 | 77 | 301,227 | 42,867 | 7 | 4.6 | 150 |
| | West of 171°W | — | 7,868,022 | 167,110 | 57 | 380 | 1,873,349 | 263,863 | 7 | 4.2 | 147 |
| | TOTAL | — | 9,257,005 | 192,170 | ND | 457 | 2,174,576 | 306,730 | 7 | 4.3 | ND |

-continued-

Table 25.—Page 2 of 6.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^c | Pots lifted | CPUE ^e | Weight ^{a,f} | Length ^{f,g} |
| 1988/89 | East of 171°W | — | 1,546,113 | 23,960 | 21 | 57 | 323,783 | 41,371 | 8 | 4.8 | 154 |
| | West of 171°W | — | 9,080,929 | 125,500 | 73 | 455 | 2,164,650 | 280,556 | 8 | 4.2 | 149 |
| | TOTAL | — | 10,627,042 | 149,460 | 76 | 512 | 2,488,433 | 321,927 | 8 | 4.3 | ND |
| 1989/90 | East of 171°W | — | 1,852,249 | 17,421 | 13 | 70 | 424,067 | 43,346 | 10 | 4.4 | 151 |
| | West of 171°W | — | 10,169,803 | 99,866 | 65 | 505 | 2,478,846 | 314,457 | 8 | 4.1 | 149 |
| | TOTAL | — | 12,022,052 | 117,287 | 68 | 575 | 2,902,913 | 357,803 | 8 | 4.1 | ND |
| 1990/91 | East of 171°W | — | 1,699,675 | 42,800 | 16 | 67 | 391,135 | 53,592 | 7 | 4.3 | 148 |
| | West of 171°W | — | 5,250,687 | 176,583 | 13 | 167 | 1,312,116 | 161,222 | 8 | 4.0 | 145 |
| | TOTAL | — | 6,950,362 | 219,383 | 23 | 234 | 1,703,251 | 214,814 | 8 | 4.1 | ND |
| 1991/92 | East of 171°W | — | 1,516,779 | 45,100 | 11 | 56 | 352,803 | 43,231 | 8 | 4.3 | 148 |
| | West of 171°W | — | 6,185,362 | 96,848 | 16 | 206 | 1,494,595 | 191,626 | 8 | 4.1 | 145 |
| | TOTAL | — | 7,702,141 | 141,948 | 19 | 258 | 1,847,398 | 234,857 | 8 | 4.2 | ND |
| 1992/93 | East of 171°W | — | 1,404,452 | 37,200 | 10 | 46 | 337,559 | 38,348 | 9 | 4.2 | 148 |
| | West of 171°W | — | 4,886,745 | 104,215 | 18 | 128 | 1,190,769 | 164,873 | 7 | 4.1 | 147 |
| | TOTAL | — | 6,291,197 | 141,415 | 22 | 174 | 1,528,328 | 203,221 | 8 | 4.1 | ND |
| 1993/94 | East of 171°W | — | 915,460 | 7,324 | 4 | 14 | 217,788 | 22,490 | 10 | 4.2 | 149 |
| | West of 171°W | — | 4,635,683 | 165,358 | 21 | 148 | 1,179,742 | 212,164 | 6 | 3.9 | 148 |
| | TOTAL | — | 5,551,143 | 172,682 | 21 | 162 | 1,397,530 | 234,654 | 6 | 4.0 | ND |
| 1994/95 | East of 171°W | — | 1,750,481 | 35,938 | 14 | 46 | 384,405 | 67,587 | 6 | 4.6 | 148 |
| | West of 171°W | — | 6,378,030 | 242,190 | 34 | 247 | 1,539,866 | 319,006 | 5 | 4.1 | 150 |
| | TOTAL | — | 8,128,511 | 278,128 | 35 | 293 | 1,924,271 | 386,593 | 5 | 4.2 | ND |

-continued-

Table 25.—Page 3 of 6.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^c | Pots lifted | CPUE ^c | Weight ^{a,f} | Length ^{f,g} |
| 1995/96 | East of 171°W | 1,500,000 | 1,993,980 | 65,156 | 17 | 42 | 431,867 | 65,030 | 7 | 4.6 | 150 |
| | West of 171°W | 5–6 million | 4,966,426 | 248,226 | 25 | 141 | 1,150,466 | 227,991 | 5 | 4.3 | 147 |
| | TOTAL | — | 6,960,406 | 313,382 | 28 | 183 | 1,582,333 | 293,021 | 5 | 4.4 | ND |
| 1996/97 | East of 174°W | 3,200,000 | 3,290,862 | 185,203 | 14 | 71 | 731,909 | 113,460 | 6 | 4.5 | ND |
| | West of 174°W | 2,700,000 | 2,524,910 | 75,506 | 13 | 99 | 602,968 | 99,267 | 6 | 4.2 | ND |
| | TOTAL | 5,900,000 | 5,815,772 | 260,709 | 18 | 166 | 1,334,877 | 212,727 | 6 | 4.4 | 147 |
| 1997/98 | East of 174°W | 3,200,000 | 3,501,055 | 131,481 | 13 | 74 | 780,610 | 106,403 | 7 | 4.5 | 147 |
| | West of 174°W | 2,700,000 | 2,444,628 | 79,564 | 9 | 96 | 569,550 | 86,811 | 7 | 4.3 | 148 |
| | TOTAL | 5,900,000 | 5,945,683 | 211,045 | 15 | 167 | 1,350,160 | 193,214 | 7 | 4.4 | 147 |
| 1998/99 | East of 174°W | 3,000,000 | 3,247,863 | 82,113 | 14 | 55 | 740,011 | 83,378 | 9 | 4.4 | 148 |
| | West of 174°W | 2,700,000 | 1,694,030 | 21,218 | 3 | 44 | 410,018 | 35,975 | 11 | 4.1 | 146 |
| | TOTAL | 5,700,000 | 4,941,893 | 103,331 | 16 | 99 | 1,150,029 | 119,353 | 10 | 4.3 | 147 |
| 1999/00 | East of 174°W | 3,000,000 | 3,069,886 | 67,574 | 15 | 60 | 709,332 | 79,129 | 9 | 4.3 | 147 |
| | West of 174°W | 2,700,000 | 2,768,902 | 104,852 | 15 | 113 | 676,558 | 107,040 | 6 | 4.1 | 147 |
| | TOTAL | 5,700,000 | 5,838,788 | 172,426 | 16 | 168 | 1,385,890 | 186,169 | 7 | 4.2 | 147 |
| 2000/01 | East of 174°W | 3,000,000 | 3,134,079 | 55,999 | 15 | 50 | 704,702 | 71,551 | 10 | 4.4 | 147 |
| | West of 174°W | 2,700,000 | 2,884,682 | 53,158 | 12 | 100 | 705,613 | 101,239 | 7 | 4.1 | 145 |
| | TOTAL | 5,700,000 | 6,018,761 | 109,157 | 17 | 149 | 1,410,315 | 172,790 | 8 | 4.3 | 146 |
| 2001/02 | East of 174°W | 3,000,000 | 3,178,652 | 50,030 | 19 | 45 | 730,030 | 62,639 | 12 | 4.4 | 147 |
| | West of 174°W | 2,700,000 | 2,740,054 | 43,519 | 9 | 90 | 686,738 | 105,512 | 7 | 4.0 | 145 |
| | TOTAL | 5,700,000 | 5,918,706 | 93,549 | 21 | 134 | 1,416,768 | 168,151 | 8 | 4.2 | 146 |

-continued-

Table 25.—Page 4 of 6.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|----------------------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^c | Pots lifted | CPUE ^e | Weight ^{a,f} | Length ^{f,g} |
| 2002/03 | East of 174°W | 3,000,000 | 2,821,851 | 55,425 | 19 | 43 | 643,886 | 52,042 | 12 | 4.4 | 148 |
| | West of 174°W | 2,700,000 | 2,640,604 | 32,101 | 6 | 73 | 664,823 | 78,979 | 8 | 4.0 | 146 |
| | TOTAL | 5,700,000 | 5,462,455 | 87,526 | 22 | 116 | 1,308,709 | 131,021 | 10 | 4.2 | 147 |
| 2003/04 | East of 174°W | 3,000,000 | 2,977,055 | 76,006 | 18 | 37 | 643,074 | 58,883 | 11 | 4.6 | 149 |
| | West of 174°W | 2,700,000 | 2,688,773 | 49,321 | 6 | 60 | 676,633 | 66,236 | 10 | 4.0 | 146 |
| | TOTAL | 5,700,000 | 5,665,828 | 125,327 | 21 | 96 | 1,319,707 | 125,119 | 11 | 4.3 | 147 |
| 2004/05 | East of 174°W | 3,000,000 | 2,886,817 | 43,576 | 19 | 32 | 637,536 | 34,848 | 18 | 4.5 | 148 |
| | West of 174°W | 2,700,000 | 2,688,234 | 43,560 | 6 | 51 | 685,465 | 56,846 | 12 | 3.9 | 146 |
| | TOTAL | 5,700,000 | 5,575,051 | 87,136 | 22 | 83 | 1,323,001 | 91,694 | 14 | 4.2 | 147 |
| 2005/06 ^h | East of 174°W | 3,000,000 | 2,866,602 | 26,962 | 7 | 39 | 623,966 | 24,569 | 25 | 4.6 | 151 |
| | West of 174°W | 2,700,000 | 2,653,716 | 30,873 | 3 | 47 | 639,370 | 30,116 | 21 | 4.2 | 148 |
| | TOTAL | 5,700,000 | 5,520,318 | 57,835 | 8 | 82 | 1,263,336 | 54,685 | 23 | 4.4 | 149 |
| 2006/07 | East of 174°W | 3,000,000 | 2,992,010 | 34,594 | 6 | 38 | 650,588 | 26,195 | 25 | 4.6 | 152 |
| | West of 174°W | 2,700,000 | 2,270,332 | 22,344 | 4 | 37 | 527,737 | 26,110 | 20 | 4.3 | 150 |
| | TOTAL | 5,700,000 | 5,262,342 | 56,938 | 7 | 74 | 1,178,325 | 52,305 | 23 | 4.5 | 150 |
| 2007/08 | East of 174°W | 3,000,000 | 2,989,997 | 21,558 | 4 | 42 | 633,253 | 22,653 | 28 | 4.7 | 153 |
| | West of 174°W | 2,700,000 | 2,518,103 | 24,870 | 3 | 39 | 600,595 | 29,950 | 20 | 4.2 | 149 |
| | TOTAL | 5,700,000 | 5,508,100 | 46,428 | 5 | 76 | 1,233,848 | 52,603 | 23 | 4.5 | 151 |
| 2008/09 | East of 174°W | 3,150,000 | 3,144,423 | 25,525 | 3 | 37 | 666,947 | 24,466 | 27 | 4.7 | 151 |
| | West of 174°W | 2,835,000 | 2,535,661 | 25,292 | 3 | 42 | 587,661 | 26,200 | 22 | 4.3 | 148 |
| | TOTAL | 5,985,000 | 5,680,084 | 50,817 | 5 | 79 | 1,254,608 | 50,666 | 25 | 4.5 | 149 |

-continued-

Table 25.—Page 5 of 6.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^c | Pots lifted | CPUE ^c | Weight ^{a,f} | Length ^{f,g} |
| 2009/10 | East of 174°W | 3,150,000 | 3,150,474 | 33,284 | 3 | 39 | 679,886 | 26,298 | 26 | 4.6 | 152 |
| | West of 174°W | 2,835,000 | 2,761,813 | 40,311 | 3 | 41 | 628,332 | 26,489 | 24 | 4.4 | 150 |
| | TOTAL | 5,985,000 | 5,912,287 | 73,595 | 5 | 80 | 1,308,218 | 52,787 | 25 | 4.5 | 150 |
| 2010/11 | East of 174°W | 3,150,000 | 3,148,188 | 71,519 | 3 | 35 | 670,981 | 25,851 | 26 | 4.7 | 153 |
| | West of 174°W | 2,835,000 | 2,820,661 | 39,727 | 3 | 38 | 626,246 | 29,944 | 21 | 4.5 | 149 |
| | TOTAL | 5,985,000 | 5,968,849 | 111,246 | 5 | 73 | 1,297,227 | 55,795 | 23 | 4.6 | 151 |
| 2011/12 | East of 174°W | 3,150,000 | 3,150,374 | 24,184 | 3 | 41 | 668,828 | 17,915 | 37 | 4.7 | 151 |
| | West of 174°W | 2,835,000 | 2,814,042 | 39,147 | 3 | 40 | 616,118 | 26,326 | 23 | 4.6 | 148 |
| | TOTAL | 5,985,000 | 5,964,416 | 63,331 | 5 | 81 | 1,284,946 | 44,241 | 29 | 4.6 | 149 |
| 2012/13 | East of 174°W | 3,310,000 | 3,315,115 | 79,434 | 3 | 45 | 687,666 | 20,827 | 33 | 4.8 | 153 |
| | West of 174°W | 2,980,000 | 2,952,644 | 55,627 | 4 | 36 | 672,916 | 32,716 | 21 | 4.4 | 150 |
| | TOTAL | 6,290,000 | 6,267,759 | 135,061 | 6 | 81 | 1,360,582 | 53,543 | 25 | 4.6 | 151 |
| 2013/14 | East of 174°W | 3,310,000 | 3,302,061 | 29,932 | 3 | 42 | 699,078 | 20,687 | 34 | 4.7 | 151 |
| | West of 174°W | 2,980,000 | 2,970,514 | 92,980 | 3 | 34 | 686,883 | 41,835 | 16 | 4.3 | 152 |
| | TOTAL | 6,290,000 | 6,272,575 | 122,912 | 5 | 76 | 1,385,961 | 62,522 | 22 | 4.5 | 151 |
| 2014/15 | East of 174°W | 3,310,000 | 3,307,016 | 29,676 | 3 | 33 | 693,474 | 16,406 | 42 | 4.8 | 152 |
| | West of 174°W | 2,980,000 | CF | CF | 2 | 44 | CF | CF | CF | CF | 148 |
| | TOTAL | 6,290,000 | CF | CF | 5 | 77 | CF | CF | CF | CF | 150 |
| 2015/16 | East of 174°W | 3,310,000 | 3,302,480 | 53,160 | 3 | 34 | 717,864 | 18,481 | 39 | 4.6 | 152 |
| | West of 174°W | 2,980,000 | CF | CF | 2 | 50 | CF | CF | CF | CF | 147 |
| | TOTAL | 6,290,000 | CF | CF | 5 | 84 | CF | CF | CF | CF | 150 |

-continued-

Table 25.—Page 6 of 6.

| Season | Location | GHL/TAC ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | | |
|---------|---------------|------------------------|------------------------|-----------------------|----------------------|----------|-------------------|-------------|-------------------|-----------------------|-----------------------|
| | | | | | Vessels ^d | Landings | Crab ^e | Pots lifted | CPUE ^e | Weight ^{a,f} | Length ^{f,g} |
| 2016/17 | East of 174°W | 3,310,000 | 3,307,162 | 65,366 | 4 | 38 | 753,332 | 23,401 | 32 | 4.4 | 152 |
| | West of 174°W | 2,235,000 | 2,236,651 | 88,769 | 3 | 37 | 543,796 | 38,118 | 14 | 4.1 | 148 |
| | TOTAL | 5,545,000 | 5,543,813 | 154,135 | 5 | 75 | 1,297,128 | 61,519 | 21 | 4.3 | 150 |
| 2017/18 | East of 174°W | 3,310,000 | 3,308,185 | 54,253 | 4 | 40 | 768,120 | 24,617 | 31 | 4.3 | 150 |
| | West of 174°W | 2,235,000 | 2,234,723 | 58,077 | 3 | 41 | 519,051 | 30,885 | 17 | 4.3 | 150 |
| | TOTAL | 5,545,000 | 5,542,908 | 112,330 | 5 | 81 | 1,287,171 | 55,502 | 23 | 4.3 | 150 |
| 2018/19 | East of 174°W | 3,856,000 | 3,854,105 | 49,918 | 3 | 47 | 896,184 | 24,481 | 37 | 4.3 | 149 |
| | West of 174°W | 2,500,000 | 2,501,344 | 52,921 | 3 | 36 | 578,221 | 29,156 | 20 | 4.3 | 151 |
| | TOTAL | 6,356,000 | 6,355,449 | 102,839 | 5 | 83 | 1,474,405 | 53,637 | 27 | 4.3 | 150 |
| 2019/20 | East of 174°W | 4,310,000 | 4,308,530 | 53,608 | 3 | 48 | 1,018,876 | 29,675 | 34 | 4.2 | 148 |
| | West of 174°W | 2,870,000 | 2,840,078 | 54,529 | 3 | 44 | 649,832 | 42,924 | 15 | 4.4 | 150 |
| | TOTAL | 7,180,000 | 7,148,608 | 108,137 | 5 | 92 | 1,668,708 | 72,599 | 23 | 4.3 | 149 |
| 2020/21 | East of 174°W | 3,650,000 | 3,650,255 | 32,004 | 3 | 47 | 863,288 | 28,833 | 30 | 4.2 | 148 |
| | West of 174°W | 2,960,000 | 2,792,835 | 62,166 | 3 | 38 | 682,107 | 46,701 | 15 | 4.1 | 150 |
| | TOTAL | 6,610,000 | 6,443,090 | 94,170 | 5 | 85 | 1,545,395 | 75,534 | 20 | 4.2 | 149 |

Note: En dashes indicate harvest limits were not set for these fisheries prior to 1995/96, and overall harvest limit was not set in 1995/96; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Guideline harvest level (GHL), total allowable catch (TAC) from 2005/06 forward.

^c Deadloss included.

^d Many vessels fished both east and west of 174° W long, and thus the total number of vessels reflects the entire Aleutian Islands.

^e Number of legal crab per pot lift.

^f Carapace length in millimeters.

^g Retained catch.

^h Crab rationalization begins.

Table 26.—Aleutian Islands golden king crab commercial fishery value and season dates, 1981/82–2020/21.

| Season | Location | Value | | Season length | | |
|---------|---------------|-----------------------|--------------|---------------|----------|------|
| | | Exvessel ^a | Total | Opened | Closed | Days |
| 1981/82 | East of 172°W | \$2.05 | \$219,274 | 11/01/81 | 01/15/82 | 76 |
| | West of 172°W | \$2.06 | \$2,414,283 | 11/01/81 | 06/15/82 | 227 |
| 1982/83 | East of 172°W | \$3.00 | \$3,412,476 | 11/01/82 | 02/15/83 | 107 |
| | West of 172°W | \$3.01 | \$23,434,448 | 11/01/82 | 04/15/83 | 166 |
| 1983/84 | East of 172°W | \$3.05 | \$5,385,400 | 11/01/83 | 02/15/84 | 107 |
| | West of 172°W | \$2.92 | \$23,234,463 | 11/10/83 | 04/15/84 | 158 |
| 1984/85 | East of 171°W | \$1.35 | \$1,958,553 | 07/01/84 | 02/15/85 | 230 |
| | West of 171°W | \$2.00 | \$8,021,350 | 11/10/84 | 07/08/85 | 241 |
| 1985/86 | East of 171°W | \$2.00 | \$3,417,310 | 07/01/85 | 10/31/85 | 123 |
| | West of 171°W | \$2.50 | \$27,067,515 | 11/01/85 | 08/15/86 | 288 |
| 1986/87 | East of 171°W | \$2.85 | \$5,300,060 | 07/01/86 | 12/31/86 | 184 |
| | West of 171°W | \$3.00 | \$37,778,469 | 11/01/86 | 08/15/87 | 288 |
| 1987/88 | East of 171°W | \$2.85 | \$3,887,181 | 07/01/87 | 09/02/87 | 64 |
| | West of 171°W | \$3.00 | \$23,102,736 | 11/01/87 | 08/15/88 | 289 |
| 1988/89 | East of 171°W | \$3.07 | \$4,672,313 | 09/01/88 | 12/04/88 | 95 |
| | West of 171°W | \$3.37 | \$30,181,208 | 11/01/88 | 08/15/89 | 288 |
| 1989/90 | East of 171°W | \$3.63 | \$6,668,081 | 09/01/89 | 02/15/90 | 168 |
| | West of 171°W | \$3.39 | \$34,140,693 | 11/01/89 | 08/15/90 | 288 |
| 1990/91 | East of 171°W | \$3.34 | \$5,530,986 | 09/01/90 | 11/09/90 | 70 |
| | West of 171°W | \$3.08 | \$15,644,967 | 11/01/90 | 08/15/91 | 288 |
| 1991/92 | East of 171°W | \$2.00 | \$2,943,358 | 09/01/91 | 11/15/91 | 76 |
| | West of 171°W | \$3.18 | \$19,338,288 | 11/01/91 | 08/15/92 | 289 |
| 1992/93 | East of 171°W | \$2.86 | \$3,906,365 | 09/01/92 | 11/17/92 | 78 |
| | West of 171°W | \$3.62 | \$17,299,699 | 11/01/92 | 08/15/93 | 288 |
| 1993/94 | East of 171°W | \$4.57 | \$4,154,668 | 09/01/93 | 03/01/94 | 182 |
| | West of 171°W | \$2.50 | \$11,175,813 | 11/01/93 | 08/15/94 | 288 |
| 1994/95 | East of 171°W | \$3.94 | \$6,757,257 | 09/01/94 | 10/28/94 | 58 |
| | West of 171°W | \$3.35 | \$20,536,409 | 11/01/94 | 08/15/95 | 288 |
| 1995/96 | East of 171°W | \$2.58 | \$4,983,453 | 09/01/95 | 10/09/95 | 39 |
| | West of 171°W | \$2.27 | \$10,715,103 | 11/01/95 | 08/15/96 | 289 |
| 1996/97 | East of 174°W | \$2.20 | \$6,835,280 | 09/01/96 | 12/25/96 | 116 |
| | West of 174°W | \$2.23 | \$5,463,529 | 09/01/96 | 08/31/97 | 365 |

-continued-

Table 26.—Page 2 of 3.

| Season | Location | Value | | Season length | | |
|----------------------|---------------|-----------------------|--------------|---------------|----------|------|
| | | Exvessel ^a | Total | Opened | Closed | Days |
| 1997/98 | East of 174°W | \$2.25 | \$7,581,682 | 09/01/97 | 11/24/97 | 85 |
| | West of 174°W | \$2.15 | \$5,087,987 | 09/01/97 | 08/31/98 | 365 |
| 1998/99 | East of 174°W | \$1.88 | \$5,952,446 | 09/01/98 | 11/07/98 | 68 |
| | West of 174°W | \$2.02 | \$3,377,002 | 09/01/98 | 08/31/99 | 365 |
| 1999/00 | East of 174°W | \$3.28 | \$9,853,086 | 09/01/99 | 10/25/99 | 55 |
| | West of 174°W | \$3.11 | \$8,284,343 | 09/01/99 | 08/14/00 | 349 |
| 2000/01 | East of 174°W | \$3.51 | \$10,806,859 | 08/15/00 | 09/24/00 | 41 |
| | West of 174°W | \$3.07 | \$8,682,694 | 08/15/00 | 05/28/01 | 287 |
| 2001/02 | East of 174°W | \$3.30 | \$10,324,453 | 08/15/01 | 09/10/01 | 27 |
| | West of 174°W | \$3.15 | \$8,487,366 | 08/15/01 | 03/30/02 | 228 |
| 2002/03 | East of 174°W | \$3.33 | \$9,199,835 | 08/15/02 | 09/07/02 | 24 |
| | West of 174°W | \$3.50 | \$9,117,906 | 08/15/02 | 03/08/03 | 206 |
| 2003/04 | East of 174°W | \$3.47 | \$10,065,228 | 08/15/03 | 09/08/03 | 25 |
| | West of 174°W | \$3.83 | \$10,109,101 | 08/15/03 | 02/06/04 | 176 |
| 2004/05 | East of 174°W | \$3.18 | \$9,039,137 | 08/15/04 | 08/29/04 | 15 |
| | West of 174°W | \$3.29 | \$8,706,763 | 08/15/04 | 01/03/05 | 142 |
| 2005/06 ^b | East of 174°W | \$2.51 | \$7,117,132 | 08/15/05 | 05/15/06 | 274 |
| | West of 174°W | \$2.12 | \$5,549,420 | 08/15/05 | 05/15/06 | 274 |
| 2006/07 | East of 174°W | \$1.71 | \$5,070,070 | 08/15/06 | 05/15/07 | 274 |
| | West of 174°W | \$1.32 | \$2,978,071 | 08/15/06 | 05/15/07 | 274 |
| 2007/08 | East of 174°W | \$2.14 | \$6,365,457 | 08/15/07 | 05/15/08 | 275 |
| | West of 174°W | \$1.79 | \$4,454,290 | 08/15/07 | 05/15/08 | 275 |
| 2008/09 | East of 174°W | \$3.42 | \$10,678,756 | 08/15/08 | 05/15/09 | 274 |
| | West of 174°W | \$1.91 | \$4,791,631 | 08/15/08 | 05/15/09 | 274 |
| 2009/10 | East of 174°W | \$1.98 | \$6,174,304 | 08/15/09 | 05/15/10 | 274 |
| | West of 174°W | \$1.96 | \$5,322,370 | 08/15/09 | 05/15/10 | 274 |
| 2010/11 | East of 174°W | \$3.03 | \$9,315,401 | 08/15/10 | 05/15/11 | 274 |
| | West of 174°W | \$3.53 | \$9,803,355 | 08/15/10 | 05/15/11 | 274 |
| 2011/12 | East of 174°W | \$3.80 | \$11,880,146 | 08/15/11 | 05/15/12 | 275 |
| | West of 174°W | \$3.72 | \$10,313,779 | 08/15/11 | 05/15/12 | 275 |
| 2012/13 | East of 174°W | \$3.47 | \$11,218,989 | 08/15/12 | 05/15/13 | 274 |
| | West of 174°W | \$3.30 | \$9,554,574 | 08/15/12 | 05/15/13 | 274 |

-continued-

Table 26.—Page 3 of 3.

| Season | Location | Value | | Season length | | |
|---------|---------------|-----------------------|--------------|---------------|----------|------|
| | | Exvessel ^a | Total | Opened | Closed | Days |
| 2013/14 | East of 174°W | \$3.48 | \$11,376,784 | 08/15/13 | 05/15/14 | 274 |
| | West of 174°W | \$3.50 | \$10,081,665 | 08/15/13 | 05/15/14 | 274 |
| 2014/15 | East of 174°W | \$3.34 | \$10,936,484 | 08/15/14 | 05/15/15 | 274 |
| | West of 174°W | CF | CF | 08/15/14 | 05/15/15 | 274 |
| 2015/16 | East of 174°W | \$3.64 | \$11,815,476 | 08/01/15 | 04/30/16 | 274 |
| | West of 174°W | CF | CF | 08/01/15 | 04/30/16 | 274 |
| 2016/17 | East of 174°W | \$4.52 | \$14,660,890 | 08/01/16 | 04/30/17 | 273 |
| | West of 174°W | \$4.50 | \$9,664,768 | 08/01/16 | 04/30/17 | 273 |
| 2017/18 | East of 174°W | \$3.59 | \$11,691,725 | 08/01/17 | 04/30/18 | 273 |
| | West of 174°W | \$3.67 | \$7,997,779 | 08/01/17 | 04/30/18 | 273 |
| 2018/19 | East of 174°W | \$4.50 | \$17,118,842 | 08/01/18 | 04/30/19 | 273 |
| | West of 174°W | \$4.49 | \$10,987,299 | 08/01/18 | 04/03/19 | 273 |
| 2019/20 | East of 174°W | \$4.64 | \$19,740,830 | 07/15/19 | 04/30/20 | 291 |
| | West of 174°W | \$4.50 | \$12,534,971 | 07/15/19 | 04/30/20 | 291 |
| 2020/21 | East of 174°W | \$4.56 | \$16,492,203 | 08/01/20 | 04/30/21 | 273 |
| | West of 174°W | \$4.51 | \$12,311,834 | 08/01/20 | 05/13/21 | 286 |

Note: CF indicates confidential data.

^a Average price per pound.

^b Crab rationalization begins.

Table 27.—Aleutian Islands golden king crab commercial fishery harvest by statistical area east of 174° W long, 2020/21.

| Statistical area | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | |
|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} |
| 705232 | 714,964 | 3,400 | 3 | 32 | 167,657 | 5,470 | 31 | 4.3 |
| 715202 | 627,428 | 5,940 | 3 | 36 | 147,940 | 5,054 | 29 | 4.2 |
| 715232 | 237,949 | 2,603 | 3 | 31 | 56,834 | 1,699 | 33 | 4.2 |
| Other ^c | 2,069,915 | 20,060 | 3 | 47 | 490,857 | 16,610 | 30 | 4.2 |
| Total | 3,650,255 | 32,004 | 3 | 47 | 863,288 | 28,833 | 30 | 4.2 |

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Combination of statistical areas (19) in which landings were made by fewer than three vessels.

Table 28.—Eastern Aleutian Islands golden king crab cost-recovery harvest data and charter length, 2013–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Charter Length | |
|--------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|--------------------------|------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Dates | Days |
| 2013 | 106,978 | 1,344 | 1 | 1 | 21,142 | 701 | 30 | 5.1 | 09/09–09/17 | 9 |
| 2014 | 119,005 | 894 | 1 | 1 | 25,590 | 596 | 43 | 4.7 | 09/07–09/17 | 11 |
| 2015 | 202,169 | 3,756 | 1 | 1 | 45,740 | 895 | 51 | 4.4 | 09/08–09/17 | 10 |
| 2016 | 172,367 | 3,448 | 1 | 1 | 40,651 | 1,069 | 38 | 4.2 | 08/28–09/07 | 11 |
| 2017 | 156,244 | 3,339 | 1 | 1 | 34,490 | 899 | 38 | 4.5 | 09/05–09/16 | 12 |
| 2018 | 180,137 | 5,312 | 1 | 1 | 44,152 | 1,072 | 41 | 4.1 | 09/04–09/14 | 11 |
| 2019 | 168,245 | 2,682 | 1 | 1 | 38,588 | 1,323 | 29 | 4.4 | 08/28–09/07 | 11 |
| 2020 | 170,863 | 1,904 | 1 | 2 | 38,833 | 1,239 | 31 | 4.4 | 09/06–09/08, 09/10–09/20 | 14 |

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

Table 29.—Eastern Aleutian District Tanner crab commercial fishery harvest data, 1973/74–2020.

| Season | Location | GHL ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | |
|-------------------|----------|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} |
| 1973/74 | — | — | 498,836 | ND | 6 | 14 | 210,539 | ND | 60 | 2.4 |
| 1974/75 | — | — | CF | CF | 2 | 2 | CF | CF | CF | CF |
| 1975/76 | — | — | 534,295 | ND | 8 | 13 | 219,166 | 4646 | 47 | 2.4 |
| 1976/77 | — | — | 1,239,569 | ND | 12 | 35 | 544,755 | 9640 | 57 | 2.3 |
| 1977/78 | — | — | 2,494,631 | ND | 15 | 198 | 1,104,631 | 29855 | 37 | 2.3 |
| 1978/79 | — | — | 1,280,115 | ND | 20 | 174 | 542,081 | 18618 | 29 | 2.4 |
| 1979/80 | — | — | 886,487 | ND | 18 | 107 | 352,819 | 18040 | 20 | 2.5 |
| 1981 | — | — | 654,514 | ND | 29 | 119 | 264,238 | 21771 | 12 | 2.5 |
| 1982 | — | — | 739,694 | ND | 31 | 138 | 332,260 | 30109 | 11 | 2.2 |
| 1983 | — | — | 547,830 | ND | 23 | 107 | 250,774 | 22168 | 11 | 2.2 |
| 1984 | — | — | 239,585 | ND | 16 | 91 | 104,761 | 11069 | 9 | 2.3 |
| 1985 | — | — | 181,407 | 60 | 7 | 56 | 78,930 | 6295 | 13 | 2.3 |
| 1986 | — | — | 167,339 | 400 | 8 | 37 | 73,187 | 10244 | 7 | 2.3 |
| 1987 | — | — | 162,097 | 115 | 8 | 65 | 72,098 | 5915 | 12 | 2.2 |
| 1988 | — | — | 309,918 | 2,000 | 20 | 130 | 129,478 | 11011 | 12 | 2.4 |
| 1989 | — | — | 326,196 | 2,300 | 12 | 108 | 144,593 | 14615 | 10 | 2.3 |
| 1990 | — | — | 155,648 | 0 | 10 | 75 | 68,859 | 6858 | 10 | 2.3 |
| 1991 | — | — | 50,038 | 0 | 5 | 27 | 21,511 | 1849 | 12 | 2.3 |
| 1992 | — | — | 98,703 | 0 | 4 | 29 | 42,096 | 2963 | 14 | 2.3 |
| 1993 | — | — | 118,609 | 0 | 7 | 34 | 51,441 | 3530 | 15 | 2.3 |
| 1994 | — | — | 166,080 | 40 | 8 | 119 | 71,760 | 6303 | 11 | 2.3 |
| 1995–2002 | | | | No Commercial Fishery | | | | | | |
| 2003 ^f | — | — | 15,138 | 9 | 3 | 10 | 6,695 | 191 | 35 | 2.3 |

-continued-

Table 29.—Page 2 of 3.

| Season | Location | GHL ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | |
|--------|-------------------|--------------------|------------------------|-----------------------|-----------------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} |
| 2004 | Unalaska Bay | 47,219 | CF | CF | 10 | 48 | CF | CF | CF | CF |
| | Makushin/Skan Bay | 87,891 | CF | CF | 6 | 13 | CF | CF | CF | CF |
| | TOTAL | 135,110 | CF | CF | 14 ^g | 61 | CF | CF | CF | CF |
| 2005 | Unalaska Bay | 35,304 | 34,022 | 0 | 25 | 79 | 14,249 | 696 | 20 | 2.4 |
| 2006 | Makushin/Skan Bay | 87,241 | CF | CF | 10 | 32 | CF | CF | CF | CF |
| 2007 | Akutan Bay | 35,000 | CF | CF | 3 | 7 | CF | CF | CF | CF |
| | Unalaska Bay | 49,000 | CF | CF | 12 | 41 | CF | CF | CF | CF |
| | TOTAL | 84,000 | CF | CF | 13 ^g | 47 | CF | CF | CF | CF |
| 2008 | Unalaska Bay | 60,000 | CF | CF | 11 | 48 | CF | CF | CF | CF |
| 2009 | Akutan Bay | 35,000 | CF | CF | 1 | 2 | CF | CF | CF | CF |
| | Makushin/Skan Bay | 35,000 | CF | CF | 1 | 3 | CF | CF | CF | CF |
| | Unalaska Bay | 58,000 | CF | CF | 10 | 83 | CF | CF | CF | CF |
| | TOTAL | 128,000 | CF | CF | 11 ^g | 88 | CF | CF | CF | CF |
| 2010 | Akutan Bay | 45,000 | CF | CF | 3 | 3 | CF | CF | CF | CF |
| | Unalaska Bay | 74,000 | CF | CF | 7 | 63 | CF | CF | CF | CF |
| | TOTAL | 119,000 | CF | CF | 8 ^g | 66 | CF | CF | CF | CF |
| 2011 | Akutan Bay | 35,000 | CF | CF | 2 | 3 | CF | CF | CF | CF |
| | Makushin/Skan Bay | 35,000 | CF | CF | 3 | 4 | CF | CF | CF | CF |
| | TOTAL | 70,000 | CF | CF | 3 ^g | 7 | CF | CF | CF | CF |
| 2012 | Makushin/Skan Bay | 35,000 | CF | CF | 1 | 6 | CF | CF | CF | CF |
| 2013 | Unalaska Bay | 35,000 | CF | CF | 6 | 28 | CF | CF | CF | CF |
| 2014 | | | | No Commercial Fishery | | | | | | |
| 2015 | Makushin/Skan Bay | 35,000 | CF | CF | 1 | 5 | CF | CF | CF | CF |

-continued-

Table 29.—Page 3 of 3.

| Season | Location | GHL ^{a,b} | Harvest ^{a,c} | Deadloss ^a | Number of | | | | Average | |
|-----------|-------------------|--------------------|------------------------|-----------------------|-----------|----------|-------------------|-------------|-------------------|-----------------------|
| | | | | | Vessels | Landings | Crab ^c | Pots lifted | CPUE ^d | Weight ^{a,e} |
| 2016 | Makushin/Skan Bay | 35,000 | CF | CF | 1 | 4 | CF | CF | CF | CF |
| 2017 | | | | No Commercial Fishery | | | | | | |
| 2018 | Makushin/Skan Bay | 35,000 | CF | CF | 2 | 8 | CF | CF | CF | CF |
| 2019–2020 | | | | No Commercial Fishery | | | | | | |

Note: En dashes indicate harvest limits and fishery management sections were not established for this fishery prior to 2004; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Guideline harvest level (GHL).

^c Deadloss included beginning 1980.

^d Number of legal crab per pot lift.

^e Retained catch.

^f January/February survey (fish ticket harvest code 15, exploratory shellfish harvest).

^g Vessel(s) participated in multiple sections.

Table 30.—Eastern Aleutian District Tanner crab commercial fishery value and season dates, 1973/74–2020.

| Season | Location | Value | | Season length | | |
|-----------|-------------------|-----------------------|-----------|---------------|----------|------|
| | | Exvessel ^a | Total | Opened | Closed | Days |
| 1973/74 | — | ND | ND | 10/01/73 | 07/31/74 | 304 |
| 1974/75 | — | CF | CF | 01/18/74 | 10/15/75 | 636 |
| 1975/76 | — | \$0.20 | \$106,859 | 01/20/75 | 10/15/76 | 635 |
| 1976/77 | — | \$0.30 | \$371,871 | 11/07/76 | 06/15/77 | 221 |
| 1977/78 | — | \$0.38 | \$947,960 | 11/01/77 | 06/15/78 | 227 |
| 1978/79 | — | \$0.52 | \$665,660 | 11/01/78 | 06/15/79 | 227 |
| 1979/80 | — | \$0.52 | \$460,973 | 11/01/79 | 06/15/80 | 228 |
| 1981 | — | \$0.58 | \$379,618 | 01/15/81 | 06/15/81 | 152 |
| 1982 | — | \$1.25 | \$924,618 | 02/15/82 | 06/15/82 | 121 |
| 1983 | — | \$1.20 | \$657,396 | 02/15/83 | 06/15/83 | 121 |
| 1984 | — | \$0.98 | \$234,793 | 02/15/84 | 06/15/84 | 122 |
| 1985 | — | \$0.96 | \$174,093 | 01/15/85 | 06/15/85 | 152 |
| 1986 | — | \$1.66 | \$277,119 | 01/15/86 | 06/15/86 | 152 |
| 1987 | — | \$2.03 | \$328,823 | 01/15/87 | 06/15/87 | 152 |
| 1988 | — | \$2.18 | \$671,261 | 01/15/88 | 04/10/88 | 87 |
| 1989 | — | \$2.72 | \$880,997 | 01/15/89 | 05/07/89 | 113 |
| 1990 | — | \$1.97 | \$306,627 | 01/15/90 | 04/09/90 | 85 |
| 1991 | — | \$1.25 | \$62,548 | 01/15/91 | 03/31/91 | 76 |
| 1992 | — | \$2.07 | \$204,315 | 01/15/92 | 03/31/92 | 77 |
| 1993 | — | \$1.70 | \$201,635 | 01/15/93 | 03/31/93 | 76 |
| 1994 | — | \$2.11 | \$350,344 | 01/15/94 | 03/31/94 | 76 |
| 1995–2003 | | No Commercial Fishery | | | | |
| 2004 | Unalaska Bay | CF | CF | 01/15/04 | 01/19/04 | 5 |
| | Makushin/Skan Bay | CF | CF | 01/15/04 | 02/03/04 | 20 |
| 2005 | Unalaska Bay | \$2.58 | \$87,777 | 01/15/05 | 01/18/05 | 4 |
| 2006 | Makushin/Skan Bay | CF | CF | 01/15/06 | 01/21/06 | 7 |
| 2007 | Akutan Bay | CF | CF | 01/15/07 | 03/31/07 | 76 |
| | Unalaska Bay | CF | CF | 01/15/07 | 01/19/07 | 5 |
| 2008 | Unalaska Bay | CF | CF | 01/15/08 | 01/29/08 | 15 |
| 2009 | Akutan Bay | CF | CF | 01/15/09 | 03/31/09 | 76 |
| | Makushin/Skan Bay | CF | CF | 01/15/09 | 03/31/09 | 76 |
| | Unalaska Bay | CF | CF | 01/15/09 | 02/11/09 | 28 |
| 2010 | Akutan Bay | CF | CF | 01/15/10 | 03/31/10 | 76 |
| | Unalaska Bay | CF | CF | 01/15/10 | 02/10/10 | 27 |

–continued–

Table 30.—Page 2 of 2.

| Season | Location | Value | | Season length | | |
|-----------|-------------------|-----------------------|-------|---------------|----------|------|
| | | Exvessel ^a | Total | Opened | Closed | Days |
| 2011 | Akutan Bay | CF | CF | 01/15/11 | 03/31/11 | 76 |
| | Makushin/Skan Bay | CF | CF | 01/15/11 | 03/18/11 | 63 |
| 2012 | Makushin/Skan Bay | CF | CF | 01/15/12 | 02/10/12 | 27 |
| 2013 | Unalaska Bay | CF | CF | 01/15/13 | 01/26/13 | 12 |
| 2014 | | No Commercial Fishery | | | | |
| 2015 | Makushin/Skan Bay | CF | CF | 01/15/15 | 02/13/15 | 30 |
| 2016 | Makushin/Skan Bay | CF | CF | 01/15/16 | 02/21/16 | 38 |
| 2017 | | No Commercial Fishery | | | | |
| 2018 | Makushin/Skan Bay | CF | CF | 01/15/18 | 03/31/18 | 76 |
| 2019–2020 | | No Commercial Fishery | | | | |

Note: En dashes indicate fishery management sections were not established for this fishery prior to 2004; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a Average price per pound.

Table 31.—Subsistence king and Tanner crab harvest from the Eastern Aleutian Islands, west of Scotch Cap Light and east of 168°W long, 1999–2020.

| Year | Permits | | | Harvest ^a | |
|-------------------|---------------|-----------------|------------------|----------------------|----------------------|
| | Number issued | Number returned | Percent returned | King crab reported | Tanner crab reported |
| 1999 | 179 | 80 | 45% | 787 | 1,432 |
| 2000 | 193 | 137 | 71% | 523 | 916 |
| 2001 | 200 | 153 | 77% | 1,149 | 1,703 |
| 2002 | 231 | 179 | 77% | 1,080 | 2,451 |
| 2003 | 229 | 160 | 70% | 387 | 4,600 |
| 2004 | 225 | 144 | 64% | 225 | 4,714 |
| 2005 | 241 | 182 | 76% | 866 | 5,447 |
| 2006 | 256 | 185 | 72% | 1,796 | 1,439 |
| 2007 | 203 | 122 | 60% | 1,359 | 1,542 |
| 2008 | 94 | 64 | 68% | 199 | 174 |
| 2009 | 72 | 51 | 71% | 186 | 634 |
| 2010 | 72 | 39 | 54% | 46 | 487 |
| 2011 | 30 | 20 | 67% | 23 | 322 |
| 2012 | 44 | 36 | 82% | 76 | 592 |
| 2013 | 161 | 102 | 63% | 537 | 1,845 |
| 2014 | 186 | 100 | 54% | 203 | 1,541 |
| 2015 | 180 | 106 | 59% | 73 | 1,570 |
| 2016 | 188 | 111 | 59% | 100 | 2,239 |
| 2017 | 152 | 94 | 62% | 29 | 1,335 |
| 2018 | 131 | 100 | 76% | 22 | 663 |
| 2019 | 140 | 77 | 55% | 32 | 428 |
| 2020 | 156 | 56 | 36% | 9 | 119 |
| 1999–2020 Average | 162 | 104 | 64% | 441 | 1,645 |

^a Reported harvest, in number of crab, from waters surrounding Unalaska Island.

^b Includes permits issued for both shellfish and salmon prior to 2008.

Table 32.—Aleutian District Dungeness crab commercial fishery data, 1974–2020.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|-----------------|------------------------|-----------------------|-----------|----------|------------------------------|-------------|-------------------|-----------------------|-----------------------|-----------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 1974 | 60,517 | ND | 3 | 13 | 24,459 | 3,399 | 7 | 2.5 | ND | ND |
| 1975 | CF | CF | 1 | 3 | CF | CF | CF | CF | CF | CF |
| 1976–1977 | | | | | No Commercial Fishing Effort | | | | | |
| 1978 | CF | CF | 2 | 9 | CF | CF | CF | CF | CF | CF |
| 1979 | CF | CF | 1 | 4 | CF | CF | CF | CF | CF | CF |
| 1980–1981/82 | | | | | No Commercial Fishing Effort | | | | | |
| 1982/83 | CF | CF | 2 | 9 | CF | CF | CF | CF | CF | CF |
| 1983/84 | CF | CF | 2 | 14 | CF | CF | CF | CF | CF | CF |
| 1984/85 | 91,739 | ND | 4 | 50 | 40,128 | 13,555 | 3 | 2.3 | \$1.33 | \$122,013 |
| 1985/86 | 17,830 | 0 | 4 | 19 | 8,590 | 1,706 | 5 | 2.1 | ND | ND |
| 1986/87 | CF | CF | 2 | 9 | CF | CF | CF | CF | CF | CF |
| 1987/88 | 26,627 | 0 | 6 | 43 | 13,247 | 2,987 | 4 | 2 | \$0.95 | \$25,296 |
| 1988/89 | 22,915 | 4 | 6 | 45 | 10,956 | 2,599 | 4 | 2.1 | \$0.81 | \$18,558 |
| 1989/90 | 11,124 | 0 | 4 | 31 | 5,165 | 2,078 | 2 | 2.2 | \$0.91 | \$10,123 |
| 1990/91 | 17,482 | 117 | 3 | 11 | 8,379 | 1,345 | 6 | 2.1 | \$1.20 | \$20,838 |
| 1991/92 | 7,412 | 0 | 4 | 14 | 3,654 | 732 | 5 | 2 | \$1.25 | \$9,265 |
| 1992/93 | 5,649 | 0 | 4 | 13 | 2,854 | 555 | 5 | 2 | \$0.83 | \$4,689 |
| 1993/94 | 7,531 | 10 | 5 | 12 | 3,448 | 797 | 4 | 2.2 | \$0.78 | \$5,866 |
| 1994/95 | | | | | No Commercial Fishing Effort | | | | | |
| 1995/96 | CF | CF | 2 | 2 | CF | CF | CF | CF | CF | CF |
| 1996/97 | | | | | No Commercial Fishing Effort | | | | | |
| 1997/98 | CF | CF | 2 | 4 | CF | CF | CF | CF | CF | CF |
| 1998/99–2000/01 | | | | | No Commercial Fishing Effort | | | | | |

-continued-

Table 32.–Page 2 of 2.

| Season | Harvest ^{a,b} | Deadloss ^a | Number of | | | | Average | | Value | |
|-----------------|------------------------|-----------------------|----------------|------------------------------|-------------------|-------------|-------------------|-----------------------|-----------------------|-------|
| | | | Vessels | Landings | Crab ^b | Pots lifted | CPUE ^c | Weight ^{a,d} | Exvessel ^e | Total |
| 2001/02 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2002/03 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2003/04–2004/05 | | | | No Commercial Fishing Effort | | | | | | |
| 2005/06 | CF | CF | 1 | 1 | CF | CF | CF | CF | CF | CF |
| 2006/07–2014 | | | | No Commercial Fishing Effort | | | | | | |
| 2015 | CF | CF | 1 ^f | 1 | CF | CF | CF | CF | CF | CF |
| 2016–2020 | | | | No Commercial Fishing Effort | | | | | | |

Note: CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds.

^b Deadloss included.

^c Number of legal crab per pot lift.

^d Retained catch.

^e Average price per pound.

^f Harvested from a dock.

Table 33.—Dutch Harbor (Area O) weathervane scallop commercial fishery harvest data, 1982–2020/21.

| Season | GHL ^{a,b} | Harvest ^a | Number of | | Meat weight CPUE ^c |
|-----------------|--------------------|----------------------|------------------------------|--------------|----------------------------------|
| | | | Vessels | Dredge hours | |
| 1982 | – | 62,105 | 5 | ND | ND |
| 1983 | – | | No Commercial Fishing Effort | | |
| 1984 | – | | No Commercial Fishing Effort | | |
| 1985 | – | 547,164 | 2 | ND | ND |
| 1986 | – | 406,642 | 5 | ND | ND |
| 1987 | – | CF | 2 | CF | CF |
| 1988 | – | CF | 1 | CF | CF |
| 1989 | – | CF | 1 | CF | CF |
| 1990 | – | CF | 1 | CF | CF |
| 1991 | – | CF | 1 | CF | CF |
| 1992 | – | CF | 1 | CF | CF |
| 1993/94 | 170,000 | 39,346 | 3 | ND | ND |
| 1994/95 | 170,000 | 1,931 | 3 | 81 | 24 |
| 1995/96 | 170,000 | 26,950 | 1 | 1,047 | 26 |
| 1996/97 | 170,000 | | No Commercial Fishing Effort | | |
| 1997/98 | 170,000 | 5,790 | 1 | 160 | 36 |
| 1998/99 | 110,000 | 46,432 | 4 | 941 | 49 |
| 1999/00 | 110,000 | 6,465 | 1 | 278 | 23 |
| 2000/01–2001/02 | | | No Commercial Fishery | | |
| 2002/03 | 10,000 | 6,000 | 1 | 184 | 33 |
| 2003/04–2007/08 | | | No Commercial Fishery | | |
| 2008/09 | 10,000 | 10,040 | 1 | 225 | 45 |
| 2009/10 | 10,000 | 6,080 | 1 | 104 | 58 |
| 2010/11 | 10,000 | 5,640 | 1 | 83 | 68 |
| 2011/12 | 10,000 | 5,570 | 1 | 77 | 72 |
| 2012/13 | 5,000 | 5,100 | 1 | 64 | 80 |
| 2013/14 | 5,000 | 5,225 | 1 | 56 | 93 |
| 2014/15 | 5,000 | 5,160 | 1 | 73 | 71 |
| 2015/16 | 10,000 | 5,040 | 1 | 157 | 32 |
| 2016/17 | 10,000 | 5,050 | 1 | 104 | 49 |
| 2017/18 | 10,000 | 285 | 1 | 24 | 12 |
| 2018/19 | 5,000 | 325 | 1 | 24 | 14 |
| 2019/20 | 5,000 | 2,625 | 1 | 130 | 20 |
| 2020/21 | 5,000 | | No Commercial Fishing Effort | | |

-continued-

Table 33.—Page 2 of 2.

Note: En dashes indicate harvest limits were not established for this fishery prior to 1993/94; CF indicates confidential data; ND indicates these data were not collected or cannot be derived.

^a In pounds of shucked scallop meat.

^b Guideline harvest level (GHL) began in 1993/94.

^c Pounds of shucked scallop meat per dredge hour.

Table 34.—Aleutian Islands commercial octopus incidental harvest in groundfish fisheries, 1996–2020.

| Season | State waters | | | State and federal waters | | | | |
|-------------------|--------------|----------|---------------------------|--------------------------|----------|---------------------------|-----------------|-----------------------------|
| | Vessels | Landings | Whole pounds ^a | Vessels | Landings | Whole pounds ^a | At-sea discards | Exvessel value ^b |
| 1996 ^c | 26 | 87 | 36,292 | 68 | 281 | 97,085 | 29,925 | \$0.39 |
| 1997 | 19 | 44 | 22,431 | 61 | 235 | 98,497 | 22,880 | \$0.33 |
| 1998 ^c | 16 | 44 | 18,375 | 46 | 195 | 54,979 | 22,554 | \$0.05 |
| 1999 | 32 | 76 | 87,420 | 60 | 268 | 152,075 | 36,292 | \$0.32 |
| 2000 | 24 | 37 | 5,911 | 69 | 286 | 71,957 | 49,832 | \$0.17 |
| 2001 | 19 | 47 | 7,120 | 69 | 254 | 99,521 | 67,936 | \$0.02 |
| 2002 | 12 | 21 | 3,063 | 56 | 199 | 96,586 | 68,752 | \$0.02 |
| 2003 | 27 | 89 | 102,104 | 69 | 329 | 288,020 | 27,011 | \$0.55 |
| 2004 ^c | 38 | 135 | 151,205 | 76 | 459 | 998,731 | 97,750 | \$0.62 |
| 2005 ^c | 22 | 82 | 57,552 | 53 | 336 | 439,500 | 37,163 | \$0.50 |
| 2006 | 33 | 114 | 133,182 | 64 | 346 | 454,847 | 92,777 | \$0.46 |
| 2007 | 31 | 96 | 46,346 | 70 | 295 | 102,291 | 13,895 | \$0.39 |
| 2008 ^c | 26 | 45 | 35,480 | 56 | 151 | 142,938 | 24,713 | \$0.40 |
| 2009 | 13 | 21 | 8,782 | 41 | 91 | 28,038 | 5,192 | \$0.23 |
| 2010 | 21 | 48 | 42,376 | 54 | 184 | 168,964 | 96,533 | \$0.20 |
| 2011 | 13 | 17 | 8,187 | 53 | 204 | 209,636 | 181,825 | \$0.23 |
| 2012 | 18 | 58 | 14,917 | 57 | 182 | 76,141 | 38,129 | \$0.12 |
| 2013 | 20 | 129 | 50,309 | 50 | 202 | 116,103 | 16,671 | \$0.01 |
| 2014 | 16 | 136 | 65,637 | 56 | 293 | 212,019 | 48,664 | \$0.26 |
| 2015 | 17 | 84 | 41,258 | 69 | 288 | 168,133 | 36,229 | \$0.26 |
| 2016 | 21 | 89 | 36,128 | 93 | 386 | 158,482 | 36,900 | \$0.18 |
| 2017 | 15 | 124 | 114,445 | 86 | 350 | 171,380 | 27,350 | \$0.03 |
| 2018 ^c | 28 | 165 | 185,615 | 93 | 395 | 494,093 | 30,617 | \$0.51 |
| 2019 | 33 | 194 | 167,465 | 85 | 376 | 283,128 | 43,505 | \$0.50 |
| 2020 | 44 | 253 | 228,716 | 105 | 465 | 334,728 | 38,744 | \$0.37 |

^a Includes discards.

^b Average price per pound, based on landed weight.

^c Includes directed octopus harvest from Commissioner's permit fishery.




Figure 11.—Aleutian Islands king crab commercial fishery Registration Area O.




Figure 12.—Eastern Aleutian Islands golden king crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1981/82–2020/21.




Figure 13.—Western Aleutian Islands golden king crab commercial fishery harvest, catch per unit effort (CPUE; number legal males per pot), and number of vessels, 1981/82–2020/21.




Figure 14.—Eastern Aleutian District Tanner crab sections of Registration Area J.

SECTION III: BERING SEA/ALEUTIAN ISLANDS COMMUNITY DEVELOPMENT QUOTA AND ADAK COMMUNITY ALLOCATION SHELLFISH FISHERIES

DESCRIPTION OF AREA

Bering Sea Community Development Quota (CDQ) crab fisheries occur within the state waters of Alaska (0–3 nmi) and the Exclusive Economic Zone (EEZ; 3–200 nmi) north of Cape Sarichef (54°36'N lat), south of Cape Prince of Wales (65°49'N lat), and east of the U.S.–Russia Maritime Boundary Line, including waters of Bristol Bay. For CDQ crab fisheries, managed by the Alaska Department of Fish and Game (ADF&G) Dutch Harbor office, Cape Romanzof (61°49'N lat) is the northern boundary (Figures 1, 3, and 5).

Aleutian Islands CDQ and Adak Community Allocation (ACA) crab fisheries encompass state waters of Alaska (0–3 nmi) and waters of the Exclusive Economic Zone (EEZ; 3–200 nmi). The CDQ crab fisheries eastern boundary is the longitude of Scotch Cap Light (164°44'W long); the northern boundary is from Cape Sarichef (54°36'N lat) to 171°W long, then north to 55°30'N lat; and the western boundary is the U.S.–Russia Maritime Boundary Agreement Line. The ACA fishery occurs west of 174°W long (Figure 11).

PROGRAM BACKGROUND

The Alaska Board of Fisheries (BOF) adopted regulations for BSAI king and Tanner crab CDQ fisheries in 1997, and the first CDQ crab fisheries took place in 1998. With the implementation of crab rationalization in 2005, the BOF adopted regulations to implement changes to the CDQ management program (5 AAC 39.690), including the addition of Aleutian Islands crab fisheries to the CDQ crab program. ADF&G manages the CDQ crab fisheries with federal oversight.

Sixty-five western Alaska coastal communities aligned into six CDQ organizations, collectively referred to as CDQ groups. The groups are as follows: Aleutian Pribilof Island Community Development Association (APICDA), Bristol Bay Economic Development Corporation (BBEDC), Central Bering Sea Fishermen's Association (CBSFA), Coastal Villages Region Fund (CVRF), Norton Sound Economic Development Corporation (NSEDG), and Yukon Delta Fisheries Development Association (YDFDA).

CDQ groups are nonprofit entities, which may have for-profit subsidiaries. Uses of CDQ funds vary widely between groups but often include fishing-related investments, scholarships, training, employment services, and other projects that are intended to benefit the communities and regions the CDQ groups represent. Some groups purchase equity in fishing vessels that harvest crab in both CDQ and individual fishing quota (IFQ) fisheries.

Each of the six CDQ groups participates in at least one CDQ fishery every year, although each group does not necessarily have an allocation for each fishery (Table 35). Groups may choose not to participate or transfer their allocation to another group. This report addresses all CDQ crab

fisheries histories and allocations except the Norton Sound CDQ red king crab fishery, which is managed by ADF&G's Arctic-Yukon-Kuskokwim Region.

In conjunction with the changes to the CDQ program during crab rationalization, the BOF adopted regulations for an Adak Community Allocation (ACA) Western Aleutian Islands golden king crab fishery. The program was established to benefit the community of Adak, who formed the Adak Community Development Corporation (ACDC). ACDC is a nonprofit entity that represents the community of Adak and has a board of directors elected by the residents of Adak. The ACA crab allocation is not a CDQ fishery because Adak is not a CDQ community. ACDC must submit a comprehensive plan to the Alaska Department of Commerce, Community, and Economic Development on the intended use of the ACA funds derived from harvesting the ACA golden king crab. The funds are intended for fisheries-related purposes and other projects to benefit the community of Adak.

The ACA is set at 10% of the TAC of the Western Aleutian Islands (west of 174°W long) golden king crab fishery. This fishery opened for the first time in August 2005.

CURRENT FISHERY

CDQ groups are required to submit preseason fishery plans to ADF&G prior to each CDQ crab fishery. Fishery plans include information such as participating vessels and their contact information, intended delivery locations, and the group's allocation including quota transfers to other CDQ entities. Almost all CDQ harvest is taken concurrently with IFQ harvest. Vessels use the same gear to harvest IFQ and CDQ crab.

TABLES

Table 35.—Community Development Quota (CDQ) and Adak Community Allocation (ACA) program percent allocation by fishery to each group.

| Fishery | Percent allocation by group ^a | | | | | | |
|--|--|-------|-------|------|-------|-------|------|
| | APICDA | BBEDC | CBSFA | CVRF | NSEDC | YDFDA | ACDC |
| Bristol Bay red king crab | 17 | 19 | 10 | 18 | 18 | 18 | 0 |
| Pribilof red & blue king crab | 0 | 0 | 100 | 0 | 0 | 0 | 0 |
| St. Matthew blue king crab | 50 | 12 | 0 | 12 | 14 | 12 | 0 |
| Norton Sound red king crab | 0 | 0 | 0 | 0 | 50 | 50 | 0 |
| Eastern Bering Sea Tanner crab | 10 | 19 | 19 | 17 | 18 | 17 | 0 |
| Western Bering Sea Tanner crab | 10 | 19 | 19 | 17 | 18 | 17 | 0 |
| Bering Sea snow crab | 8 | 20 | 20 | 17 | 18 | 17 | 0 |
| Aleutian Islands red king crab (west of 179°W long) ^b | 8 | 18 | 21 | 18 | 21 | 14 | 0 |
| Eastern Aleutian Islands golden king crab (east of 174°W long) ^b | 8 | 18 | 21 | 18 | 21 | 14 | 0 |
| Western Aleutian Islands golden king crab (west of 174°W long) | 0 | 0 | 0 | 0 | 0 | 0 | 100 |

^a APICDA (Aleutian Pribilof Island Community Development Association).

BBEDC (Bristol Bay Economic Development Corporation).

CBSFA (Central Bering Sea Fishermen's Association).

CVRF (Coastal Villages Region Fund).

NSEDC (Norton Sound Economic Development Corporation).

YDFDA (Yukon Delta Fisheries Development Association).

ACDC (Adak Community Development Corporation).

^b Aleutian Islands red king crab west of 179°W long and Eastern Aleutian Islands golden king crab east of 174°W long were not part of the CDQ program until the initiation of crab rationalization in the 2005/06 season.