2018 Southeast Alaska Herring Spawn-On-Kelp Pound Fishery Management Plan

by

Eric Coonradt

Dave Harris

Troy Thynes

and

Scott Walker

March 2018

Alaska Department of Fish and Game



Division of Commercial Fisheries

Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative C	ode AAC	all standard mathematical	
deciliter	dL	all commonly accepted		signs, symbols and	
gram	g	abbreviations	e.g., Mr., Mrs.,	abbreviations	
hectare	ha		AM, PM, etc.	alternate hypothesis	H_A
kilogram	kg	all commonly accepted		base of natural logarithm	e
kilometer	km	professional titles	e.g., Dr., Ph.D.,	catch per unit effort	CPUE
liter	L		R.N., etc.	coefficient of variation	CV
meter	m	at	@	common test statistics	$(F, t, \chi^2, etc.)$
milliliter	mL	compass directions:		confidence interval	CI
millimeter	mm	east	Е	correlation coefficient	
		north	N	(multiple)	R
Weights and measures (English)		south	S	correlation coefficient	
cubic feet per second	ft ³ /s	west	W	(simple)	r
foot	ft	copyright	©	covariance	cov
gallon	gal	corporate suffixes:		degree (angular)	0
inch	in	Company	Co.	degrees of freedom	df
mile	mi	Corporation	Corp.	expected value	E
nautical mile	nmi	Incorporated	Inc.	greater than	>
ounce	OZ	Limited	Ltd.	greater than or equal to	≥
pound	lb	District of Columbia	D.C.	harvest per unit effort	HPUE
quart	qt	et alii (and others)	et al.	less than	<
yard	yd	et cetera (and so forth)	etc.	less than or equal to	≤
		exempli gratia		logarithm (natural)	ln
Time and temperature		(for example)	e.g.	logarithm (base 10)	log
day	d	Federal Information		logarithm (specify base)	\log_{2} , etc.
degrees Celsius	°C	Code	FIC	minute (angular)	'
degrees Fahrenheit	°F	id est (that is)	i.e.	not significant	NS
degrees kelvin	K	latitude or longitude	lat or long	null hypothesis	H_{O}
hour	h	monetary symbols		percent	%
minute	min	(U.S.)	\$, ¢	probability	P
second	S	months (tables and		probability of a type I error	
		figures): first three		(rejection of the null	
Physics and chemistry		letters	Jan,,Dec	hypothesis when true)	α
all atomic symbols		registered trademark	®	probability of a type II error	
alternating current	AC	trademark	TM	(acceptance of the null	
ampere	A	United States	*** 0	hypothesis when false)	β
calorie	cal	(adjective)	U.S.	second (angular)	"
direct current	DC	United States of	***	standard deviation	SD
hertz	Hz	America (noun)	USA	standard error	SE
horsepower	hp	U.S.C.	United States	variance	
hydrogen ion activity	pН	IIC -4-4-	Code	population	Var
(negative log of)		U.S. state	use two-letter abbreviations	sample	var
parts per million	ppm		(e.g., AK, WA)		
parts per thousand	ppt,		(c.g., AIX, WA)		
	‰				
volts	V				
watts	W				

REGIONAL INFORMATION REPORT NO. 1J18-01

2018 SOUTHEAST ALASKA HERRING SPAWN-ON-KELP POUND FISHERY MANAGEMENT PLAN

by

Eric Coonradt Alaska Department of Fish and Game, Division of Commercial Fisheries, Sitka

Dave Harris Alaska Department of Fish and Game, Division of Commercial Fisheries, Douglas

Troy Thynes
Alaska Department of Fish and Game, Division of Commercial Fisheries, Petersburg

and

Scott Walker Alaska Department of Fish and Game, Division of Commercial Fisheries, Ketchikan

> Alaska Department of Fish and Game Division of Commercial Fisheries, Publications Section 802 3rd Street, Douglas, Alaska, 99824

> > March 2018

The Regional Information Report Series was established in 1987 and was redefined in 2007 to meet the Division of Commercial Fisheries regional need for publishing and archiving information such as project operational plans, area management plans, budgetary information, staff comments and opinions to Board of Fisheries proposals, interim or preliminary data and grant agency reports, special meeting or minor workshop results and other regional information not generally reported elsewhere. Reports in this series may contain raw data and preliminary results. Reports in this series receive varying degrees of regional, biometric and editorial review; information in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author or the Division of Commercial Fisheries if in doubt of the level of review or preliminary nature of the data reported. Regional Information Reports are available through the Alaska State Library and on the Internet at: http://www.adfg.alaska.gov/sf/publications/

Eric Coonradt, Alaska Department of Fish and Game, Division of Commercial Fisheries, 304 Lake St. Rm. 103 Sitka, AK 99835 USA

Dave Harris, Alaska Department of Fish and Game, Division of Commercial Fisheries, 802 3rd Street Douglas, AK 99824 USA

Troy Thynes, Alaska Department of Fish and Game, Division of Commercial Fisheries, 16 Sing Lee Alley Petersburg, AK 99833 USA

and

Scott Walker, Alaska Department of Fish and Game, Division of Commercial Fisheries, 2039 Sea level Drive Suite 205, Ketchikan, AK 99901, USA

This document should be cited as follows:

Coonradt, E., D. Harris, T. Thynes, and S. Walker. 2018. 2018 Southeast Alaska herring spawn-on-kelp pound fishery management plan. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 1J18-01, Douglas.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526 U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203 Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers: (VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648, (Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

TABLE OF CONTENTS

	Page
LIST OF TABLES	ii
LIST OF FIGURES	ii
ABSTRACT	1
INTRODUCTION	1
HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE	2
Methods of Forecasting Herring Biomass Craig-Klawock (Section 3-B) Ernest Sound (District 7) Tenakee Inlet (Section 12-A) Hoonah Sound (Section 13-C)	2 2 3 3
CALENDAR OF EVENTS	
REGULATIONS	
General Spawn-On-Kelp Regulations	
Closure of some pound types	5
Other Regulations	5
HARVEST AND ALLOCATION OF KELP FOR 2018	6
FISHERY CONDUCT AND MANAGEMENT	6
Open Waters	7
Closed Waters	7
OTHER AGENCY REQUIREMENTS	8
Alaska Department Of Natural Resources U.S. Forest Service U.S. Coast Guard	8
LIST OF MANAGEMENT CONTACTS	9
TABLES AND FIGURES	10

LIST OF TABLES

Table		Page
1.	Craig/Klawock stock size and winter food and bait harvests, 1987–2017.	11
2.	Craig/Klawock herring spawn-on-kelp fishery summary, 1998–2017	12
3.	A comparison of Craig and Ernest Sound herring spawning dates for years 1997–2017	
4.	Ernest Sound miles of spawn, stock size and harvest, 1969–2017	14
5.	Ernest Sound herring SOK fishery summary, 2004, 2008–2009, 2011, and 2013–2014	16
6.	Tenakee Inlet herring seasonal spawning dates, mileage, biomass estimate, and harvest, 1978–2017.	17
7.	Tenakee Inlet herring spawn-on-kelp fishery summary, 2003–2005, 2009, and 2014	18
8.	A comparison of Hoonah Sound and Tenakee Inlet herring spawn dates for years 1996–2017	19
9.	Hoonah Sound herring spawning stock and fishery performance, 1971–2017.	
10.	Hoonah Sound herring spawn-on-kelp fishery summary, 2003–2017.	21
	LIST OF FIGURES	
Figure		Page
1.	Updated Open area for Craig/Klawock (Section 3-B) herring SOK fishery.	22
2.	Diagram of a herring pound showing two alternatives methods of marking herring pounds	
3.	Coast Guard Requirements for marking pounds.	24

ABSTRACT

This management plan provides an overview of the management approach, permit requirements, and regulations for the 2018 herring spawn-on-kelp pound fisheries in Southeast Alaska. A spawn-on-kelp pound fishery will only occur in Craig-Klawock. Staff biologists listed at the end of this document are available to answer questions regarding this plan. Pound operators are also advised to review the section of this plan that describes requirements of other agencies.

Key words: Pacific herring, Clupea pallasii, herring pound, Macrocystis kelp, allocation, management plan, spawn on kelp

INTRODUCTION

This plan provides an overview of the 2018 management approach, permit requirements, and regulations for the Southeast Alaska herring (*Clupea pallasii*) spawn-on-kelp fisheries. 5 AAC 27.185 *Management Plan for Herring Spawn on Kelp in Southeastern Alaska* establishes the regulatory framework for the Southeast Alaska spawn-on-kelp (SOK) fisheries and provides for fisheries in Sections 3-B (Craig/Klawock), 12-A (Tenakee Inlet), 13-C (Hoonah Sound), and in District 7 (Ernest Sound).

A *closed-pound-fishery* involves capturing sexually mature herring and releasing them into a net impoundment in which kelp is suspended. The herring are released from the pound after they spawn on the kelp and the kelp with eggs is then sold. An *open-pound-fishery* involves suspending kelp from a floating frame structure in an area where herring are spawning. The herring are not impounded but instead are allowed to naturally spawn on the suspended kelp. The kelp blades with eggs are removed from the water then sold.

In the Southeast Alaska herring SOK fisheries, a closed or an open pound may be operated by one or more Commercial Fisheries Entry Commission (CFEC) permit holders.

The 2017–2018 herring guideline harvest level (GHL) for the Craig/Klawock area stock is 2,312 tons of herring. Forty percent (40%) or 925 tons is allocated to the SOK fishery plus any unharvested portion of the winter food and bait quota. The 2017–2018 winter food and bait fishery closed on February 28, 2018, with a final harvest of 645 tons leaving 742 tons to add to the SOK GHL. This unharvested amount plus the 925 tons puts the GHL for the Craig/Klawock spawn-on-kelp fishery at 1,666 tons, placing the fishery in the 1,500 or more ton kelp allocation range. Fishermen are advised that new regulations are in place for the Craig/Klawock fishery for the 2018 season (see Regulations section).

No fishery will occur in Ernest Sound during the 2017–2018 season. In 2017, a spawn deposition survey was not conducted due to a lack of samples being obtained and the low mileage of herring spawn observed. Future assessment and fisheries are dependent on available funding.

No fishery will occur in Hoonah Sound during the 2017–2018 season. In 2017, no spawning biomass was observed therefore no forecast was generated for the 2017–2018 season. Future assessment and fisheries are dependent on available funding.

No fishery will occur in Tenakee Inlet during the 2017–2018 season. Approximately 2.1 nautical miles of spawn was observed in 2017, but a minimal number of samples were taken and no forecast was generated. Future assessment and fisheries are dependent on available funding.

HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE

METHODS OF FORECASTING HERRING BIOMASS

The Biomass Accounting (BA) method of forecasting is used to determine the expected prefishery mature spawning biomass and to establish the fishery GHL in Hoonah Sound and Ernest Sound. The BA method uses the most recent year's spawn deposition estimate of eggs, the age composition of the spawning biomass, and weights-at-age to project the following year's return of mature herring. The Hoonah Sound projection uses the estimated survival and maturity estimates from the age-structured-analyses (ASA) for the nearby Sitka Sound herring stock. A median historical level of recruitment of age-3 herring specific to Hoonah Sound is also applied to forecast biomass. The Ernest Sound projection uses the estimated survival and maturity estimates from the ASA for the Craig herring stock and a median historical recruitment of age-3 herring specific to Ernest Sound is applied to forecast biomass.

This BA method is unlike the ASA method used for forecasting herring biomass for several of the larger stocks in Southeast Alaska, including Craig/Klawock and Tenakee Inlet. The ASA method also uses the spawn deposition estimate of the eggs and the age composition to project the following year's return of mature herring. However, the ASA model calculates survival and maturation rates specific to the spawning stock. The ASA model utilizes a long time series of spawn deposition and age composition information to provide an estimate of the most recent biomass from which the forecast biomass for the next year is determined. ADF&G will continue to consider converting to use of the preferred ASA method for forecasting once there is an adequate time series of data to do so.

Once a forecast of the season's biomass is calculated, a variable harvest rate formula allows for a harvest rate of 10–20% of the forecast of mature spawning biomass. When the spawning biomass forecast for an area equals the threshold, the exploitation rate is 10% of the estimated spawning biomass. For each incremental increase in the spawning biomass equal to the threshold, the exploitation rate increases by 2%.

CRAIG-KLAWOCK (SECTION 3-B)

<u>Fishermen are advised that new regulations are in place for the Craig/Klawock fishery for the 2018 season (see Regulations section).</u>

Winter food and bait herring fisheries have occurred in Section 3-B (in the Mears Passage and Bocas de Finas areas) since the 1960s. Seasonal landings from the 1960s through 1985 were small, averaging approximately 210 tons. From the start of the herring SOK fishery in 1992, the Craig/Klawock herring GHLs have averaged 1,902 tons, ranging from a low of 626 tons in 2000 to a high of 6,847 tons in 2012 (Table 1). Spawn-on-kelp fishing effort, harvest, spawning dates, fishery dates, and product values are summarized in Table 2.

Annual harvest levels are based on a formula that allows for higher harvest rates as the herring population increases relative to the threshold level. No harvest is allowed if the biomass estimate for the stock is less than the threshold level. The established threshold level for the Craig/Klawock stock is 5,000 tons. The 2018 forecast for the Craig area is 16,039 tons. The 14.4% harvest rate will allow a combined quota of 2,312 tons for the winter food and bait and the

SOK fisheries. The GHL split between is 1,387 tons to the winter food and bait fishery and 925 tons to the SOK fishery. The forecast anticipates a strong return of age-4 herring (60%).

The GHL for the Section 3-B stock is allocated between both the winter food and bait fishery and the herring SOK fishery. When the fishery was created, the GHL allocation was 85% for the winter food and bait fishery and 15% for the SOK fishery. In 1998, the allocation was modified so that the winter food and bait fishery is allocated 60% of the GHL with the remaining 40% and any unharvested portion of the food and bait fishery going to the herring SOK fishery.

The 2017–2018 winter food and bait fishery closed on February 28, 2018. All unharvested winter food and bait quota remaining will be added to the GHL for the SOK fishery. The original GHL of 925 tons along with the remainder of the food and bait fishery will allow for the kelp allocation in the Section 3-B SOK fishery to fall within the maximum allocation of 1,500 or more tons.

Herring spawning normally occurs in the Craig/Klawock area between mid-March and early April. The earliest spawning observed since the mid-1970s was March 9 and the latest date of initial spawning was April 9. Figure 1 shows the new open area for the Craig/Klawock SOK fishery.

ERNEST SOUND (DISTRICT 7)

The Ernest Sound SOK pound fishery was created in January 2003 by the BOF. Additionally, the BOF created a herring bait pound fishery that is allocated 10% of the area's GHL and is similar to other herring fisheries in that its allocation is based upon the GHL remaining after the herring food and bait fisheries occur. Any remaining GHL from the winter food and bait fishery, or the bait pound fishery after March 15, is allocated to the SOK fishery. SOK fisheries in Ernest Sound have occurred intermittently and at various levels of effort and harvest since the first fishery occurred in 2004. From 2004 through 2017, there have been SOK GHLs in 6 years. Effort has varied from 0 participants in 2011 to 129 in 2014 (Table 5).

The Ernest Sound herring stock has a threshold level of 2,500 tons. A forecast was not developed for 2018 due to no spawn deposition survey being conducted. Historical spawning biomass, forecast, GHLs, spawning dates, harvest, and fishery dates are summarized in Tables 3, 4, and 5.

No commercial herring fisheries will occur in Ernest Sound during the 2017-2018 season. Monitoring herring spawn, collecting samples of spawning herring, and spawn deposition estimate in 2018 will be dependent on available funds.

TENAKEE INLET (SECTION 12-A)

The Tenakee Inlet stock has been utilized for the winter food and bait fishery since the 1978–1979 season. During seasons that the estimated spawning biomass was above the 3,000 ton threshold, the GHL has ranged from a low of 200 tons in the initial season to a peak of 1,700 tons in 1985–1986 (Table 5). Regulations adopted by the BOF in January 2003, provide for a SOK fishery in Tenakee Inlet if sufficient GHL remains at the close of the winter food and bait fishery. The SOK fishery occurred for the first time in April 2003. Summary results of the Tenakee Inlet SOK fisheries are presented in Table 6.

ADF&G has been conducting aerial surveys in Tenakee Inlet since the early 1970s; documenting the total miles of spawn each season to provide an indication of herring stock size or biomass. Aerial surveys were supplemented with hydroacoustic surveys from 1979 through 1986. Spawn-

deposition dive surveys have been used starting in 1987 as the most reliable and accurate means to assess the spawning biomass, and have not been conducted in Tenakee Inlet since 2015.

The Tenakee Inlet spawning stock has historically exhibited cycles of abundance. After a decade of fisheries, the stock declined below threshold in the early 1990s and no fisheries took place until 1996. Good recruitment led to nearly a decade of harvestable surplus until the forecasted biomass again declined below threshold in 2006. Aerial spawn surveys and spawn deposition dive surveys conducted in 2008 indicated a significant increase in spawning biomass to the levels seen in 1997–1999. However, recent surveys conducted since 2009 once again indicated a decreasing trend in mature spawning biomass. The herring spawn mileage observed in 2014, 2015, and 2017 was approximately 2.0 nauctical miles (nmi) each year. No herring spawn was documented within the inlet in 2016.

Spawning in Tenakee Inlet has generally occurred between the last week in April and the first week in May (Tables 5 and 7). The 2017 spawning dates were the latest observed to date with peak spawning occurring on May 13 and May 16. Traditionally, herring spawn primarily along the south shoreline of Tenakee Inlet between Saltery Bay and South Passage Point, with the core areas centered east and west of the Kadashan River flats. The 2015 spawn was centered much further east from the core area in the vicinity of Trap Bay. The 2017 spawn occurred in the traditional core area on either side of the Kadashan River flats. In addition, spawn has been documented intermittently along the Chatham Strait shoreline from South Passage Point to Basket Bay.

No commercial herring fisheries will occur in Tenakee Inlet during the 2017-2018 season. Aerial surveys will begin in mid-April of 2018.

HOONAH SOUND (SECTION 13-C)

ADF&G began monitoring the Hoonah Sound herring population in 1971. Since that time, the herring spawning stock has averaged 7.5 nmi of spawn and an estimated average escapement of 4,025 tons. Since 1990, the year the SOK fishery started, the stock has averaged 9.9 nmi of spawn and 5,922 tons of estimated escapement (Table 9). The highest recorded escapement biomass occurred in 2008 with an estimated 14.5 nmi of spawn and an escapement of 19,975 tons based on the spawn deposition survey.

In 1990, when Hoonah Sound became an SOK fishery, the minimum threshold at which a fishery could occur was reduced from 2,000 tons to 1,000 tons. In 2015, to be more consistent with similar sized stocks around the region, the threshold in Hoonah Sound was increased to 2,000 tons. A summary of historic herring spawn timing and herring SOK harvest in Hoonah Sound is available in Tables 8 and 10, respectively.

In 2017, no herring spawn was observed; because of this, no estimate of spawning biomass was generated. The biomass has been decreasing in this area since 2008. The reason for this decrease is not known, however, there are many oceanographic and biological influences on herring populations that are not well understood. There will be no formal forecast of herring biomass for the 2018 season. Therefore, no commercial fishery will occur in Hoonah Sound in 2018.

CALENDAR OF EVENTS

The following is a calendar of events to be considered by pound operators for the 2018 herring season.

March 2	News Release announcing the 2018 Ernest Sound, Hobart Bay/Port Houghton, Hoonah Sound, and Tenakee Inlet closures, and the 2018 Craig/Klawock GHL.
March 10	2018 Southeast Alaska Herring Spawn-On-Kelp Pound Fishery Management Plan will be available at all Southeast Alaska area offices.
March 17	The Craig/Klawock fishery will be open to seining of herring for placement in pounds effective 12:00 noon.
May 31	Pounds and all associated equipment in support of the fishery must be completely removed from the waters of the herring pound fishing area in Section 3-B. This includes the area covered by extreme high tide.

REGULATIONS

GENERAL SPAWN-ON-KELP REGULATIONS

The BOF met in Sitka from January 11 to 23, 2018, and adopted two regulatory changes to the SOK management plan. These changes include: an expansion to the open pounding area 5 AAC 27.185 (f)(1) and adoption of a regulation that allows the department to close fishing for some pound types listed in 5 AAC 27.185 (c) and 5 AAC 27.185 (dd)(2) if necessary to stay within the GHL.

The regulatory framework for the SOK fishery is found in 5 AAC 27.185. Management Plan for Herring Spawn on Kelp in Pounds in Sections 3-B, 12-A, 13-C, and District 7.

Expanded Open Pounding Area

Pounds for the taking of herring spawn-on-kelp and seines for the taking of herring for placement in pounds may be operated in the following locations, in Section 3-B, in the waters of the Gulf of Esquibel, San Alberto Bay, Shinaku Inlet, and San Christoval Channel south of the latitude of the northernmost tip of Saint Phillips Island at 55°39.31' N. latitude, east of a line from the northernmost tip of Saint Phillips Island to the northernmost tip of Point Garcia at 55°33.65′ N. latitude, 133°26.47′ W. longitude, and north of a line from Entrance Point to the southernmost tip of Clam Island to the southernmost tip of Fern Point and east of 133°20' W. longitude (Figure 1). The closed waters under 5 AAC 27.185 (f)(1) remain in effect.

Closure of some pound types

The department may close fishing for some pound types if necessary to avoid exceeding the GHL. In years when the GHL is low, the department instead of closing the fishery, may instead close the fishery to certain pound types, such as single-closed, double-closed. There will be no restrictions on pound types for the 2018 Craig/Klawock SOK fisherv.

Other Regulations

Additional regulations pertaining to the pound fisheries can be found in the 2017/2018 Statewide Commercial Herring Fishing Regulations booklet under CHAPTER 27, ARTICLE 4, SOUTHEAST ALASKA AREA under the following sections: 5 AAC 27.110 Fishing Seasons

for Southeastern Alaska Area (f), 5 AAC 27.130 Lawful Gear for Southeastern Alaska Area (d), and (e); 5 AAC 27.185 Management Plan for Herring Spawn on Kelp in Pounds (a) through (dd); 5 AAC 27.187 Buyer and Processors Reporting Requirements for Spawn on Kelp in Pounds for the Southeastern Alaska Area; harvesting requirements for Macrocystis kelp in 5 AAC 37.100 Permits; and 5 AAC 37.300 Harvesting Requirements for Macrocystis.

Under 5 AAC 27.185 (w) all pounds and associated equipment used in these fisheries must be removed from the water by a specific date for a specific period of time. ADF&G and Alaska Wildlife Troopers (AWT) are advising permit holders that any pounds, nets, buoys, lines, and anchors left on the grounds will be removed and impounded or destroyed.

It is the responsibility of permit holders to carefully review and follow these regulations.

HARVEST AND ALLOCATION OF KELP FOR 2018

A permit issued by ADF&G is required to harvest kelp for use in pounds (5 AAC 37.900). Kelp harvest permits may be obtained from local department offices. Kelp blades will be allocated equally among permit holders fishing the same type of gear. The amount of kelp allowed for harvest for each permit holder is based on the kelp allocation table as indicated under regulation 5 AAC 27.185 (d) plus an allowance for breakage and loss during transport.

Section 3-B (Craig/Klawock):

- Single permit closed pound—600 blades of *Macrocystis* kelp.
- Double permit closed pound—900 blades of *Macrocystis* kelp.
- Triple permit closed pound—1,000 blades of *Macrocystis* kelp.
- Quadruple permit or more closed pound—1,000 blades of *Macrocystis* kelp.
- Single permit open pounds—3,000 blades of *Macrocystis* kelp;
- Multiple permit open pounds—9,000 blades of *Macrocystis* kelp.

FISHERY CONDUCT AND MANAGEMENT

The Craig/Klawock herring pound fishery will be the only SOK fishery in southeast Alaska for 2018.

ADF&G will be closely monitoring herring activity using vessel and aerial surveys. Results of aerial surveys will be announced by department news release or in fishery updates.

ADF&G will continue to monitor the practice of *top-off-fishing*. Regulation 5 AAC 27.185 (q) prevents a permit holder from releasing any herring from their pound when they are adding fresh herring.

To avoid mortality, the transport of herring to the pound site should be done with the pound itself or a pushable/towable net pen. Towable net pens used only for transporting herring must be marked "Tow Pound" and include the CFEC permit number of a participating permit holder. Transporting with a purse seine is discouraged except for very short distances. Pound operators should slowly push pounds or tow the pound alongside to avoid prop wash and prevent crushing

herring against the net. Permit holders are asked to avoid making and holding large sets intended to fill multiple pounds to avoid herring mortality and stress.

Although regulations determine the maximum allowable number of kelp blades that can be harvested and placed in each permit holder's pound, fishermen are encouraged to fish the number of blades which will provide the maximum overall quality and value of their product rather than simply to fish the total amount allowed. Other measures have successfully been used in the fishery that may be considered when trying to maximize spawn on kelp quality and value including the following:

- 1) Pound nets may be shaped with internal frames to provide the full net volume;
- 2) The kelp depths in the pound may be matched with the depth of active spawning by testing spawn deposition with a weighted string;
- 3) Fishing and transferring herring to pounds should only occur once herring are fully mature;
- 4) Top-off sets may be added over a 4 day period;
- 5) The herring density in the net should be limited since spawning is retarded by excessive crowding;
- 6) Web depth may be adjusted (consistent with specifications under 5 AAC 27.130 (e)(1)(C) or 5 AAC 27.185(cc)) to provide good water exchange.

Open Waters

The open waters for Section 3-B were modified during the 2018 BOF meeting in Sitka and now include: the waters of the Gulf of Esquibel, San Alberto Bay, Shinaku Inlet, and San Christoval Channel south of the latitude of the northernmost tip of Saint Phillips Island at 55°39.31′ N. latitude, east of a line from the northernmost tip of Saint Phillips Island to the northernmost tip of Point Garcia at 55°33.65′ N. latitude, 133°26.47′ W. longitude, and north of a line from Entrance Point to the southernmost tip of Clam Island to the southernmost tip of Fern Point and east of 133°20′ W. longitude. (Figure 1).

Closed Waters

In Section 3-B, certain areas are closed to the operation of herring pounds and seines for taking of herring for placement in pounds. Those areas are shown in Figure 1 and include:

- Klawock Inlet and Big Salt Lake;
- Those waters of San Christoval Channel in the main channel enclosed by a line from 55°35.62′ N. latitude, 133°20′ W. longitude to 55°35.17′ N. latitude, 133°20′ W. longitude to 55°33.37′ N. latitude, 133°17.52′ W. longitude to 55°33.50′ N. latitude, 133°17.28′ W. longitude;
- Those waters of Fish Egg and Ballena Islands south of 55°31′ N. latitude and north of the southernmost tip of Cape Suspiro and east of the longitude of Ballena Island Shoal Light.

OTHER AGENCY REQUIREMENTS

Prospective pound operators are advised to consider other agency requirements for constructing and operating pounds in Craig/Klawock, Ernest Sound, Tenakee Inlet, and Hoonah Sound. Pound operators are urged to contact the Alaska Department of Natural Resources, U.S. Forest Service, the U.S. National Marine Fisheries Service, and the U.S. Coast Guard to determine other regulations and requirements. Phone numbers for those agencies are listed below.

ALASKA DEPARTMENT OF NATURAL RESOURCES

The Alaska Department of Natural Resources (907-465-3400) manages the use of tidelands and submerged lands seaward of mean high water.

U.S. FOREST SERVICE

In the Hoonah Sound and Tenakee Inlet areas, the U.S. Forest Service has jurisdiction over and manages most of the lands above mean high tide. People who plan to use National Forest land in connection with the fishery must apply for a special use permit from the U.S. Forest Service prior to any occupancy. Special use permit applications are available at the Sitka Ranger District Office, 204 Siginaka Way, Sitka, Alaska 99835, (907-747-6671). Completed applications should be submitted to the Sitka Ranger District well in advance of operations to ensure that a permit is received in time for the fishery. Examples of use needing a permit include (but are not limited to): camping on National Forest land in conjunction with the commercial fishery and storage of gear on the National Forest.

U.S. COAST GUARD

Structures such as floating fish pens are subject to the requirements of the Code of Federal Regulations, Title 33, Part 64. This regulation requires an owner to apply for a U.S. Coast Guard permit and to install and maintain a light or other private aid to navigation if the U.S. Coast Guard determines it to be necessary to protect maritime navigation.

Herring pounds used in the SOK pound fishery do not require permits for private aids to navigation at this time, provided the owners:

Place two signs on opposite corners of the structure. These signs will be worded "Danger, Fish Pens" (Figure 3).

Place a single, all-points white light on one corner of structures less than 400 square feet in size.

Place a single, all-points white light on every corner of structures larger than 400 square feet in size.

Anchor fish pens within the boundary areas specified in ADF&G regulation 5 AAC 27.185 (f) (Figure 1).

If all these conditions are not met, the permit holder must apply to the U.S. Coast Guard for an individual "Private Aids to Navigation Permit." If you have questions, call the U.S. Coast Guard Aids to Navigation office, at (907) 463-2254.

LIST OF MANAGEMENT CONTACTS

Following are ADF&G Division of Commercial Fisheries contacts regarding this management plan:

Name and Title	Address and Phone Number
Lowell Fair	802 3rd Street
Southeast Alaska Regional Supervisor	Douglas, Alaska 99824
	(907) 465-4250
Dan Gray	304 Lake St., Rm. 103
Southeast Alaska Regional Management Biologist	Sitka, Alaska 99835
	(907) 747-6688
Kyle Hebert	802 3rd Street
Herring Research Biologist	Douglas, Alaska 99824
-	(907) 465-4250
Scott Walker	2030 Sea Level Dr. Ste. 205
Area Management Biologist	Ketchikan, Alaska 99901
	(907) 225-5195
Bo Meredith or Justin Breese	
Assistant Management Biologists	
Troy Thynes	16 C' I All
Area Management Biologist	16 Sing Lee Alley Petershyra: AV 00822 USA (007) 772 2801
Kevin Clark	Petersburg, AK 99833 USA (907) 772-3801
Assistant Management Biologist	
Tom Kowalske	215 F
Assistant Management Biologist	215 Front Street
Assistant Management Diologist	Wrangell, AK 99929-0200
-	(907) 874-3822
Eric Coonradt	304 Lake St., Rm. 103
Area Management Biologist	Sitka, Alaska 99835
A . D .	(907) 747-6688
Aaron Dupuis	
Assistant Management Biologist	
Dave Harris	802 3rd Street
Area Management Biologist	Douglas, Alaska 99824
Scott Forbes	(907) 465-4250
Assistant Management Biologist Jim Craig	802 3rd Street
Southeast Alaska Regional Publications Manager	Douglas, Alaska 99824
Southeast Alaska Regional I ubilications ividilagei	(907) 465-4236
	(701) 703-7230

TABLES AND FIGURES

Table 1.-Craig/Klawock stock size and winter food and bait harvests, 1987-2017.

Season	Miles Spawn	Forecasted Pre- fishery Biomass	Total GHL Bait and SOK ^c (Tons)	Bait Quota (Tons)	Bait Harvest (Tons)	Number of Permits	Ex-Vessel Value
87–88 ^d	27.0	16,550	N/A	2,200	2,014	21	\$564,000
88–89	31.7	16,350	N/A	1,810	1,730	22	\$465,000
89–90	30.0	19,800	N/A	3,150	3,221	23	\$ 998,200
90-91	22.0	18,350	N/A	2,841	3,272	27	\$981,600
91–92	23.0	17,800	2,684	2,281	2,295	28	\$619,650
92–93 ^e	8.4	12,350	1,602	1,362	623	10	\$150,960
93-94	8.0	7,996	895	760	636	6	\$187,578
94–95	5.5	6,778	725	617	***	***	***
95–96	9.9	6,262	658	558	***	***	***
96–97	13.2	6,755	715	615	517	4	\$137,788
97–98 ^f	11.0	7,018	755	455	***	***	***
98–99	15.4	6,951	750	450	***	***	***
99-00	12.9	6,013	626	376	***	***	***
00-01	16.7	9,091	1,058	635	***	***	***
01-02	18.0	8,387	952	571	***	***	***
02-03	11.2	6,045	630	378	***	***	***
03-04	12.0	13,204	1,754	1,052	***	***	***
04-05	18.0	15,577	2,217	1,330	553	3	\$199,012
05-06	8.2	14,262	1,955	1,173	689	3	\$247,934
06-07	22.3	13,768	1,860	1,116	576	3	\$139,000
07-08	11.0	14,213	1,945	1,167	565	3	\$133,300
08-09	17.0	14,213	1,945	1,167	142	3	\$51,304
09-10	18.7	14,870	2,074	1,244	***	***	***
10-11	14.8	17,886	2,710	1,140	***	***	***
11-12	14.9	34,235	6,847	4,060	309	3	\$113,784
12-13	15.3	23,391	4,060	2,436	321	3	\$116,000
13–14	13.6	26,085	4,808	2,884	***	***	***
14–15	11.5	15,803	2,362	1,367	964	3	\$396,205
15–16	12.3	12,303	1,590	954	898	3	\$369,660
16–17	22.8	7,833	872	523	527	3	\$210,800
17–18	_	16,083	2,312	1,387	_	3	_
Average	15.9	13,749	1,902	1,357	800	7	\$220,106

^{***} confidential

^a Spawn year is beginning year of regulatory season listed in the adjacent year column.

b Forecasted pre-fishery biomass values were estimated with hydroacoustics for 86/87, spawn deposition surveys for 87/88 to 92/93, and age-structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured models for 93/94 to 14/15.

Construction of the structured m

^e First year bait quota was split between pound fishery 85%:15%.

f Herring allocation changed to 60% for the winter food and bait fishery, 40% to the pound fishery.

^g Updated forecast based on updated scale ages.

Table 2.-Craig/Klawock herring spawn-on-kelp fishery summary, 1998–2017.

Statistic	2000	2001	2002	2003	2004	2005
Herring Quota (tons)	280	914	852	528	1,579	1,667
Total harvest SOK (tons)	0	26.9	41.7	69.2	50	115.2
Exvessel value	\$0	\$146,859	\$218,700	\$423,000	\$325,000	\$603,723
Average Price/lb	\$0.00	\$2.70	\$3.10	\$3	\$3.25	\$2.62
Average Income	\$0	\$2,880	\$2,460	\$3,385	\$3,420	\$9,011
Number of pounds	50	31	50	61	50	42
Number of landings	0	51	89	118	95	67
Blade allocation	d	e	e	e	d	f
Total kelp harvest (tons)	2.0	3.2	8.2	7.5	14.0	4.9
Herring spawning dates	3/22-4/5	4/1-4/7	3/31-4/7	3/31-4/7	3/26-4/7	4/9-4/14
Miles of spawn	12.9	16.7	18.4	11.2	12.0	18.0
Forecasted Pre-Fishery biomass (tons)	9,951	8,042	8,387	6,045	13,204	15,577
	2006	2007	2008	2009	2010	2011
Herring Quota (tons)	1266	1,284	1,380	1,802	1,953	2,710
Total harvest SOK (tons)	28.9	44.5	148.5	137.3	116.7	70
Exvessel value	\$298,575	\$1,087,532	\$3,066,788	\$1,256,777	\$884,715	\$728,147
Average Price/lb	\$5.15	\$12.08	\$10.33	\$4.58	\$3.80	\$5.13
Average Income	\$8,782	\$23,139	\$25,138	\$9,107	\$8,268	\$14,003
Number of pounds	50	52	66	96	63	34
Number of landings	34	47	122	137	107	52
Blade allocation	g	g	g	g	g	g
Total kelp harvest (tons)	4.6	5.6	12.2	7.3	8.2	4.6
Herring spawning dates	3/30-4/3	4/3-4/12	4/3-4/12	4/3-4/10	4/5-4/14	4/1-4/7
Miles of spawn	8.2	22.3	11.0	17.0	18.7	14.8
Forecasted Pre-Fishery biomass (tons)	14,262	13,768	14,213	14,213	14,870	17,886
(1311)	2012	2013	2014	2015	2016	2017
Herring Quota (tons)	6,847	4,060	4,808	2,362	1,590	872
Total harvest SOK (tons)	98.1	137.7	confidential	confidential	confidential	69.9
Exvessel value	\$2,099,002	\$3,099,002	confidential	confidential	confidential	\$932,917
Average Price/lb	\$10.69	\$12.00	confidential	confidential	confidential	\$6.68
Average Income	\$32,795	\$23,656	confidential	confidential	confidential	\$8,042
Number of pounds	35	80	75	76	46	19
Number of landings	64	131	136	135	133	116
Blade allocation	g	g	g	g	h	i
Total kelp harvest (tons)	5.3	9.3	19.2	19.2	9.2	10.2
Herring spawning dates	4/3-4/8	3/31–4/3	4/1-4/5	3/27-4/1	3/25-3/31	3/24–4/2 & 4/6–4/8
Miles of spawn	14.9	15.3	13.6	11.5	12.3	22.8
Forecasted Pre-Fishery biomass (tons)	34,235	23,391	26,085	15,803	12,303	7,833

⁷⁰ blades for a single closed pound, 210 for a multiple pound permit holder.

²⁰⁰ blades for a single closed pound, 600 blades for a multiple pound permit holder

³⁵⁰ blades for a single closed pound, 750 blades for a double closed pound, 1,125 blades for a triple closed pound. 600 blades for a single closed pound, 750 blades for a double closed pound, 1,125 blades for a triple closed pound.

³⁰⁰ blades for a single closed pound, 400 blades for a double closed pound, 700 blades for a triple closed pound.

⁵⁰⁰ blades per permit with six permit holders per pound structure.

Table 3.-A comparison of Craig and Ernest Sound herring spawning dates for years 1997–2017.

<u>Table</u>	3	A com	parı	ISOI	n of	t C	rai	g a	ınd	<u>E</u>	rne	est	S	ou	nd	h	eri	nr	g s	pa	lWI	nır	ıg	da	tes	s to	or	ye	ars	3 I	<u>99</u>	/-/	<u> 201</u>	/.														
				N	Mai	rch	l																										Αj	pril														
		16 17 1	18 19	20	21	22	23	2	1 2:	5 2	6 2	27	28	29	3	0 :	31	1	2	3	3	4	5	6	7	7 :	8	9 1	10	11	12	13	14	15	16	17	7 18	3 1	9 2	0 2	1 2	22	23	24	25	26	27	28
1997 Craig																																																
Ernes																																																
1998 Craig	2																																								Т							
Ernes																																									\perp							
1999 Craig	<u>g</u>																				Т																				Т							
Ernes																																																
2000 Craig																																									Т							
Ernes																																									\perp							
2001 Craig	2																																								Т							
Ernes	st																																								\perp							
2002 Craig																																									Τ							
Ernes																																																
2003 Craig																																									Т							
Ernes	st																																															
2004 Craig	g																																								Т							
Ernes																																									\perp							
2005 Craig	3																				Т																				Т							
Ernes	st									Т					Т						Т	Т					Г											Т										
2006 Craig																																									Т							
Ernes	st																																								\perp							
2007 Craig																																									Т							
Ernes	st																																															
2008 Craig	g																																															
Ernes																																																
2009 Craig																																									Т							
Ernes																																																
2010 Craig																																																
Ernes																																																
2011 Craig	2																																								Т							
Ernes																																																
2012 Craig																																																
Ernes																																																
2013 Craig																																									\perp							
Ernes	st																																															
2014 Craig																																									I							
Ernes																																									\perp							
2015 Craig	3																																															
Ernes																																																
2016 Craig	3																																															
Ernes																																									I							
2017 Craig	g																																								Т							П
Ernes																					Т						Т												Т		Т				\exists			
			_	_	_				_	_				_	_	_			_	_	_	_				_	_	_	_					_					_		_	_			-	-		

Note: Black bars indicate dates of active spawning for the Craig stock and gray bars indicate dates of active spawning for the Ernest Sound stock.

Table 4.-Ernest Sound miles of spawn, stock size and harvest, 1969-2017.

Season	Date of first spawn ^a	Nautical miles of spawn ^b	Forecast Used for GHL Determination ^c (tons)	Spawning Biomass (tons) ^d	Guideline Harvest Level (tons) ^e	Bait harvest (tons) ^f	SOK Harvest (lbs)	Sac Roe Harvest (tons)	SOK GHL (tons)
1969/1970	_	_	-	_	_	17	_	_	_
1970/1971	_	3	_	13,100	_	206	_	_	_
1971/1972	_	_	13,100	3,650	_	967	_	_	_
1972/1973	_	_	3,650	450	_	775	_	_	_
1973/1974	_	_	450	400	_	535	_	_	_
1974/1975	_	_	400	2,900	_	593	_	_	_
1975/1976	_	3	2,900	4,350	580	708	_	_	0
1976/1977 ^g	_	3	4,350	3,035	870	901	_	49	0
1977/1978	3-May	_	3,035	1,505	455	340	_	_	115
1978/1979	16-Apr	2.6	1,505	255	_	_	_	_	_
1979/1980	2-May	4	255	500	_	_	_	_	_
1980/1981	_	3.5	500	410	-	_	_	_	_
1981/1982	_	_	410	160	-	_	_	_	_
1982/1983	_	_	160	1,640	-	_	_	_	_
1983/1984	11-Apr	_	1,640	1,000	_	_	_	_	_
1984/1985	_	4.5	1,000	1,000	_	_	_	-	_
1985/1986	_	_	1,000	1,000	_	_	_	_	_
1986/1987	_	1	1,000	_	_	_	_	_	_
1987/1988	21-Apr	2	_	_	_	_	_	_	_
1988/1989	17-Apr	2.4	_	500	_	_	_	_	_
1989/1990	_	2.1	500	1,000	_	_	_	_	_
1990/1991	_	_	1,000	3,000	_	_	_	_	-
1991/1992	16-Apr	9.1	3,000	2,650	_	_	_	_	-
1992/1993	23-Apr	9	2,650	684	200	8	_	_	192
1993/1994	26-Apr	8.4	684	2,544	0	_	_	-	-
1994/1995	25-Apr	6.5	2,544	2,470	255	111	_	_	144
1995/1996	16-Apr	6.9	2,744	2,665	280	220	_	_	60
1996/1997	16-Apr		4,852	0	377	6	-	-	371
1997/1998	9-Apr	11.8	-	5,998	0	_	_	_	-
1998/1999	5-Apr	1.8	5,381	No survey	662	96	_	_	566
1999/2000	8-Apr	9.1	-	920	0	_	_	_	-
2000/2001	11-Apr	6.9	-	2,052	0	_	-	-	-
2001/2002	15-Apr	4.8	1,653	2,406	0				_

-continued-

Note: Table footnotes are at the bottom of the following page.

Table 4.–Page 2 of 2.

Season	Date of first spawn ^a	Nautical miles of spawn ^b	Forecast Used for GHL Determination ^c (tons)	Spawning Biomass (tons) ^d	Guideline Harvest Level (tons) ^e	Bait harvest (tons) ^f	SOK Harvest (lbs)	Sac Roe Harvest (tons)	SOK GHL (tons)
2002/2003	10-Apr	8.5	2,407	5,509	0	_	-	_	_
2003/2004	11-Apr	7.1	6,592	2,413	875	44	112,286	_	831
2004/2005	22-Apr	10.1	1,906	3,268	0	_	_	_	_
2005/2006	6-Apr	7.9	2,284	2,538	0	_	_	_	_
2006/2007	19-Apr	11.3	1,955	7,353	0	_	_	_	_
2007/2008	20-Apr	15.4	9,060	4,846	1,382	**	19,650	_	>700
2008/2009	17-Apr	6.6	4,545	2,862	529	**	4,911	_	100-299
2009/2010	14-Apr	7.8	2,879	3,523	297	**	_	_	< 50
2010/2011	17-Apr	8.1	5,080	2,559	476	**	0	_	100-299
2011/2012	16-Apr	8.9	2,682	3,193	272	**	_	_	< 50
2012/2013	16-Apr	9.7	3,509	7,556	379	**	129,265	_	100-299
2013/2014	14-Apr	3.7	7,613	2,631	1,073	**	**	_	>700
2014/2015	8-Apr	5.5	1,991	562	0	_	_	_	_
2015/2016	10-Apr	4.4	1,207	346	0	_	_	_	_
2016/2017	14-Apr	4.4	_	_	_	_	_	_	_
10-yr Average	16-Apr	7.5	4,285	3,120	490	**	**	_	578
Total Average	16-Apr	6.1	2,590	2,591	332	368	**	49	353

Note: ** indicates data is confidential.

^a Since 1997/1998 the first spawn and the major spawn have been within five days of each other.

b 1996/1997 No survey, fish all spawned (7.5 miles) along Ship Island; 1998/1999 No survey, only 1.8 miles of spawn observed, surveys probably missed main spawn; spawn year is the beginning year of regulatory season.

c 1971/1972 through 1984/1985 forecasts were based on hydroacoustical-computer generated estimates; 1985/1986 through 1991/1992 forecasts were based on visual estimates; 1992/1993 through 1994/1995 were based on spawn deposition estimates; 1995/1996 through 2006/2007 were biomass accounting forecasts.

d 1969/1970 through 1983/1984 biomass estimates were hyrdoacoustical-computer generated estimates; 1984/1985 through 1990/1991 were visual estimates; and 1991/1992 through 2005/2006 were spawn deposition estimates. 1975/1976 & 1976/1977 GHLs are based upon 20% of the acoustical estimate. 1977/1978 GHL is based upon 15% of the acoustical estimate.

e 2003/2004 GHL includes 90 tons rolled over from the bait pound fishery.

f 1973/1974,1974/1975, 1976/1977 also include harvests from Fools and Menefee Inlets. Does not include harvests from statistical area 107-40.

Since 1976/1977 season the threshold for a fishery has been 2,500 tons.

Table 5.-Ernest Sound herring SOK fishery summary, 2004, 2008–2009, 2011, and 2013–2014.

Statistic	2004	2008	2009
Ernest Sound GHL (tons)	875	1,382	529
GHL Available for SOK (tons)	775	**	**
SOK Harvest (tons)	56.1	9.8	2.5
Exvessel Value	\$514,912	\$131,059	\$18,359
Average Price/lb	\$4.59	\$6.87	\$3.67
Average Income/permit	\$8,046	\$10,091	\$4,590
Number of Permits participating	64	13	4
Number of Pounds	51/6/0/1 ^a	1/6/0/0 a	0/2/0/0 a
Number Permits Landing Product	64	13	4
Kelp Allocation (blades)	c	d	e
Kelp Blade Harvest (lbs)	4,600	29,400	2,400
Fishery Open—Closed	4/1-5/10	4/1-5/10	4/1-5/10
Fishing Occurred	4/10-4/17	4/16/-4/24	4/17-4/21
Harvest Occurred	4/14–1/17	24-Apr	21-Apr
Statistic	2011	2013	2014
Ernest Sound GHL (tons)	476	379	1,073
GHL Available for SOK (tons)	**	**	**
SOK Harvest (tons)	0	64.3	**
Exvessel Value	\$0	\$1,574,729	**
Average Price/lb	\$0	\$12.25	**
Average Income/permit	\$0	\$19,441	**
Number of Permits participating	0	81	129
Number of Pounds	0	$1/5/1/22/0^{b}$	25/52/0 a
Number Permits Landing Product	0	81	129
Kelp Allocation (blades)	e	e	d
Kelp Blade Harvest (lbs)	0	6,400	29,000
Fishery Open—Closed	4/1-5/10	4/1-5/10	4/1-5/10
Fishing Occurred	NA	4/10-4/16	4/12-4/20
Harvest Occurred	NA	4/15-4/16	4/18-4/20

Note: ** indicates data is confidential.

^a single/double/triple/open

b single/double/combined double/triple/open

c 1,000 blades single-closed/1,000 blades double-closed/1,000 blades triple-closed/2,500 blades single-open/7,500 multiplepermit-open d 1,000 blades single-closed/2,000 blades double-closed/1,000 blades triple-closed/2,500 blades single-open/7,500 multiple-

permit-open e 200 blades single-closed/400 blades double-closed/500 blades triple-closed/1,500 blades single-open/4,500 multiple-permit-

Table 6.—Tenakee Inlet herring seasonal spawning dates, mileage, biomass estimate, and harvest, 1978–2017.

			Spawning	Food/Bait	
	Major Spawning	Nautical Miles of	Biomass ^a	GHL	Food/Bait
Season	Dates	Spawn (nmi)	(tons)	(tons)	Harvest (tons)
1978/1979	5/9-5/11	3.3	2,500	200	0
1979/1980	4/28-5/2	3.9	4,485	400	504
1980/1981	4/27-5/5	9.3	7,500	750	847
1981/1982	4/25-5/7	11.1	6,650	650	687
1982/1983	4/25-5/6	13.1	8,870	875	749
1983/1984	4/20-4/26	8.3	12,100	850	619
1984/1985	4/24-5/1	9.9	11,000	1,400	1,406
1985/1986	4/27-5/1	8.3	12,500	1,700	2,040
1986/1987	4/22-4/30	7.9	6,600	800	1,275
1987/1988	4/22-4/27	9.1	6,000	1,450	1,577
1988/1989	4/26-4/29	10.3	5,360	720	655
1989/1990	4/25-5/6	2.9	2000	650	595
1990/1991	4/25-5/4	2.1	400	No t	fishery
1991/1992	5/5	trace	200	No t	fishery
1992/1993	4/21-4/23	6.4	904	No t	fishery
1993/1994	4/24-4/26	0.25	400	No f	fishery
1994/1995	4/26	0.05	200	No f	fishery
1995/1996	5/4-5/14	18.1	4,560	No f	fishery
1996/1997	4/26-5/7	14.4	9,926	300	98
1997/1998	4/24-4/29	12.4	10,419	825	586
1998/1999	4/25-4/28	11.0	11,049	1,023	835
1999/2000	4/26-5/3	13.8	9,425	542	494
2000/2001	4/21-5/1	12.2	7,576	906	775
2001/2002	4/23-4/27	15.4	4,084	840	355
2002/2003	4/25-4/28	12.2	3,529	528	328
2003/2004	4/28-5/3	13.0	4,728	399	confidential
2004/2005	4/26-5/2	8.9	3,036	476	0
2005/2006	5/2-5/6	5.9	5,110	No f	fishery
2006/2007	4/23-4/26	4.4	3,346		fishery
2007/2008	4/30; 5/7–5/8	11.4	11,252		fishery
	4/25–4/26; 4/29–4/30	6.9	5,283	875	254
2009/2010	5/7-5/9	2.7	1,437	583	confidential
2010/2011	5/9	1.0	N/A		ishery
2011/2012	4/20–4/23	4.6	5,119		ishery
2012/2013	5/7–5/10	5.4	4,936		ishery
2013/2014	4/29	2.0	927	557	confidential
2014/2015	4/22–4/25	2.3	2,223		ishery
2015/2016	none	0.0	No forecast		ishery
2016/2017	5/13–5/16	2.1	No forecast		ishery

Spawning biomass estimates were calculated from hydro-acoustical surveys from 1979 through 1986, and from spawn deposition surveys from 1987 through 2015 – bolded values were derived from ASA models.

Table 7.-Tenakee Inlet herring spawn-on-kelp fishery summary, 2003-2005, 2009, and 2014.


Statistic	2003	2004	2005
Tenakee Inlet GHL (tons)	528	399	476
GHL Available for SOK (tons)	180	confidential	476
SOK Harvest (tons)	47.6	100.7	101.4
Exvessel Value	\$580,500	\$981,464	\$512,900
Average Price/lb	\$6.10	\$4.68	\$2.53
Average Income/permit	\$10,555	\$11,684	\$5,636
Number of Permits participating	55	85	98
Number of Pounds	$1/15/8/0^{a}$	1/32/6/2/2 ^b	1/29/13/3 ^a
Number Permits Landing Product	55	85	91
Kelp Allocation (blades)	200/400/550/0 ^a	$300/500/500/2000^a$	$300/500/500/2000^a$
Kelp Blade Harvest (lbs)	35,375	39,000	53,850
Fishery Open—Closed	4/6-5/6	4/6–5/6	4/6-5/5
Fishing Occurred	4/25-4/28	4/28 - 5/1	4/27-4/30
Harvest Occurred	4/30-5/4	5/3-5/6	5/2-5/4
Statistic	2009	2014	
Tenakee Inlet GHL (tons)	875	557	
GHL Available for SOK (tons)	621	confidential	
SOK Harvest (tons)	64.1	84.4	
Exvessel Value	\$558,900	\$1,155,276	
Average Price/lb	\$4.36	\$6.85	
Average Income/permit	\$6,499	\$16,271	
Number of Permits participating	86	78	
Number of Pounds	11/27/7/0 ^a	2/25/5/1°	
Number Permits Landing Product	86	71	
Kelp Allocation (blades)	$400/500/500/0^{a}$	300/500/500/0 ^a	
Kelp Blade Harvest (lbs)	42,600	41,250	
Fishery Open—Closed	4/6–5/5	4/6–5/4	
Fishing Occurred	4/28-5/1	4/27-5/1	
Harvest Occurred	5/2-5/5	5/2-5/4	

Note: No fishery occurred from 2006 to 2008 since the biomass forecast was below the 3,000-ton threshold.

^a single/double/triple/test.

b single/double/triple/long line/test. c single/double/triple/quadruple.

Table 8.–A comparison of Hoonah Sound and Tenakee Inlet herring spawn dates for years 1996–2017.



Note: Black bars indicate spawning dates for Hoonah Sound and gray bars indicate spawning dates for Tenakee Inlet.

Table 9.–Hoonah Sound herring spawning stock and fishery performance, 1971–2017.

Year	Spawn Dates	Nautical Miles Spawn	Estimated Escapement (tons)	SOK Harvested (tons)
1971	5/10-5/17	2.5	833	-
1972	5/11-5/12	1.5	666	-
1973	N/A	1.0	333	_
1974	14-May	3.0	999	-
1975	N/A	N/A		_
1976	5-May	1.0	333	-
1977	N/A	3.5	1,166	_
1978	N/A	5.3	1,765	_
1979	N/A	0.5	167	_
1980	N/A	N/A	_	
1981	4/30-5/01	2.3	750	_
1982	4/29-5/01	1.5	500	_
1983	1-May	1.0	333	_
1984	4/26–5/01	3.0	540	
1985	5/01-5/03	3.5	1,166	_
1986	4/28–5/01	3.8	1,249	_
1987	4/28–5/02	3.8	740	_
1988	4/30–5/01	5.0	1,665	_
1989	4/16–4/20	17.0	4,000	_
1990	4/13-4/28	10.0	2,350	11.9
1991	4/19–4/24	8.7	2,175	13.3
1992	4/22–4/24	10.8	5,714	23.1
1993	4/27–4/29	5.7	1,099	14.0
1994	4/21–4/23	9.0	2,450	32.7
1995	4/20–4/21	4.5	274	27.4
1996	5/02–5/9	10.1	4,023	27.4
1997	4/25–4/28	14.5	5,884	65.2
1998	4/23-4/27	14.5	6,472	85.6
1999	4/27–5/1	13.8	4,426	71.6
2000	4/27–4/30	13.0	3,635	35.7
2000	4/27–4/30	13.7	8,538	66.2
2001	4/25–4/27	11.9	4,936	136.6
2002	4/23-4/27	16.7	9,423	141.5
2003	4/23–4/27	11.1	7,502	237.4
2004	4/18–4/25	10.3	6,924	190.6
2005	4/23–4/26	9.0	6,028	
2007	4/46–5/2	16.5	10,946	162.1 159.4
2007	4/46–5/2 4/23–4/30	14.5	19,975	202.3
2008		10.3		
2009	4/22-4/27 4/22 - 4/28	12.4	15,829 15,264	234.7 290.4
			4 4 5 4 5	
2011 2012	4/20–5/5 4/20–4/23	12.6 4.2	14,215 923	193.7 186.0
2012	4/20 -4 /23 5/7-5/9	2.4	412	0.0
	5/5–5/8			0.0
2014		3.2	444	
2015	4/24–5/6	4.2	23	0.0
2016 2017	N/A N/A	N/A N/A	_	0.0 0.0
verage	1971–2017	7.5	4,025	N/A
verage	1971–2017	7.5 9.9	4,025 5,922	N/A 95.6

Note: Shaded estimated escapements are based on average spawn density from years 1989 to 2002.

Table 10.-Hoonah Sound herring spawn-on-kelp fishery summary, 2003–2017.

-					
Statistic	2003	2004	2005	2006	2007
Herring Quota (tons)	427	1,207	728	669	681
Harvest Quota (tons)	NA	NA	NA	NA	NA
Harvest (tons)	141.6	237.4	190.6	162.1	144.5
Exvessel Value	\$1,922,50	\$2,071,347	\$1,117,568	\$1,943,422	\$4,491,070
Average Price/lb	\$6.79	\$4.36	\$2.93	\$6.00	\$14.09
Average Income	\$17,800	\$19,541	\$11,889	\$24,600	\$49,352
Number of Applicants	NA	NA _.	NA	NA	NA
Number of Pounds	49/1/3 ^a	92/12/2 ^b	81/5/3 ^c	17/45 ^a	67/12 ^a
Number Selling Product	108	106	94	79	91
	500/300/750	1,000/1,000/	1,000/1,000/	2,500/1,000/	2,500/1,000/
Kelp Allocation (blades)	a	$3,000^{b}$	$1,500^{c}$	$1,500^{a}$	$1,500^{a}$
Kelp Blade Harvest	60,301	126,000	118,450	136,698	122,565
Fishery Open—Closed	4/6-4/25	4/6-4/28	4/6–4/28	4/6-4/27	4/6–5/4
Fishing Occurred	4/19–4/24	4/20-4/25	4/19–4/28	4/18–4/23	4/23-4/29
Harvest Occurred	4/24-4/27	4/26–4/28	4/25–4/28	4/23–4/27	4/30–5/4
Statistic	2008	2009	2010	2011	2012
Herring Quota (tons)	2,238	2,238	3,182	3,015	2,139
Harvest Quota (tons)	NA	NA	NA	NA	NA
Harvest (tons)	223	234.7	290.4	193.7	186.5
Exvessel Value	\$5,115,45	\$2,332,514	\$2,580,517	\$1,820,952	\$4,033,078
Average Price/lb	\$11.47	\$4.97	\$4.44	\$4.70	\$10.81
Average Income/Landing	\$51,155	\$23,094	\$25,550	\$20,460	\$55,248
Number of Applicants	NA	NA	NA	NA	NA
Number of Pounds	98/3 ^a	99/4 ^a	97/2 ^a	85/4 ^a	83/4 ^a
Number Selling Product	100	101	101	89	73
	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/
Kelp Allocation (blades)	$1,500^{a}$	$1,500^{a}$	$1,500^{a}$	$1,500^{a}$	$1,500^{a}$
Kelp Blade Harvest	201,262	196,492	178,898	169,922	155,104
Fishery Open–Closed	4/6-5/02	4/6-4/30	4/6–4/30	4/6–4/30	4/6–5/15
Fishing Occurred	4/22–4/27	4/22-4/25	4/21-4/25	4/20-4/24	4/19–4/21
Harvest Occurred	4/27-5/1	4/26–4/29	4/22–4/28	4/26–4/29	4/26–4/27
Statistic	2013	2014	2015	2016	2017
Herring Quota (tons)	130	0	0	0	0
Harvest Quota (tons)	NA	NA	NA	NA	NA
Harvest (tons)	0	0	0	0	0
Exvessel Value	\$0	\$0	\$0	\$0	\$0
Average Price/lb	\$0	\$0	\$0	\$0	\$0
Average Income/Landing	\$0	\$0	\$0	\$0	\$0
Number of Applicants	NA	NA	NA	NA	NA
Number of Pounds	3 open	0	0	0	0
Number Selling Product	0	0	0	0	0
Kelp Allocation (blades)	600 open	0	0	0	0
Kelp Blade Harvest	7,940	0	0	0	0
Fishery Open—Closed	4/6–5/15	NA	NA	NA	NA
Fishing Occurred	NA	NA	NA	NA	NA
Harvest Occurred	NA	NA	NA	NA	NA
	2122	212.2	1111	1111	1111

Note: No fishery occurred in 1996 since the biomass forecast was below the 1,000-ton threshold

^a Double closed pounds/single closed pounds/triple closed pounds.

^b Double closed pounds/single closed pounds/open pounds.

^c Single-permit closed pound/double-permit closed pound/triple-permit closed pounds.




Figure 1.—Updated Open area for Craig/Klawock (Section 3-B) herring SOK fishery.




Figure 2.—Diagram of a herring pound showing two alternatives methods of marking herring pounds.

Note: Regulations require vertical signs with the permit holder's first and last name and five-digit CFEC permit number (5 AAC 27.185(k)). Letters and numbers must be at least six inches high and at least one-half inch wide, must contrast with the background and must be above the waterline at all times.




Figure 3.—Coast Guard Requirements for marking pounds.