

2022 Southeast Alaska Herring Spawn-On-Kelp Pound Fishery Management Plan

by

Aaron Dupuis

Dave Harris

Bo Meredith

and

Paul Salomone

February 2022

Alaska Department of Fish and Game

Division of Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code		all standard mathematical signs, symbols and abbreviations	
deciliter	dL		AAC		
gram	g	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H _A
hectare	ha			base of natural logarithm	e
kilogram	kg			catch per unit effort	CPUE
kilometer	km	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	coefficient of variation	CV
liter	L			common test statistics	(F, t, χ^2 , etc.)
meter	m	at	@	confidence interval	CI
milliliter	mL	compass directions:		correlation coefficient (multiple)	R
millimeter	mm	east	E	correlation coefficient (simple)	r
Weights and measures (English)		north	N	covariance	cov
cubic feet per second	ft ³ /s	south	S	degree (angular)	°
foot	ft	west	W	degrees of freedom	df
gallon	gal	copyright	©	expected value	E
inch	in	corporate suffixes:		greater than	>
mile	mi	Company	Co.	greater than or equal to	≥
nautical mile	nmi	Corporation	Corp.	harvest per unit effort	HPUE
ounce	oz	Incorporated	Inc.	less than	<
pound	lb	Limited	Ltd.	less than or equal to	≤
quart	qt	District of Columbia	D.C.	logarithm (natural)	ln
yard	yd	et alii (and others)	et al.	logarithm (base 10)	log
Time and temperature		et cetera (and so forth)	etc.	logarithm (specify base)	log ₂ , etc.
day	d	exempli gratia (for example)	e.g.	minute (angular)	'
degrees Celsius	°C	Federal Information Code	FIC	not significant	NS
degrees Fahrenheit	°F	id est (that is)	i.e.	null hypothesis	H ₀
degrees kelvin	K	latitude or longitude	lat or long	percent	%
hour	h	monetary symbols (U.S.)	\$, ¢	probability	P
minute	min	months (tables and figures): first three letters	Jan,...,Dec	probability of a type I error (rejection of the null hypothesis when true)	α
second	s	registered trademark	®	probability of a type II error (acceptance of the null hypothesis when false)	β
Physics and chemistry		trademark	™	second (angular)	"
all atomic symbols		United States (adjective)	U.S.	standard deviation	SD
alternating current	AC	United States of America (noun)	USA	standard error	SE
ampere	A	U.S.C.	United States Code	variance	
calorie	cal			population sample	Var var
direct current	DC	U.S. state	use two-letter abbreviations (e.g., AK, WA)		
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

REGIONAL INFORMATION REPORT NO. 1J22-04

**2022 SOUTHEAST ALASKA HERRING SPAWN-ON-KELP POUND
FISHERY MANAGEMENT PLAN**

by

Aaron Dupuis

Alaska Department of Fish and Game, Division of Commercial Fisheries, Sitka

Dave Harris

Alaska Department of Fish and Game, Division of Commercial Fisheries, Douglas

Bo Meredith

Alaska Department of Fish and Game, Division of Commercial Fisheries, Ketchikan

and

Paul Salomone

Alaska Department of Fish and Game, Division of Commercial Fisheries, Petersburg

Alaska Department of Fish and Game
Division of Commercial Fisheries, Publications Section
802 3rd Street
Douglas, AK 99824-5412

February 2022

The Regional Information Report Series was established in 1987 and was redefined in 2007 to meet the Division of Commercial Fisheries regional need for publishing and archiving information such as area management plans, budgetary information, staff comments and opinions to Alaska Board of Fisheries proposals, interim or preliminary data and grant agency reports, special meeting or minor workshop results and other regional information not generally reported elsewhere. Reports in this series may contain raw data and preliminary results. Reports in this series receive varying degrees of regional, biometric and editorial review; information in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author or the Division of Commercial Fisheries if in doubt of the level of review or preliminary nature of the data reported. Regional Information Reports are available through the Alaska State Library and on the Internet at: <http://www.adfg.alaska.gov/sf/publications/>.

Product names used in this publication are included for completeness and do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

*Aaron Dupuis,
Alaska Department of Fish and Game, Division of Commercial Fisheries,
304 Lake St. Rm. 103
Sitka, AK 99835 USA*

*Dave Harris,
Alaska Department of Fish and Game, Division of Commercial Fisheries,
802 3rd Street
Douglas, AK 99824 USA*

*Bo Meredith,
Alaska Department of Fish and Game, Division of Commercial Fisheries,
2030 Sea level Drive Suite 205,
Ketchikan, AK 99901, USA*

and

*Paul Salomone,
Alaska Department of Fish and Game, Division of Commercial Fisheries,
16 Sing Lee Alley
Petersburg, AK 99833 USA*

This document should be cited as follows:

Dupuis, A., D. Harris, B. Meredith, and P. Salomone. 2022. 2022 Southeast Alaska herring spawn-on-kelp pound fishery management plan. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 1J22-04, Douglas.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,
(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	ii
LIST OF FIGURES	ii
ABSTRACT	1
INTRODUCTION.....	1
HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE	2
Methods of Forecasting Herring Biomass	2
Craig-Klawock (Section 3-B).....	2
Ernest Sound (District 7)	3
Tenakee Inlet (Section 12-A).....	4
Hoonah Sound (Section 13-C).....	4
CALENDAR OF EVENTS	6
REGULATIONS	6
General Spawn-On-Kelp Regulations	6
Placement and Release of Herring in Pounds	6
Post-Harvest Requirements.....	6
Harvest and Production.....	6
Requirements for Buyers	7
Closure of some pound types.....	7
Other Regulations	7
HARVEST AND ALLOCATION OF KELP FOR 2022.....	7
Section 3-B (Craig/Klawock)	8
FISHERY CONDUCT AND MANAGEMENT	8
Open Waters	9
Closed Waters.....	9
OTHER AGENCY REQUIREMENTS.....	9
Alaska Department of Natural Resources.....	9
U.S. Forest Service	9
U.S. National Marine Fisheries Service.....	10
U.S. Coast Guard	10
LIST OF CONTACTS.....	11
TABLES AND FIGURES.....	13

LIST OF TABLES

Table	Page
1. Craig/Klawock herring stock size and fishery summary, 1990–2021.....	14
2. Craig/Klawock herring SOK detailed fishery summary, 2004–2021.....	15
3. Craig and Ernest Sound herring spawning dates comparison, 2002–2021.....	16
4. Ernest Sound herring stock and fishery summary, 1969–2021.....	17
5. Ernest Sound SOK detailed fishery summary, 2004, 2008–2009, 2011, and 2013–2014.....	19
6. Tenakee Inlet herring stock and fishery summary, 1978–2021.....	20
7. Tenakee Inlet herring SOK detailed fishery summary, 2003–2005, 2009, and 2014.....	22
8. Hoonah Sound and Tenakee Inlet herring spawn dates comparison, 2001–2021.	23
9. Hoonah Sound herring spawning stock and fishery summary, 1971–2021.....	24
10. Hoonah Sound herring SOK detailed fishery summary, 2003–2017.	25

LIST OF FIGURES

Figure	Page
1. Craig/Klawock (Section 3-B) herring SOK fishery open area.	26
2. Diagram of a herring pound showing two alternative methods of marking herring pounds.	27
3. U.S. Coast Guard requirements for marking pounds.....	28

ABSTRACT

This management plan provides an overview of the management approach, permit requirements, and regulations for the 2022 herring spawn-on-kelp (SOK) pound fisheries in Southeast Alaska. A herring spawn-on-kelp pound fishery will only occur in the Section 3-B (Craig-Klawock) area. The forecasted mature spawning biomass in Section 3-B is 63,250 tons of herring for the 2021/22 season. This biomass estimate allows for a 20% harvest rate and a combined GHL of 12,650 for the winter food and bait and the SOK fishery. The minimum GHL available to the Section 3-B SOK fishery is 5,060 tons, allowing for full kelp allocation.

Keywords: Pacific herring, *Clupea pallasii*, herring pound, *Macrocystis* kelp, allocation, management plan, spawn on kelp

INTRODUCTION

This plan provides an overview of the 2022 management approach, permit requirements, and regulations for Pacific herring (*Clupea pallasii*) harvest in the Southeast Alaska spawn-on-kelp fisheries. 5 AAC 27.185 *Management Plan for Herring Spawn on Kelp in Southeastern Alaska* establishes the regulatory framework for the Southeast Alaska spawn-on-kelp (SOK) fisheries and provides for fisheries in Sections 3-B (Craig/Klawock), 12-A (Tenakee Inlet), 13-C (Hoonah Sound), and in District 7 (Ernest Sound).

A *closed-pound fishery* involves capturing sexually mature herring and releasing them into a net impoundment in which kelp is suspended. The herring are released from the pound after they spawn on the suspended kelp and the kelp with eggs attached is then sold. An *open-pound fishery* involves suspending kelp from a floating frame structure in an area where herring are spawning. The herring are not impounded but instead are allowed to naturally spawn on the suspended kelp. The kelp blades with eggs are removed from the water then sold. In the Southeast Alaska herring SOK fisheries, a closed or an open pound may be operated by one or more Commercial Fisheries Entry Commission (CFEC) permit holders.

The 2021/22 season (October 1–September 30) herring guideline harvest level (GHL) for the Craig/Klawock area stock is 12,650 tons of herring. Forty percent (40%) or 5,060 tons is allocated to the SOK fishery plus any unharvested portion of the winter food and bait quota. The current GHL for the Craig/Klawock spawn-on-kelp fishery will put the fishery in the 1,500 or more tons kelp allocation range.

No fishery will occur in Ernest Sound during the 2021/22 season. A forecast was not generated due to the low mileage of herring spawn observed in 2021. In 2022, Ernest Sound will be monitored throughout the duration of historical spawn timing. If enough herring spawn is observed and funding allows, a spawn deposition survey may be conducted.

No fishery will occur in Hoonah Sound during the 2021/22 season. In 2021, no herring or herring spawn was observed therefore no forecast was generated for the 2021/22 season. Future assessment and fisheries are dependent on available funding.

No fishery will occur in Tenakee Inlet during the 2021/22 season. In 2021, no herring or herring spawn was observed inside Tenakee Inlet, and approximately 1.7 nautical miles (nmi) of spawn was observed on the Chatham Strait shoreline from north of Basket Bay south to Point Craven (just inside Peril Strait). No forecast was generated, and future assessment and fisheries are dependent on available funding.

HERRING STOCK STATUS AND HISTORICAL FISHERY PERFORMANCE

METHODS OF FORECASTING HERRING BIOMASS

The age-structured analysis (ASA) method is used for forecasting herring biomass for several of the larger stocks in Southeast Alaska, including Craig/Klawock and Tenakee Inlet. The ASA method primarily uses a time series of estimates of total egg deposition, age compositions of spawning and commercially caught herring, and commercial harvest to project the following year's return of mature herring. The ASA model utilizes a three-to-four-decade long time series of these data sets to estimate the current biomass, historical biomass, and forecast biomass for the next year. The ASA model also calculates survival and maturation rates specific to the spawning stock and is the preferred model when data sets are adequate.

The biomass accounting (BA) method of forecasting is simpler alternative to the ASA method and used for areas with historical datasets that are shorter or less complete. This method is used to forecast the pre-fishery mature biomass and to establish the fishery GHL in Hoonah Sound and Ernest Sound. The BA method uses only the current year's estimate of egg deposition, the current age compositions of the spawning and commercially caught biomass, weight at age, and fecundity estimates to project the following year's return of mature herring. The Hoonah Sound forecast uses the estimated survival and maturity estimates derived from the ASA model for the nearby Sitka Sound herring stock. A median historical level of recruitment of age-3 herring specific to Hoonah Sound is also applied to forecast biomass. The Ernest Sound forecast uses the estimated survival and maturity estimates derived from the ASA model for the Craig herring stock and a median historical recruitment of age-3 herring specific to Ernest Sound is applied to forecast biomass.

Once a forecast of the season's biomass is calculated, a variable harvest rate formula allows for a harvest rate of 10–20% of the forecast of mature spawning biomass. When the spawning biomass forecast for an area equals the threshold, the exploitation rate is 10% of the estimated spawning biomass. For each incremental increase in the spawning biomass equal to the threshold, the exploitation rate increases by 2%.

CRAIG-KLAWOCK (SECTION 3-B)

Herring fisheries have occurred in Section 3-B since the 1960s. The first commercial harvest was in the form of a wild SOK fishery in the Craig/Klawock area beginning in the early 1960s that ended in 1967. A small winter bait herring fishery began in 1965. Initial harvests were small, less than five tons through 1973, and a formal GHL was never established. Beginning in 1973, hydroacoustic surveys began and the winter bait harvest increased with an average harvest of 1,147 tons from 1973–1991. The Section 3-B SOK fishery was established by the board in 1992. The GHL for the Section 3-B stock was allocated between both the winter food and bait fishery and the herring SOK fishery. When the fishery was created in 1992, the GHL allocation was 85% for the winter food and bait fishery and 15% for the SOK fishery. In 1998, the allocation was modified so that the winter food and bait fishery is allocated 60% of the GHL with the remaining 40% and any unharvested portion of the food and bait fishery allocated to the herring SOK fishery. Since 1992, the Craig/Klawock herring GHL has averaged 2,864 tons, ranging from a low of 626 tons in 2000 to a high of 19,456 tons in 2021 (Table 1). The herring spawn-on-kelp fishery effort, harvest, spawning dates, fishery dates, and product values are summarized in Table 2.

Annual harvest levels are based on a formula that allows for higher harvest rates as the herring population increases relative to the threshold level. No harvest is allowed if the biomass estimate for the stock is less than the threshold level. The established threshold level for the Craig/Klawock stock is 5,000 tons.

Herring biomass information collected in 2019 indicated a pronounced recruitment pulse of herring, not just in Craig, but throughout Southeast Alaska and the Gulf of Alaska. Specifically, samples from both cast net and the commercial catch during the 2019 spawning event in Craig indicate that the mature biomass was dominated by age-3 fish, 81% of cast net samples and 85% of the commercial harvest. This large proportion of age-3 fish was observed in several Southeast Alaska herring stocks, including Ernest Sound, Revilla Channel, Seymour Canal and Sitka Sound. The 2021 forecast of mature spawning biomass for the Craig/Klawock stock was 97,282 tons of herring, the highest forecast on record. The decrease in the 2022 forecasted mature spawning biomass is due to low survival and low forecasted age-3 recruitment compared to 2019 and 2021. Model-estimated age-3 recruitment in 2021 is the third highest recruitment on record and the population is now supported by two of the highest year classes (2016 and 2018 year classes) seen since the start of the time series in 1988.

The forecasted mature spawning biomass for 2021/22 is 63,250 tons of herring. This forecast allows for a 20% harvest rate and a resultant GHL of 12,650 for the winter food and bait and the SOK fisheries. Therefore, the GHL is 7,590 tons to the winter food and bait fishery and 5,060 tons to the SOK fishery. The department anticipates the spawning biomass in 2022 to be comprised primarily (70%) of age-6 herring, based on the age structure of the 2021 spawning biomass.

The 2021/22 winter food and bait fishery will close by regulation on February 28, 2022. All unharvested winter food and bait quota will be added to the GHL for the SOK fishery. The allocated SOK GHL of 5,060 tons will allow for the kelp allocation in the Section 3-B SOK fishery to fall above the maximum allocation of 1,500 or more tons.

Herring spawning normally occurs in the Craig/Klawock area between mid-March and early April. The earliest spawning observed since the mid-1970s was March 9 and the latest date of initial spawning was April 9 (Table 3).

ERNEST SOUND (DISTRICT 7)

The Ernest Sound SOK pound fishery was created in January 2003 by the BOF. Additionally, the BOF created a herring bait pound fishery that is allocated 10% of the area's GHL and is similar to other herring fisheries in that its allocation is based upon the GHL remaining after the herring food and bait fisheries occur. Any remaining GHL from the winter food and bait fishery, or the bait pound fishery after March 15, is allocated to the SOK fishery. SOK fisheries in Ernest Sound have occurred intermittently and at various levels of effort and harvest since the first fishery occurred in 2004 (Table 4). From 2004 through 2021, there have been SOK GHLs in 6 years. Effort varied from 0 participants in 2011 to 129 in 2014 (Table 5).

The Ernest Sound herring stock has a threshold level of 2,500 tons. A forecast was not developed for 2022 because no spawn deposition survey was conducted. Historical spawning biomass, forecast, GHLs, spawning dates, harvest, and fishery dates are summarized in Tables 3, 4, and 5.

No commercial herring fisheries will occur in Ernest Sound during the 2021/22 season. The department plans to monitor herring spawn in 2022 and if herring spawn is substantial, samples of spawning herring may be collected, and spawn deposition survey may be conducted.

TENAKEE INLET (SECTION 12-A)

The Tenakee Inlet stock has been utilized for the winter food and bait fishery since the 1978/79 season. During seasons that the estimated spawning biomass was above the 3,000-ton threshold, the GHJ ranged from a low of 200 tons in the initial season to a peak of 1,700 tons in the 1985/86 season (Table 5). Regulations adopted by the BOF in January 2003, provide for a SOK fishery in Tenakee Inlet if sufficient GHJ remains at the close of the winter food and bait fishery. The SOK fishery occurred for the first time in April 2003. Summaries of the Tenakee Inlet SOK fisheries are presented in Table 6.

The department has conducted aerial surveys in Tenakee Inlet since the early 1970s, documenting the total miles of spawn each season to provide an indication of herring stock size or biomass. Aerial surveys were supplemented with hydroacoustic surveys from 1979 through 1986. Spawn-deposition surveys began in 1987 as the most reliable and accurate means to assess the spawning biomass and have not been conducted in Tenakee Inlet since 2015.

With respect to historical stock assessment, the Tenakee Inlet stock includes spawn documented within the inlet itself and any spawn documented along the Chichagof Island shoreline in Chatham Strait between Tenakee Inlet and Pt. Hayes. This spawning stock has historically exhibited cycles of abundance. After a decade of fisheries, the stock declined below threshold in the early 1990s and no fisheries took place until the 1996/97 season. Good recruitment led to nearly a decade of harvestable surplus until the forecasted biomass again declined below threshold in 2006. Aerial spawn surveys and spawn deposition dive surveys conducted in 2008 indicated a significant increase in spawning biomass to the levels seen in 1997–1999. However, recent surveys conducted since 2009 once again indicated a decreasing trend in mature spawning biomass. The herring spawn mileage observed in 2014, 2015, and 2017 was approximately 2.0 nmi each year, but no herring spawn was documented within the inlet in 2016. Since 2017, observed spawn mileage within Tenakee Inlet has shown a continual decline: 1.4 nmi in 2018, 0.5 nmi in 2019, 0.15 nmi in 2020, and none observed in the spring of 2021. The only herring spawn observed in the area in 2021 was approximately 1.7 nmi of spawn along the Chichagof Island shoreline of Chatham and mouth of Peril Straits with concentrations just north of Basket Bay and around Point Craven.

Spawning in Tenakee Inlet has generally occurred between the last week in April and the first week in May (Tables 5 and 7). Traditionally, herring spawn primarily in Tenakee Inlet along the southern shoreline between Saltery Bay and South Passage Point, with the core areas centered east and west of the Kadashan River flats. Herring spawn has also been documented intermittently along the Chatham Strait shoreline from South Passage Point to Point Hayes, and the only significant spawn observed in the last two seasons (2020 and 2021) has been along this shoreline. The 2021 Chatham/Peril Straits major spawning events were observed from April 20–23, among the earliest timing documented for this stock in the last 30 years.

No commercial herring fisheries will occur in Tenakee Inlet during the 2021/22 season. Aerial surveys will begin in mid-April of 2022.

HOONAH SOUND (SECTION 13-C)

ADF&G began monitoring the Hoonah Sound herring stock in 1971. From 1971–2020, the herring spawning stock has averaged 6.8 nmi of spawn and an estimated average spawning biomass of 3,633 tons. From 1990, the year the SOK fishery started, through 2020 the stock has averaged 8.7 nmi of spawn and 5,103 tons of estimated spawning biomass (Table 9). The highest recorded

spawning biomass occurred in 2008 with an estimated spawning biomass of 19,975 tons based on the spawn deposition survey.

In 1990, when Hoonah Sound became a SOK fishery, the minimum threshold at which a fishery could occur was reduced from 2,000 tons to 1,000 tons. In 2015, to be more consistent with similar sized stocks around the region, the threshold in Hoonah Sound was increased to 2,000 tons. A summary of historic herring spawn timing and herring SOK harvest in Hoonah Sound is available in Tables 8 and 10.

In 2021, no herring spawn was observed; because of this, no estimate of spawning biomass was generated. The biomass in this area sharply decreased in 2012 and, based on aerial surveys, appears to have remained at a very low level since that time. The reason for this decrease is not known, however, there are many oceanographic and biological influences on herring populations that are not well understood. There was no forecast for the 2021/22 season; therefore, no commercial fishery will occur in Hoonah Sound in 2022.

CALENDAR OF EVENTS

The following is a calendar of events to be considered by pound operators for the 2022 herring season.

- | | |
|----------------|---|
| By February 28 | 2022 Southeast Alaska Herring Spawn-On-Kelp Pound Fishery Management Plan will be available at all Southeast Alaska area offices. |
| March 3 | Advisory announcement of the 2022 Ernest Sound, Hoonah Sound, and Tenakee Inlet closures, and the 2022 Craig/Klawock GHL. |
| March 17 | The Craig/Klawock fishery will open to seining of herring for placement in pounds effective 12:00 noon. |
| May 31 | Pounds and all associated equipment in support of the fishery must be completely removed from the waters of the herring pound fishing area in Section 3-B. This includes the area covered by extreme high tide. |

REGULATIONS

GENERAL SPAWN-ON-KELP REGULATIONS

The regulatory framework for the SOK fishery is found in 5 AAC 27.185. *Management Plan for Herring Spawn on Kelp in Pounds in Sections 3-B, 12-A, 13-C, and District 7.*

Placement and Release of Herring in Pounds

Herring may be placed in or added to a pound for four days starting with the initial placement of herring in a pound. After 11:59 p.m. on the fourth day, no additional herring may be added to the pound (5 AAC 27.185 [q]). All herring in the pound must be released by 12:00 noon on the seventh day after the initial placement of herring in a pound (5 AAC 27.185 [s]). Under 5 AAC 27.185 (s), the “*first day*” is defined as the day herring are first placed into a pound. Once herring have been released or SOK product has been harvested, no additional herring or kelp may be introduced into a pound (5 AAC 27.185 [q]). Fishermen must take responsibility to ensure that when adding herring to a pound that herring are not at the same time swimming out of the pound as this would be a violation of 5 AAC 27.185 (q). When releasing herring, at least one full side of the pound's webbing must be lowered a minimum of two feet below the surface of the water (5 AAC 27.185 [s]).

Post-Harvest Requirements

After a permit holder releases herring and harvests product from the pound, the permit holder must maintain the webbing in place for at least four weeks. To optimize hatching success, the permit holder must position egg-covered webbing in the original size and configuration of the pound structure with adequate water circulation on all sides. The webbing support system must be above the surface of the water and clearly marked as per 5 AAC 27.185(k).

Harvest and Production

Each permit holder's SOK blades must remain separate from other permit holder's SOK blades until after processing and grading is completed. Permit holders will be allowed to harvest all SOK produced in their pounds. A permit holder's fish ticket must report only the SOK they harvested from their pound. Each permit holder fishing a jointly operated pound shall be issued a fish ticket

and the sum of the weights of those tickets shall equal the total weight of product produced in the jointly operated pound. All permit holders and any vessel carrying commercial SOK product from the fishing grounds must first contact the Alaska Department of Fish and Game (ADF&G) with the estimated amount of SOK product harvested and indicate the intended time and location of the delivery. For any product that has been delivered on the grounds to a licensed processor, the processor (not the permit holder) will be required to contact the department with delivery weight for each landing on board.

REQUIREMENTS FOR BUYERS

Reporting requirements for buyers and processors of SOK product from Southeast Alaska SOK fisheries can be found in 5 AAC 27.187 *Buyer and Processors Reporting Requirements for Spawn-On-Kelp in Pounds for the Southeastern Alaska Area*. Buyers, processors, and permit holders should read and become familiar with these reporting requirements.

Operators of floating processing vessels, tender vessels, and catcher-processors will be required to report in person, by VHF radio, or by telephone, to the local ADF&G office or directly to department area management biologists on the grounds before the start of processing operations in Southeast Alaska. These reporting requirements are specified by regulation 5 AAC 39.130 (f) and (g).

CLOSURE OF SOME POUND TYPES

The department may close fishing for some pound types, if necessary, to avoid exceeding the GHL. In years when the GHL is low, the department, instead of closing the fishery, may instead close the fishery to certain pound types, such as single closed and double closed. There will be no restrictions on pound types for the 2022 Craig/Klawock SOK fishery.

OTHER REGULATIONS

Additional regulations pertaining to the pound fisheries can be found in the 2021–2022 Statewide Commercial Herring Fishing Regulations booklet under CHAPTER 27, ARTICLE 4, SOUTHEAST ALASKA AREA under the following sections: 5 AAC 27.110 *Fishing Seasons for Southeastern Alaska Area* (f), 5 AAC 27.130 *Lawful Gear for Southeastern Alaska Area* (d), and (e); 5 AAC 27.185 *Management Plan for Herring Spawn on Kelp in Pounds* (a) through (dd); 5 AAC 27.187 *Buyer and Processors Reporting Requirements for Spawn on Kelp in Pounds for the Southeastern Alaska Area*; harvesting requirements for *Macrocystis* kelp in 5 AAC 37.100 *Permits*; and 5 AAC 37.300 *Harvesting Requirements for Macrocystis*.

Under 5 AAC 27.185 (w) all pounds and associated equipment used in these fisheries must be removed from the water by a specific date for a specific period of time. ADF&G and Alaska Wildlife Troopers are advising permit holders that any pounds, nets, buoys, lines, and anchors left on the grounds will be removed and impounded or destroyed.

It is the responsibility of permit holders to carefully review and follow these regulations.

HARVEST AND ALLOCATION OF KELP FOR 2022

A permit issued by ADF&G is required to harvest kelp for use in pounds (5 AAC 37.900). Kelp harvest permits may be obtained from local department offices. Kelp blades will be allocated equally among permit holders fishing the same type of gear. The amount of kelp allowed for

harvest for each permit holder is based on the kelp allocation table as indicated under regulation 5 AAC 27.185 (c) plus an allowance for breakage and loss during transport.

SECTION 3-B (CRAIG/KLAWOCK)

- Single permit closed pound—600 blades of *Macrocystis* kelp
- Double permit closed pound—900 blades of *Macrocystis* kelp
- Triple permit closed pound—1,000 blades of *Macrocystis* kelp
- Quadruple permit or more closed pound—1,000 blades of *Macrocystis* kelp
- Single permit open pound—3,000 blades of *Macrocystis* kelp
- Multiple permit open pounds—9,000 blades of *Macrocystis* kelp

FISHERY CONDUCT AND MANAGEMENT

The Craig/Klawock herring pound fishery will be the only SOK fishery in Southeast Alaska for 2022.

ADF&G will be closely monitoring herring activity using vessel and aerial surveys. Results of aerial surveys will be announced by department advisory announcements on the day that an aerial survey occurs and may be updated on the grounds on Marine VHF Channel 10. Herring survey data collected by the department from aerial and boat surveys will be available online through the interactive herring spawn maps located at the department's website at: <http://www.adfg.alaska.gov/index.cfm?adfg=commercialbyareasoutheast.herring#maps>

ADF&G will continue to monitor the practice of *top-off fishing*. Regulation 5 AAC 27.185 (q) prevents a permit holder from releasing any herring from their pound when they are adding fresh herring.

To avoid mortality, the transport of herring to the pound site should be done with the pound itself or a pushable/towable net pen. Net pens used only for transporting herring must be marked "Tow Pound" and include the CFEC permit number of a participating permit holder. Transporting with a purse seine is discouraged except for very short distances. Pound operators should slowly push pounds or tow the pound alongside to avoid prop wash and prevent crushing herring against the net. Permit holders are asked to avoid making and holding large sets to avoid herring mortality and stress.

Although regulations determine the maximum allowable number of kelp blades that can be harvested and placed in each permit holder's pound, fishermen are encouraged to fish the number of blades which will provide the maximum overall quality and value of their product rather than simply to fish the total amount allowed. Other measures have successfully been used in the fishery that may be considered when trying to maximize spawn on kelp quality and value including the following:

- 1) Pound nets may be shaped with internal frames to provide the full net volume.
- 2) Kelp depths in the pound may be matched with the depth of active spawning by testing spawn deposition with a weighted string.
- 3) Fishing and transferring herring to pounds should only occur once herring are fully mature.

- 4) Top-off sets may be added over a 4-day period.
- 5) The herring density in the pound should be closely monitored since successful spawning is inhibited by excessive crowding.
- 6) Web depth may be adjusted (consistent with specifications under 5 AAC 27.130 (e)(1)(C) or 5 AAC 27.185(cc)) to provide good water exchange.

OPEN WATERS

In Section 3-B, pounds for the taking of herring spawn on kelp and seines for the taking of herring for placement in pounds may be operated in the waters of the Gulf of Esquibel, San Alberto Bay, Shinaku Inlet, and San Christoval Channel south of the latitude of the northernmost tip of Saint Philip Island at 55°39.31' N lat, 133°25.12' W long, east of a line from the northernmost tip of Saint Philip Island to the northernmost tip of Point Garcia at 55°33.65' N lat, 133°26.47' W long, and north of a line from Entrance Point at 55°31.10' N lat, 133°09.00' W long to the southernmost tip of Clam Island at 55°30.96' N lat, 133°09.43' W long to the southernmost tip of Fern Point 55°30.05' N lat, 133°16.97' W long (Figure 1).

CLOSED WATERS

In Section 3-B, certain areas are closed to the operation of herring pounds and seines for taking of herring for placement in pounds. Those areas are shown in Figure 1 and include:

- Klawock Inlet and Big Salt Lake
- those waters of San Christoval Channel in the main channel enclosed by a line from 55°35.62' N lat, 133°20.00' W long to 55°35.17' N lat, 133°20.00' W long to 55°33.37' N lat, 133°17.52' W long to 55°33.50' N lat, 133°17.28' W long
- those waters of Fish Egg and Ballenas Islands south of 55°31.00' N lat, north of the southernmost tip of Cape Suspiro at 55°27.47' N lat, 133°08.40' W long, and east of the longitude of Ballena Island Shoal Light at 133°13.25' W long

OTHER AGENCY REQUIREMENTS

Prospective pound operators are advised to consider other agency requirements for constructing and operating pounds in Craig/Klawock, Ernest Sound, Tenakee Inlet, and Hoonah Sound. Pound operators are urged to contact the Alaska Department of Natural Resources (ANDR), U.S. Forest Service (USFS), the U.S. National Marine Fisheries Service (NMFS), and the U.S. Coast Guard (USCG) to determine other regulations and requirements. Phone numbers for those agencies are listed below.

ALASKA DEPARTMENT OF NATURAL RESOURCES

The Alaska Department of Natural Resources manages the use of tidelands and submerged lands seaward of mean high water. Contact ADNR at (907) 465-3400.

U.S. FOREST SERVICE

The U.S. Forest Service has jurisdiction over and manages most of the lands above mean high tide. People who plan to use National Forest land in connection with the fishery must apply for a special use permit from the USFS prior to any occupancy. Special use permit applications are available

from local USFS offices. Completed applications should be submitted well in advance of operations to ensure that a permit is received in time for the fishery. Examples of use needing a permit include (but are not limited to): camping on National Forest land in conjunction with the commercial fishery and storage of gear on the National Forest.

U.S. NATIONAL MARINE FISHERIES SERVICE

Marine mammal interactions should be reported to the U.S. National Marine Fisheries Service at (907) 586-7221.

U.S. COAST GUARD

Structures such as floating fish pens are subject to the requirements of the Code of Federal Regulations, Title 33, Part 64. This regulation requires an owner to apply for a USCG permit and to install and maintain a light or other private aid to navigation if the USCG determines it to be necessary to protect maritime navigation.

Herring pounds used in the SOK pound fishery do not require permits for private aids to navigation at this time, provided the owners:

- Place two signs on opposite corners of the structure. These signs will be worded “Danger, Fish Pens” (Figure 3).
- Place a single, all-points white light on one corner of structures less than 400 square feet in size.
- Place a single, all-points white light on every corner of structures larger than 400 square feet in size.
- Anchor fish pens within the boundary areas specified in ADF&G regulation 5 AAC 27.185(f) (Figure 1).

If all these conditions are not met, the permit holder must apply to the USCG for an individual “Private Aids to Navigation Permit.” If you have questions, call the USCG Aids to Navigation office, at (907) 463-2254.

LIST OF CONTACTS

The following ADF&G, Division of Commercial Fisheries personnel may be contacted regarding this management plan:

Bo Meredith, Justin Breese, and Whitney Crittenden Area Management Biologists Ketchikan Area Office	2030 Sea Level Dr., Suite 205 Ketchikan, Alaska 99901 (907) 255-5195
Aaron Dupuis and Jason Jones Area Management Biologists Sitka Area Office	304 Lake St. Rm. 103 Sitka, Alaska 99835 (907) 747-6688
Dave Harris and Scott Forbes Area Management Biologists Douglas Regional Office	P.O. Box 110024 Juneau, Alaska 99811 (907) 465-4250
Paul Salomone, Tom Kowalske, and Katie Taylor Area Management Biologists Petersburg Area Office	P.O. Box 667 Petersburg, Alaska 99833 (907) 772-3801
Kyle Hebert Herring Research Biologist Douglas Regional Office	P.O. Box 110024 Douglas, Alaska 99811 (907) 465-4250
Troy Thynes Region I Management Coordinator Petersburg Area Office	P.O. Box 667 Petersburg, Alaska 99833 (907) 772-3801
Lowell Fair Region I Supervisor Douglas Regional Office	P.O. Box 110024 Douglas, Alaska 99811 (907) 465-4250

TABLES AND FIGURES

Table 1.—Craig/Klawock herring stock size and fishery summary, 1990–2021.

Season	Miles Spawn	Forecasted	Total GH L Bait and SOK ^c (Tons)	Bait GH(L)(Tons)	Bait Harvest (Tons)	Number of Permits	Exvessel Value
		Pre- fishery Biomass					
1990/91	22.0	18,350	N/A	2,841	3,272	27	\$981,600
1991/92	23.0	17,800	2,684	2,281	2,295	28	\$619,650
1992/93 ^e	8.4	12,350	1,602	1,362	623	10	\$150,960
1993/94	8.0	7,996	895	760	636	6	\$187,578
1994/95	5.5	6,778	725	617	***	***	***
1995/96	9.9	6,262	658	558	***	***	***
1996/97	13.2	6,755	715	615	517	4	\$137,788
1997/98 ^f	11.0	7,018	755	455	***	***	***
1998/99	15.4	6,951	750	450	***	***	***
1999/00	12.9	6,013	626	376	***	***	***
2000/01	16.7	9,091	1,058	635	***	***	***
2001/02	18.0	8,387	952	571	***	***	***
2002/03	11.2	6,045	630	378	***	***	***
2003/04	12.0	13,204	1,754	1,052	***	***	***
2004/05	18.0	15,577	2,217	1,330	553	3	\$199,012
2005/06	8.2	14,262	1,955	1,173	689	3	\$247,934
2006/07	22.3	13,768	1,860	1,116	576	3	\$139,000
2007/08	11.0	14,213	1,945	1,167	565	3	\$133,300
2008/09	17.0	14,213	1,945	1,167	142	3	\$51,304
2009/10	18.7	14,870	2,074	1,244	***	***	***
2010/11	14.8	17,886	2,710	1,140	***	***	***
2011/12	14.9	34,235	6,847	4,060	309	3	\$113,784
2012/13	15.3	23,391	4,060	2,436	321	3	\$116,000
2013/14	13.6	26,085	4,808	2,884	***	***	***
2014/15	11.5	15,803	2,362	1,367	964	3	\$396,205
2015/16	12.3	12,303	1,590	954	898	3	\$369,660
2016/17	22.8	7,833	872	523	527	3	\$210,800
2017/18	17.3	16,083	2,312	1,387	527	5	\$310,000
2018/19	28.9	22,810	3,906	2,344	710	6	\$301,750
2019/20	56.1	55,072	11,014	6,608	995	4	\$410,000
2020/21	34.2	97,282	19,456	11,674	540	5	\$225,000
2021/22		63,250	12,650	7,590			
Average	16.9	19,123	3,174	1,791	824	6	\$240,000

Note: *** indicates data is confidential.

^a Spawn year is second year of regulatory season listed in the adjacent year column.

^b Forecasted pre-fishery biomass values were estimated with hydroacoustics for 86/87, spawn deposition surveys for 87/88 to 92/93, and age-structured models for 93/94 to 18/19.

^c Spawn-On-Kelp (SOK).

^d Bait quota reduced to 1,600 tons on the grounds.

^e First year bait quota was split between pound fishery 85%:15%.

^f Herring allocation changed to 60% for the winter food and bait fishery, 40% to the pound fishery.

^g Updated forecast based on updated scale ages.

Table 2.—Craig/Klawock herring SOK detailed fishery summary, 2004–2021.

Statistic	2004	2005	2006	2007	2008	2009
Herring SOK Quota (tons) ^k	1,597	1,743	1,266	1,283	1,369	1,802
Total harvest SOK (tons)	50	115.2	28.9	44.5	148.5	137.3
Exvessel value	\$325,000	\$603,723	\$298,575	\$1,087,532	\$3,066,788	\$1,256,777
Average Price/lb	\$3.25	\$2.62	\$5.15	\$12.08	\$10.33	\$4.58
Average Income	\$3,420	\$9,011	\$8,782	\$23,139	\$25,138	\$9,107
Number of pounds	50	42	50	52	66	96
Number of landings	95	67	34	47	122	137
Blade allocation	d	f	g	g	g	g
Total kelp harvest (tons)	14.0	4.9	4.6	5.6	12.2	7.3
Herring spawning dates	3/26–4/7	4/9–4/14	3/30–4/3	4/3–4/12	4/3–4/12	4/3–4/10
Miles of spawn	12.0	18.0	8.2	22.3	11.0	17.0
Forecasted Pre-Fishery biomass (tons)	13,204	15,577	14,262	13,768	14,213	14,213
	2010	2011	2012	2013	2014	2015
Herring SOK Quota (tons) ^k	1,953	2,402	6,538	3,739	4,544	1,308
Total harvest SOK (tons)	116.7	70	98.1	137.7	confidential	confidential
Exvessel value	\$884,715	\$728,147	\$2,099,002	\$3,099,002	confidential	confidential
Average Price/lb	\$3.80	\$5.13	\$10.69	\$12.00	confidential	confidential
Average Income	\$8,268	\$14,003	\$32,795	\$23,656	confidential	confidential
Number of pounds	63	34	35	80	75	76
Number of landings	107	52	64	131	136	135
Blade allocation	g	g	g	g	g	g
Total kelp harvest (tons)	8.2	4.6	5.3	9.3	19.2	19.2
Herring spawning dates	4/5–4/14	4/1–4/7	4/3–4/8	3/31–4/3	4/1–4/5	3/27–4/1
Miles of spawn	18.7	14.8	14.9	15.3	13.6	11.5
Forecasted Pre-Fishery biomass (tons)	14,870	17,886	34,235	23,391	26,085	15,803
	2016	2017	2018	2019	2020	2021
Herring SOK Quota (tons) ^k	692	349	1,602	2,895	10,119	18,916
Total harvest SOK (tons)	confidential	69.9	205.3	202.4	283.9	262.6
Exvessel value	confidential	\$932,917	\$3,262,900	\$3,300,000	\$3,256,000	\$2,150,000
Average Price/lb	confidential	\$6.68	\$7.95	\$8.15	\$5.73	\$4.77
Average Income	confidential	\$8,042	\$24,533	\$23,358	\$21,853	\$15,468
Number of pounds	46	19	66	73	75	80
Number of landings	133	116	133	141	149	158
Blade allocation	i	h	j	j	j	j
Total kelp harvest (tons)	9.2	10.2	9.4	9.8	11.4	12.5
Herring spawning dates	3/25–3/31	3/24–4/2 & 4/6–4/8	3/29–4/4	3/27–4/6	4/3–4/13	4/5–4/14
Miles of spawn	12.3	22.8	17.3	28.9	56.1	34.2
Forecasted Pre-Fishery biomass (tons)	12,303	7,833	16,039	22,810	86,087	97,282

^a 100 blades for single-closed pound, 150 blades for multiple pound permit holder.^b 120 blades for a single closed pound, 180 blades for a multiple pound permit holder.^c 155 blades for a single closed pound, 235 blades for a multiple pound permit holder.^d 70 blades for a single closed pound, 210 for a multiple pound permit holder.^e 200 blades - single closed pound, 600 blades/permit holder multiple closed pound.^f 350 blades for a single closed pound, 750 blades for a double closed pound, 1,125 blades for a triple closed pound.^g 600 blades for a single closed pound, 750 blades for a double closed pound, 1,125 blades for a triple closed pound.^h 500 blades per permit with six permit holders per pound structureⁱ 300 blades for a single closed pound, 400 blades for a double closed pound, 700 blades for a triple closed pound.^j 600 blades for a single closed pound, 900 blades for a double closed pound, 1,000 blades for a triple closed pound.^k The quota was updated to include the total SOK quota which includes the carry-over from unharvested winter bait.

16




Table 4.—Ernest Sound herring stock and fishery summary, 1969–2021.

Season	Date of First Spawn ^a	Nautical Miles of Spawn ^b	Forecast used for GHL Determination ^{c,d} (tons)	Spawning Biomass (tons) ^e	GHL (tons) ^f	Bait Harvest (tons) ^g	SOK Harvest (lbs)	Sac Roe Harvest (tons)	Remaining GHL (tons)
1969/70	—	—	—	—	—	17	—	—	—
1970/71	—	3	—	13,100	—	206	—	—	—
1971/72	—	—	13,100	3,650	—	967	—	—	—
1972/73	—	—	3,650	450	—	775	—	—	—
1973/74	—	—	450	400	—	535	—	—	—
1974/75	—	—	400	2,900	—	593	—	—	—
1975/76	—	3	2,900	4,350	580	708	—	—	—
1976/77 ^g	—	3	4,350	3,035	870	901	—	49	0
1977/78	5/3	—	3,035	1,505	455	340	—	—	0
1978/79	4/16	2.6	1,505	255	—	—	—	—	115
1979/80	5/2	4	255	500	—	—	—	—	—
1980/81	—	3.5	500	410	—	—	—	—	—
1981/82	—	—	410	160	—	—	—	—	—
1982/83	—	—	160	1,640	—	—	—	—	—
1983/84	4/11	—	1,640	1,000	—	—	—	—	—
1984/85	—	4.5	1,000	1,000	—	—	—	—	—
1985/86	—	—	1,000	1,000	—	—	—	—	—
1986/87	—	1	1,000	—	—	—	—	—	—
1987/88	4/21	2	—	—	—	—	—	—	—
1988/89	4/17	2.4	—	500	—	—	—	—	—
1989/90	—	2.1	500	1,000	—	—	—	—	—
1990/91	—	—	1,000	3,000	—	—	—	—	—
1991/92	4/16	9.1	3,000	2,650	—	—	—	—	—
1992/93	4/23	9	2,650	684	200	8	—	—	192
1993/94	4/26	8.4	684	2,544	0	—	—	—	—
1994/95	4/25	6.5	2,544	2,470	255	111	—	—	144
1995/96	4/16	6.9	2,744	2,665	280	220	—	—	60
1996/97	4/16	0	4,852	0	377	6	—	—	371
1997/98	4/9	11.8	—	5,998	0	—	—	—	—
1998/99	4/5	1.8	5,381	No survey	662	96	—	—	566
1999/00	4/8	9.1	—	920	0	—	—	—	—
2000/01	4/11	6.9	—	2,052	0	—	—	—	—
2001/02	4/15	4.8	1,653	2,406	0	—	—	—	—

-continued-

Table 4.–continued (Page 2 of 2).

Season	Date of First Spawn ^a	Nautical Miles of Spawn ^b	Forecast Used for GHL Determination ^{c,d} (tons)	Spawning Biomass (tons) ^e	GH L (tons) ^f	Bait harvest (tons) ^g	SOK Harvest (lbs)	Sac Roe Harvest (tons)	Remaining GH L (tons)
2002/03	4/10	8.5	2,407	5,509	0	—	—	—	—
2003/04	4/11	7.1	6,592	2,413	875	44	112,286	—	831
2004/05	4/22	10.1	1,906	3,268	0	—	—	—	—
2005/06	4/6	7.9	2,284	2,538	0	—	—	—	—
2006/07	4/19	11.3	1,955	7,353	0	—	—	—	—
2007/08	4/20	15.4	9,060	4,846	1382	***	19,650	—	>700
2008/09	4/17	6.6	4,545	2,862	529	***	4911	—	100-299
2009/10	4/14	7.8	2,879	3,523	297	***	—	—	<50
2010/11	4/17	8.1	5,080	2,559	476	***	0	—	100-299
2011/12	4/16	8.9	2,682	3,193	272	***	—	—	<50
2012/13	4/16	9.7	3,509	7,556	379	***	129,265	—	100-299
2013/14	4/14	3.7	7,613	2,631	1073	***	***	—	>700
2014/15	4/8	5.5	1,991	562	0	—	—	—	—
2015/16	4/10	4.4	1,207	346	0	—	—	—	—
2016/17	4/14	4.4	—	—	—	—	—	—	—
2017/18	4/24	3.5	—	—	—	—	—	—	—
2018/19	4/14	1	—	—	—	—	—	—	—
2019/20	4/19	0.3	—	—	—	—	—	—	—
2020/21	4/15	0.5	—	—	—	—	—	—	—
Recent 10-yr Average.	4/15	4.2	3,400	2,858	345	**	**	—	317
Long term Average	4/16	5.6	2,852	2,591	331	44	**	—	253

Note: En dashes indicate no data and *** indicates data is confidential.

^a Since the 1997/1998 season, the first spawn and the major spawn have been within five days of each other.

^b 1996/1997 No spawn deposition survey conducted; 7.5 miles of spawn observed along Ship Island;

1998/1999 No spawn deposition survey conducted; 1.8 miles of spawn observed

^c from the 1971/1972 season through the 1984/1985 season forecasts were based on computer generated estimates derived using hydroacoustic techniques; from 1985/1986 through 1991/1992 forecasts were based on visual estimates; from 1992/1993 through 1994/1995 forecasts were based on spawn deposition estimates; from 1995/1996 through 2006/2007 the method used was biomass accounting forecasts.

^d beginning with the 1976/1977 season fishery threshold established at 2,500 tons.

^e from 1969/1970 through 1983/1984 biomass estimates were -computer generated estimates derived using hydroacoustic techniques; from 1984/1985 through 1990/1991 estimates were based on visual observations and from 1991/1992 through 2005/2006 were spawn deposition estimates. 1975/1976 & 1976/1977 GH Ls are based upon 20% of the hydroacoustic estimate. 1977/1978 GH L is based upon 15% of the hydroacoustic estimate.

^f 2003/2004 GH L includes 90 tons rolled over from the bait pound fishery.

^g 1973/1974, 1974/1975, 1976/1977 also include harvests from Fools and Menefee Inlets. Does not include harvests from statistical area 107-40.

Table 5.—Ernest Sound SOK detailed fishery summary, 2004, 2008–2009, 2011, and 2013–2014.

Statistic	2004	2008	2009
GHL (tons)	875	1,382	529
GHL Available for SOK (tons)	775	***	***
SOK Harvest (tons)	56.1	9.8	2.5
Exvessel Value	\$514,900	\$131,100	\$18,400
Average Price/lb	\$4.59	\$6.87	\$3.67
Average Income/permit	\$8,046	\$10,091	\$4,590
Number of Permits	64	13	4
Number of Pounds	51/6/0/1 ^a	1/6/0/0 ^a	0/2/0/0 ^a
Number of Landings	64	13	4
Kelp Allocation (blades)	b	c	d
Kelp Blade Harvest (lbs)	4,600	29,400	2,400
Fishery Open—Closed	4/1–5/10	4/1–5/10	4/1–5/10
Fishing Occurred	4/10–4/17	4/16–4/24	4/17–4/21
Harvest Occurred	4/14–1/17	24-Apr	21-Apr
	2011	2013	2014
GHL (tons)	476	379	1,073
GHL Available for SOK (tons)	***	***	***
SOK Harvest (tons)	0	64.3	***
Exvessel Value	\$0	\$1,574,700	***
Average Price/lb	\$0	\$12.25	***
Average Income/permit	\$0	\$19,441	***
Number of Permits	0	81	129
Number of Pounds	0	1/5/1/22/0 ^e	25/52/0 ^a
Number of Landings	0	81	129
Kelp Allocation (blades)	d	d	c
Kelp Blade Harvest (lbs)	0	6,400	29,000
Fishery Open—Closed	4/1–5/10	4/1–5/10	4/1–5/10
Fishing Occurred	NA	4/10–4/16	4/12–4/20
Harvest Occurred	NA	4/15–4/16	4/18–4/20

Note: *** indicates data is confidential.

^a Single/double/triple/open.

^b 1,000 blades single-closed/1,000 blades double-closed/1,000 blades triple-closed/2,500 blades single-open/7,500 multiple-permit-open.

^c 1,000 blades single-closed/2,000 blades double-closed/1,000 blades triple-closed/2,500 blades single-open/7,500 multiple-permit-open.

^d 200 blades single-closed/400 blades double-closed/500 blades triple-closed/1,500 blades single-open/4,500 multiple-permit-open.

^e Single/double/combined double/triple/open.

Table 6.—Tenakee Inlet herring stock and fishery summary, 1978–2021.

Season	Major Spawning Dates	Nautical Miles of Spawn	Spawning Biomass ^a (tons)	Food/Bait GHL (tons)	Food/Bait Harvest (tons)
1978/79	5/9–5/11	3.3	2,500	200	0
1979/80	4/28–5/2	3.9	4,485	400	504
1980/81	4/27–5/5	9.3	7,500	750	847
1981/82	4/25–5/7	11.1	6,650	650	687
1982/83	4/25–5/6	13.1	8,870	875	749
1983/84	4/20–4/26	8.3	12,100	850	619
1984/85	4/24–5/1	9.9	11,000	1,400	1,406
1985/86	4/27–5/1	8.3	12,500	1,700	2,040
1986/87	4/22–4/30	7.9	6,600	800	1,275
1987/88	4/22–4/27	9.1	6,000	1,450	1,577
1988/89	4/26–4/29	10.3	5,360	720	655
1989/90	4/25–5/6	2.9	2000	650	595
1990/91	4/25–5/4	2.1	400	—	—
1991/92	5/5	trace	200	—	—
1992/93	4/21–4/23	6.4	904	—	—
1993/94	4/24–4/26	0.25	400	—	—
1994/95	4/26	0.05	200	—	—
1995/96	5/4–5/14	18.1	4,560	—	—
1996/97	4/26–5/7	14.4	9,926	300	98
1997/98	4/24–4/29	12.4	10,419	825	586
1998/99	4/25–4/28	11	11,049	1,023	835
1999/00	4/26–5/3	13.8	9,425	542	494
2000/01	4/21–5/1	12.2	7,576	906	775
2001/02	4/23–4/27	15.4	4,084	840	355
2002/03	4/25–4/28	12.2	3,529	528	328
2003/04	4/28–5/3	13	4,728	399	***
2004/05	4/26–5/2	8.9	3,036	476	0
2005/06	5/2–5/6	5.9	5,110	—	—
2006/07	4/23–4/26	4.4	3,346	—	—
2007/08	4/30; 5/7–5/8	11.4	11,252	—	—
2008/09	4/25–4/26; 4/29–4/30	6.9	5,283	875	254
2009/10	5/7–5/9	2.7	1,437	583	***
2010/11	5/9	1	—	—	—
2011/12	4/20–4/23	4.6	5,119	—	—
2012/13	5/7–5/10	5.4	4,936	—	—

-continued-

Table 6.—continued (Page 2 of 2).

Season	Major Spawning Dates	Nautical Miles of Spawn	Spawning Biomass ^a (tons)	Food/Bait GHL (tons)	Food/Bait Harvest (tons)
2013/14	29-Apr	2	927	557	***
2014/15	4/22–4/25	2.3	2,223	—	—
2015/16	—	0	—	—	—
2016/17	5/13–5/16	2.1	—	—	—
2017/18	5/7–5/9	1.4	—	—	—
2018/19	5/12–5/14	0.5	—	—	—
2019/20	5/5–5/6	1.9	—	—	—
2020/21	4/20–4/21	1.7	—	—	—

Note: En dashes indicate no data and *** indicates data is confidential.

^a Spawning biomass estimates were calculated from hydro-acoustical surveys from 1979 through 1986, and from spawn deposition surveys from 1987 through 2015—bolded values were derived from ASA models.

Table 7.—Tenakee Inlet herring SOK detailed fishery summary, 2003–2005, 2009, and 2014.

Statistic	2003	2004	2005
GHL (tons)	528	399	476
GHL Available for SOK	180	***	476
SOK Harvest (tons)	47.6	100.7	101.4
Exvessel Value	\$580,500	\$981,500	\$512,900
Average Price/lb	\$6.10	\$4.68	\$2.53
Average Income/permit	\$10,560	\$11,680	\$5,640
Number of Permits	55	85	98
Number of Pounds	1/15/8/0 ^a	1/32/6/2/2 ^b	1/29/13/3 ^a
Number Permits Landing	55	85	91
Kelp Allocation (blades)	200/400/550/0 ^a	300/500/500/2000 ^a	300/500/500/2000 ^a
Kelp Blade Harvest (lbs)	35,375	39,000	53,850
Fishery Open—Closed	4/6–5/6	4/6–5/6	4/6–5/5
Fishing Occurred	4/25–4/28	4/28–5/1	4/27–4/30
Harvest Occurred	4/30–5/4	5/3–5/6	5/2–5/4
Statistic	2009	2014	
GHL (tons)	875	557	
GHL Available for SOK	621	***	
SOK Harvest (tons)	64.1	84.4	
Exvessel Value	\$558,900	\$1,155,300	
Average Price/lb	\$4.36	\$6.85	
Average Income/permit	\$6,500	\$16,270	
Number of Permits	86	78	
Number of Pounds	11/27/7/0 ^a	2/25/5/1 ^c	
Number Permits Landing	86	71	
Kelp Allocation (blades)	400/500/500/0 ^a	300/500/500/0 ^a	
Kelp Blade Harvest (lbs)	42,600	41,250	
Fishery Open—Closed	4/6–5/5	4/6–5/4	
Fishing Occurred	4/28–5/1	4/27–5/1	
Harvest Occurred	5/2–5/5	5/2–5/4	

Note: *** indicates data is confidential.

^a Single/double/triple/test.

^b Single/double/triple/long line/test.

^c Single/double/triple/quadruple.

Table 8.–Hoonah Sound and Tenakee Inlet herring spawn dates comparison, 2001–2021.



Note: Black bars indicate spawning dates for Hoonah Sound and gray bars indicate spawning dates for Tenakee Inlet.

Table 9.—Hoonah Sound herring spawning stock and fishery summary, 1971–2021.

Year	Spawn Dates	Nautical Miles of Spawn	Spawning Biomass (tons)	SOK Harvest (tons)
1971	5/10–5/17	2.5	833	—
1972	5/11–5/12	1.5	666	—
1973	N/A	1.0	333	—
1974	14-May	3.0	999	—
1975	N/A	N/A	—	—
1976	5-May	1.0	333	—
1977	N/A	3.5	1,166	—
1978	N/A	5.3	1,765	—
1979	N/A	0.5	167	—
1980	N/A	N/A	—	—
1981	4/30–5/01	2.3	750	—
1982	4/29–5/01	1.5	500	—
1983	1-May	1.0	333	—
1984	4/26–5/01	3.0	540	—
1985	5/01–5/03	3.5	1,166	—
1986	4/28–5/01	3.8	1,249	—
1987	4/28–5/02	3.8	740	—
1988	4/30–5/01	5.0	1,665	—
1989	4/16–4/20	17.0	4,000	—
1990	4/13–4/28	10.0	2,350	12.0
1991	4/19–4/24	8.7	2,175	13.6
1992	4/22–4/24	10.8	5,714	23.1
1993	4/27–4/29	5.7	1,099	14.8
1994	4/21–4/23	9.0	2,450	33.3
1995	4/20–4/21	4.5	274	28.7
1996	5/02–5/9	10.1	4,023	—
1997	4/25–4/28	14.5	5,884	64.3
1998	4/23–4/27	14.5	6,472	85.6
1999	4/27–5/1	13.8	4,426	93.8
2000	4/27–4/30	13.0	3,635	36.0
2001	4/27–5/1	13.7	8,538	66.2
2002	4/25–4/27	11.9	4,936	136.6
2003	4/23–4/27	16.7	9,423	141.6
2004	4/22–4/29	11.1	7,502	236.1
2005	4/18–4/25	10.3	6,924	181.7
2006	4/23–4/26	9.0	6,028	162.1
2007	4/46–5/2	16.5	10,946	159.4
2008	4/23–4/30	14.5	19,975	228.1
2009	4/22–4/27	10.3	15,829	234.7
2010	4/22 – 4/28	12.4	15,264	290.4
2011	4/20–5/5	12.6	14,215	193.7
2012	4/20–4/23	4.2	923	186.5
2013	5/7–5/9	2.4	412	—
2014	5/5–5/8	3.2	444	—
2015	4/24–5/6	4.2	22	—
2016	N/A	N/A	—	—
2017	N/A	N/A	—	—
2018	N/A	N/A	—	—
2019	N/A	N/A	—	—
2020	N/A	N/A	—	—
2021	N/A	N/A	—	—
Average	1971–2020	6.8	3,689	N/A
Average	1990–2020	8.7	5,158	87.4

Note: Shaded estimated escapements are based on average spawn density from years 1989 to 2002; due to funding, comprehensive aerial surveys have not been conducted since 2016. En dashes indicate no data and N/A indicates no applicable since no spawn occurred.

Table 10.—Hoonah Sound herring SOK detailed fishery summary, 2003–2017.

Statistic	2003	2004	2005	2006	2007
Herring GHL (tons)	427	1,207	728	669	681
Harvest (tons)	141.6	237.4	190.6	162.1	144.5
Exvessel Value	\$1,922,500	\$2,071,347	\$1,117,568	\$1,943,422	\$4,491,070
Average Price/lb	\$6.79	\$4.36	\$2.93	\$6.00	\$14.09
Average Income	\$17,800	\$19,541	\$11,889	\$24,600	\$49,352
Number of Pounds	49/1/3 ^a	92/12/2 ^b	81/5/3 ^c	17/45 ^a	67/12 ^a
Number Selling Product	108	106	94	79	91
		1,000/1,000/	1,000/1,000/	2,500/1,000/	2,500/1,000/
Kelp Allocation (blades)	500/300/750 ^a	3,000 ^b	1,500 ^c	1,500 ^a	1,500 ^a
Kelp Blade Harvest	60,301	126,000	118,450	136,698	122,565
Fishery Open—Closed	4/6–4/25	4/6–4/28	4/6–4/28	4/6–4/27	4/6–5/4
Fishing Occurred	4/19–4/24	4/20–4/25	4/19–4/28	4/18–4/23	4/23–4/29
Harvest Occurred	4/24–4/27	4/26–4/28	4/25–4/28	4/23–4/27	4/30–5/4
Statistic	2008	2009	2010	2011	2012
Herring Quota (tons)	2,238	2,238	3,182	3,015	2,139
Harvest (tons)	223	234.7	290.4	193.7	186.5
Exvessel Value	\$5,115,459	\$2,332,514	\$2,580,517	\$1,820,952	\$4,033,078
Average Price/lb	\$11.47	\$4.97	\$4.44	\$4.70	\$10.81
Average Income/Landing	\$51,155	\$23,094	\$25,550	\$20,460	\$55,248
Number of Pounds	98/3 ^a	99/4 ^a	97/2 ^a	85/4 ^a	83/4 ^a
Number Selling Product	100	101	101	89	73
	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/	3,000/2,000/
Kelp Allocation (blades)	1,500 ^a	1,500 ^a	1,500 ^a	1,500 ^a	1,500 ^a
Kelp Blade Harvest	201,262	196,492	178,898	169,922	155,104
Fishery Open—Closed	4/6–5/02	4/6–4/30	4/6–4/30	4/6–4/30	4/6–5/15
Fishing Occurred	4/22–4/27	4/22–4/25	4/21–4/25	4/20–4/24	4/19–4/21
Harvest Occurred	4/27–5/1	4/26–4/29	4/22–4/28	4/26–4/29	4/26–4/27
Statistic	2013	2014	2015	2016	2017
Herring Quota (tons)	130	0	0	0	0
Harvest (tons)	0	0	0	0	0
Exvessel Value	\$0	\$0	\$0	\$0	\$0
Average Price/lb	\$0	\$0	\$0	\$0	\$0
Average Income/Landing	\$0	\$0	\$0	\$0	\$0
Number of Pounds	3 open	0	0	0	0
Number Selling Product	0	0	0	0	0
Kelp Allocation (blades)	600 open	0	0	0	0
Kelp Blade Harvest	7,940	0	0	0	0
Fishery Open—Closed	4/6–5/15	NA	NA	NA	NA
Fishing Occurred	NA	NA	NA	NA	NA
Harvest Occurred	NA	NA	NA	NA	NA

Note: Fisheries did not occur in 1996 and 2014–2021 since the biomass forecast was either below the 1,000-ton threshold or no survey was conducted due to minimum herring spawn observed.

^a Double closed pounds/single closed pounds/triple closed pounds.

^b Double closed pounds/single closed pounds/open pounds.

^c Single-permit closed pound/double-permit closed pound/triple-permit closed pounds.




Figure 1.—Craig/Klawock (Section 3-B) herring SOK fishery open area.




Figure 2.—Diagram of a herring pound showing two alternative methods of marking herring pounds.

Note: Regulations require vertical signs with the permit holder's first and last name and five-digit CFEC permit number (5 AAC 27.185(k)). Letters and numbers must be at least six inches high and at least one-half inch wide, must contrast with the background and must be always above the waterline.




Figure 3.—U.S. Coast Guard requirements for marking pounds.