

**YUKON RIVER SALMON 2021 SEASON SUMMARY  
AND 2022 SEASON OUTLOOK**

Prepared by

THE UNITED STATES AND CANADA  
YUKON RIVER JOINT TECHNICAL COMMITTEE

March 2022

Regional Information Report 3A22-01

Alaska Department of Fish and Game

333 Raspberry Road

Anchorage, AK 99518, USA



## Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

| Weights and measures (metric)           |                    | General                                  |   | Mathematics, statistics                                    |                         |
|---|--------------------|--|---|--|-------------------------|
| centimeter                              | cm                 | Alaska Administrative Code               |   | all standard mathematical signs, symbols and abbreviations |                         |
| deciliter                               | dL                 |  | AAC   |  |                         |
| gram                                    | g                  | all commonly accepted abbreviations      | e.g., Mr., Mrs., AM, PM, etc.               | alternate hypothesis                                       | H <sub>A</sub>          |
| hectare                                 | ha                 |  |   | base of natural logarithm                                  | <i>e</i>                |
| kilogram                                | kg                 | all commonly accepted                    |   | catch per unit effort                                      | CPUE                    |
| kilometer                               | km                 | professional titles                      | e.g., Dr., Ph.D., R.N., etc.                | coefficient of variation                                   | CV                      |
| liter                                   | L                  |  |   | common test statistics                                     | (F, t, $\chi^2$ , etc.) |
| meter                                   | m                  | at                                       | @   | confidence interval  | CI                      |
| milliliter                              | mL                 | compass directions:                      |   | correlation coefficient (multiple)                         | R                       |
| millimeter                              | mm                 | east                                     | E   | correlation coefficient (simple)                           | r                       |
| <b>Weights and measures (English)</b>   |                    | north                                    | N   | covariance   | cov                     |
| cubic feet per second                   | ft <sup>3</sup> /s | south                                    | S   | degree (angular )  | °                       |
| foot                                    | ft                 | west                                     | W   | degrees of freedom   | df                      |
| gallon                                  | gal                | copyright                                | ©   | expected value   | <i>E</i>                |
| inch                                    | in                 | corporate suffixes:                      |   | greater than   | >                       |
| mile                                    | mi                 | Company                                  | Co.   | greater than or equal to                                   | ≥                       |
| nautical mile                           | nmi                | Corporation                              | Corp.                                       | harvest per unit effort                                    | HPUE                    |
| ounce                                   | oz                 | Incorporated                             | Inc.  | less than  | <                       |
| pound                                   | lb                 | Limited                                  | Ltd.  | less than or equal to                                      | ≤                       |
| quart                                   | qt                 | District of Columbia                     | D.C.  | logarithm (natural)  | ln                      |
| yard                                    | yd                 | et alii (and others)                     | et al.                                      | logarithm (base 10)  | log                     |
|   |                    | et cetera (and so forth)                 | etc.  | logarithm (specify base)                                   | log <sub>2</sub> etc.   |
| <b>Time and temperature</b>             |                    | exempli gratia                           |   | minute (angular)   | '                       |
| day                                     | d                  | (for example)                            | e.g.  | not significant  | NS                      |
| degrees Celsius                         | °C                 | Federal Information Code                 | FIC   | null hypothesis  | H <sub>0</sub>          |
| degrees Fahrenheit                      | °F                 | id est (that is)                         | i.e.  | percent  | %                       |
| degrees kelvin                          | K                  | latitude or longitude                    | lat or long                                 | probability  | P                       |
| hour                                    | h                  | monetary symbols                         |   | probability of a type I error                              |                         |
| minute                                  | min                | (U.S.)                                   | \$, ¢                                       | (rejection of the null hypothesis when true)               | $\alpha$                |
| second                                  | s                  | months (tables and figures): first three |   | probability of a type II error                             |                         |
| <b>Physics and chemistry</b>            |                    | letters                                  | Jan,...,Dec                                 | (acceptance of the null hypothesis when false)             | $\beta$                 |
| all atomic symbols                      |                    | registered trademark                     | ®   | second (angular)   | "                       |
| alternating current                     | AC                 | trademark                                | ™   | standard deviation   | SD                      |
| ampere                                  | A                  | United States                            |   | standard error   | SE                      |
| calorie                                 | cal                | (adjective)                              | U.S.  | variance   |                         |
| direct current                          | DC                 | United States of America (noun)          | USA   | population   | Var                     |
| hertz                                   | Hz                 | U.S.C.                                   | United States Code                          | sample   | var                     |
| horsepower                              | hp                 |  |   |  |                         |
| hydrogen ion activity (negative log of) | pH                 | U.S. state                               | use two-letter abbreviations (e.g., AK, WA) |  |                         |
| parts per million                       | ppm                |  |   |  |                         |
| parts per thousand                      | ppt, ‰             |  |   |  |                         |
| volts                                   | V                  |  |   |  |                         |
| watts                                   | W                  |  |   |  |                         |

***REGIONAL INFORMATION REPORT NO. 3A22-01***

**YUKON RIVER SALMON 2021 SEASON SUMMARY  
AND 2022 SEASON OUTLOOK**

The United States and Canada  
Yukon River Joint Technical Committee

Alaska Department of Fish and Game  
Division of Commercial Fisheries  
333 Raspberry Road, Anchorage, Alaska, 99518-1565

March 2022

The Regional Information Report Series was established in 1987 and was redefined in 2007 to meet the Division of Commercial Fisheries regional need for publishing and archiving information such as area management plans, budgetary information, staff comments and opinions to Alaska Board of Fisheries proposals, interim or preliminary data and grant agency reports, special meeting or minor workshop results and other regional information not generally reported elsewhere. Reports in this series may contain raw data and preliminary results. Reports in this series receive varying degrees of regional, biometric and editorial review; information in this series may be subsequently finalized and published in a different department reporting series or in the formal literature. Please contact the author or the Division of Commercial Fisheries if in doubt of the level of review or preliminary nature of the data reported. Regional Information Reports are available through the Alaska State Library and on the Internet at: <http://www.adfg.alaska.gov/sf/publications/>.

Product names used in this publication are included for completeness and do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

*This document should be cited as follows:*

*JTC (Joint Technical Committee of the Yukon River U.S./Canada Panel). 2022. Yukon River salmon 2021 season summary and 2022 season outlook. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 3A22-01, Anchorage.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

# TABLE OF CONTENTS

|  | Page |
|--|------|
| TABLE OF CONTENTS .....  | i    |
| LIST OF TABLES.....  | iii  |
| LIST OF FIGURES .....  | iii  |
| LIST OF APPENDICES .....   | iv   |
| 1.0 ABSTRACT .....   | 1    |
| 2.0 INTRODUCTION .....   | 1    |
| Yukon River Salmon Agreement Management Performance Summary .....                                      | 2    |
| 2021 Total Run Size, Harvest, and Escapement of Canadian-origin Chinook Salmon .....                   | 2    |
| 2021 Total Run Size, Harvest, and Escapement of Canadian-origin Fall Chum Salmon .....                 | 2    |
| 2022 Outlooks.....   | 3    |
| 2022 Escapement Goals.....   | 3    |
| 3.0 ALASKA MANAGEMENT OVERVIEW .....   | 3    |
| 3.1 Chinook and Summer Chum Salmon .....   | 3    |
| Preseason Management Strategy Planning .....   | 4    |
| Chinook and Summer Chum Salmon Inseason Management .....   | 4    |
| 3.2 Fall Chum And Coho Salmon.....   | 6    |
| Fall Chum Salmon Management Overview .....   | 7    |
| Coho Salmon Management Overview .....  | 8    |
| 4.0 ALASKA HARVEST SUMMARIES .....   | 8    |
| 4.1 Subsistence Salmon Fishery .....   | 8    |
| 4.2 Commercial Fishery .....   | 10   |
| Summer Season Harvest .....  | 10   |
| Fall Season Harvest .....  | 10   |
| 4.3 Sport Fishery.....   | 10   |
| 4.4 Personal Use Fishery .....   | 11   |
| 5.0 CANADIAN MANAGEMENT OVERVIEW .....   | 11   |
| 6.0 CANADIAN HARVEST SUMMARIES .....   | 16   |
| 7.0 TOTAL RUN, ESCAPEMENT, AND HARVEST SHARE ASSESSMENTS FOR 2021.....                                 | 18   |
| 7.1 Chinook Salmon .....   | 18   |
| 7.2 Summer Chum Salmon Alaska (U.S. Only) .....  | 20   |
| 7.3 Fall Chum Salmon .....   | 21   |
| Mainstem Yukon River Canadian-origin Fall Chum Salmon.....   | 22   |
| Porcupine River (Including the Fishing Branch River) Canadian-origin Fall Chum Salmon.....             | 22   |
| 8.0 PROJECT SUMMARIES.....   | 22   |
| 8.1 Alaska, U.S.....   | 22   |
| Lower Yukon Test Fishery .....   | 23   |
| Pilot Station Sonar .....  | 24   |
| <i>Ichthyophonus</i> investigations.....   | 25   |
| Chinook Salmon Genetic Sampling, 2021 .....  | 25   |
| Mixed Stock Analysis of Yukon River Chinook Salmon Sampled at the Pilot Station Sonar, 2005–2021 ..... | 25   |
| Mixed Stock Analysis of Yukon River Chinook Salmon Harvested in Alaska, 2021 .....                     | 26   |

## TABLE OF CONTENTS (Continued)

|   | Page |
|---|------|
| Yukon River Chum Salmon Mixed Stock Analysis, 2021 .....  | 27   |
| Environmental Conditions Report .....   | 27   |
| 8.2 Eagle Sonar.....  | 28   |
| 8.3 Yukon, Canada .....   | 29   |
| Yukon River (Mainstem) Adult Chinook Salmon Assessment .....  | 29   |
| Big Salmon Sonar.....   | 29   |
| Pelly River Sonar .....   | 29   |
| Klondike River Sonar.....   | 30   |
| Whitehorse Rapids Fishway Chinook Salmon Enumeration .....  | 30   |
| Whitehorse Hatchery Operations .....  | 30   |
| Porcupine River Investigations .....  | 31   |
| Porcupine River Chinook Salmon Sonar.....   | 31   |
| Fishing Branch River Chum Salmon Weir.....  | 32   |
| Aerial Surveys .....  | 33   |
| Kluane River Aerial Survey .....  | 33   |
| Mainstem Yukon River Aerial Survey .....  | 33   |
| Genetic Stock Identification and Stock Composition of Canadian Yukon River Chinook and Fall Chum Salmon ..... | 33   |
| Environmental Conditions Report .....   | 34   |
| November 2020 to April 2021.....  | 34   |
| May 2021 to July 2021 .....   | 35   |
| August 2021 to November 2021.....   | 35   |
| Summary .....   | 36   |
| 9.0 MARINE FISHERIES INFORMATION .....  | 36   |
| 10.0 RUN OUTLOOKS 2022 .....  | 37   |
| 10.1 Yukon River Chinook Salmon.....  | 37   |
| Canadian-origin Brood Table .....   | 37   |
| Canadian-origin Yukon River Chinook Salmon.....   | 37   |
| Dynamic Sibling Model .....   | 38   |
| Spawner-recruit Model.....  | 38   |
| Juvenile-based Forecast.....  | 38   |
| 2022 Canadian-origin Chinook Salmon Forecast.....   | 39   |
| 10.2 Yukon River Summer Chum Salmon .....   | 39   |
| 10.3 Yukon River Fall Chum Salmon .....   | 40   |
| Drainagewide Fall Chum Salmon.....  | 40   |
| Canadian-origin Upper Yukon River Fall Chum Salmon.....   | 42   |
| Canadian-origin Porcupine River Fall Chum Salmon .....  | 42   |
| 10.4 Yukon River Coho Salmon.....   | 43   |
| 11.0 STATUS OF ESCAPEMENT GOALS.....  | 43   |
| 11.1 Spawning Escapement Target Options in 2021 .....   | 43   |
| Canadian-origin Mainstem Yukon River Chinook Salmon .....   | 43   |
| Canadian-origin Mainstem Yukon River Fall Chum Salmon .....   | 44   |
| Fishing Branch River Fall Chum Salmon .....   | 44   |
| 12.0 REFERENCES CITED .....   | 45   |

## TABLE OF CONTENTS (continued)

|   | <b>Page</b> |
|---|-------------|
| TABLES AND FIGURES .....  | 47          |
| APPENDIX A: TABLES .....  | 85          |
| APPENDIX B: TABLES .....  | 101         |
| APPENDIX C: BERING SEA-ALEUTIAN ISLANDS BYCATCH SUMMARY AND IMPACT ON YUKON<br>RIVER CANADIAN-ORIGIN SALMON ..... | 169         |

## LIST OF TABLES

| <b>Table</b>  | <b>Page</b> |
|---|-------------|
| 1 Yukon Area regulatory subsistence salmon fishing schedule, in U.S. waters. ....   | 48          |
| 2 Yukon River drainage fall chum salmon management plan overview, in U.S. waters. ....  | 48          |
| 3 Inseason fishery management decision matrix for Yukon River mainstem Chinook salmon in Canada,<br>2021 .....  | 49          |
| 4 Inseason fishery management decision matrix for mainstem Yukon River fall chum salmon in Canada,<br>2021 .....  | 49          |
| 5 Cumulative fish passage estimates by species with 90% confidence intervals (CI), at the Pilot Station<br>sonar in 2021 .....  | 50          |
| 6 Yukon River Chinook salmon age and female percentage estimated from samples collected at the Pilot<br>Station and Eagle sonar projects, 2021 .....                                    | 50          |
| 7 Summary of 2021 Chinook salmon escapement estimates in U.S. (Alaska) tributaries compared to<br>existing escapement goals. ....   | 51          |
| 8 Summary of 2021 summer chum salmon escapement estimates in U.S. (Alaska) compared to existing<br>escapement goals. ....   | 51          |
| 9 Summary of 2021 fall chum salmon escapement estimates in U.S. (Alaska) compared to existing<br>escapement goals. ....   | 52          |
| 10 Summary of 2021 preliminary fall chum salmon escapement counts to Canada in comparison with<br>existing international interim management escapement goals (IMEG) .....               | 52          |
| 11 Pilot Station sonar Chinook salmon passage and Canadian-origin proportion by strata, 2005–2021. ....   | 53          |
| 12 Genetic baseline (microsatellite, 37 populations) used for stock separation of chum salmon sampled in<br>the Pilot Station sonar drift gillnet test fishery, 2021. ....              | 55          |
| 13 Preseason Canadian-origin Yukon River Chinook salmon outlooks for 2000–2022 and the observed<br>run sizes for 2000–2021. ....  | 56          |
| 14 Forecasted 2022 total run size of fall chum salmon based on parent year escapement for each brood<br>year and predicted return per spawner (R/S) rates, Yukon River, 2016–2019. .... | 57          |
| 15 Preseason Yukon River drainagewide fall chum salmon outlooks 1998–2022 and estimated run sizes<br>for 1998–2021. ....  | 57          |
| 16 Preseason Canadian-origin mainstem Yukon River fall chum salmon outlooks for 1998–2022 and<br>observed run sizes for 1998–2021. ....   | 58          |
| 17 Preseason Fishing Branch River fall chum salmon outlooks for 1998–2022 and observed run sizes for<br>1998–2021. ....   | 59          |

## LIST OF FIGURES

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| 1 Map of the Alaska (U.S.) portion of the Yukon River drainage showing communities and fishing<br>districts. ....  | 60          |
| 2 Primary assessment projects operated in the U.S. and Canada used to assess Chinook and fall chum<br>salmon run strength or escapement. ....                  | 61          |
| 3 Daily (top) and cumulative (bottom) catch per unit effort (CPUE) for Chinook salmon in the Lower<br>Yukon set gillnet test fishery at Big Eddy in 2021. .... | 62          |

## LIST OF FIGURES (Continued)

| Figure  | Page |
|---|------|
| 4 U.S. (Alaska) harvest of Chinook salmon, Yukon River, 1961–2021.....  | 63   |
| 5 U.S. (Alaska) harvest of fall chum salmon, Yukon River, 1961–2021.....  | 64   |
| 6 U.S. (Alaska) harvest of coho salmon, Yukon River, 1961–2021.....   | 65   |
| 7 Commercial fishing boundaries, tributaries, and major towns within the Yukon Territory, Canada. ....  | 66   |
| 8 Canadian harvest of Chinook salmon, Yukon River, 1961–2021. ....  | 67   |
| 9 Estimated spawning escapement estimates and escapement goals (minimum or range) for Canadian-<br>origin Yukon River mainstem Chinook salmon, 1982–2021. ....  | 68   |
| 10 Chinook salmon ground-based escapement estimates for selected tributaries in the U.S. (Alaska)<br>portion of the Yukon River drainage, 1986–2021. ....   | 69   |
| 11 Estimated drainagewide escapement of fall chum salmon, Yukon River, 1974–2021.....   | 71   |
| 12 Fall chum salmon escapement estimates for selected spawning areas in the U.S. (Alaska) portion of the<br>Yukon River drainage, 1972–2021.....  | 72   |
| 13 Canadian harvest of fall chum salmon, Yukon River, 1961–2021.....  | 73   |
| 14 Estimated spawning escapement and escapement goals (minimum or range) for Canadian-origin fall<br>chum salmon for the mainstem Yukon River and Fishing Branch River, 1972–2021. ....   | 74   |
| 15 Daily passage estimates of Chinook salmon at the Pilot Station sonar in 2021 (top) and cumulative<br>passage estimate, including 90% confidence intervals (bottom), 2021 compared to historical average. ....  | 75   |
| 16 Daily passage estimates of chum salmon at the Pilot Station sonar in the fall season in 2021 (top),<br>cumulative passage estimates, including 90% confidence intervals (bottom), compared to median<br>passages.....  | 76   |
| 17 Daily passage estimates of coho salmon at the Pilot Station sonar in 2021 (top), cumulative passage<br>estimates, including 90% confidence intervals (bottom), compared to median passages. ....   | 77   |
| 18 Lower Yukon daily water temperatures, comparing 2021 to historical minimum, maximum, and<br>average temperatures.....  | 78   |
| 19 Expected versus observed number of Canadian-origin Chinook salmon returning to spawn, 2000–2021....  | 79   |
| 20 Yukon River Canadian-origin Chinook salmon recruits versus spawners, Ricker curve (thick line), and<br>1:1 replacement line (thin line). Brood years 1982–2015 are included.....   | 80   |
| 21 Juvenile abundance estimates of Canadian-origin Chinook salmon from the Yukon River based on<br>pelagic trawl research surveys in the northern Bering Sea (2003–2021). ....  | 81   |
| 22 The relationship between juvenile abundance estimated from surface trawl surveys and adult returns<br>for Canadian-origin Chinook salmon from the Yukon River. Data labels indicate juvenile year, gray<br>shaded area indicates the 80% confidence interval, and black dashed lines indicate the 80% prediction<br>interval. Data include 2003–2017, excluding 2005, 2008, and 2014. .... | 82   |
| 23 Historic run size estimates of Canadian-origin Chinook salmon in the Yukon River (gray bars) 1982–<br>2021 and preliminary projected run size for 2022 based on juvenile abundance (gray hashed bar). ....   | 83   |

## LIST OF APPENDICES

| Appendix  | Page |
|---|------|
| A1 Passage estimates from the Pilot Station sonar, 1995 and 1997–2021.....  | 86   |
| A2 Alaska commercial salmon sales (number of fish) by district and subdistrict, 2021.....   | 88   |
| A3 Yukon River Canadian-origin Chinook salmon total run by brood year and escapement by year.....   | 89   |
| A4 Chinook salmon age and sex percentages from selected Yukon River monitoring projects operated in<br>U.S. (Alaska), 2021. ....  | 91   |
| A5 Yukon River Chinook salmon age, female percentage, and mean length from Eagle sonar project,<br>2005–2021.....   | 92   |
| A6 Yukon River Chinook salmon harvest percentage by stock group for U.S. harvest, U.S. and Canada<br>harvest combined, and the percentage of the upper stock group harvest by each country, 1981–2021. .... | 93   |
| A7 Stock group percentage by major stock and by country, from chum salmon beginning July 19 at the<br>Pilot Station sonar, Yukon River, 1999–2021.....  | 95   |
| A8 Drainagewide Yukon River fall chum salmon estimated brood year production and return per spawner<br>estimates 1974–2021.....   | 96   |



## LIST OF APPENDICES (Continued)

| Appendix   | Page |
|--|------|
| A9 Escapement, rebuilding and interim goals for Canadian-origin Chinook and fall chum salmon stocks, 1985–2022.....  | 98   |
| A10 Fall chum salmon age and sex percentages with average lengths from selected Yukon River monitoring projects, 2021. ....  | 100  |
| B1 Alaskan and Canadian total utilization of Yukon River Chinook, chum, and coho salmon, 1961–2021....   | 102  |
| B2 Alaska harvest of Yukon River Chinook salmon, 1961–2021. ....   | 105  |
| B3 Alaska harvest of Yukon River summer chum salmon, 1970–2021. ....   | 108  |
| B4 Alaska harvest of Yukon River fall chum salmon, 1961–2021.....  | 111  |
| B5 Alaska harvest of Yukon River coho salmon, 1961–2021. ....  | 114  |
| B6 Alaska (U.S.) and Canada total utilization of Yukon River Chinook and fall chum salmon, 1961–2021...  | 117  |
| B7 Canadian harvest of Yukon River Chinook salmon, 1961–2021. ....   | 119  |
| B8 Canadian harvest of Yukon River fall chum salmon, 1961–2021.....  | 122  |
| B9 Chinook salmon aerial survey indices for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1961–2021.....   | 125  |
| B10 Chinook salmon escapement counts and percentage females counted for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1986–2021. ....                    | 127  |
| B11 Estimated run size and spawning escapement of Canadian-origin Yukon River mainstem Chinook salmon, 1982–2021. ....   | 129  |
| B12 Chinook salmon escapement counts for selected spawning areas in the Canadian (Yukon) portion of the Yukon River drainage, 1961–2021.....   | 131  |
| B13 Summer chum salmon escapement counts for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1973–2021.....  | 134  |
| B14 Fall chum salmon abundance estimates or escapement estimates for selected spawning areas in the U.S. (Alaska) portions of the Yukon River drainage, 1971–2021.....                       | 139  |
| B15 Fall chum salmon escapement estimates for selected spawning areas in Canadian (Yukon) portions of the Yukon River drainage, 1971–2021.....   | 142  |
| B16 Estimated spawning escapement of Canadian-origin mainstem Yukon River fall chum salmon, 1980–2021.....   | 144  |
| B17 Coho salmon passage estimates or escapement estimates for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1972–2021. ....                              | 146  |
| B18 Yukon River Salmon Agreement specified obligations for harvest shares, border passage and spawning escapement for mainstem Canadian-origin Yukon River Chinook salmon, 2001–2021.....    | 149  |
| B19 Summary of management and conservation measures implemented in the U.S. (Alaska) and Canada for Chinook salmon, 2001–2021. ....  | 151  |
| B20 Yukon River Salmon Agreement specified obligations for harvest shares, border passage and spawning escapement for mainstem Canadian-origin fall chum salmon, 2001–2021.....              | 161  |
| B21 Summary of management and conservation measures implemented in the U.S. (Alaska) fall season fisheries and Canada Yukon mainstem for fall chum salmon fisheries, 2001–2021.....          | 163  |
| B22 Yukon River Salmon Agreement specified obligations for spawning escapement for Fishing Branch River fall chum salmon, 2001–2021. ....  | 166  |
| B23 Summary of management and conservation measures implemented for fall chum salmon in the U.S. (Alaska) and Canada on the Porcupine River, 2001–2021.....                                  | 168  |
| C1 Bering Sea-Aleutian Islands Bycatch Summary and impact on Yukon river Canadian-origin salmon. Prepared by NOAA, in coordination with ADF&G, at the request of the Yukon River Panel. .... | 170  |



## 1.0 ABSTRACT

The Yukon River Joint Technical Committee (JTC) of the United States and Canada meets twice a year to analyze and discuss harvest and escapement goals, management trends, postseason reviews, preseason outlooks, and results of cooperative research projects for Canadian-origin Yukon River salmon. This report summarizes the status of Chinook *Oncorhynchus tshawytscha*, coho *O. kisutch*, and summer and fall chum salmon *O. keta* stocks in 2021, presents a 2022 season outlook, and provides data about salmon harvests in commercial, subsistence, First Nations, personal use, domestic, and sport or public angling fisheries. Summaries of Yukon River research projects are also included. For 2021, the preliminary estimate of Chinook salmon (mainstem) spawning escapement in Canada was 31,452 fish, which was below the interim management escapement goal (IMEG) range of 42,500–55,000 fish. A preliminary estimate of the total Canadian-origin Chinook salmon run was 32,972 fish. The preliminary estimate of fall chum salmon spawning escapement in the Canadian mainstem Yukon River was approximately 23,170 fish, which was below the IMEG range of 70,000–104,000 fish. The preliminary estimate of fall chum salmon spawning escapement in the Fishing Branch River (Porcupine River), obtained from a weir count was 2,413 fish, which was below the IMEG range of 22,000–49,000 fish. Recommended interim management escapement goals for Canadian-origin mainstem fall chum salmon and Fishing Branch (Porcupine River) fall chum salmon in 2022 remain the same as for 2021. The JTC recommended a new escapement goal for Canadian-origin Yukon River Chinook salmon of 52,500 with an acceptable deviation range of 42,500–62,500. The Yukon River Panel will be presented with the new Chinook salmon escapement goal recommendation for consideration at their 2022 pre-season meeting.

**Key words:** Chinook salmon *Oncorhynchus tshawytscha*, chum salmon *O. keta*, coho salmon *O. kisutch*, Yukon River, Yukon River Salmon Agreement, Joint Technical Committee, escapement, escapement goal, interim management escapement goal IMEG, management strategy, season outlook.

## 2.0 INTRODUCTION

The purpose of this annual *Yukon River Season Summary and Season Outlook* report is to present data for the Canadian-origin Yukon River salmon stocks subject to the *Yukon River Salmon Agreement* (YRSA). After 16 years of negotiations, Canada and the United States reached a consensus on the elements of a draft YRSA, which was finalized and signed in December 2002. The YRSA continues to represent an international commitment to the restoration, conservation, and management of Canadian-origin Yukon River salmon. The YRSA also established the Yukon River Panel (YRP) as the main instrument to implement the Treaty and the Joint Technical Committee (JTC) as the body responsible for acquiring the best science and management expertise possible to support the YRP.

The JTC was established as an international advisory committee to evaluate management plans and escapement goals for the transboundary stocks of salmon within the Yukon River drainage. The JTC is comprised of representatives from both State, Territorial, and Federal agencies, and local and regional organizations in the U.S. and Canada. The JTC meets twice a year and is charged with various tasks related primarily to Yukon River salmon stock assessment and management, including reporting on preseason outlooks and postseason reviews, examining management regimes and recommending how they may be improved to achieve management and escapement goals, and evaluating the status of Canadian-origin salmon stocks and making recommendations for adjustments to rebuilding programs. This report fulfills several of the JTC's functions outlined in the YRSA and serves as a repository for important data related to Canadian-origin Yukon River salmon stocks. This repository is used by fisheries managers, Tribal and Yukon First Nation governments, fishers, and other stakeholders as the primary record for Yukon River salmon.

This report focuses on Chinook *Oncorhynchus tshawytscha*, fall chum *O. keta*, and coho salmon *O. kisutch* stocks that originate in Canadian waters and are covered by the *Yukon River Salmon Agreement*. Summer chum salmon occur entirely within the U.S. portion of the Yukon River drainage and have overlapping run timing with Chinook salmon and fall chum salmon. Where they

overlap, the management of summer chum salmon is affected by the management of Chinook salmon and vice versa. As such, this report contains information about summer chum salmon to provide context for fisheries assessment and management decisions that affect Canadian-origin Chinook and fall chum salmon. Few coho salmon are bound for the upper reaches of the Yukon River in Canada, therefore discussion of coho salmon is primarily limited to the Porcupine River population. This annual report covers salmon fishery and management topics addressed by the JTC following the 2021 season and preceding the 2022 season.

## **YUKON RIVER SALMON AGREEMENT MANAGEMENT PERFORMANCE SUMMARY**

The following is a summary of information contained in the main body of the report, tables, figures, and appendices. This information is provided at the request of the YRP to summarize specific outcomes of the 2021 season, size of the 2022 salmon runs, and 2022 escapement goal recommendations related to the YRSA.

### **2021 Total Run Size, Harvest, and Escapement of Canadian-origin Chinook Salmon**

The preliminary estimate of the 2021 Canadian-origin Chinook salmon run in the mainstem Yukon River was 32,972 fish and was below the 2021 preseason outlook range of 42,000–77,000 fish. There was no total allowable catch for Canadian-origin Chinook in 2021. The harvest of Canadian-origin Chinook salmon in the U.S. was estimated to be 1,214 fish, which was above the U.S. harvest share of 0 fish. The estimated U.S./Canada border passage of Chinook salmon was 31,758 fish. The mainstem harvest of Chinook salmon in Canada was estimated to be 306 fish, which was above the Canada harvest share of 0 fish. The spawning escapement of mainstem Canadian-origin Yukon River Chinook salmon was estimated to be 31,452 fish, which was below the lower end of the interim management escapement goal (IMEG) range of 42,500–55,000 fish.

### **2021 Total Run Size, Harvest, and Escapement of Canadian-origin Fall Chum Salmon**

The preliminary estimate of the 2021 Canadian-origin fall chum salmon run in the mainstem Yukon River was approximately 23,000 fish and was substantially lower than the preseason outlook range of 136,000–191,000 fish. The preliminary harvest estimate of mainstem Canadian-origin fall chum salmon in the U.S. was approximately 176 fish. The U.S. harvest is not known with certainty and was approximated as 25% of the total U.S. harvest of fall chum salmon ( $703 \times 0.25 = 176$  fish). The estimated U.S./Canada border passage of mainstem fall chum salmon was 23,170 fish. The harvest of mainstem fall chum salmon in Canada was 0. The spawning escapement of mainstem Canadian-origin fall chum salmon was estimated to be 23,170 fish and was well below the IMEG range of 70,000–104,000 fish.

The total run size estimate for 2021 Fishing Branch fall chum salmon was 2,500 fish and is highly uncertain. Total harvest of Fishing Branch fall chum salmon in the U.S. was approximately 28 fish and assumed that 4% of the total U.S. harvest of fall chum salmon were bound for the Fishing Branch River. The total harvest of Porcupine River fall chum salmon in Canada was reported as 21, of which 63% (13 fish) were estimated to be bound for the Fishing Branch River. Escapement past the Fishing Branch River weir was 2,413 fall chum salmon and was well below the IMEG range of 22,000–49,000 fish.

## 2022 Outlooks

The preseason outlook range presented by the JTC for Canadian-origin salmon stocks:

- Chinook salmon: 41,000–62,000
- Mainstem fall chum salmon: 20,000–37,000
- Fishing Branch fall chum salmon: 3,000–6,000

## 2022 Escapement Goals

Pertaining to stocks subject to the *Yukon River Salmon Agreement*, the JTC recommends a revision to the Canadian-origin Chinook salmon mainstem stock goal and no changes to the either of the Yukon River fall chum salmon stock goals. Recommendations for the 2022 season are:

- Chinook salmon: 52,500 with an acceptable range of 42,500–62,500<sup>1</sup>
- Mainstem fall chum salmon: IMEG of 70,000–104,000
- Fishing Branch fall chum salmon: IMEG of 22,000–49,000

# 3.0 ALASKA MANAGEMENT OVERVIEW

## 3.1 CHINOOK AND SUMMER CHUM SALMON

The Yukon River drainage in Alaska (Yukon Area) is divided into fishery districts and subdistricts for management purposes (Figure 1). Management of the Yukon Area summer season commercial salmon fisheries is in accordance with 5 AAC 39.222 *Policy for the Management of Sustainable Salmon Fisheries*, 5 AAC 05.360 *Yukon River Drainage King Salmon Management Plan*, and 5 AAC 05.362 *Yukon River Summer Chum Salmon Management Plan*. The summer chum salmon management plan establishes run size thresholds needed to allow subsistence, commercial, sport, and personal use fishing, prioritizing subsistence among uses, and prioritizing escapement over consumptive uses. Because summer chum and Chinook salmon migrate concurrently, regulations in the management plans allow for using selective gear types that target summer chum salmon during times of Chinook salmon conservation and allow immediate, live release of Chinook salmon back to the water.

During the “summer season” (early May–July 15 in District 1), management and research staff focus on assessing and managing the summer chum and Chinook salmon runs. After July 15, in Emmonak, Chinook salmon are nearly done entering the river and the summer chum salmon run transitions to the fall chum salmon run. On July 16, management transitions to the “fall season” and assessment and management become focused on fall chum and coho salmon runs.

Throughout most of the fishing season, the Yukon River Drainage Fisheries Association (YRDFA) facilitated weekly teleconferences to provide managers, fishermen, tribal/traditional council representatives, and other stakeholders the opportunity to share information, provide input, and discuss inseason management options. During these weekly teleconferences, Alaska Department of Fish and Game (ADF&G) and U.S. Fish and Wildlife Service (USFWS) staff provided inseason run assessment information from various assessment projects (Figure 2). Managers also relayed information about management strategies and subsistence fishermen reported on river conditions in their respective communities along the river.

---

<sup>1</sup> The decision to recommend that the management entities in both countries implement this revised escapement goal for the 2022 season is at the discretion of the YRP, as is consideration of whether the goal recommendation would be considered an interim goal.

## **Preseason Management Strategy Planning**

The 2021 JTC preseason forecast for Canadian-origin Chinook salmon was for a run of approximately 42,000–77,000 fish, and the ADF&G preseason forecast for the Yukon River drainagewide run (U.S. and Canada stocks combined) was 102,000–189,000 fish. For Canadian origin Chinook salmon, the IMEG range recommended by the YRP was 42,500–55,000 fish.

The summer chum salmon outlook was projected to be approximately 1.2 million fish (80% CI  $\pm$  500,000), which was a run size sufficient to meet escapement and subsistence needs and provide a harvestable surplus for commercial fisheries.

Additional considerations in 2021 included travel limitations related to COVID-19. Under State of Alaska health mandates, commercial and subsistence fishing activities were considered essential, however activities were impacted by reduced airline and freight services, local travel guidelines, and concerns for crew and community health and safety. However, key projects such as the Pilot Station and Eagle sonars operated successfully and provided estimates of salmon passage for the entirety of the 2021 season (Tables 7 and 8). The Lower Yukon test fishery (LYTF) was operated with modified operations to reduce Chinook salmon mortality and provided indices of relative abundance throughout the 2021 season. The Gisasa River weir did not operate in 2021 due to COVID-19 related travel restrictions and funding concerns.

Prior to the annual preseason planning meeting, YRDFA hosted additional teleconferences with participants from each district in mid-April. Discussion topics included how to prepare for low salmon runs, importance and ability to harvest nonsalmon species, gear types, and how to adapt to changing conditions.

YRDFA hosted the preseason planning meeting with board members attending in person, and the public invited via teleconference. This meeting was held in late April and funded by the Yukon River Panel. The purpose of this meeting was to present the preseason outlook and management strategies and answer questions from participants. Fishermen from throughout the drainage discussed management options and raised additional concerns about environmental factors, bycatch, fish diseases, food security, and project operations. Updates on COVID-19 contingency plans and changes to project operations and were also given. An annual informational flyer detailing the outlooks for Chinook, chum, and coho salmon and fishery management strategies was mailed preseason to approximately 2,700 Yukon River households and distributed as an advisory announcement on May 13.

## **Chinook and Summer Chum Salmon Inseason Management**

Based on the forecasts, managers expected to restrict salmon fishing early in the run while there was uncertainty about the size of the run. Due to the late timing of summer chum salmon in 2020 and 2019, fishing for summer chum salmon with selective gear also remained closed at the start of the season.

During the 2018 Board of Fisheries meeting, the regulation requiring full fishing closures during the first pulse of Chinook salmon in Districts 1 and 2 was removed when projected run sizes are adequate to meet escapements. Due to the low forecast in 2021, and low returns in 2019 and 2020, managers took additional precautionary measures and closed salmon fishing shortly after the first Chinook salmon was counted at Pilot Station sonar. Closures went into place on June 2 in District 1 and the Coastal Area and were implemented in upriver districts based on run timing (Table 1;

Appendix 21). This closure also protected the early-arriving fish, as well as the entire first pulse of fish.

Fishing for nonsalmon remained open with 4-inch or smaller mesh gillnets restricted to 60 feet or less in length and other nonsalmon gear types (fyke net, longline, hook and line, etc.). Fishermen could also use selective gear types such as beach seines and dip nets however all Chinook and chum salmon were required to be released alive. Pink and sockeye salmon retention was allowed, however the abundance of sockeye salmon is low in the Yukon River, and pink salmon abundance is low in odd years (Appendix 21).

Around the typical midpoint of the run (June 23) at the Pilot Station sonar, the Chinook salmon run projection indicated that the drainagewide run size was too weak to meet escapement goals and provide any harvestable surplus. Once the second stratum of genetic samples were analyzed, the Canadian-origin run estimate was approximately 64,000 Chinook salmon (with an 80% confidence interval of 58,000 to 70,000 fish). While there was a higher-than-average proportion of Canadian-origin fish, the overall run was too weak to support harvest, and a conservative approach was warranted based on the lower-than-expected passage of Canadian-origin Chinook salmon at the border the last two years.

Cumulative summer chum salmon counts at the Pilot Station sonar were the lowest ever observed in the history of the project (from 1995–2021). At the midpoint of the summer chum salmon run at the Pilot Station sonar (July 6), season cumulative counts were 79,138 fish. Season total counts of summer chum salmon were approximately 153,497 fish (with a 90% confidence interval of 137,200 to 169,800 fish), which was well below the historical median from years with late run timing of 1.6 million fish. Season total counts of summer chum salmon at the Pilot Station sonar were the lowest in all the years of project operations (1995–2021) and were well below the previous lowest counts of 442,546 and 448,665 in 2001 and 2000 respectively. Forecasting models, even with some adjustments to account for poor age class returns, did not accurately predict this weak run. No escapement goals were met and project counts were below historical medians (Tables 7 and 8). Summer season aerial surveys were not conducted due to poor weather and increased rain and water levels impacting survey visibility during the normal survey dates. To protect summer chum and Chinook salmon, all salmon fishing remained closed, and selective gear types were only open for nonsalmon species

As run size estimates were refined inseason, the management team subtracted the IMEG (42,500–55,000) from the inseason estimate of Canadian-origin Chinook salmon and multiplied that result by the midpoint of the U.S. harvest share (77%) to estimate a harvest range of Canadian-origin fish available for Alaskan fishermen. Near the midpoint of the Chinook salmon run at Pilot Station sonar, ADF&G estimated the U.S. harvest share of Canadian-origin Chinook salmon to be approximately 3,100–13,000 fish. However, due to the difference between inseason projections and the abundance estimated at the border in 2019 and 2020, managers continued with a precautionary strategy and maintained salmon fishing closures.

Despite very conservative management, inseason passage counts at the Eagle sonar project indicated that like 2019 and 2020, fewer Canadian-origin Chinook salmon were going to make it to the border than were projected by the Pilot Station sonar genetic estimates. Historically, the midpoint of late Chinook salmon runs at Eagle sonar is around July 28. In 2021, Chinook salmon

passage was only 15,900 fish on this date, which was well below average. More detail on management and conservation measures implemented<sup>2</sup> are summarized in Appendix B19.

It is not certain why the 2019–2021 inseason projections of Canadian-origin Chinook salmon based on Pilot Station sonar passage and application of genetics did not align well with the estimates at the Eagle sonar. In recent years (2014–2018), inseason projection methods have provided enough information to enable managers to restrict harvest sufficiently to achieve or exceed both the lower end of the border escapement IMEG and provide for the Canadian harvest share. The U.S. harvest alone does not account for the difference between inseason projections and the abundance estimated at the border in 2019–2021. Based on preliminary harvest estimates and genetic analysis, an estimated 1,214 Canadian-origin Chinook salmon were harvested in the U.S. in 2021 (Appendix B18). A large portion of this harvest was salmon from test fishery projects that were distributed to communities.

In 2021, water temperatures were close to average and water levels were low during most of the summer season. However, increased levels of *Ichthyophonus* infections were observed, and infections were noted in small Chinook salmon harvested incidentally in 4-inch or smaller mesh gear (Stan Zuray, fisherman, Tanana; personal communication). Preliminary results of samples taken from a first-year study at the Pilot Station sonar indicate that the prevalence of *Ichthyophonus* infections in 2021 were much higher than other periods in the past when baseline sampling was in place and above the threshold level identified by the JTC of 25% to establish a monitoring program. *Ichthyophonus* infections may have contributed to an increased level of en route mortality of Chinook salmon headed for Canada. Though the number of fish that die during migration before making it to the border cannot currently be measured, more research is being planned in this area.

### 3.2 FALL CHUM AND COHO SALMON

Management of the Yukon Area fall season salmon fisheries is in accordance with 5 AAC 39.222 *Policy for the Management of Sustainable Salmon Fisheries*, 5 AAC 01.249 *Yukon River Drainage Fall Chum Salmon Management Plan*, 5 AAC 05.369 *Yukon River Coho Salmon Management Plan*, and 5 AAC 05.367 *Tanana River Salmon Management Plan*. The intent of these plans is to align management objectives with the established escapement goals, provide flexibility in managing subsistence harvests when stocks are low, and bolster salmon escapement as run abundance increases (Table 2). The sustainable escapement goal (SEG) range for the entire Yukon River drainage is 300,000–600,000 fall chum salmon (Fleischman and Borba 2009). The threshold number of fall chum salmon needed to allow for a fall chum salmon directed commercial fishery is 550,000 fish, and commercial fishing is considered only on the surplus projected above that level.

Management also incorporates conditions found in the *Yukon River Salmon Agreement*. Those conditions include treaty objectives for border passages that are based on the IMEG into Canada, and harvest shares of fall chum salmon. The IMEG for Canadian-origin mainstem Yukon River is 70,000–104,000 fall chum salmon, and the IMEG for Fishing Branch River is 22,000–49,000 fall chum salmon.

---

<sup>2</sup> To look up advisory announcements for Yukon River fisheries in the U.S. go to the following website:  
<http://www.adfg.alaska.gov/index.cfm?adfg=cfnews.search>



The *Yukon River Coho Salmon Management Plan* allows for a coho salmon-directed commercial fishery if the fall chum salmon run is assessed to be more than 500,000 fish, incidental catch of fall chum salmon remains above the 500,000 fish threshold, and a harvestable surplus of coho salmon is identified, or a commercial fishery will not have a significant impact on fall chum salmon escapement and allocation. The Tanana River plan specifies that commercial fishing in Subdistrict 5-A and District 6 are based on the assessment and timing of salmon stocks bound for the Tanana River drainage as both areas are considered terminal harvest areas.

## **Fall Chum Salmon Management Overview**

By regulation, the fall season began in District 1 on July 16. Assessment information collected from projects located in the lower river were used to inform management decisions. The projects included two lower river drift gillnet test fisheries that provided run timing and relative abundance information, and a mainstem Yukon River sonar, located near the community of Pilot Station, that provided fish abundance estimates. Stock composition information for chum salmon was provided by genetic samples collected at the mainstem Yukon River sonar.

Upriver projects that monitored escapement consisted of a mainstem Yukon River sonar operated at Eagle near the U.S./Canada border; a weir/video project operated in the Fishing Branch River, a Porcupine River headwater; sonars in the Teedriinjik River and in the Canadian portion of the Porcupine River near Old Crow; foot surveys conducted in the Delta River, a tributary of the Tanana River; boat surveys in the Delta Clearwater River, a tributary of the Tanana River; and aerial surveys in the Tanana River drainage, Kluane River and mainstem Yukon River between Tatchun Creek and the Pelly River. Age, sex, and length information was collected at the lower river test fisheries, the Eagle sonar near the U.S./Canada border, and from the Fishing Branch and Delta rivers.

The preseason forecast was revised to a preseason run size projection in mid-July, using the relationship between historical summer and fall chum salmon run size estimates. Based on the critically low 2021 summer chum salmon, the preseason drainage projection for fall chum salmon was a run size of less than 300,000 fish.

Preseason management strategies included the following:

- Concurrent with the fall chum salmon migration upriver, all Yukon Area districts and subdistricts would remain closed to subsistence fishing unless the run projection exceeded 300,000 fish.
- To improve fall chum salmon escapement to the spawning grounds, the department anticipated implementing a complete closure of subsistence salmon fishing in the Alaska portion of the mainstem Porcupine River when the fall chum salmon migration reached that area.
- Commercial salmon fishing would not be allowed unless the inseason drainagewide fall chum salmon run projection exceeded 550,000 fish, and a commercial surplus was identified.

According to the *Yukon River Drainage Fall Chum Salmon Management Plan*, the preseason projection did not meet the threshold of 300,000 fish needed to allow subsistence, personal use, and commercial salmon fishing. Based on inseason assessment projects at the midpoint of the run, the projection indicated a run size of approximately 100,000 fall chum salmon. All Yukon Area districts remained closed to fall chum salmon fishing for the duration of the season. Gillnets of 4-

inch or smaller mesh were allowed to target non-salmon. However, due to the conservation concern for Chinook and chum salmon, 4-inch or smaller mesh gillnets were restricted to a maximum length of 60-feet. Subsistence fishing opportunity was provided with selective gears (dip nets and hook and line) for pink, sockeye, and coho salmon that are present in the Lower Yukon Area through Subdistrict 4-A Lower. While using selective gear, all Chinook and chum salmon were required to be released alive (Appendix 23).

As the season progressed, it became apparent that the body size of fall chum and coho salmon was the smallest observed in the historical datasets and the percentage of female fall chum salmon was tracking about 10% below average in Lower Yukon assessment projects for most of the season. Due to the higher probability of encountering smaller bodied salmon and females, 4-inch or smaller mesh gillnets that are used to target non-salmon species were placed on a reduced schedule to allow more salmon to reach their spawning grounds. To provide more fishing opportunities during this time, subsistence fishing opened with fish wheels (manned) for non-salmon, while fall chum salmon were required to be released alive immediately, and coho salmon were strongly recommended to be released as well (Appendix 23).

Starting October 1, subsistence salmon fishing restrictions were lifted in the Coastal District and District 1. Restrictions were subsequently lifted in upriver districts and subdistricts as the tail end of the fall chum salmon run reached those areas. To protect spawning salmon, important spawning areas for fall chum and coho salmon in Yukon River drainage tributaries remained closed to subsistence salmon fishing through the end of December (Appendix 23).

### **Coho Salmon Management Overview**

The coho salmon run overlaps with much of the fall chum salmon run. While subsistence fishing for fall chum salmon was closed for most of the season, fishermen in the Lower Yukon Area through Subdistrict 4-A could use selective gear such as dip nets and hook and line for coho salmon. On August 28, selective gear opportunity closed and further restrictions to 4-inch or smaller mesh gillnets were implemented due to concerns for both fall chum and coho salmon.

The coho salmon run appeared to be weak and late, and information from lower river assessment projects showed a record low coho salmon run. The preliminary coho salmon run size was estimated to be 45,000 fish, which was below the historical average of 240,000 fish.

## **4.0 ALASKA HARVEST SUMMARIES**

### **4.1 SUBSISTENCE SALMON FISHERY**

Subsistence salmon fishing activities in the Yukon River drainage typically begin in late May and continue through mid-October (Jallen et al. 2017). Fishing opportunity in the Lower Yukon Area (Districts 1–3) in May and the Upper Yukon Area (Districts 4–6) in October is highly dependent upon river ice conditions. Throughout the drainage, most Chinook salmon harvested for subsistence use are dried, smoked, or frozen for later human consumption. Summer chum, fall chum, and coho salmon harvested in the lower Yukon River are primarily utilized for human consumption. In the Upper Yukon Area, summer chum, fall chum, and coho salmon are also an important human food source, but a larger portion of the harvest is fed to dogs used for recreation and transportation (Andersen 1992).

Documentation of the subsistence salmon harvest is necessary to determine if sufficient salmon are returning to the Yukon Area and enough fishing opportunities are being provided to meet

subsistence needs. In years with fishery restrictions, estimates of harvest can be used to assess the effect of the management actions taken to meet escapement goals to maintain future salmon production. The primary method of estimating subsistence harvest is voluntary participation in the annual subsistence salmon harvest survey program conducted by ADF&G, Division of Commercial Fisheries. The survey is conducted in 33 communities (including the 2 coastal communities of Hooper Bay and Scammon Bay) during the fall, after most households have completed fishing for salmon. Additional information about harvest timing is obtained from harvest calendars that are sent to households and filled out voluntarily. Fishing permits also provide information about harvest timing for areas of the river where permits are required (District 6 and portions of District 5 and the Koyukuk River).

In 2021, subsistence harvest surveys identified approximately 2,570 households in the Yukon Area in 33 communities. Of these, an estimated 222 households fished for salmon. Permits are not required for subsistence fishing throughout most of the Yukon Area, except for the urban areas around Fairbanks and other areas accessible by road. Therefore, the largest share of subsistence harvest in the Yukon Area is estimated from the postseason survey results. A total of 213 salmon fishing permits were issued in 2021, approximately 95% of the subsistence salmon permits had been returned at the time of this publication, and 46 salmon permits reported fishing.

All 2021 subsistence harvest data are considered preliminary as of the publication date of this report. Final results will be included in an ADF&G Fishery Data Series publication after the analysis is completed and reviewed. Based on survey and permit data, the 2021 preliminary subsistence salmon harvest in the Alaska portion of the Yukon River drainage was estimated to be 1,945 Chinook; 1,253 summer chum; 703 fall chum; and 293 coho salmon (Appendices B2–B5). For comparison, recent 2016–2020 average subsistence salmon harvest estimates (including test fish donations) were 32,501 Chinook; 71,633 summer chum; 60,855 fall chum; and 5,912 coho salmon (Appendices B2–B5) from communities in the Alaska portion of the Yukon River drainage. In 2021, Chinook, summer chum, fall chum, and coho salmon harvests all fell below their respective ranges of Amounts Reasonably Necessary for Subsistence (ANS) as defined by the Alaska Board of Fisheries (Brown and Jallen 2012).

For a second year in a row, due to the COVID-19 pandemic, subsistence salmon harvest surveys were conducted remotely via telephone, mail, and internet. An electronic version of the survey was employed to provide subsistence users an avenue to self-report harvests online. To improve survey response rate, all known households were attempted to be contacted. The survey questions largely remained the same as previous years. The 2021 estimates and 95 % CI were  $735 \pm 134$  Chinook;  $730 \pm 206$  summer chum ;  $123 \pm 65$  fall chum; and  $93 \pm 60$  coho salmon. It is important to restate the estimates and 95% CI provided here are preliminary and will change as additional mail surveys are entered and quality control measures are conducted. Survey estimates are a subtotal of the overall subsistence estimates provided above and 95% CI only apply to survey estimates.

## **4.2 COMMERCIAL FISHERY**

### **Summer Season Harvest**

There was no commercial fishing in the Yukon Area during the 2021 summer season. Historical commercial harvest information of Chinook and summer chum salmon can be found in Figure 4 and Appendices B2 and B3.

### **Fall Season Harvest**

There was no commercial fishing in the Yukon Area during the 2021 fall season. Historical commercial harvest information of fall chum and coho salmon can be found in Figures 5 and 6, and Appendices B4 and B5.

## **4.3 SPORT FISHERY**

Since 2011, sport fishing effort for wild salmon in the Yukon River drainage has been directed primarily at Chinook, chum, and coho salmon, with lesser numbers of sockeye and pink salmon targeted in the lower Yukon River. Over the past decade, Chinook salmon stocks have experienced periods of low productivity with subsequent restrictions to subsistence fishing opportunities. As a result, Chinook salmon sport fishing restrictions and closures have been implemented each year during this period in the ADF&G Division of Sport Fish Yukon Management Area (YMA, excludes the Tanana River drainage) and similarly in the Tanana River Management Area (TRMA), except for 2017. All chum salmon harvested in the sport fishery are categorized as summer chum salmon because these fish are mostly caught incidental to Chinook salmon during midsummer in clearwater tributaries. Some harvest of fall chum salmon occurs after Chinook salmon spawning concludes but is considered negligible relative to summer chum salmon harvests. Coho salmon are targeted primarily in the fall.

Alaska sport fishing effort and harvests are monitored annually through the Statewide Harvest Survey (SWHS)<sup>3</sup>. The SWHS is an annual survey of households where at least one person (resident or nonresident) purchased a sport fishing license. Harvest estimates are not available until approximately one calendar year after the fishing season; therefore, 2021 estimates were not available for this report. Total sport harvest of salmon during 2020 in the Alaska portion of the Yukon River drainage (YMA and TRMA) was estimated to be 49 Chinook, 1,684 chum, and 1,337 coho salmon (Appendices B2, B3, and B5). The 2016–2020 average sport salmon harvest was estimated to be 65 Chinook, 474 chum, and 583 coho salmon and that for 2011–2020 was estimated to be 132 Chinook, 493 chum, and 622 coho salmon (Appendices B2, B3, and B5).

Most sport fishing effort for the Yukon River occurs in the Tanana River along the road system (Baker 2018) due to the proximity of major population centers such as Fairbanks, North Pole, and Delta Junction. On average, 62% and 94% of Chinook salmon harvested during 2011–2020 and 2016–2020 respectively occurred in the Tanana River. During 2016–2020, average sport harvests for chum and coho salmon in the Tanana River represented 2% and 52% of the total for these species respectively for the Yukon River. In the Tanana River, most Chinook and chum salmon

---

<sup>3</sup> Alaska Sport Fishing Survey database [Internet]. 1996–2020. Anchorage, AK: Alaska Department of Fish and Game, Division of Sport Fish (cited December 9, 2021). Available from: <http://www.adfg.alaska.gov/sf/sportfishingsurvey/>.

sport fishing effort occurs in the Chena and Salcha rivers, whereas most coho salmon are harvested from the Delta Clearwater and Nenana Rivers. The majority of sport fishing effort for Chinook, chum, and coho salmon for the rest of the Yukon River drainage takes place in the Anvik and Andreafsky rivers.

During 2006–2016, all freshwater sport fishing guides and guide businesses operating in Alaska were required to be licensed and to report harvest and released (numbers of fish captured and released) in logbooks. From 2012–2016, guided sport harvests in the Yukon River drainage (YMA and TRMA) averaged 34 Chinook and 356 coho salmon.

For 2021, all waters of the YMA and TRMA were closed to sport fishing of Chinook salmon effective May 10, 2021 and June 24, 2021, respectively. Sport fishing for chum and coho salmon were also closed in both the YMA and TRMA effective July 1, 2021 and August 26, 2021, respectively. These closures were a result of below average counts past Pilot Station sonar and subsequent restrictions to the subsistence fishery.

#### **4.4 PERSONAL USE FISHERY**

The Fairbanks Nonsubsistence Area, located in the middle portion of the Tanana River, contains the only personal use fishery within the Yukon River drainage. Subsistence or personal use permits have been required in this portion of the drainage since 1973. Personal use fishing regulations were in effect from 1988 until July 1990 and from 1992 until April 1994. In 1995, the Joint Board of Fisheries and Game established the Fairbanks Nonsubsistence Area which has subsequently been managed consistently under personal use regulations. Historical harvest data must account for these changes in status. Subdistrict 6-C is completely within the Fairbanks Nonsubsistence Area and therefore falls under personal use fishing regulations. Personal use salmon or whitefish/sucker permits, and a valid resident sport fishing license are required to fish within the Fairbanks Nonsubsistence Area. The harvest limit for a personal use salmon household permit is 10 Chinook, 75 summer chum, and 75 fall chum and coho salmon combined. The personal use salmon fishery in Subdistrict 6-C has a subdistrict harvest limit of 750 Chinook; 5,000 summer chum; and 5,200 fall chum and coho salmon combined.

In 2021, a total of 45 personal use salmon permits were issued. The 2021 preliminary harvest, based on 100% of the personal use salmon permits returned in Subdistrict 6-C is 0 salmon. The 2016–2020 average personal use harvest was 148 Chinook, 297 summer chum, 274 fall chum, and 149 coho salmon (Appendices B2–B5) in the Alaska portion of the Yukon River drainage.

### **5.0 CANADIAN MANAGEMENT OVERVIEW**

#### **5.1 CHINOOK SALMON**

The 2021 pre-season outlook range for Canadian-origin mainstem Yukon River Chinook salmon was 42,000 to 77,000. This range was well below historically-observed run sizes (average 153,411, 1982–1997) and also below the average run size (82,894) observed from 1998–2020. When accounting for uncertainty and past forecast performance it was recognized that the run size was unlikely to meet the upper end of this range.

New to 2021 in Canadian Yukon River fishery management was applying the concept of run size probabilities to pre-season fishery management planning and communications with First Nation

governments and stakeholders. This approach takes into account the inherent uncertainty of the outlook, addresses the reality that some run sizes are more probable than others and provides separate probabilities for different run sizes (e.g. there is a 75% chance that the run size will be at least 48,500 and a 50% chance that the run size will be at least 57,000). This approach is useful in fishery and harvest management planning.

Prior to the season, Fisheries and Oceans Canada (DFO) hosted and/or participated in virtual meetings with the Yukon Salmon Subcommittee (YSSC), Yukon First Nation governments, Renewable Resources Councils, and the public to discuss the 2021 forecast and potential management scenarios.

Each year, in advance of the salmon season, DFO develops an Integrated Fisheries Management Plan<sup>4</sup> (IFMP) for Yukon River Chinook, fall chum and coho salmon. The IFMP, which is in effect from July 1 of the current year to June 30 of the subsequent year, serves to identify the primary objectives (i.e. YRSA) and requirements for the management of Canadian salmon fisheries in the Yukon River, as well as the management measures to be used to achieve these objectives in the commercial, domestic (non-aboriginal food fishery) and licensed public angling fisheries.

In accordance with Yukon First Nation self-governing agreements, First Nation fisheries are managed by First Nation governments. In support of this, DFO includes First Nation advisors in Yukon River Panel processes, and provides scientific information and management updates to the First Nations on a weekly basis (more frequently if/when requested) throughout the season.

Canadian management decisions in 2021 were guided by the YRSA, YSSC recommendations, implementation of the precautionary approach, obligations as set out in the Final Land Claim Agreements with Yukon First Nations and the application of inseason assessment information to the *inseason fishery management decision matrix* (a component of the IFMP) and the following management recommendations from the YRP for the 2021 season:

1. Consistent with the 2021 Chinook salmon forecast and in consideration of the probability of expected low run size, 2021 directed Chinook salmon fisheries should be closed through at least the mid-point of the run. Chinook salmon harvest opportunity after the mid-point should be considered conservatively to ensure escapement falls within the 2021 IMEG range (42,500–55,000) goals and that Yukon River Salmon Agreement harvest share obligations are achieved. In-season fishery management decisions will account for the reality that the Canadian-origin run size past the Pilot Station sonar program will not be known with certainty, as such, should rely on the lower bound of in-stream run size projections.
2. To provide for Canadian-origin Chinook salmon conservation, limit use of gillnets to 6” mesh or smaller upstream of the Tanana River / Yukon River Mainstem confluence for the duration of the Chinook salmon migration consistent with the regulatory structures in both countries.
3. Contingent on status of border passage and achievement of Treaty objectives, restrictions on Chinook salmon fishing should be maintained into the Fall Season through the approximate 95% point or greater of the Chinook salmon run timing.

---

<sup>4</sup> The IFMP is available online at <https://waves-vagues.dfo-mpo.gc.ca/Library/40801445.pdf>

4. Environmental conditions, in particular extreme events, should be considered in-season to inform fishery management measures implemented and resulting harvest opportunities.
5. In the event that inseason assessment programs are unable to operate in 2020 due to circumstances beyond Agency control, fishery harvest opportunities should be provided conservatively based on 2021 pre-season outlooks and associated Total Allowable Catch and harvest share allocations.
6. U.S. and Canadian Managers shall collaborate closely throughout the season to evaluate and discuss in-season run assessment, management actions and expected outcomes resulting from management actions.

Based on the preseason forecast and a 75% probability of the run size being at least 48,500, DFO commenced the 2021 season with limited harvest opportunities for the First Nation fisheries. By the time that Chinook entered the Canadian portion of the Yukon River, First Nations governments were advised that inseason information indicated that the run size would be below or near the lower end of the pre-season forecast of 42,000. The public angling fishery was prohibited from catching or retaining Chinook salmon and similarly, the commercial and domestic fisheries remained closed (no allocation).

Allocations to the commercial, domestic and public angling fisheries are subject to run abundance, and opportunities (i.e. allocation) may only be provided if there is sufficient confidence that the abundance of Chinook salmon will meet the upper end of the IMEG (55,000), and Canada's harvest share exceeds the number required for a full allocation to the First Nation fishery.

In consideration of the YRP management recommendations, conditions of licence in the commercial and domestic salmon fisheries would restrict harvesters to a maximum allowable gillnet mesh size of six (6) inches and mandated the release of incidentally caught Chinook salmon in the chum salmon commercial and domestic fisheries.

As confidence in inseason abundance improved, fishery management actions proceeded according to the *inseason fishery management decision matrix*. The decision matrix provides guidance for the management of fisheries, is linked to specific inseason run abundance levels, summarizes the management reference points, general allocation plans, and anticipated management responses under different run size scenarios (Table 3).

### **Inseason Management Yukon River Mainstem Chinook Salmon**

DFO commenced the season with an allocation available for Yukon First Nations (managed by First Nation governments) while public angling, commercial and domestic fisheries were closed.

Early in the 2021 season, information from the ADF&G's Lower Yukon Test Fishery (LYTF) near Emmonak and the Pilot Station sonar in the Lower Yukon Area suggested a low run and late run timing. By mid-July, the run at Pilot Station sonar was nearly complete with a cumulative passage estimate of around 65,000 Canadian-origin Chinook salmon (with an estimated Canadian harvest share of 2,300 to 5,000 fish). Although this estimate was within the pre-season forecast, Canadian managers considered uncertainty, past inseason forecast performance, low run sizes observed at other assessment sites in Alaska, and Yukon River fisher reports of low Chinook abundance among other factors in planning.

The first Chinook salmon were counted at Eagle sonar (located near the international border) on June 28. Cumulative passage estimates at Eagle during the early part of the run were low and slow to increase. As the season progressed the mismatch between the information from Pilot Station and the observed run size at Eagle sonar became increasingly apparent and inseason estimates at Eagle sonar indicated that the IMEG was not likely to be achieved.

First Nation Governments were advised to adjust harvest strategies accordingly in the First Nation fishery and DFO maintained the closures in the public angling, commercial and domestic fisheries. DFO enacted a complete salmon angling closure on the Yukon River and its tributaries for the duration of the season. The YSSC recommended to the First Nation governments to plan to not harvest Chinook salmon. First Nation governments were responsive to inseason information and advised their citizens to not harvest.

There were no available allocations for the commercial, domestic, and public angling fisheries. First Nation governments maintained a conservative approach throughout the Canadian season and were responsive to inseason information. Throughout the season DFO provided weekly email updates to First Nations and harvesters and hosted bi-weekly inseason meetings with the YSSC and First Nation Lands and Resources managers as a means to provide a forum to exchange management and assessment updates. DFO staff also participated in weekly inter-agency meetings with ADF&G summer season staff and provided updates during the weekly YRDFA teleconferences.

The public angling fishery daily catch and possession limits were reduced to 0 for 2021–2022 season. The public angling fishery along the Yukon River was closed to salmon fishing from July 1 to September 29. Chinook salmon commercial and domestic fisheries in Canada remained closed throughout the 2021 season. A summary of management and conservation measures implemented in Canada are presented in Appendix B19.

### **Inseason Management Porcupine River Chinook Salmon**

In the absence of stock-specific information about Porcupine River Chinook salmon in Canada, the early season management of this stock is based on information and management of mainstem Yukon River Chinook salmon. Given the below-average outlook for mainstem Chinook salmon in 2021, it was advised that the First Nation Fishery refrain from harvest. Public angling on the Porcupine River was closed from July 1 to September 8.

By late July, the inseason assessment of run strength at the Porcupine River sonar indicated that the 2021 Chinook salmon run was approximately 10% of project average (2014–2019). The Vuntut Gwitchin Government, which directs the First Nation fishery in accordance with Yukon First Nation Self-Governing Agreements and is guided by the *Porcupine River Salmon Plan*, directed First Nation harvesters not to fish for Chinook.

## **5.2 FALL CHUM SALMON**

### **Mainstem Yukon River**

The 2021 preseason forecast for the Canadian-origin fall chum salmon run to the mainstem Yukon River was 136,000 to 191,000 fish. The preseason forecast was preliminary and was revised in mid-July, following the summer chum run. The IMEG range recommended by the YRP remained at 70,000–104,000 for Canadian-origin fall chum salmon.



Throughout the season DFO provided weekly email updates to First Nations and harvesters and hosted bi-weekly inseason meetings with the YSSC and First Nation Lands and Resources managers as a means to provide a forum to exchange management and assessment updates. DFO staff also participated in regular inter-agency meetings with ADF&G fall season staff and provided updates during the weekly YRDFA teleconferences

Canadian management decisions were based on the application of inseason assessment information to the management decision matrix – a component of the IFMP. The decision matrix provides detailed guidance for the management of fisheries linked to specific inseason run abundance levels. The 2021 decision matrix summarized the management reference points, general allocation plans, and anticipated management responses under different run size scenarios (Table 4). A summary of management and conservation measures implemented in Canada is presented in Appendix B21.

### **Inseason Management Mainstem Yukon Fall Chum Salmon**

Inseason decisions about fishery openings and closures in Canada for fall chum salmon were made in a similar way as those for Chinook salmon. 2021 saw the lowest summer chum run on record which resulted in a revised drainagewide fall chum salmon projection that would be well below the pre-season forecast and unlikely to meet spawning escapement goals.

Inseason projections of the Canadian component of the fall chum salmon run were based on cumulative passage estimates and genetic apportionment of Canadian-origin fall chum salmon from the Pilot Station sonar and assessment information from the LYTF. As early as July 22, the revised projection for fall chum was well below the pre-season forecast and unlikely to meet spawning escapement. As fall chum salmon approached and entered Canada in early September, Canadian managers began considering passage estimates from Eagle sonar.

In consideration and implementation of the YRP's management recommendations, in the event that allocations were available to the commercial and domestic fisheries the conditions of license would include the following.

- All incidentally caught Chinook salmon in the chum salmon commercial and domestic fisheries must be released, and;
- The maximum allowable gillnet mesh size is 6 inches in both the commercial and domestic chum salmon fisheries.

The intention of management actions in 2021 was to ensure that the IMEG range of 70,000–104,000 fall chum salmon was achieved. However, the revised projection and observed low run size at Pilot indicated that the low fall chum run would not be sufficiently abundant to provide for spawning escapement and resulting in no available Canadian-origin chum salmon allocation.

By early August, information from the lower river in Alaska indicated that the total run would be the lowest on record, which was later supported by Eagle sonar passage estimates that indicated that the run into Canada would not meet the IMEG. Given the poor run, First Nation governments were advised that there would not be a Canadian allocation and to adjust their management plans accordingly. First Nation governments advised their citizens to forgo chum harvest.

### **Fishing Branch (Porcupine) River Fall Chum Salmon**

The 2021 preseason forecast estimate for Fishing Branch-origin fall chum salmon was 22,000–30,000 fish. The preseason forecast was preliminary and was revised in mid-July, following the

summer chum run. The IMEG for the Fishing Branch River recommended by the YRP was 22,000–49,000 adult fall chum salmon.

Considering that the IMEG has only been achieved in 6 of the last 10 years, a precautionary approach was warranted. The IFMP recommended that, until an inseason projection for Fishing Branch chum exceeded 26,000 fish, a conservative approach to harvest be taken in the Porcupine River First Nation fishery. Important to note is that in accordance with Yukon First Nation Self-Governing Agreements, the Vuntut Gwitchin Government directs the First Nation fishery. A summary of management and conservation measures implemented on the Porcupine River in Canada is presented in Appendix B23.

### **Inseason Management Fishing Branch (Porcupine) Fall Chum Salmon**

Canadian fishery management considered early season information from the LYTF and Pilot Station sonar. Estimates of fall chum salmon passage in combination with genetic mixed stock analysis (MSA) cannot be reliably used to project the run to Fishing Branch River. Because the Fishing Branch River component at the Pilot Station sonar is such a small part of the total run, the uncertainty associated with these estimates is very high; therefore, management decisions cannot be based on this information.

Inseason fishery management decisions are largely based on information from the Porcupine River sonar located near the community of Old Crow. The Porcupine River sonar passage projection is the primary indicator used to inform inseason management decisions, however harvest in Alaska before the fish reach Canada is also considered when making management decisions.

As the season progressed the fall chum salmon run projections were reduced to levels that would not support meeting the Fishing Branch River spawning escapement goal, at which time the Vuntut Gwitchin First Nation asked their citizens to refrain from fall chum salmon harvest.

In 2021, escapement to the Fishing Branch River was monitored by a combined weir and video counter/sonar. Only a portion of the fall chum salmon that return to the Canadian Porcupine River are destined for the Fishing Branch River. Based on 2021 Porcupine River sonar counts and Fishing Branch River weir counts approximately 63% of Canadian-origin Porcupine River fall chum salmon were considered Fishing Branch River origin.

## **6.0 CANADIAN HARVEST SUMMARIES**

### **6.1 FIRST NATION SUBSISTENCE FISHERIES**

Harvest estimates of salmon in the First Nation fisheries on the Yukon and Porcupine rivers are determined from locally-conducted inseason interviews and postseason reports. For additional ease in reporting, DFO provides harvest calendars and harvest reporting forms to First Nation governments' Lands and Resources staff for distribution among harvesters.

#### **Mainstem Yukon River Chinook Salmon**

Based on a preseason outlook for a below-average run of 42,000–77,000 Canadian-origin Yukon Chinook salmon, and the probability that the run size would not be at the upper end of the range, several recommendations and conservation measures were proposed for early fishing opportunities in the First Nation fisheries. These included initiating harvest activities in a conservative manner and to direct harvest at smaller (younger) fish by using selective gear and release of larger (older) fish. Following a slow start to the season, inseason information from the LYTF and Pilot Station

sonar projects indicated that the run was within the preseason forecast range, which would provide for a limited First Nation fishery. Ultimately, inseason Eagle sonar passage data did not align with Pilot Station sonar projections. As the run progressed, the Eagle sonar passage indicated that the IMEG was deemed unlikely to be met in 2021. Yukon First Nation governments were responsive to inseason information and followed conservative management plans throughout the 2021 season, resulting in a significantly reduced harvest compared to long term historical averages. The First Nation harvest in the Canadian Yukon River mainstem drainage in 2021 was estimated to be 306 fish (Appendix B7). For comparison, the First Nations long-term (1961–2020) average harvest is 4,890 fish; the most recent 10-year average (2011–2020) is 2,374; and the most recent 5-year average (2016–2020) is 2,837 fish (Appendix B7).

### **Mainstem Yukon River Fall Chum Salmon**

The preseason outlook for Canadian-origin fall chum salmon in 2021 suggested a below average run of 136,000 to 191,000 fish. The preseason forecast was preliminary and was revised in mid – July, following the summer chum run. By July 21, the inseason projection was revised, and the projected run size was not expected to meet the minimum spawning escapement of 70,000 Canadian-origin fall chum salmon. Inseason Eagle sonar counts suggested that border passage would be insufficient to meet border passage obligations under the YRSA. First Nations abstained from harvest in the First Nation fishery on the Yukon River mainstem. For the second year in a row there was 0 fall chum salmon harvest reported in the First Nation fishery on the mainstem Yukon River drainage in 2021 (Appendix B8). For comparison, the long-term (1961–2020) average First Nation subsistence harvest is 2,173 fish; the most recent 10-year average (2011–2020) is 775 and 5-year average (2016–2020) is 800 fish (Appendix B8).

### **Porcupine River Chinook, Fall Chum, and Coho Salmon**

An estimated harvest of 16 Chinook salmon occurred in the in 2021 First Nation subsistence fishery near Old Crow (Appendix B7). For comparison, the long-term (1961–2020) average harvest is 249 fish; the most recent 10-year average (2011–2020) is 208 fish; and, the most recent 5-year average (2016–2020) is 227 fish (Appendix B7).

An estimated harvest of 21 fall chum salmon occurred in the 2021 First Nation subsistence fishery near Old Crow (Appendix B8). For comparison, the long-term (1961–2020) average harvest is 4,104 fish; the most recent 10-year average (2011–2020) is 1,808 fish; and the most recent 5-year average (2016–2020) is 1,658 fish (Appendix B8).

There was no reported harvest of coho salmon on the Porcupine River in 2021.

## **6.2 COMMERCIAL FISHERY**

The commercial Chinook, fall chum, and coho salmon fisheries remained closed throughout the 2021 fishing season (Appendices B7 and B8). The long-term (1961–2006, 2009) average commercial harvest of Chinook salmon is 5,717 fish, and there have been 0–4 Chinook salmon harvested annually in the past 12 years (2010–2021; Appendix B7).

The long-term (1961–2020) average commercial harvest of fall chum salmon is 9,193 fish, and the most recent 5-year average (2016–2020) is 1,567 fish (Appendix B8). Since 1997, there has been a marked decrease in commercial catches of Upper Yukon River fall chum salmon as a result of a limited market. Between 1961 and 2019, the commercial fall chum salmon catch ranged from a low of 293 fish in 2009, when the run was late and the fishery had been closed for most of season

due to conservation concerns, to a high of 40,591 fish in 1987. Note that commercial harvest of coho salmon in the mainstem Yukon River in Canada rarely occurs. This is thought to be due to a combination of low abundance and their late migration timing which limits availability of this species.

### **6.3 DOMESTIC SUBSISTENCE FISHERY**

The domestic fishery was closed during the Chinook and fall chum salmon season (Appendices B7 and B8); there were no salmon harvested in the domestic fishery in 2021. Openings in the domestic salmon fisheries are concurrent with commercial fishery openings. For comparison, with respect to harvest of Chinook salmon in the domestic fishery the long-term (1961–2020) average is 393 fish. Domestic harvest of Chinook salmon has been 0 since 2010 (Appendix B7). With respect to domestic harvest of fall chum salmon, the long-term (1961–2020) average is 405 fish; the most recent ten-year average (2011–2020) is 10 fish; and the most recent five-year average (2016–2020) is 6 fish (Appendix B8).

### **6.4 LICENSED PUBLIC ANGLING FISHERY**

In 1999, the YSSC introduced a mandatory Yukon Salmon Conservation Catch Card to improve harvest estimates and to serve as a statistical base to ascertain the importance of salmon to the Yukon River public angling fishery. Anglers are required to report their catch and harvest by November 30. The information reported includes the number, species, fate (kept or released), sex, size, date, and location of all salmon caught. From preliminary catch card information received at the time of this publication, no Chinook salmon were caught or retained in the Yukon River or its tributaries in the 2021 public angling fishery, which is consistent with the angling restrictions and closures which were in place for the duration of the 2021 Chinook and chum salmon season.

Over the last 10 years retention (harvest) of Chinook salmon in the public angling fishery was only permitted in 2009 and 2011 (Appendix B7). For the 2021 season, the daily catch and possession limits of fall chum salmon in the public angling fishery were varied to 0 prior to the start of the season which was followed by a complete angling closure to salmon on the Yukon River and its tributaries (Appendix B19).

## **7.0 TOTAL RUN, ESCAPEMENT, AND HARVEST SHARE ASSESSMENTS FOR 2021**

### **7.1 CHINOOK SALMON**

In 2021, the total Chinook salmon passage at the Pilot Station sonar was approximately 124,845 fish  $\pm$ 10,831 (90% CI, Table 5, Appendix A1). This is considered an index of the drainagewide Chinook salmon run, rather than a total run size estimate, because some salmon are harvested or enter spawning areas below this sonar site. This passage was below the historical average<sup>5</sup> of 181,967 fish (Appendix A1). Most of the Chinook salmon entered the river in four pulses consisting of approximately 22,550 fish; 53,670 fish; 20,760 fish; and 3,790 fish. However, similar to 2020, the front end of the run had an unusually long and consistent flow of ‘tricklers’ that lasted for almost two weeks before the more distinctive first pulse arrived. The first quarter point,

---

<sup>5</sup> Average includes years 1995, 1997, 2000, 2002–2008, and 2010–2019. The sonar did not operate in 1996 and project difficulties occurred in 1998–1999, 2001, and 2009.

midpoint, and third quarter point for Chinook salmon at the Pilot Station sonar project were on June 19, June 29, and July 4, respectively. The 2021 Chinook salmon run appears to have been six days later than average based on the midpoint at the sonar project.

Chinook salmon passage estimated at Eagle sonar in 2021 was 31,796 fish (Appendix B11). The estimated mainstem border passage into Canada was 31,758 fish, which is calculated by subtracting the harvest upriver from the Eagle sonar site (Appendices B11, B18). The estimated spawning escapement of Canadian-origin Yukon River Chinook salmon (mainstem) was 31,452 fish, which is calculated by subtracting Canadian harvest (Figures 8 and 9; Appendices B11 and B18). This escapement was below the lower end of the IMEG of 42,500–55,000 fish. Combining the spawning escapement estimate with the U.S. and Canadian harvests of Canadian-origin Chinook salmon indicates the total mainstem Canadian-origin run size was approximately 32,972 Chinook salmon (Appendix B18).

Based on a total run size estimate of 32,972 Chinook salmon, and prescriptions outlined in the YRSA, the TAC for 2021 was 0 fish (Appendix B18). The U.S. harvest of 1,214 exceeded the harvest share of 0. The number of Chinook salmon that passed into Canada (31,758) was 10,742 fewer fish than what was needed to meet the lower end of the IMEG range (42,500 fish). The Canadian mainstem harvest of 306 Chinook salmon exceeded the Canadian harvest share of 0.

Age, sex, and length (ASL) composition of Chinook salmon were assessed at both mainstem sonar sites and in various escapement projects (Table 6; Appendices A4–A5). The ASL samples collected at the Pilot Station sonar are thought to be representative of all Chinook salmon stocks passing the site and include both U.S. and Canadian stocks. The ASL samples collected at the Eagle sonar are exclusively from Canadian-origin fish. Gillnet mesh sizes used to sample the runs differ at each location. The Chinook salmon age composition from 646 samples that were aged from the drift gillnet test fishery at the Pilot Station sonar project (all mesh sizes combined) was 2% age-4, 48% age-5, 45% age-6, 5% age-7, and less than 1% age-8 fish (Appendix A4). Females comprised 50% of all fish sampled (including un-ageable samples; Table 6). The age composition for age-4 and age -5 fish were below the recent 10-year average. However, all other age classes were above the recent 10-year average with percent female also above average. It is important to note that while the Pilot Station sonar test fishery uses a wide range of gillnet mesh sizes, and likely captures a representative sample across sizes and age classes, the sex is determined visually, and this method has reduced accuracy compared to internal inspection (Table 6; Appendix A4).

The Chinook salmon age composition from 327 samples that were aged from the test fishery at the Eagle sonar project was 2% age-4, 45% age-5, 49% age-6, and 4% age-7 fish (Appendix A4). The age composition for age-4 was below the recent 10-year average. However, all other age classes were similar or above the recent 10-year average with percent female also similar to the recent 10-year average. (Table 6). Slight modifications have been made to the drift gillnet mesh sizes used at the Eagle sonar during the first three years of operation (2005–2007); however, mesh sizes measuring 5.25, 6.5, 7.5, and 8.5-inch have been used consistently since 2007. Small fish may be underrepresented in the samples, due to not fishing gillnets smaller than 5.25-inch mesh.

Chinook salmon escapement in U.S. tributaries was assessed at two weirs and two counting towers (Table 7; Figure 10). In 2021, all U.S. tributary Chinook salmon escapement goals were not met and escapements for systems without goals were below average (Liller and Savereide 2018; Table 7; Appendix B10). River conditions were favorable on the Chena and Salcha River systems this year, with below average water levels during the summer season passage dates (late-June to mid-

August). Most assessment projects were able to get successful counts for nearly all days of operation. However, aerial surveys of the East and West Forks of the Andreafsky River, Anvik River, Nulato River, Gisasa River, Henshaw Creek, etc. were not conducted due to record levels of rain and sustained poor weather in western and interior Alaska during early August survey dates. Due to logistical challenges resulting from the COVID-19 pandemic, the Gisasa River weir did not operate.

Passage of Chinook salmon to tributaries in Canada was assessed at the Whitehorse Rapids Fishway and sonars operated on the Porcupine, Klondike, Pelly, Big Salmon and Takhini rivers and at a weir on Tatchun Creek (Appendix B12). The 2021 estimate for Chinook salmon passage on the Porcupine River was 409 fish, much lower than the 2014–2019 average of 3,896. On the Klondike River, 855 Chinook salmon were counted, which was lower than in the previous operating years, 2009–2011 and 2020 (average of 1,900). On the Pelly River, Chinook salmon passage was estimated at 4,980 fish, which was lower than the 2017–2020 average of 7,859 fish<sup>6</sup>. On the Big Salmon River, 1,958 Chinook salmon were counted, which was below the 2011–2020 average count of 4,620 fish. At Tatchun Creek, 2021 passage was estimated at 17 Chinook salmon, well below the 1970–2000 average of 243 fish. At the Whitehorse Rapids Fishway, 274 Chinook salmon were counted, which was below the ten-year average count of 1,074 fish, and among the lowest on record. Hatchery-produced fish accounted for 36% of the fish that returned to the Whitehorse Fishway in 2021, compared to 2011–2020 average of 47%.

## **7.2 SUMMER CHUM SALMON ALASKA (U.S. ONLY)**

In 2021, an estimated 153,718 summer chum salmon  $\pm 16,149$  (90% CI) passed the Pilot Station sonar (Table 5, Appendix A1), which was well below the 1995–2020 (excluding 1996, 1998, 1999, 2001 and 2009) median of 1.9 million fish for the project and were well below the previous lowest counts of 442,546 and 448,665 in 2001 and 2000 respectively. The first quarter point, midpoint, and third quarter point were June 30, July 6, and July 11, respectively, which was likely 9 days later than average and the latest on record based on the midpoint at the sonar project. Three pulses of summer chum salmon were detected at the sonar project with the largest group consisting of approximately 83,045 fish and passed between July 6 and July 18. A summer chum salmon drainagewide biological escapement goal (BEG) with a range of 500,000–1,200,000 was adopted in 2016 (Liller and Savereide 2018; Table 8), and the 2021 estimated escapement of 153,000 fish fell below the lower end of the goal and was the lowest on record.

In addition to the drainagewide biological escapement goal, escapement goals exist for summer chum salmon on the East Fork Andreafsky River and the Anvik River (Table 8). None of these goals were met in 2021 and counts at the other projects (Henshaw Creek weir, Chena and Salcha River sonars) were well below the historic medians.

The Gisasa River weir did not operate in 2021 due to COVID-19 travel restrictions and staffing concerns, and aerial surveys were not conducted because of poor weather. Carcass sampling on the Salcha River was canceled due to low abundance of fish and high water during the peak spawning and carcass sampling periods.

---

<sup>6</sup> Average excludes sonar estimate from 2016 feasibility study.

### 7.3 FALL CHUM SALMON

The initial method of determining total drainagewide (i.e., U.S.-origin and Canadian-origin) fall chum salmon run size inseason was based on the Pilot Station sonar passage estimate and the estimated inriver harvest of fall chum salmon downstream of the sonar site. The inseason run size model primarily uses the commercial fishery, which is the largest harvest component below the sonar site, to produce overall projections of abundance used to manage the fishery. In 2021 due to the low run of fall chum salmon, the commercial fishery was not prosecuted and the subsistence harvest was minimal due to fishery closures; therefore, no harvest was added to the passage estimate. Genetic mixed stock analysis (MSA) was used inseason to account for the strictly fall chum salmon component of the run, which transitions from summer to fall runs in mid-July. The inseason total run size using these methods was estimated to be near 100,000 chum salmon.

Postseason, a Bayesian state-space model was used to estimate drainagewide escapement (Fleischman and Borba 2009). The total drainagewide run size is then derived by adding the estimated total harvest (U.S. and Canada) to the estimate of drainagewide escapement. In 2021 this method resulted in a total drainagewide run size estimate of 95,000 fall chum salmon, which was well below the 2021 forecast of 542,000–762,000 fish. The total run size ended up even lower than the inseason projection of 184,000 fall chum salmon, based on the relationship between summer and fall chum salmon total run sizes.

The drainagewide escapement estimate produced by the Bayesian state-space model was 94,000 fall chum salmon, which was well below the escapement goal range of 300,000–600,000 fall chum salmon (Liller and Savereide 2018; Table 8; Figure 14). The model utilized historical escapement data from the Toklat, Delta, Teedriinjik (Chandalar), Sheenjek, Fishing Branch, and Canadian mainstem Yukon rivers, as well as mark–recapture estimates of abundance from the upper Tanana, and Kantishna projects (Appendices B14–B16). All of the individual fall chum salmon escapements were well below their respective goal ranges (Liller and Savereide 2018; Table 8; Figure 15; Appendices B14, B16).

In 2021, the proportions by age class for fall chum salmon caught in both the LYTF and MVTF were weighted by project, then combined and used to represent the drainagewide run and included <1% age-3, 92% age-4, 7% age-5, and <1% age-6 fish. The age-3, age-5, and age-6 components were all below average, while the age-4 was well above average (ranked second highest on record) when compared to LYTF weighted averages for all years 1977–2020. The unweighted proportions of fall chum salmon samples from LYTF included 2% age-3, 87% age-4, 10% age-5, and 1% age-6 (Appendix A10). The fall chum salmon samples collected from the test fishery operated at Mountain Village included 1% age-3, 94% age-4 and 5% age-5. Fall chum salmon ASL composition estimates from collections in the Delta River included 8% age-3, 76% age-4, and 16% age-5. Samples were also collected from fall chum salmon for the escapement into Canada based on test fishing near the Eagle sonar site and included 91% age-4 and 9% age-5. Fall chum salmon sampled at the weir on the Fishing Branch River included 2% age-3, 84% age-4 and 14% age-5. All projects reported proportions of age-4 well above average. The proportion of females was lower than males in all escapement projects (Appendix A10). Fall chum salmon were the longest in the Fishing Branch River at 571 mm, measured from mid eye to tail fork, here referred to as fork length (MEFL), and the shortest in the Delta River at 558 mm MEFL.

## **Mainstem Yukon River Canadian-origin Fall Chum Salmon**

The U.S./Canada border passage estimate for fall chum salmon was the lowest on record, at 23,170 fish. There was no reported fall chum salmon harvest in the U.S. or Canada upstream of Eagle sonar in 2021; the border passage and spawning escapement estimates for Canadian-origin Yukon River mainstem fall chum salmon are also 23,170 (Figure 13; Appendices B8 and B16). For comparison, the 10-year average (2011–2020) escapement is 163,317 (Appendix B16). The 2021 spawning escapement of Canadian-origin Yukon River mainstem fall chum salmon was well below the IMEG of 70,000–104,000 fish (Figure 14, Table 10).

The preliminary reconstruction of the total 2021 Canadian-origin Yukon River mainstem fall chum salmon run was just over 23,000 fish (Appendix B20). Total run size was approximated using the expanded estimate of fall chum salmon that passed the Eagle sonar near the U.S./Canada border (23,170 fish) plus 25% of the U.S. harvest of fall chum salmon that occurred downstream of Eagle sonar ( $703 \times 0.25 = 176$  fish) and then rounded to the nearest 1,000. This run size estimate was well below both the preseason outlook range of 136,000–191,000 Canadian-origin Yukon River mainstem fall chum salmon. The final run size, however, approached the upper confidence limit of the estimate based on Pilot Station Sonar and genetic stock identification (17,300; 90% CI 10,000–24,000).

## **Porcupine River (Including the Fishing Branch River) Canadian-origin Fall Chum Salmon**

In 2021 DFO and Vuntut Gwitchin Government operated the Porcupine River sonar immediately downstream of the community of Old Crow. An estimated 3,486 fall chum salmon passed by the sonar (Appendix B15). An estimated 21 fish were harvested in the Old Crow fishery (Appendix B8; details are presented in Section 8.3).

DFO and Vuntut Gwitchin Government also operated the Fishing Branch River weir in 2021. Counts of fall chum salmon passing a constrained opening in the weir were made using a video counter from 7–16 September, when the video system was replaced with a sonar that operated for the remainder of the season. The 2021 spawning escapement estimate for fall chum salmon above the Fishing Branch River weir was 2,413 fish, the lowest recorded since the program began in 1972 (Figure 14, Table 10 and Appendix B15). The Canadian harvest of Fishing Branch River fall chum salmon in 2021 was estimated at 13 fish (of 21 total chum salmon harvested). This assumes that 63% of the fall chum salmon in the Porcupine River drainage are destined for Fishing Branch River, based on the slope of the regression between Fishing Branch River weir counts and Porcupine sonar estimates (2015–2017, 2019). The total run size estimate for 2021 Fishing Branch fall chum salmon was 2,500 fish (Appendix B22). This was calculated as the sum of the weir passage (2,413 fish), the estimated Canadian harvest (13 fish), and the estimated U.S. harvest of Fishing Branch fall chum salmon (4% of the total U.S. fall chum salmon harvest,  $703 \times 0.04 = 28$  fish) and then rounded to the nearest 500.

# **8.0 PROJECT SUMMARIES**

## **8.1 ALASKA, U.S.**

Salmon assessment programs operated throughout the U.S. portion of the Yukon River drainage are collaborative. This report summarizes salmon run, harvest, and escapement monitoring results from numerous projects. Data were provided by various entities including the Mountain Village



Test Fishery (G. Sandone Consulting, LLC) and the chum salmon genetic stock identification (USFWS). Other project results were provided by ADF&G Division of Commercial Fisheries and Division of Sport Fisheries. Due to COVID-19 travel restrictions and staffing concerns the Gisasa River weir (USFWS) did not operate and aerial surveys were not conducted because of poor weather. Partner organizations that assisted with data collection Yukon Delta Fisheries Development Association, Yukon River Drainage Fisheries Association, and DFO. A more in-depth overview of select stock assessment programs are described in the following sections of this report.

### **Lower Yukon Test Fishery**

The LYTF program is designed to assess salmon run timing and relative abundance and typically consists of two Chinook salmon test fisheries; an 8.5-inch mesh set gillnet test fishery operated in the South and Middle mouths of the Yukon River, and an 8.25-inch mesh drift gillnet operated at Big Eddy in the South Mouth, near Emmonak. The LYTF also has a summer chum salmon-directed drift gillnet test fishery using 5.5-inch mesh gear operated in the South and Middle mouths. These test fisheries provide catch per unit effort (CPUE), which gives an index of abundance and indicates the presence of large groups of fish, or “pulses”, entering the mouths of the river.

The LYTF operated at normal effort at the South Mouth (Big Eddy) drift and set gillnet sites starting on May 22 and June 3, respectively. The Middle Mouth Chinook set gillnet site did not operate for the 2021 summer season because of restrictions due to COVID-19 and logistical complications of operating a field camp. However, an additional 8.25-inch mesh drift gillnet was fished in the Middle Mouth allowing the crew to effectively commute to and from Emmonak while still providing test fishing indices of the run from that mouth of the river. Furthermore, the use of a drift gillnet reduced the incidental mortality of Chinook salmon in a low abundance year and streamlined fish donations for a logistically challenging location. The 5.5-inch drift net operations for summer chum salmon also returned to the Middle Mouth following a one year suspension.

The LYTF 8.5-inch set gillnet concluded operations on July 12 in the South Mouth. The cumulative Chinook salmon CPUE for the Big Eddy set gillnet was 29.82. The first quarter point, midpoint, and third quarter point of the set net was on June 14, June 20, and June 25, respectively. The 8.25-inch drift gillnet projects for Chinook salmon operated in the South Mouth and Middle Mouths until July 15 and provided valuable supplemental run timing information for Chinook salmon entering the Yukon River. The combined cumulative Chinook salmon CPUE for the South and Middle Mouth drift gillnet sites was 64.89. The combined first quarter point, midpoint, and third quarter points of the drift gillnets were on June 5, June 18, and June 30, respectively. The 5.5-inch drift gillnets for summer chum salmon at both the South and Middle Mouth sites also concluded operations on July 15. The combined cumulative chum salmon CPUE for the South and Middle Mouth drift gillnet sites was 191.90, which was below the historical median CPUE of 7,265.63. The first quarter point, midpoint, and third quarter point were June 21, June 28, and July 4, respectively.

The LYTF project continues in the fall season after switching to 6-inch drift gillnets on July 16 and completed operations on September 10. The cumulative CPUE for fall chum salmon of 125.72 was well below the historical median of 1,588.86 and the cumulative CPUE for coho salmon of 14.09 was also well below the historical median of 414.56.

Chinook, chum, and coho salmon caught in the LYTF were released alive if healthy enough to do so, otherwise they were kept, sampled, and distributed to local community. Fish kept and distributed are included in the subsistence harvest estimates. The fish donation program was coordinated with village tribal councils and with the assistance of Yukon Delta Fisheries Development Association.

### **Pilot Station Sonar**

The goal of the Pilot Station sonar project is to estimate daily upstream passage of Chinook (Figure 15), summer and fall chum (Figure 16), and coho salmon (Figure 17). The project has been in operation since 1986 but data is only reported back to 1995. Due to changes in methodology, data from 1995 to present are the most consistent (Appendix A1). Both split-beam and Adaptive Resolution Imaging Sonar (ARIS)<sup>7</sup> are used to estimate total fish passage, and CPUE from the drift gillnet test fishing portion of the project is used to estimate species composition. The project's sonar equipment and apportionment methodologies have evolved over time (Pfisterer et al. 2017; Dreese and Lozori 2019).

Fish passage estimates at the Pilot Station sonar project are based on a sampling design in which sonar equipment is operated daily in three 3-hour periods and drift gillnets 25 fathoms long with mesh sizes ranging from 7.0 cm to 21.6 cm (2.75- to 8.5-inch), approximately 4.3 fathoms in depth, that are fished twice each day between sonar periods to apportion the sonar counts to species. During the 2021 season, both banks were fully operational on May 31 and continued operations through September 7. The ice went out on the mainstem Yukon River near Pilot Station on May 13, based on National Weather Service (NWS) data<sup>8</sup>. Test fishing began on May 31; the first Chinook salmon was caught May 31, the first chum salmon on June 7, and the first coho salmon was caught on August 2.

An estimated 1,040,660 fish passed through the sonar sampling area between May 31 and September 7 (Table 5). Drift gillnetting resulted in a catch of 4,681 fish including 761 Chinook; 453 summer chum; 739 fall chum; and 411 coho salmon. A total of 2,317 fish of other species were also caught. Chinook salmon were sampled for ASL; while only sex (external) and length were collected from chum, pink *O. gorbuscha*, sockeye *O. nerka*, and coho salmon without aging structures; for all other non-salmon species, only length was collected. Genetic samples were taken from Chinook and chum salmon. Any captured fish that were not successfully released alive were distributed daily to residents in Pilot Station. Overall in 2021, there were no significant operational problems. Both sonars performed well throughout the season.

River discharge recorded by the NWS near Pilot Station was below the 2011–2020 mean at the beginning of the season until June 21, then remained above the mean until July 21. During late July the NWS equipment at Pilot Station experienced issues related to sediment burial, and gage data are not available through the remaining field season<sup>9</sup>. The NWS estimated the discharge at Pilot Station using upstream data from the Steven's Village monitoring station. The estimated

---

<sup>7</sup> Product names used in this report are included for scientific completeness, but do not constitute a product endorsement.

<sup>8</sup> <https://www.weather.gov/aprfc/breakupDB?site=488>

<sup>9</sup> [USGS Current Conditions for USGS 15565447 YUKON R AT PILOT STATION AK](#)

values suggest the discharge again rose above the mean approximately Aug 25 and remained through the end of the project.

In 2021, all project goals were met, and passage estimates were provided to fisheries managers daily during the season. Information generated at the Pilot Station sonar was also discussed weekly through multi-agency international teleconferences that included stakeholders from the lower Yukon River to the headwater communities in Canada. Preliminary daily salmon passage estimates were available online<sup>10</sup> and disseminated daily to the general public via a listserv.

### ***Ichthyophonus* investigations**

The prevalence and severity of *Ichthyophonus* disease in Yukon River Chinook salmon was investigated in 2021. Preliminary results of 200 samples taken from the first-year feasibility study at the Pilot Station sonar indicate that the prevalence of *Ichthyophonus* infections in 2021 was 44%. This prevalence was similar to peak levels observed in the past when baseline sampling was in place, and above the threshold level identified by the JTC of 25% to establish a monitoring program. Of the 185 samples that were genetically grouped to country of origin, 47% were Alaskan stocks and 45% were of Canadian stocks. In addition to quantifying the prevalence of infections this study is also estimating the level of intensity or severity of those infections and preliminary results indicate the intensity levels may be unprecedented or higher than previously documented for this system. Samples were also collected from subsistence-caught Chinook salmon at Rapids and of the 68 fish sampled, level of prevalence was 39% and all samples were taken from 4-inch mesh or less gillnets. Full results will be available post season and efforts are continuing to establish a monitoring program for next season.

### **Chinook Salmon Genetic Sampling, 2021**

In 2021, ADF&G and collaborators collected 1,600 genetic tissue samples from adult Chinook salmon caught in Alaskan test fisheries on the Yukon River. Samples included 755 fish from the Pilot Station sonar test fishery (PSTF), 376 fish from the Eagle sonar test fishery (ETF), and 469 fish from the Lower Yukon Test Fishery (LYTF). Additionally, a total of 77 adult Chinook salmon genetic baseline samples were collected from Henshaw Creek.

### **Mixed Stock Analysis of Yukon River Chinook Salmon Sampled at the Pilot Station Sonar, 2005–2021**

The ADF&G Gene Conservation Laboratory (GCL) uses mixed stock analysis (MSA) to estimate inseason stock compositions of Chinook salmon passage at the Pilot Station sonar using genotypes of samples collected from the PSTF. These data provide fishery managers an important “first look” at the Canadian-origin Chinook salmon run strength and timing before those fish migrate through most Alaska fisheries. Without genetic MSA of the PSTF samples, fishery managers would have no information about the Canadian-origin run until fish arrive at Eagle sonar, when most of the run has already passed through 1,900 km of fisheries. Knowledge of relative abundance and migration timing from this project has aided in inseason projections of total run size of Canadian-origin Chinook salmon and more refined management strategies to meet border passage goals.

Genetic MSA is conducted to provide insight on stock-specific run dynamics and has proven to be a critical component of inseason management of salmon fisheries in Alaska. Pilot Station sonar

---

<sup>10</sup> <https://www.adfg.alaska.gov/index.cfm?adfg=commercialbyareayukon.main>

project data has been used to estimate the total proportion of Canadian-origin Chinook salmon each year since 2005. The weighted postseason estimates from this project indicate that on average (2005–2020) the Canadian stock makes up 40% of the total run and has ranged from 31%–50% (Table 11). Over this 16-year timeframe, the contribution of the Canadian-origin stock to the total run has been relatively stable; however, this project has highlighted a considerable amount of within-year variability in the relative abundance of Canadian-origin Chinook salmon (Table 11). In nearly all years (2005–2020), the proportion of Canadian-origin stocks has been highest, often exceeding 50%, during the early portion of the run, but typically decreases as the run progresses. This project, combined with the Pilot Station sonar passage estimates, has shown that while the proportion of Canadian-origin stocks are typically highest in the early portion of the run, the abundance (i.e., numbers of fish) of Canadian-origin fish is generally higher during the middle part of the run (Table 11). Analysis of the 2021 PSTF samples conforms to this typical pattern.

Tissue samples taken from Chinook salmon caught in the 2021 PSTF were analyzed in 3 strata for genetic MSA. The 3 strata periods were May 31–June 22 (number analyzed (n) = 252), June 23–July 06 (n = 379), and July 07–August 6 (n = 113). Genetic MSA indicated the proportion of the total Chinook salmon passage at the Pilot Station sonar that were Canadian-origin was 62% (approximately 28,000 fish) in stratum 1, 54%, (approximately 32,000 fish) in stratum 2, and 35%, (approximately 7,000 fish) in stratum. The total season Canadian percentage was 54% (weighted by passage) which is the highest total season Canadian percentage observed within the 2005–2021 time series (Table 11).

### **Mixed Stock Analysis of Yukon River Chinook Salmon Harvested in Alaska, 2021**

Three broad-scale stock (reporting) groups are used to apportion Chinook salmon harvest by Alaska fisheries within the Yukon River drainage. The Lower and Middle Yukon River stock groups spawn in Alaska and the Upper Yukon River stock group spawns in the Canadian mainstem. Scale pattern analysis, age composition estimates, and geographic distribution were used by ADF&G from 1981–2003 to estimate Chinook salmon stock composition in Yukon River harvests. From 2004 to present, genetic analysis has been the primary method for stock identification (e.g., DuBois 2018). Harvest percentages by stock group for 2014–2021 include the harvest from the Coastal District, whereas the Coastal District was not included in years prior to 2014.

An estimate of the 2021 total U.S. harvest of Chinook salmon by stock of origin required information about the genetic stock composition of the subsistence harvest, test fish giveaways, and incidental commercial harvest in each district. The Canadian-origin harvests from each district were then summed for a total estimated U.S. harvest of Canadian-origin stocks (e.g., DuBois 2018). There was no directed subsistence harvest sampling program in place for 2021. Therefore, genetic MSA results from prior year (2006–2018) subsistence harvest sampling programs, samples taken from the 2021 LYTF, and samples collected from the Pilot Station test fishery (PSTF) in mesh sizes 5.25-inch or less were used to inform the 2021 subsistence harvest composition. A total of 180 samples were collected from the LYTF and were used to determine the stock composition of the test fish giveaway. The subsistence fishery was closed and restricted to 4-inch mesh gillnets or less to target non-salmon. In order to represent the stock composition of fish harvested incidentally from 4-inch mesh or less, the 123 samples collected in mesh sizes 5.25-inch or less from the PSTF were applied to harvests from the Coastal District through District 3. Genetic MSA results from prior year (2006–2018) subsistence harvest sampling programs were used to inform the 2021 subsistence harvest composition for Districts 4 through 5. Chinook salmon harvested in

the Black River, Koyukuk drainage, Teedriinjik (Chandalar River), Birch Creek, and District 6 (Tanana River) are presumed to be U.S.-origin. Similarly, sport fishery harvests typically occur in Alaskan tributaries and assumed to harvest few if any Canadian-origin fish. Stock apportionment information and assumptions were applied to the total U.S. harvest of Chinook salmon (all stocks) of 1,945 (Appendix B2). An estimate of 1,214 Canadian-origin Chinook salmon were harvested in the U.S in 2021 (Appendix B18). Subsistence harvest and stock composition estimates for 2021 are still considered preliminary as of the publication date of this report.

Genetic MSA results for 2021 indicate that the weighted U.S. harvest of Yukon River Chinook salmon was comprised of 15% Lower, 23% Middle, and 62% Upper (Canadian-origin) stock groups. U.S. harvest composition for 2021 was above the 2016–2020 average for the Lower and Upper stock groups and below the 2016–2020 average for the Middle stock group (Appendix A6).

### **Yukon River Chum Salmon Mixed Stock Analysis, 2021**

Chum salmon were sampled from the Pilot Station sonar from June 7 through September 7 and analyzed by the USFWS gene lab to provide stock composition estimates for most of the summer and fall chum salmon runs. Populations in the baseline are reported in aggregated stock groups (Table 12). Results from analysis of these samples were reported for each pulse or time stratum and distributed by email to fishery managers within 24–48 hours of receiving the samples. For summer chum salmon, the lower river stock group comprised 75% of the run and the middle river stock group comprised 25%. The Tanana component of the middle river stock group comprised 5% of the total summer chum salmon run and peaked in passage at the Pilot Station sonar during the sampling period of July 19–August 5. During the fall management season in 2021 the summer component was 31%, which is above the 2006–2020 average (Appendix A7). Due to low abundance of chum salmon, sample sizes were the low for both the summer and fall seasons with only 2 strata (normally 4) in the summer and only 3 strata in the fall (normally 5). The first stratum (July 19–August 5) included the 2 largest pulses and contained a large portion of U.S. border (Teedriinjik-Chandalar, Sheenjek, and Draanjik-Black rivers) stocks. The run transition from summer to fall chum salmon was nearly completed in the second stratum of the fall season (August 6–22) when 93% of the mixture was comprised of fall chum salmon. For fall chum salmon, 82% of the run was of U.S.-origin and 18% of Canadian-origin. The fall chum salmon composition of the U.S. contributions was 42% Tanana and 40% U.S. border. The composition of the Canadian contribution was 11% mainstem Yukon, 6% White, and both Teslin and Porcupine rivers were <1%. Preparations are underway to continue the project for the 2022 season.

### **Environmental Conditions Report**

This U.S. environmental conditions report was added for the first time in 2019. This report differs from the Canadian environmental conditions report, which is much more detailed and was requested by the YRP. Instead, this addition was a first step to document environmental conditions relevant to adult salmon migrating through the U.S. portion of the Yukon River drainage. Currently, environmental monitoring within the U.S. portion of the Yukon River is limited and existing assessment programs are inadequate to quantify environmental impacts to migrating and spawning salmon. Climate change is bringing warming conditions to northern latitudes and in some years water temperatures in the mainstem Yukon River may have exceeded the tolerances of adult salmon. Research has indicated that adult salmon exposed to temperatures of >21–22°C, can experience increased mortality (McCullough et al. 2001).

Water temperature records from LYTF and Pilot Station sonar project sites remain the most reliable and consistent historical inseason data available for the mainstem Yukon River. However, there has been a multi-year effort by ADF&G to expand the spatial distribution of temperature loggers throughout the Yukon River drainage. In 2021, there were 16 temperature loggers deployed representing 8 locations along the Yukon River mainstem. There were 2 located at the LYTF sites at Big Eddy and Middle Mouth, 5 at Pilot Station sonar, 2 near the community of Nulato, 2 near the community of Galena, 1 near the Dalton highway bridge, 2 near Fort Yukon, and 2 deployed by Eagle sonar.

Loggers at LYTF in 2021 recorded below average water temperatures much of the summer and fall seasons. Water temperature were near record low on June 13 and again July 17. The highest water temperatures (by time-of-season) occurred for 1 week in late June and similar temperatures for 3 days in mid-July (16–17°C). The historically warmest parts of the summer in mid-July were most often below historic average temperatures in 2021 (Figure 18). The maximum water temperature reached was 17.6°C during the entirety of LYTF operations from late May into September.

Pilot Station sonar temperature loggers encountered similar temperatures to LYTF and the maximum water temperature reached was 18.5°C (July 19) during the entirety of operations which were of a similar time frame to LYTF. Eagle sonar temperature loggers were deployed from June 30 through October 6 and generally experienced temperatures below (time-of-season) historical averages and the maximum water temperature reached was 19.5°C on July 23, 2021. Other temperature loggers, while not all deployed early in the season, had no indications of prolonged elevated temperatures occurring during the salmon migration. Other available information includes USGS temperature data<sup>11</sup> for Yukon River at Pilot Station, Tanana River at Nenana, Chena River near Two Rivers, and Salcha River near Salchaket. LYTF project also provides daily temperature data that is reported throughout the season using handheld monitors, these readings match closely with the loggers since the Yukon River waters are generally well mixed.

Inseason ADF&G takes water levels (discharge in cubic feet per second) into account when tracking groups of adult salmon as they migrate up the Yukon River as it affects their travel time, debris loads, and effectiveness of fishing gear. ADF&G provides inseason daily updates of water levels to the public for the following sites: Yukon River near Eagle, Stevens Village, and Pilot Station as well as locations on the Tanana River near Nenana, Chena River near Fairbanks, and Salcha River near Salchaket. In 2021 water levels were generally average to below average throughout the season in each of the sites monitored.

## **8.2 EAGLE SONAR**

ADF&G and DFO collaborate to jointly assess the passage of Yukon River mainstem Chinook and chum salmon just downstream of the international border (project is referred to as Eagle sonar). Since 2006, Chinook and fall chum salmon passage has been estimated using split-beam and imaging sonar operated near the community of Eagle, Alaska (McDougall and Brodersen 2020). There are effectively two separate fishing efforts at the project. The first is for collecting ASL and genetic samples from Chinook salmon and utilizes 5.25, 6.5, 7.5, and 8.5-inch mesh fished in a rotating schedule. These drifts are conducted twice a day (two fishing periods) until August 1 when one period is discontinued and, in its place, drifts are conducted to determine the crossover date

---

<sup>11</sup> <https://waterdata.usgs.gov/ak/nwis/current>

between the Chinook and fall chum salmon runs. The crossover drifts utilize only the 5.25 and 7.5-inch nets and incorporate a beach walk to ensure fall chum salmon are adequately represented in the catches. The drifts for collecting Chinook salmon samples are discontinued August 15 with the crossover drifts continuing through September 30. Although there is some minor overlap, Chinook and fall chum salmon runs are largely discrete in time based on test fishery results, local knowledge of catches, and data collected in Canada. There was a brief period when the ARIS was not operational on the right bank. This resulted in two days with partial counts and one day with no counts for the right bank. Estimates reported include interpolation of the missing data. The 2021 Chinook salmon passage estimate at the project was 31,796 fish  $\pm$  341 (90% CI) for the dates June 28 through August 31 (Appendix B11). The fall chum salmon passage estimate was 19,668 fish  $\pm$  400 (90% CI) for the dates September 1 through October 6. Because of continued passage at the termination of the project, the fall chum salmon estimate was subsequently adjusted to 23,170 fish (Appendix B16). This expansion was calculated using a second order polynomial for each day through October 18.

## **8.3 YUKON, CANADA**

### **Yukon River (Mainstem) Adult Chinook Salmon Assessment**

#### ***Big Salmon Sonar***

An ARIS Explorer 1800 multi-beam sonar was used to enumerate the Chinook salmon escapement to the Big Salmon River in 2021. This was the seventeenth year of escapement monitoring at a site approximately 1.5 km upstream of the confluence with the Yukon River. Sonar operation began on July 13 and continued without interruption through August 20, producing a count of 1,909 Chinook. An expansion was used to estimate the end of the run to September 1, using an exponential equation based on daily counts of the previous 10 days. The extrapolation resulted in the addition of 49 Chinook and a total passage estimate of 1,958 Chinook salmon (Appendix B12). This is the third lowest Big Salmon Chinook escapement recorded and was below the 10-year average (2011–2020) estimate of 5,048 fish. The peak daily count of 109 fish occurred on August 6 at which point 70% of the run had passed the sonar site. Approximately 50% of the run had passed the sonar by August 2, 2 days earlier than the 10-year average (2011–2020) midpoint (August 4). The 2021 Big Salmon sonar project report will be publicly available through the YRP website<sup>12</sup> after submission to the Pacific Salmon Commission R&E Fund Administrator. Carcass sample collection efforts in 2021 were truncated due to high water levels and turbidity and consequently no samples were collected.

#### ***Pelly River Sonar***

On the Pelly River, an ARIS Explorer 1800 multi-beam (left bank), and an ARIS Explorer 1200 multi-beam (right bank) sonar system were used to estimate the 2021 Chinook salmon passage. This was the sixth year of assessment undertaken by the Selkirk First Nation in collaboration with EDI Environmental Dynamics Inc., (EDI) at a site approximately 20 km upstream of the confluence of the Pelly and Yukon rivers. Sonar operation began on July 1 and concluded on August 25, counting 4,802 Chinook salmon. A postseason expansion to September 1 brought the total estimate to 4,980 fish (Appendix B12). The peak daily count of 197 fish on July 21 occurred when 27% of the run had passed. Correcting for leap year, approximately 50% of the run had

---

<sup>12</sup> <https://www.yukonriverpanel.com/restoration-enhancement-fund/r-e-project-reports/>



passed by July 28, matching the 2016–2020 average. No fish were captured during test netting, due to high water and equipment failures. Project reports will be publicly available through the YRP website after submission to the Pacific Salmon Commission R&E Fund Administrator.

### ***Klondike River Sonar***

A single ARIS Explorer 1200 multi-beam sonar system was installed on the right bank of the Klondike River to estimate the 2021 Chinook salmon passage. The 2021 season was the second year of assessment undertaken by the Tr'ondëk Hwëch'in First Nation and EDI following a trial year in 2019. This project is a continuation of sonar work conducted in 2009–2011 by Mercer and Associates, as supported by the R&E Fund. The 2021 sonar site was located near the Klondike River bridge, and approximately 2.6 km downstream of the 2009–2011 site and 2.1 km from the confluence of Klondike River with the Yukon River. Sonar operation began on July 1 and concluded on August 13, counting 843 Chinook salmon. A postseason expansion to August 22 brought the total estimate to 855 fish (Appendix B12). The peak daily count of 48 fish on July 21 occurred when 59% of the run had passed. Correcting for leap year, approximately 50% of the run had passed by July 20, two days earlier than the 2009–2011 and 2020 average (July 22). Project reports will be publicly available through the YRP website after submission to the Pacific Salmon Commission R&E Fund Administrator.

### **Whitehorse Rapids Fishway Chinook Salmon Enumeration**

The Whitehorse Rapids Fishway is a fish ladder, owned and operated by Yukon Energy Corporation, that bypasses the Whitehorse dam. It has an observation window into a chamber with upstream and downstream gates. The viewing window allows visual enumeration of migrating adult Chinook salmon. In 2021, Fishway staff counted 274 adult Chinook salmon at the Whitehorse Rapids Fishway between August 2 and September 5 (Appendix B12). This escapement was well below the 2011–2020 average of 1,075 Chinook salmon, and the second lowest count recorded since 1976 (the lowest count was in 2020.) Of these salmon, 98 (36% of run) were of hatchery origin and 176 (64% of run) were considered to be wild origin. The hatchery component included 8 females and 90 males. The wild component included 41 females and 135 males. Female Chinook salmon made up 18% of the total run to the Fishway (1988–2020 average 33%, range 18–56%, 2011–2020 average 31%, range 18–51%; DFO files).

The Whitehorse Rapids Fishway enumeration program is a joint initiative of the Yukon Fish and Game Association and Yukon Energy Corporation, with support from DFO. Students count all adult salmon migrating through the Fishway, record the sex and size category (small, medium, or large) of each salmon, identify hatchery-origin fish based on the absence of the adipose fin, and describe tags present on migrating salmon. Fishway staff also assist the Whitehorse Rapids Hatchery with broodstock collection at the Fishway.

### **Whitehorse Hatchery Operations**

The Whitehorse Rapids Hatchery, owned and operated by Yukon Energy Corporation, has released Chinook salmon fry upstream of the dam since 1985. The current annual release target of 150,000 (2.0 gram) fry has been in place since 2002; releases since that time have ranged from 85,306 fry in 2008 to 176,648 fry in 2003. The recent 10-year average (2011–2020) is 137,803 fry clipped and released upstream of the dam (unpublished data on file with Trix Tanner, Restoration Coordinator, DFO, Whitehorse, YT).



In 2021, all Chinook salmon fry released from the Whitehorse Rapids Hatchery into the Yukon River were marked. Fish had their adipose fin removed and were released upstream of the dam. This marking facilitates visual determination of the hatchery contribution to the run during observation of adult Chinook salmon migrating upstream through the viewing chamber at the Whitehorse Rapids Fishway; it also allows hatchery managers to identify hatchery-origin fish during broodstock collection. Fin clipping also enables researchers to distinguish hatchery fry from wild fry when investigating juvenile Chinook salmon habitat use. Marked fish are also recovered in marine studies, in river stock assessment of juvenile and adult Yukon River Chinook salmon, and in harvests. As in 2020, tagging under COVID-19 pandemic conditions posed safety concerns and logistical difficulties; no coded wire tags were applied in 2021. However, genetic samples were collected from parent broodstock to initiate a parentage based tagging program, which, once established, is expected to enable identification of Whitehorse Rapids hatchery release groups through genetic sampling of returning Chinook salmon.

A total of 124,547 Chinook salmon fry from the 2020 brood year were reared and marked (adipose fin-clipped) at the Whitehorse Rapids Hatchery and then released to two locations upstream of the Whitehorse Rapids hydroelectric dam (one site in Michie Creek, and one in M'Clintock River) on June 15, 2021. Average weight of all tagged fish at the time of release was 3.2 grams, while release groups average weights ranged from 2.9 grams to 3.4 grams.

Additionally, 665 fry from Whitehorse Rapids Hatchery eggs grown in the Stream to Sea classroom incubation program, were marked and released to Wolf Creek, tributary to the Yukon River upstream of the dam, between May 11 and June 7, 2021.

Brood stock collection in 2021 began on August 11, after 22 Chinook salmon had migrated through the Whitehorse Rapids Fishway and ended on September 5. A total of 59 males, including 37 wild and 22 adipose-clipped (hatchery) Chinook salmon, were removed from the Fishway for the brood stock program. The hatchery removed 26.2% of the total 225 Chinook salmon males. In total, 28 female Chinook salmon (57.1% of the total 49 female Chinook salmon), including 24 wild and 4 adipose-clipped (hatchery) salmon were removed for hatchery brood stock. Eggs were taken between August 30 and September 11, 2021 from 24 full (or nearly full) ripe females, and 1 partially spent female. Fecundity estimates ranged from 3,520 to 6,559 eggs, with a preliminary average, excluding partial spawns, estimated at 4,788 eggs.

The total estimated egg take in 2021 was 113,263 green eggs. Preliminary fertilization rate was estimated to be 99%. Egg removals prior to the eyed stage included 250 eggs to assess development and 3,459 mortalities; green egg to eyed egg survival was estimated at 97%. Thereafter, removals included 3,072 mortalities (between October 19 and October 27), and 850 eyed eggs donated to the Stream to Sea classroom incubation program; eyed egg to hatch survival was estimated at 97%. Since hatching, 1,121 dead alevins have been removed, resulting in an estimated 104,511 Chinook salmon alevins in incubators on December 14, 2021.

## **Porcupine River Investigations**

### ***Porcupine River Chinook Salmon Sonar***

In 2021, the Vuntut Gwitchin First Nation Government and DFO collaborated to enumerate Chinook salmon on the Porcupine River near Old Crow using multi-beam ARIS Explorer 1200 (right bank) and Explorer 1800 (left bank) sonars. Both sonars alternated every 30 minutes between

inshore ranges (1–20 m) and offshore ranges (20–40 m) 24 hours a day. On July 13, the right bank offshore range was increased to 20–70 m. The range was reset to 20–40 m on August 25. Set gill nets were deployed throughout the run to assess species composition and collect ASL data from Chinook salmon. This was the seventh year of Chinook salmon sonar enumeration on the Porcupine River.

Chinook salmon operations occurred from July 1 to August 14, producing a passage estimate of 409 Chinook salmon, including interpolated estimates for short periods of sonar downtime (Appendix B12). August 15 was selected as the crossover date based on daily passage estimates, with a 7-day rolling average showing an upward inflection beginning 15 August. This inflection point compares well to past crossover dates and corresponds with an increase in proportion of passage on left bank (typical bank used by migrating chum), and is shortly before a reported increase in 500 – 600 mm salmon-behaving targets viewed on sonar. Peak daily passage of 30 Chinook occurred on July 11, when 36% of the run had passed the sonar site. Approximately 50% of the run had passed the sonars on July 15 (the average midpoint of the run from 2014–2019 is July 23). Most Chinook enumerated (63%) migrated along the right bank. Approximately 46% of Chinook, enumerated along the right bank, migrated within 10 m of the sonar. Approximately 73% of Chinook, enumerated along the left bank, migrated within 10 m of the sonar. Passage rates were higher during the first six hours of the day with approximately 45% of inshore fish and 35% of offshore fish counted during this time.

The estimated passage of Chinook salmon was by far the lowest on record. Subtracting the local harvest estimate of 16 (14 harvested downstream of sonar and 2 harvested upstream) results in an escapement of 393 Chinook salmon to the upper Porcupine River drainage.

### ***Porcupine River Chum Salmon Sonar***

In 2021, the Vuntut Gwitchin First Nation Government and DFO collaborated to enumerate fall chum salmon on the Porcupine River near Old Crow using multi-beam ARIS Explorer 1200 (right bank) and Explorer 1800 (left bank) sonars. Both sonars alternated every 30 minutes between inshore ranges (1–20 m) and offshore ranges (20–40 m) 24 hours a day. This was the ninth year of Porcupine fall chum sonar enumeration (2011–2017, 2019, 2021).

The first chum salmon was caught in a set net on August 31, and a crossover date of August 15 was determined based on daily passage estimates, with a 7-day rolling average showing an upward inflection beginning 15 Aug. The final day of sonar operation was September 29. A second order polynomial equation (Crane and Dunbar 2011) postseason expansion was applied from Sept 29–Oct 11, adding 163 additional chum. The final total season passage estimate was 3,486 fall chum.

The run had three minor peaks; September 10 (206 fish, 54% of the run passed), September 17 (150 fish, 79% of the run passed) and September 27 (79 fish, 98% of the run passed). Approximately 50% of the run had passed by the sonars on September 9; the average midpoint of the run (2011–2017, 2019) is September 15.

The estimated passage of chum salmon was the lowest on record. Subtracting the local harvest of 21 fall chum (all harvested downstream of sonar) resulted in an escapement of 3,465 fall chum to the upper Porcupine River drainage.

### ***Fishing Branch River Chum Salmon Weir***

Fall chum salmon runs to the Fishing Branch River have been assessed annually since 1971. A weir has been used in most years, aerial surveys were used in some years, and in 2013 and 2014

estimates were based on proportion of radio tag recoveries combined with the sonar-based passage estimate on the Porcupine River mainstem (Appendix B15). Previous spawning escapement estimates for the Fishing Branch River have ranged from 4,795–353,282 fall chum salmon in 2020 and 1975, respectively (Appendix B15). In 2021, Fishing Branch River enumeration of fall chum salmon was conducted using a combination of a weir and video counter/sonar.

Weir installation began September 2 and was completed September 6. Video enumeration began September 7 and continued until September 16. Sonar enumeration occurred from September 16 to October 22. The video counting system was replaced with an above-weir sonar after chum were observed pooling at the entrance of the aluminum box at night. No preseason or postseason expansion was applied. Passage numbers at the start and end of the monitoring period were sufficiently low that expansion formulae would have provided no additional estimates. The final passage estimate of 2,413 fall chum salmon (Appendix B15) was below the Fishing Branch River interim escapement goal range of 22,000–49,000 fish. This escapement was the lowest estimate in 38 years of weir operation, and 51 years of assessment.

The fall chum salmon run peaked on September 21 with a maximum daily count of 130 fish (39% of the run had passed). Approximately 50% of the run had passed the weir by September 24. The average midpoint of the run from the past 10 years of weir operation (2008–2012 and 2015–2020) is September 25.

ASL data were collected from 195 fall chum (live and carcasses) between September 10 and October 20. The mean MEFL was 570 mm for sampled fall chum salmon (567 mm for females and 573 mm for males). Of the 184 samples that were successfully aged, 2% were age-3, 84% were age-4, 14% were age-5, and less than 1% were age-6. The sex composition of the combined video assessment (September 7–16) and ASL samples (n=411) was 51% female.

## **Aerial Surveys**

### ***Kluane River Aerial Survey***

An aerial survey of the Kluane River was conducted on October 25, 2021. Annual surveys of Kluane River were conducted 1972–2006, and were restarted in 2017 following a river piracy event at the headwaters of Kluane Lake (Shugar et al. 2017). The Kluane River index for 2021 was 64 fall chum salmon. Fish countability was considered fair due to moderate water clarity. This was among the lowest aerial counts on record, with counts reaching a maximum of 39,347 in 2003 (Appendix B15).

### ***Mainstem Yukon River Aerial Survey***

An aerial survey of the Yukon River mainstem index area (from Tatchun River confluence to Pelly River confluence) was conducted on October 19, 2021. Prior aerial surveys of this area occurred in 1973, 1975, 1983–1998, 2000–2006 and 2020. Historical fall chum salmon index counts ranged from 383 (1973) to 16,425 (2005). The 2021 index was 1,131 fish, among the lowest on record (Appendix B15).

## **Genetic Stock Identification and Stock Composition of Canadian Yukon River Chinook and Fall Chum Salmon**

Genetic stock composition of the 2021 Chinook salmon migration bound for Canada was estimated using genetic samples collected from the gillnet test fishing program conducted in conjunction with Yukon River sonar operations near Eagle. Genetic stock identification was conducted using

single nucleotide polymorphisms (SNPs). Of the 376 Chinook sampled at Eagle sonar in 2021, 373 returned usable genetic stock identifications. Chinook from the 2021 sample were identified to mid-mainstem Yukon River including Teslin River (33.5%), Carmacks area tributaries (22.5%), Pelly River and tributaries (13.1%), Stewart River and tributaries (8.8%), Teslin watershed above Teslin Lake (7.7%), northern Yukon River and tributaries (6.4%), upper Yukon River and tributaries (5.1%), and the White River and tributaries (3.0%; Appendix B24).

Relative stock composition estimates in 2021 were considerably lower than the 2008–2020 average for Teslin watershed (26.2%), and higher than average for Carmacks area (17.4%) and mid-mainstem Yukon River (21.1%; Appendix B24). Estimates for remaining stock groups were closer to historical values.

Genetic stock composition for fall chum salmon passing Eagle sonar was determined in a similar fashion to Chinook salmon. Genetic samples from 85 fall chum salmon captured in the gillnet test fishing program at Eagle sonar in 2021 were analyzed; all 85 samples returned usable genetic stock identifications using SNPs. Fall chum from the 2021 sample were identified to Yukon River mainstem including Minto area, Tatchun Creek area, Big Creek and Pelly River (65.0%), White River drainage including Kluane River (34.3%), Teslin River (0.7%) and the Yukon early stock group including Chanindu River (<0.1%; Appendix B25).

Relative stock composition estimates in 2021 were higher than the 2009–2020 average for Yukon mainstem (53.5%), and lower for White River drainage average (44.8%), though both were within the historically observed range of stock proportions (Appendix B25).

## **Environmental Conditions Report**

This annual summary describes environmental conditions influencing salmon habitat in the Canadian sub-basin of the Yukon River, including the Yukon and Porcupine rivers. The sub-basin encompasses over 100 documented spawning streams and many more rearing streams.

Due to the spatial scale, specific salmon habitat information is not collected extensively from year to year; the following information is a regional synopsis of what was experienced in the Canadian sub-basin during a given year. Weather records and stream discharge data are examined and compared with historic records to identify anomalies and/or unusual events, and their implications for salmon are considered. This report on environmental conditions is based on scientific evidence, field observations of the public, fishers, consultants, and DFO, and professional judgment.

### ***November 2020 to April 2021***

The 2020–2021 winter involved a range of conditions throughout the territory. In southern and central Yukon, precipitation events between November and March led to higher than average snowpack accumulation<sup>13</sup>. Meanwhile northern Yukon saw lower than average precipitation most of the winter until March and April. By May, the snowpack accumulation in southern Yukon was above average (up to 265% of the historical median), while central Yukon was slightly above average, and northern Yukon was below average (70%).

Air temperatures were colder than average in November, but unseasonably warm in December and January. Colder than average air temperatures occurred in February, March and April<sup>14</sup>. A cold

---

<sup>13</sup> Yukon Snow Survey and Water Forecast Bulletin <https://yukon.ca/en/yukon-snow-survey-bulletin-water-supply-forecast-may-2021>

<sup>14</sup> Environment Canada Monthly Climate Data <https://dd.weather.gc.ca/climate/observations/monthly/csv/YT/>

spring helped delay the spring melt, but ultimately the melting of the substantial snowpack would lead to a historic year of flooding in the Southern Lakes region. Yukon University hydrometric summaries<sup>15</sup> documented that the flooding of houses in the Whitehorse area began as early as April and May. Water temperature data, while limited in 2021, suggests spring temperatures were generally cooler than average, and similar to 2020 conditions (von Finster 2021).

Conditions in this period (November to April) align with Chinook and chum salmon incubation and emergence, and the beginning of outmigration of age-1+ Chinook salmon

### ***May 2021 to July 2021***

Yukon University hydrometric summaries noted that a combination of record snowpack, a late spring snowmelt, and above average temperatures at the end of June were conducive to flooding in southern Yukon. The Southern Lakes reached record heights at the end of June, in an event expected to occur only once every 200 years. Teslin Lake, Teslin River, and the Yukon River at Carmacks reached 50 year returns levels (levels expected to occur only once every 50 years). As a result, the summer saw a succession of flood watches, flood advisories, and evacuation notices<sup>16</sup>.

Air temperatures between May and July were average across the territory<sup>17,18</sup> and precipitation was below average<sup>19</sup>. Water temperatures in May were generally colder than average. After June water temperature data was only available from six stations, and all six stations were warmer than 2020 and periodically above average in June and July (von Finster 2021).

On Porcupine River conditions were more consistent over the season. The slow melt of a lower than average snowpack resulted in a relatively low spring flow. The low volume of water was relatively warm during the summer. The average monthly surface temperature measured at DFO and Vuntut Gwitchin Government's Porcupine Sonar Project near Old Crow in July was 18 °C, and similar to the 2017 and 2019 seasons.

For juvenile salmon, May through July corresponds with the downstream migration of age-1+ Chinook salmon, emergence and dispersal to rearing tributaries of age-0+ Chinook salmon, and emergence and downstream migration of chum salmon (age-0+). High water levels and discharge could promote early outmigration of age-1+ Chinook salmon fry, as well as the downstream displacement of newly emerged age-0+ juveniles of both species. Adult Chinook salmon enter the Yukon River in late May/early June and reach the mainstem Canadian border at the beginning of July. Chinook salmon spawning activity peaks in July in the Klondike River and starts in July in many Canadian Yukon River tributaries. Canadian-origin fall chum salmon enter the Yukon River mouth during this time. High water levels may have slowed the adult Chinook salmon migration. Warm water conditions are less favorable for migrating adult salmon.

### ***August 2021 to November 2021***

Flood level conditions in the Southern Lakes region persisted into late summer and the level of Atlin lake reached its peak on August 17. By early September levels had returned to normal

---

<sup>15</sup> Benoit Turcotte Hydrometric Blog <http://scholar.yukonu.ca/bturcotte/blog/significant-hydrological-events-2021-northwestern-canada>

<sup>16</sup> Government of Yukon Active flood warnings and advisories [Find out water levels in Yukon lakes and rivers | Government of Yukon](https://www.yukon.ca/en/flood-warnings)

<sup>17</sup> Environment Canada Canadian Climate Normals [https://climate.weather.gc.ca/climate\\_normals/index\\_e.html](https://climate.weather.gc.ca/climate_normals/index_e.html)

<sup>18</sup> Environment Canada Monthly Climate Data <https://dd.weather.gc.ca/climate/observations/monthly/csv/YT/>

<sup>19</sup> Environment Canada Seasonal Forecast\_ [https://weather.gc.ca/saisons/charts\\_e.html?season=mjj&year=2021&type=p](https://weather.gc.ca/saisons/charts_e.html?season=mjj&year=2021&type=p)

heights and all high water advisories for the Southern Lakes were ended. A late rain event in the Pelly and Stewart Rivers produced high flows in August, but they started to return to normal as winter approached. On the White River relatively high flows were experienced for the season, and a sharp runoff event on August 9 resulted in a large increase in discharge. On the Porcupine River, runoff events in August brought the water level up to average.

Average daily air temperatures during August to October were largely consistent with historical monthly normals. The end of September was colder than average in Old Crow leading to an early freeze up of the Porcupine River. The beginning of October was warmer than average in Dawson and Old Crow. November started off warmer than average, but the month ended with colder than average temperatures. While this pattern was seen across the territory, it was more evident in Old Crow where the mean daily temperature ranged from -4 to -38 °C.

Water temperatures from the few stations monitored showed warmer than average temperatures in early August, cooling by the end of the month, and variable temperatures into September.

This period (August to November) corresponds to Chinook and chum salmon migration, spawning, and early egg incubation. High water may have resulted in slower travel speeds, and contributed to late run timing of Chinook salmon to southern spawning areas. Warmer than average water temperatures could negatively affect spawning salmon. For juveniles warmer temperatures can potentially speed up Chinook salmon egg development if temperatures remain favorable throughout the winter. Chum salmon spawning sites in Yukon are dominated by groundwater; fall chum salmon are generally less susceptible than Chinook salmon to thermal effects on development due to moderating groundwater influences.

### ***Summary***

Migration, spawning, and rearing conditions in the Canadian sub-basins of the Yukon River were varied throughout the drainage in 2020–2021, but were dominated by flooding conditions in southern Yukon. How these conditions influence salmon varies with age and season. High water could accelerate the downstream migration of ocean bound juveniles, and disperse newly emerging Chinook to downstream habitat. High water may also delay the adult Chinook salmon migration in the Yukon River mainstem. While high water levels may allow adults to enter otherwise inaccessible small channels, this can negatively impact eggs by reducing water quality through increased sediment load. Cold conditions in the spring could delay emergence, and slow juvenile growth, while warmer water in the summer and fall are less favorable for adult migrating salmon.

Limited information is available for the Porcupine River watershed, but weather patterns suggest this region differed from other areas of the Yukon. After a cool spring the Porcupine River level remained low and warm for much of the summer. Warm temperatures in July and August may have adversely affected migrating Chinook salmon.

With increased climate variability, increased habitat monitoring and assessment in the Yukon River Canadian Sub-basin is encouraged to inform management, research, restoration strategies, and habitat considerations for Yukon River Pacific salmon populations.

## **9.0 MARINE FISHERIES INFORMATION**

Yukon River salmon migrate into the Bering Sea during the spring and summer after spending 0, 1, or 2 winters rearing in fresh water. Information about stock of origin from tagging, scale patterns, parasites, and genetic analysis indicate that Yukon River salmon are present throughout

the Bering Sea, in regions of the North Pacific Ocean, south of the Aleutian Chain, and the Gulf of Alaska during their ocean migration (Healey 1991; Salo 1991). Yukon River salmon have the potential to be captured by fisheries that harvest mixed stocks of salmon, other species of fish (bycatch), and by illegal fishing activities throughout their oceanic distribution. Coded-wire tag recoveries in these fisheries and in research surveys provide a key descriptor of the oceanic distribution of Yukon River Chinook salmon. However, genetic stock identification has become the primary tool for identifying Yukon River Chinook salmon in marine habitats (Larson et al. 2013; Guthrie et al. 2016). The U.S. groundfish trawl fisheries in the Gulf of Alaska (GOA) and Bering Sea-Aleutian Islands (BSAI) management areas are managed to limit the incidental harvest (bycatch) of salmon.

Appendix C was prepared by NOAA in coordination with ADF&G at the request of the YRP. It provides background information on BSAI fisheries, bycatch regulations, and information to understand bycatch impacts on Canadian-origin salmon. Recent year and historical bycatch information is provided and will be updated annually as new information becomes available. Estimated adult equivalent bycatch of Yukon River Canadian-origin Chinook salmon from the BSAI pollock fisheries are available from 1994–2017 (Ianelli and Stram 2018).

## **10.0 RUN OUTLOOKS 2022**

### **10.1 YUKON RIVER CHINOOK SALMON**

Over the years, the JTC has used a range of methods to produce an annual preseason outlook of Canadian-origin Chinook salmon run abundance. Run outlooks are used by fishery managers and stakeholders as a tool for guiding the development of preseason harvest strategies (Table 13, Figure 19).

#### **Canadian-origin Brood Table**

The brood table for Canadian-origin Yukon River Chinook salmon (Appendix A3) is the basis of the current spawner-recruitment model (Figure 20) which is one of the models used to forecast returns in future years. Age-specific returns have been estimated from border passage, harvest and escapement data. Because assessment methods have changed over time, the brood table is constructed from a variety of data sources. For the years 1982–2001, initial border passage estimates were derived from the DFO Chinook salmon mark–recapture program, but information from several sources, reviewed in 2008, indicated that these data were biased low. Subsequently, the 1982–2001 Canadian spawning escapement estimates were reconstructed using a linear regression of the estimated total spawning escapements for 2002–2007 against a 3-area aerial survey index of combined counts from Big Salmon, Little Salmon, and Nisutlin rivers. Spawning escapement estimates for years 2002–2004 were based on radiotelemetry studies. Since 2005, spawning escapement estimates have been derived by subtracting both Canadian and U.S. harvests that occurred upriver from the sonar project site from the passage estimates at Eagle sonar. A standardized age dataset for Chinook salmon passage at the U.S./Canada border (Hamazaki 2018) was adopted by the JTC in 2019 and used to update the brood table (JTC 2020).

#### **Canadian-origin Yukon River Chinook Salmon**

The JTC forecast subcommittee has been in the process of updating the Canadian-origin Chinook salmon run-size forecast model to improve the forecast accuracy and to improve methods used to account for uncertainty. The 2022 preseason forecast for Canadian-origin Chinook salmon is based

on three independent models weighted by forecast performance within a Bayesian framework. The three models include a dynamic sibling model, spawner-recruitment model, and juvenile abundance model based on Northern Bering Sea surface trawl surveys. The common time period over which performance of these three models is evaluated for weighting purposes is: 2010, 2013-2021.

### ***Dynamic Sibling Model***

The dynamic sibling model predicts the 2022 run size of Canadian-origin Chinook salmon will be approximately 47,000 (80% credible interval 30,000–66,000) fish. This model predicts age class returns based on prior years sibling (younger) returns and accounts for change in age at maturity over time. Age-5, age-6, and age-7 predictions were based on the dynamic sibling model using model fits from 1982–2021; whereas age-3, age-4, and age-8 predictions were based on the recent 10-year average return. Age class predictions were summed to produce the total estimated run size.

### ***Spawner-recruit Model***

The spawner-recruitment model predicts the 2022 run size of Canadian-origin Chinook salmon will be approximately 79,000 (80% credible interval 49,000–116,000) fish. This model uses a Ricker relationship based on the number of spawners and recruits from 1982–2015 to calculate the total expected returns from each brood year escapement. Run size predictions for 2022 were based on the predicted recruitment from the appropriate brood years, multiplied by the 5-year average (2017–2021) proportions for age-5, age-6, and age-7 fish. Predictions for the subdominant age classes, age-3, age-4, and age-8 are the recent 10-year average of abundance. The current formulation of this model does not account for changes in productivity over time. Over the last 10 years, the spawner-recruitment model has been on average 71% different (i.e., (observed – forecast) / observed) compared to observed run sizes. The JTC forecast subcommittee intends to continue exploring the appropriateness of the Ricker model as a forecast tool and consider options to account for changes in productivity to improve performance. However, the fully integrated Bayesian forecast method accounts for the uncertainty and heavily penalizes the Ricker component by weighting it less compared to other information sources.

### ***Juvenile-based Forecast***

Surface trawl surveys in the northern Bering Sea (Murphy et al. 2021) are used to estimate the abundance of Yukon River salmon stocks during their first summer at sea (juvenile life-history stage). Since 2003, juvenile Chinook salmon catch from the trawl surveys, coupled with genetic MSA, has been used to provide stock-specific juvenile abundance estimates (Figure 21; Murphy et al. 2017, Howard et al. 2019, Howard et al. 2020, Murphy et al. 2021). Juvenile Chinook salmon have experienced relatively stable marine survival following their first summer in the northern Bering Sea. As a result of this stable marine survival, the relationship between juvenile Chinook salmon abundance in the northern Bering Sea correlates to adult returns to the Yukon River (Figure 22). This relationship is pivotal to the juvenile-based forecast model used to predict adult returns up to 3 years in advance. Juvenile abundance-based forecasts of Canadian-origin Chinook salmon have been provided to the JTC and YRP since 2013. Beginning in 2018, the JTC decided to explicitly incorporate the juvenile-based forecast as part of the formal outlook.

Juvenile Chinook salmon in the Bering Sea in 2018 and 2019 (returning as age-6 and age-5, respectively) will be the primary contributors to the 2022 adult run. Both the 2018 and 2019 juvenile abundances were below average, continuing the downward trend in juvenile abundance



that began in 2017 (Figure 21). Juvenile abundance models indicate that the projected run size of Canadian-origin Chinook salmon in 2022 should be between 25,000–59,000 fish (point estimate of 42,000 fish, Figure 23). The juvenile forecast ranges are based on an 80% prediction interval calculated from the relationship between juvenile abundance and adult returns. The run-size forecasts and ranges are estimated from predicted returns using a three-year window of average maturity.

### ***2022 Canadian-origin Chinook Salmon Forecast***

The final forecast for 2022 Canadian-origin Chinook salmon run was developed using an integrated Bayesian approach to better account for uncertainty and to weight each individual component model (i.e., dynamic sibling, spawner recruit and juvenile models) by its performance in the recent past. To weight each component model the empirical standard deviation was calculated based on predictions for the comparable time period of 2010 and 2013–2021<sup>20</sup>. Standard deviation was calculated as the standard deviation of the log of predicted divided by observed. A random variable for the combined run size prediction was estimated with an uninformative prior, and log-normal likelihoods aligned this estimate with the prediction from each forecast model, in proportion to past performance. This integrated Bayesian estimation procedure results in forecast component models with low relative standard deviation being given a higher weight in the integrated model (and vice versa). The Ricker model had the poorest fit to the observed run sizes (i.e., greatest amount of uncertainty), and an empirical standard deviation<sup>21</sup> of 0.330 was applied. The dynamic sibling and juvenile models tended to fit similarly well to prior observed run sizes and were assigned an empirical standard deviation of 0.284 and 0.223 respectively. The result is a posterior distribution for the integrated forecast that resulted in a combined point estimate of 51,000 with a 80% credible interval of 41,000–62,000.

The JTC recommends using an 80% credible interval as the basis for an operational forecast range of 41,000–62,000 Canadian-origin Chinook salmon for 2022 (Table 13). The 80% credible interval implies a 20% chance (1 in 5) that the 2022 run size will fall outside the forecast range based on past model performance. The lower end of the 2022 outlook range suggests a possible run size similar to the record low run size observed in 2021 (Table 13). The upper end of the outlook is for a run size smaller than the recent 10-year average (2012–2021) of 64,000 Chinook salmon (Appendix B18), and well below the 1982–1997 average of 153,000 Chinook salmon (Appendix B11).

The Chinook salmon runs on the Yukon River are typically dominated by age-5 and age-6 fish. The brood years producing these age classes in 2022 are 2016 (age-6) and 2017 (age-5). The Canadian-origin Yukon River Chinook salmon spawning escapement in 2016 of 68,798 fish and 2017 escapement of 68,315 fish were above the 1982–2014 average escapement of 47,000 fish (Appendix A3; Figure 9). The age-4 (725) and age-5 (14,940) estimated returns in 2021 were below the long-term average brood year return of 5,683 and 32,020 fish, respectively and the age-4 return was the lowest on record (Appendix A3).

## **10.2 YUKON RIVER SUMMER CHUM SALMON**

The strength of the summer chum salmon run in 2022 will be dependent on production from the 2018 (age-4 fish) and 2017 (age-5 fish) escapements, because these age classes generally dominate

---

<sup>20</sup> The years 2011 and 2012 could not be included because juvenile forecasts were not available.

<sup>21</sup> Empirical deviation is reported in log-space.

the run. The drainagewide spawning escapement in 2017 and 2018 was approximately 3.0 million and 1.4 million summer chum salmon, respectively. The return of age-4 and age-5 fish in 2021 were the second smallest and smallest, respectively, observed since 1978. The overall return of 154,000 summer chum in 2021 was the smallest on record and 93% smaller than the 1978-2020 average of 2.5 million. Below average returns of age-4 and age-5 chum salmon were also observed in Yukon fall chum salmon, other wild chum salmon stocks throughout Alaska, as well as hatchery stocks of chum salmon in Alaska. The spatial extent of observations in 2020 and 2021 is evidence that common ocean conditions contributed to the poor run of age-4 and age-5 chum salmon which indicates the return of age-5 and age-6 summer chum salmon in 2022 may be poor.

Historically, the drainagewide summer chum salmon forecast was developed by forecasting the run size of the Anvik River component, based on projections of brood year returns and sibling relationships, and then scaling up based on historical contribution of the Anvik River to the total run. Unfortunately, Anvik sonar did not operate in 2020 due to COVID-19 related travel restrictions. However, a drainagewide run reconstruction model was developed in 2016 (Hamazaki and Conitz 2015), and the resulting model estimates of escapement and total return (1978–2020) were used to develop a drainagewide brood table and forecast the 2021 summer chum salmon run. This method was again used to develop the 2022 forecast. The expected 2022 summer chum salmon run is forecast to be 330,000 (80% CI range of 160,000 to 540,000) fish, which is slightly larger than the 2021 run of approximately 154,000 fish. The relatively wide forecast range is representative of the uncertainty associated with the poor 2021 age-4 and age-5 returns and implications for the 2022 run.

The 2022 summer chum salmon run is highly uncertain and may not provide for a normal subsistence harvest or a surplus for a commercial harvest. The upper end of the forecast range is near the lower end of the drainage wide escapement goal of 500,000 summer chum salmon. Unless the 2022 summer chum run is stronger than predicted, no summer chum are available for harvest.

## **10.3 YUKON RIVER FALL CHUM SALMON**

### **Drainagewide Fall Chum Salmon**

The preseason forecast is determined using estimates of escapement and resulting production (spawner-recruit). The brood table for the drainagewide fall chum salmon is the basis of the current spawner-recruitment model. The age-specific returns have been estimated based on the samples collected in the lower Yukon River which is primarily gillnet fisheries applied to the escapement and harvests throughout the drainage. Yukon River drainagewide estimated escapement of fall chum salmon for the period 1974 through 2015 has ranged from approximately 224,000 (2000) to 2,200,000 (1975) fish, based on Bayesian analysis of escapement assessments to approximate overall abundance (Fleischman and Borba 2009). Escapements between 1974 and 2015 resulted in subsequent brood year returns that ranged in size from approximately 313,000 (1996 production) to 2,900,000 (2001 production) fish. Corresponding return per spawner rates ranged from 0.3–9.0, averaging 1.8 for all years combined (1974–2015; Appendix A8).

A considerable amount of uncertainty has been associated with these run forecasts, particularly in the last two decades, because of unexpected run failures (1998–2002) followed by strong runs from 2003 through 2008. Weakness in these salmon runs prior to 2003 was generally attributed to reduced productivity in the marine environment and not to low levels of parental escapement. The 2020 and 2021 run failures also appear to be attributed to the marine environment as it was observed to be widespread in chum salmon throughout western Alaska.

Beginning in 1999, Yukon River fall chum salmon preseason forecasts have been presented as a range, to better represent uncertainty in the expected run size. In most years, the expected run size (point estimate) was forecast using estimates of brood year escapement, estimates of returns per spawner (production), and maturity schedules developed for even and odd years based on historical averages. In 1998, the forecast method overestimated run size due to an unexpected poor return. To account for this, the point estimate was used as the upper bound of the forecast range in subsequent years (1999–2005; Brenner et al. 2020). The lower end of the forecast range was generated by adjusting the point estimate based on the average forecast performance (i.e., ratio of observed to predicted). Forecast performance from 1998–2003 were used to inform the 1999–2004 outlooks. As run sizes increased over the early to mid-2000s, the forecast performance improved, and in 2005 the lower bound of the forecast range was based on the 2001–2004 average forecast performance. Beginning in 2006, adjustments to the point estimate were no longer applied. Instead, the forecast range was based on a statistical confidence interval around the point estimate. Since 2006, the annual forecasts have been informed by different odd- and even-year maturity schedules based on the historical averages available at the time and assumptions of stock productivity. For example, in 2006 and 2007 average age composition from years 1974–1983 were used to represent high productivity years, whereas in 2008–2012 data from 1984–2012 was used to represent low productivity years. Since 2013, the average odd- and even-year maturity schedules have been calculated from the complete historical dataset.

The majority of fall chum salmon return at age-4 (2018) and age-5 (2017), and a smaller proportion return at age-3 and age-6 (Appendix A8). As such, the 2022 run will be composed of brood years 2016–2019 (Table 14). The escapements in both 2017 and 2018 were above the upper end of the drainagewide escapement goal range of 300,000–600,000 fall chum salmon. It is anticipated that the 2022 return will be dominated by age-4 fish (Table 14), as the age-5 components from both 2016 and 2017 resulted in new record low brood year returns (Appendix A8). Estimates of returns per spawner (R/S) were used to estimate production for 2016 and 2017, and typically a Ricker spawner-recruit model was used to predict returns from 2018 and 2019. The average odd and even year maturity schedule was calculated from the complete historical dataset since 1974. That maturity schedule was applied to the estimated production (i.e., returns) for each contributing brood year and summed to estimate the total number of fall chum salmon that are expected to return in the coming year. The result from the Ricker model was a forecast point estimate of 643,000 fall chum salmon returning drainagewide.

The sibling model predicts the drainagewide run size will be approximately 415,000 fall chum salmon. The model predicts age class returns based on prior years sibling returns. Ages-3, age-4, age-5, and age-6 predictions were based on the sibling model using 1974-2015. Age class predictions were summed to produce the total estimated run size. Brood year returns of age-3 fish range from 0 to 198,000 fall chum salmon. Returns of age-4 fish from even-numbered brood years during the time period 1974–2017 average 487,000 fall chum salmon with a range from a low of 89,000 for brood year 2016 to a high of 1,200,000 for brood year 2012. Returns of age-5 fish from the same time period for odd-numbered brood years average 254,000 fall chum salmon with a range from a low of 67,000 fish for brood year 1975 to a high of 719,000 fish for brood year 2001. Considering the sibling relationship described, and the record low returns of age-5 fish in the last two years the contribution of age-5 fish is expected to be well below the odd-numbered year average. The age-4 fish are also expected to be well below the even-numbered year average until there is an observed improvement in production in this dominant age class.

Forecast models rarely predict extreme changes in production. The fluctuations observed in fall chum salmon run sizes (postseason run size estimates) in comparison with the expected run sizes (preseason outlooks) are reflected in the outlook performance, i.e., proportions of the expected run size, observed for the 1998–2021 period (Table 15). In attempts to produce a more credible forecast, considering the recent drastic run failures, the 2022 forecast point estimate of 110,000 fall chum salmon was developed based on the average forecast performance (i.e., ratio of observed to predicted) for 2020 and 2021. To provide the range of possibilities around the estimate both the individual proportions were used and resulted in the forecast range of 78,100 to 148,000 fall chum salmon. This forecasted drainagewide fall chum salmon run size is well below average (1998–2021; Table 15).

During the 2022 fall fishing season, estimated strength of the projected run of fall chum salmon will be adjusted using the relationship to summer chum salmon run abundance and assessed based on various inseason monitoring project data. With a forecasted run size range of 78,100–148,000 fall chum salmon (point estimate 110,000 fish; Table 14), it is anticipated that escapement goals will not be met, and the run will not support normal subsistence fishing activities. The forecast suggests no surplus of fall chum salmon will be available for commercial harvest. However, harvestable surpluses for Yukon River fisheries will be determined inseason and applied to the guidelines outlined in the fall chum salmon management plan. The first inseason projection will refine the forecast based on the relationship between the summer and fall chum salmon runs in mid-July at the beginning of the fall season.

### **Canadian-origin Upper Yukon River Fall Chum Salmon**

To develop an outlook for the 2022 Canadian-origin Yukon River fall chum salmon, the drainagewide outlook range of 78,100–148,000 fall chum salmon (point estimate 110,000) was multiplied by 25% (the estimated contribution of mainstem Yukon River Canadian-origin fall chum salmon), producing an outlook range of 20,000–37,000 fish with a midpoint of 28,000 fish (rounded to the nearest 1,000; Table 16). Recent genetic stock identification analyses have indicated that the assumption of 25% is reasonable.

### **Canadian-origin Porcupine River Fall Chum Salmon**

In the Canadian section of the Porcupine River, a majority of the production of fall chum salmon originates from the Fishing Branch River. Canadian-origin Porcupine River stocks have been estimated to comprise approximately 5% of the drainagewide run. Fishing Branch River fall chum salmon are estimated to comprise between 40% and 80% of the Canadian-origin Porcupine River stocks, and approximately 4% of the drainagewide run, though estimates have ranged from 1%–7%. Applying the 4% average estimate to the drainagewide outlook range of 78,100–148,000 fish (point estimate 110,000) results in a Fishing Branch River outlook of 3,000–6,000 fish, with a midpoint of 4,000 fish (rounded to the nearest 1,000 fish; Table 17). This outlook is considered uncertain due to the high variation in contributions of Fishing Branch River fall chum salmon to drainagewide stocks.

Though the models used to develop forecasts have varied from year-to-year, the postseason run size estimates of Fishing Branch River fall chum salmon have been consistently below preseason outlooks since 1998, except for 2003–2005, 2016, and 2017.

## **10.4 YUKON RIVER COHO SALMON**

Although there is little comprehensive escapement information for Yukon River drainagewide coho salmon, it is known that coho salmon primarily return as age-2.1 fish (4-year-old, age in European notation) and overlap in run timing with fall chum salmon. The major contributor to the 2022 coho salmon run will be age-4 fish returning from the 2018 parent year. Based on the run reconstruction index (1995–2021, excluding 1996 and 2009), the 2018 escapement was estimated to be 143,000 coho salmon, which was below the average (163,000). In 2018, a relatively large amount of coho salmon was harvested incidentally in the directed fall chum salmon commercial fisheries (exploitation estimate at 45%). Subsistence harvest in 2018 was well below the 2013–2017 average of 13,000 coho salmon (Appendix B5). The returns from 2014 through 2018 have been high abundance years (averaging over 300,000 fish) which may indicate good productivity which typically cycles for several years in succession. However, the run sizes have been declining since 2016 with run sizes less than 200,000 coho salmon in both 2019 and 2020, followed by a record low return in 2021 which confirms a transition to a cycle of lower productivity.

Escapements are primarily monitored in the Tanana River drainage. The Delta Clearwater River (DCR) is a major producer of coho salmon in the upper Tanana River drainage and has comparative escapement monitoring data since 1972 (Appendix B17). The DCR parent year escapement of 2,884 fish in 2018 was well below the SEG range of 5,200–17,000 coho salmon. Six other locations in the Tanana River drainage were surveyed for coho salmon specifically; half of them were above average when compared to the 2016–2020 average escapements. Very informal coho salmon outlooks are made preseason based on average survival of the primary parent year escapement estimate, which in 2022 would indicate that the return would be average to below average.

## **11.0 STATUS OF ESCAPEMENT GOALS**

### **11.1 SPAWNING ESCAPEMENT TARGET OPTIONS IN 2021**

Canadian-origin mainstem Yukon River Chinook, and mainstem and Fishing Branch fall chum salmon, are managed under the umbrella of the YRSA. The YRP meets annually and recommends escapement goals for Canadian-origin stocks to the Canadian and U.S. management agencies.

#### **Canadian-origin Mainstem Yukon River Chinook Salmon**

In 2010, the YRP adopted an IMEG range of 42,500–55,000 Chinook salmon. In the absence of a biological escapement goal, i.e., a goal based on a production or population model, the IMEG has been retained each year since then. Beginning in 2019, the JTC undertook a comprehensive bilateral effort to model the spawner-recruit dynamics for this stock aggregate and estimate biological reference points and probability profiles that could be used to recommend a biological escapement goal to the YRP. Model results were peer-reviewed in January of 2022 through the Canadian Science Advice Secretariat and found to be appropriate for informing management decisions. At its March 2022 meeting, the JTC used the model results to review the current IMEG, consider alternative goal options, and formulate an updated recommendation for the YRP<sup>22</sup>.

---

<sup>22</sup> At the time of publication of this JTC report, detailed reports describing the model development, data inputs, results, and conclusions are pending along with results of the peer review proceedings and science advice. Summaries of the JTC considerations and rationale pertaining to its recommendation will be provided to the YRP in April 2022 via presentations and written documentation.

The JTC reached bilateral consensus to recommend an escapement goal objective of 52,500 with an acceptable deviation range of 42,500–62,500. The JTC recommendation to revise the goal stemmed from a desire to equally balance yield and recruitment objectives. Review of goal range options that focused only on future yield or recruitment revealed inherent trade-offs, such that one could not be optimized without compromising the other. Due to limitations in guidance within the Pacific Salmon Treaty and guidance provided by the Panel, the JTC was not able to clearly prioritize future harvest (i.e., yield) or future run sizes (i.e., recruitment).

The JTC considered a variety of strategies to equally prioritize yield- and recruitment-based objectives. The JTC recognized that even when consistent criteria were applied to both yield and recruitment probabilities, inherent trade-offs at the lower and upper bounds remained. The JTC agreed to identify a single escapement value that had the same probability of maximizing yield and recruitment as a preferred option to demonstrate that both objectives were of equal priority. The value of 52,500 was identified as the escapement where the probability of maximum yield and maximum recruitment were as equal as possible. This resulted in a 72% and 70% probability of achieving at least 90% of maximum yield and recruitment respectively.

Notwithstanding the desire to achieve the stated target of 52,500, the JTC acknowledged that management to precisely achieve a single escapement value was not realistic, and an acceptable level of deviation from the target was required to account for assessment and management imprecision. A value of  $\pm 20\%$  of the target escapement objective was agreed to as reasonable. This identified maximum acceptable deviations from the 52,500 target of  $\pm 10,500$  fish, resulting in an acceptable range of 42,000–63,000. The acceptable range was adjusted to 42,500–62,500 in consideration of the current IMEG, which was intended to emphasize that the JTC did not recommend lowering the escapement goal without additional rationale or guidance.

### **Canadian-origin Mainstem Yukon River Fall Chum Salmon**

In 2010, the YRP adopted an IMEG range of 70,000–104,000 Canadian-origin mainstem Yukon River fall chum salmon. This range was developed as 0.8–1.2 times the estimated spawners at maximum sustained yield (86,600 fish), which was derived prior to the returns from the exceptional 2005 spawning escapement of over 437,000 fall chum salmon. Based on prior recommendations by the JTC, the YRP extended this IMEG for the 3-year period of 2020–2022.

### **Fishing Branch River Fall Chum Salmon**

An IMEG range of 22,000–49,000 fall chum salmon for the Fishing Branch River has been extended for 3-year periods since 2008 (Appendix B15). Based on prior recommendations by the JTC, the YRP extended this IMEG for the 3-year period of 2020–2022.

## 12.0 REFERENCES CITED

- Andersen, D. B. 1992. The use of dog teams and the use of subsistence-caught fish for feeding sled dogs in the Yukon River drainage. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 210, Juneau.
- Baker, B. 2018. Fishery management report for recreational fisheries in the Tanana River management area, 2017. Alaska Department of Fish and Game, Fishery Management Report No. 18-33, Anchorage.
- Brenner, R. E., S. J. Larsen, A. R. Munro, and A. M. Carroll, editors. 2020. Run forecasts and harvest projections for 2020 Alaska salmon fisheries and review of the 2019 season. Alaska Department of Fish and Game, Special Publication No. 20-06, Anchorage.
- Brown, C., and D. Jallen. 2012. Options for amounts reasonably necessary for subsistence uses of salmon: Yukon Management Area; prepared for the January 2013 Anchorage Alaska Board of Fisheries meeting. Alaska Department of Fish and Game, Division of Subsistence, Special Publications No. BOF 2012-08, Fairbanks.
- Dreese, L. M., and J. D. Lozori. 2019. Sonar estimation of salmon passage in the Yukon River near Pilot Station, 2018. Alaska Department of Fish and Game, Fishery Data Series No. 19-16, Anchorage.
- DuBois, L. 2018. Origins of Chinook salmon in the Yukon Area fisheries, 2014. Alaska Department of Fish and Game, Fishery Data Series No. 18-25, Anchorage.
- Fleischman, S. J., and B. M. Borba. 2009. Escapement estimation, spawner-recruit analysis, and escapement goal recommendation for fall chum salmon in the Yukon River drainage. Alaska Department of Fish and Game, Fishery Manuscript No. 09-08, Anchorage.
- Guthrie, C. M., H. T. Nguyen, and J. R. Guyon. 2016. Genetic stock composition analysis of the Chinook salmon bycatch from the 2014 Bering Sea walleye pollock (*Gadus chalcogrammus*) trawl fishery. U.S. Dep. Commerce NOAA Tech. Memo. NMFS-AFSC-310. 25 p.
- Hamazaki, T. 2018. Estimation of U.S.-Canada border age-composition of Yukon River Chinook salmon, 1982–2006. Alaska Department of Fish and Game, Fishery Data Series No. 18-21, Anchorage.
- Hamazaki, T., and J. M. Conitz. 2015. Yukon River summer chum salmon run reconstruction, spawner-recruitment analysis, and escapement goal recommendation. Alaska Department of Fish and Game, Fishery Manuscript Series No. 15-07, Anchorage.
- Healey, M. C. 1991. Life history of Chinook salmon (*Oncorhynchus tshawytscha*). [In]: Groot, C. and L. Margolis, editors, Pacific Salmon Life Histories. UBC Press, Vancouver, B.C., Canada, pp. 311-394.
- Howard, K.G., S. Garcia, J. Murphy, and T.H. Dann. 2019. Juvenile Chinook salmon abundance index and survey feasibility assessment in the Northern Bering Sea, 2014–2016. Alaska Department of Fish and Game, Fishery Data Series No. 19-04, Anchorage.
- Howard, K. G., S. Garcia, J. Murphy, and T. H. Dann. 2020. Northeastern Bering Sea juvenile Chinook salmon survey, 2017 and Yukon River adult run forecasts, 2018–2020. Alaska Department of Fish and Game, Fishery Data Series No. 20-08, Anchorage.
- Ianelli, J. N. and D. L. Stram. 2014. Estimating impacts of the pollock fishery bycatch on western Alaska Chinook salmon. *ICES J. Mar. Sci.* **72**: 1159-1172. doi:10.1093/icesjms/fsu173.
- Ianelli, J. N., and D. L. Stram. 2018. Chinook Bycatch Mortality Update. Discussion paper presented to the North Pacific Fishery Management Council, April 2018. Available online at: <http://npfmc.legistar.com/gateway.aspx?M=F&ID=e172520e-fc22-46e8-b5aa-72ba233f129e.pdf>
- Jallen, D. M., S. K. S. Decker, and T. Hamazaki. 2017. Subsistence and personal use salmon harvests in the Alaska portion of the Yukon River drainage, 2015. Alaska Department of Fish and Game, Fishery Data Series No. 17-39, Anchorage.
- JTC (Joint Technical Committee of the Yukon River U.S./Canada Panel). 2020. Yukon River salmon 2019 season summary and 2020 season outlook. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 3A20-01, Anchorage.

- Larson, W. A., F. M. Utter, K. W. Myers, W. D. Templin, J. E. Seeb, C. M. Guthrie, A. V. Bugaev, and L. W. Seeb. 2013. Single-nucleotide polymorphisms reveal distribution and migration of Chinook salmon (*Oncorhynchus tshawytscha*) in the Bering Sea and North Pacific Ocean. *Canadian Journal of Fisheries and Aquatic Sciences* 70:128-141.
- Liller, Z. W., and J. W. Savereide. 2018. Escapement goal recommendations for select Arctic-Yukon-Kuskokwim Region salmon stocks, 2019. Alaska Department of Fish and Game, Fishery Manuscript No. 18-08, Anchorage.
- McCullough, D. S. Spalding, D. Sturdevant, and M. Hicks. 2001. Summary of technical literature examining the physiological effects of temperature on salmonids. Issue Paper 5. U.S. Environmental Protection Agency. EPA-910-D-01-005.
- McDougall, M. J., and N. B. Brodersen. 2020. Sonar estimation of Chinook and fall chum salmon passage in the Yukon River near Eagle, Alaska, 2019. Alaska Department of Fish and Game, Fishery Data Series No. 20-26, Anchorage.
- Murphy, J.M. E.V. Farley, J.N. Ianelli, and D.L. Stram. 2017. Distribution, diet, and bycatch of chum salmon in the Eastern Bering Sea. *N. Pac. Anadr. Fish. Comm. Bull.* 6:219-234. doi: 10.23849/npafcb6/219.234
- Murphy, J. M., K. G. Howard, J. C. Gann, K. C. Cieciel, W. D. Templin, and C. M. Guthrie. 2017. Juvenile Chinook salmon abundance in the northern Bering Sea: Implications for future returns and fisheries in the Yukon River. *Deep Sea Research Part II: Topical Studies in Oceanography* 135:156-167.
- Murphy, J., S. Garcia, J. Dimond, J. Moss, F. Sewall, W. Strasburger, E. Lee, T. Dann, E. Labunski, T. Zeller, A. Gray, C. Waters, D. Jallen, D. Nicolls, R. Conlon, K. Cieciel, K. Howard, B. Harris, N. Wolf, and E. Farley Jr. 2021. Northern Bering Sea surface trawl and ecosystem survey cruise report, 2019. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-423, 124 p.
- NPAFC (North Pacific Anadromous Fish Commission). 2001. Plan for NPAFC Bering-Aleutian Salmon International Survey (BASIS) 2002-2006. North Pacific Anadromous Fish Commission Doc. 579. 27 pp. Available at: [www.npafc.org](http://www.npafc.org).
- Pfisterer, C. T., T. Hamazaki, and B. C. McIntosh. 2017. Updated passage estimates for the Pilot Station sonar project, 1995-2015. Alaska Department of Fish and Game, Fishery Data Series No. 17-46, Anchorage.
- Salo, E. O. 1991. Life history of chum salmon, *Oncorhynchus keta*. [In]: Groot, C., and L. Margolis, editors. *Pacific Salmon Life Histories*. UBC Press, Vancouver, B.C., Canada, pp. 231-309.
- Shugar, D.H., J. J. Clage, J. L. Best, C. Schoof, M. J. Willie, L. Copeland and G. H. Roe. 2017. River piracy and drainage basin reorganization led by climate driven glacier retreat. *Nature Geoscience* 10:3701-376.
- von Finster, A. 2021. Water temperature monitoring of Yukon River Chinook salmon spawning and migration habitats in Canada – annual project report – 2021. Report to the Yukon River Panel, Project CRE-20-21. 39 pp.



## **TABLES AND FIGURES**

Table 1.—Yukon Area regulatory subsistence salmon fishing schedule, in U.S. waters.

| Area                      | Regulatory subsistence fishing periods | Open fishing times                          |
|---------------------------|--|---|
| Coastal District          | 7 days per week                        | M/T/W/TH/F/SA/SU - 24 hours/day             |
| District 1                | Two 36-hour periods per week           | Mon 8 pm to Wed 8 am / Thu 8 pm to Sat 8 am |
| District 2                | Two 36-hour periods per week           | Wed 8 pm to Fri 8 am / Sun 8 pm to Tue 8 am |
| District 3                | Two 36-hour periods per week           | Wed 8 pm to Fri 8 am / Sun 8 pm to Tue 8 am |
| District 4                | Two 48-hour periods per week           | Sun 6 pm to Tue 6 pm / Wed 6 pm to Fri 6 pm |
| Koyukuk and Innoko Rivers | 7 days per week                        | M/T/W/TH/F/SA/SU - 24 hours/day             |
| Subdistricts 5-A, -B, -C  | Two 48-hour periods per week           | Tue 6 pm to Thu 6 pm / Fri 6 pm to Sun 6 pm |
| Subdistrict 5-D           | 7 days per week                        | M/T/W/TH/F/SA/SU - 24 hours/day             |
| Subdistrict 6             | Two 42-hour periods per week           | Mon 6 pm to Wed Noon / Fri 6 pm to Sun Noon |
| Old Minto Area            | 5 days per week                        | Friday 6 pm to Wednesday 6 pm               |

*Note:* In the Upper Yukon, fishing times are longer by regulation to help account for longer travel times and lower numbers of fish available as fish leave the mainstem Yukon River to spawn in U.S. tributaries. This schedule was not used in 2021; salmon fishing closed as Chinook salmon moved up river and remained closed all season.

Table 2.—Yukon River drainage fall chum salmon management plan overview, in U.S. waters.

| Run size estimate <sup>b</sup><br>(point estimate) | Recommended management action <sup>a</sup><br>Fall chum salmon directed fisheries |                      |                      |  | Targeted<br>drainagewide<br>escapement |
|--|---|----------------------|----------------------|--|--|
|  | Commercial  | Personal use         | Sport                | Subsistence                              |  |
| 300,000<br>or Less                                 | Closure   | Closure              | Closure              | Closure <sup>c</sup>                     | 300,000<br><br>to<br><br>600,000       |
| 300,001<br>to<br>550,000                           | Closure   | Closure <sup>c</sup> | Closure <sup>c</sup> | Possible<br>Restrictions <sup>c, d</sup> |  |
| Greater than<br>550,001                            | Open <sup>e</sup>   | Open                 | Open                 | No<br>restrictions                       |  |

<sup>a</sup> Considerations for the Canadian mainstem interim management escapement goal may require more restrictive management actions.

<sup>b</sup> Alaska Department of Fish and Game will use the best available data, including preseason projections, mainstem river sonar passage estimates, test fisheries indices, subsistence and commercial fishing reports, and passage estimates from escapement monitoring projects.

<sup>c</sup> The fisheries may be opened or less restrictive in areas where indicator(s) suggest the escapement goal(s) in that area will be achieved.

<sup>d</sup> Subsistence fishing will be managed to achieve a minimum drainagewide escapement goal of 300,000 fall chum salmon.

<sup>e</sup> Drainagewide commercial fisheries may be open and the harvestable surplus above 550,000 fall chum salmon will be distributed by district or subdistrict (in proportion to the guidelines harvest levels established in 5 AAC 05.365 and 5 AAC 05.367).

Table 3.—Inseason fishery management decision matrix for Yukon River mainstem Chinook salmon in Canada, 2021.

| Canada total run size | Border passage projection <sup>a</sup> | Canada allowable harvest (CAH) <sup>b</sup> | Projected escapement <sup>b</sup> | Fishery allocations <sup>c</sup> |                      |                       |
|-----------------------|--|---|-----------------------------------|----------------------------------|----------------------|-----------------------|
|                       |  |   |                                   | First Nation                     | Public angling       | Commercial & domestic |
| 0–42,500              | 0–42,500                               | 0   | 0–42,500                          | 0                                | 0                    | 0                     |
| 42,501–96,848         | 42,501–55,000                          | 1–6,250                                     | 42,500–48,750                     | 1–6,250                          | 0                    | 0                     |
| 96,849–141,196        | 55,001–65,200                          | 6,251–10,200                                | 48,750–55,000                     | 6,251–10,000                     | 0 – 200 <sup>d</sup> | 0                     |
| 141,197–143,804       | 65,201–65,800                          | 10,201–10,800                               | 55,000                            | 10,000                           | 201–800              | 0                     |
| 143,805–150,761       | 65,801–67,400                          | 10,801–12,400                               | 55,000                            | 10,000                           | 801–1,260            | 0–1,140 <sup>d</sup>  |
| 150,762–259,891       | 67,401–92,500                          | 12,401–37,500                               | 55,000                            | 10,000                           | 1,260–2,515          | 1,141–24,985          |
| 259,892–292,500       | 92,501–100,000                         | 37,501–45,000                               | 55,000                            | 10,000                           | 2,515–2,890          | 24,986–32,110         |

<sup>a</sup> Border passage projection is Eagle Sonar estimate plus estimated U.S. harvest between sonar and U.S./Canada border.

<sup>b</sup> Canadian allowable harvest and projected escapement levels may vary within the First Nation fishery depending on the trade-offs between the two; this is influenced by the priority that First Nations may place on escapement or harvest in any given year.

<sup>c</sup> Allocations to fisheries are depicted categories of opportunity, with dark grey representing no fishery opportunities, light grey as limited fishery opportunities, and unshaded as extensive fishery opportunities.

<sup>d</sup> This fishery allocation represents the level of management precision for that fishery and is the threshold required before considering harvest opportunities.

Table 4.—Inseason fishery management decision matrix for mainstem Yukon River fall chum salmon in Canada, 2021.

| International border passage<br>(based on Eagle sonar estimate) | Fishery   |   |   |   |
|---|---|---|---|---|
|   | First Nation  | Public angling  | Commercial  | Domestic  |
| < 40,000<br>(Red Zone)  | <b>Closed</b><br>Removal of allocation for conservation purposes              | <b>Closed</b><br>No retention permitted                               | <b>Closed</b>   | <b>Closed</b>   |
| 40,000 to 73,000<br>(Yellow Zone)                               | <b>Varies <sup>a</sup></b><br>Catch target to vary with abundance within zone | <b>Closed</b><br>No retention permitted                               | <b>Closed</b>   | <b>Closed</b>   |
| > 73,000<br>(Green Zone)  | <b>Open</b><br>Unrestricted   | <b>Open <sup>a</sup></b><br>Retention permitted. No catch anticipated | <b>Open <sup>a</sup></b><br>Allocation varies with run size | <b>Open <sup>a</sup></b><br>Allocation varies with run size |

<sup>a</sup> Allocations (harvest opportunities) are subject to run abundance and international harvest sharing provisions (Yukon River Salmon Agreement).

Table 5.—Cumulative fish passage estimates by species with 90% confidence intervals (CI), at the Pilot Station sonar in 2021.

| Species                                | Total passage | 90% CI  |         |
|--|---------------|---------|---------|
|  |               | Lower   | Upper   |
| Large Chinook <sup>a</sup>             | 104,267       | 93,928  | 114,606 |
| Small Chinook <sup>b</sup>             | 20,578        | 17,346  | 23,810  |
| All Chinook subtotal                   | 124,845       | 114,014 | 135,676 |
| Summer chum                            | 153,718       | 137,569 | 169,867 |
| Fall chum                              | 146,197       | 134,511 | 157,883 |
| Coho                                   | 37,255        | 33,376  | 41,134  |
| Pink                                   | 22,181        | 16,349  | 28,013  |
| Cisco                                  | 195,566       | 170,302 | 220,830 |
| Broad whitefish <i>C. nasus</i>        | 264,160       | 242,969 | 285,351 |
| Humpback whitefish <i>C. pidschian</i> | 23,859        | 19,888  | 27,830  |
| Sheefish <i>Stenodus leucichthys</i>   | 34,820        | 29,446  | 40,194  |
| Other <sup>c</sup>                     | 38,059        | 33,213  | 42,905  |
| Total <sup>d</sup>                     | 1,040,660     |         |         |

<sup>a</sup> Large Chinook salmon >655 mm.

<sup>b</sup> Small Chinook salmon ≤655 mm.

<sup>c</sup> Includes sockeye salmon, burbot *Lota lota*, long nose sucker *Catostomus catostomus*, Dolly Varden, *Salvelinus malma*, and northern pike *Esox lucius*.

<sup>d</sup> All Chinook subtotal not included in total passage sum.

Table 6.—Yukon River Chinook salmon age and female percentage estimated from samples collected at the Pilot Station and Eagle sonar projects, 2021.

| Age/sex | Chinook salmon age and sex composition (percentage of test fishery samples) |       |                                   |       |
|---------|---|-------|-----------------------------------|-------|
|         | Pilot Station sonar   |       | Eagle sonar                       |       |
|         | Historical average<br>(2011–2020)   | 2021  | Historical average<br>(2011–2020) | 2021  |
| Age-4   | 11.1%   | 2.0%  | 6.7%                              | 2.1%  |
| Age-5   | 50.2%   | 47.9% | 41.3%                             | 45.3% |
| Age-6   | 36.2%   | 45.3% | 48.8%                             | 48.6% |
| Female  | 42.8%   | 49.9% | 45.3%                             | 44.3% |

*Note:* Sampling at the Pilot Station sonar uses 6 gillnets that range in mesh sizes from 2.75–8.5 inch whereas sampling at Eagle sonar uses 4 gillnets that range in mesh size from 5.25–8.5 inch. This difference in gillnet mesh sizes can possibly affect the difference in observed age classes. In addition, sex is determined only through visual inspection of external body characteristics at both projects. Sexual dimorphism is more pronounced by the time fish reach Eagle, making sex identification more accurate at that site. These factors need to be considered when comparing between projects. Percent female was calculated using all sampled Chinook salmon including fish that were unable to be aged successfully.

Table 7.—Summary of 2021 Chinook salmon escapement estimates in U.S. (Alaska) tributaries compared to existing escapement goals.

| Location                | Assessment method | Escapement goal (type) | 2021 Chinook salmon |
|-------------------------|-------------------|------------------------|---------------------|
| East Fork Andreafsky    | Weir              | 2,100–4,900 (SEG)      | 1,418               |
| West Fork Andreafsky    | Aerial survey     | 640–1,600 (SEG)        | No Survey           |
| Anvik (drainagewide)    | Aerial survey     | 1,100–1,700 (SEG)      | No Survey           |
| Nulato (forks combined) | Aerial survey     | 940–1,900 (SEG)        | No Survey           |
| Gisasa                  | Weir              | none                   | Not operated        |
| Henshaw                 | Weir              | none                   | 130                 |
| Chena                   | Tower/Sonar       | 2,800–5,700 (BEG)      | 1,417               |
| Salcha                  | Tower/Sonar       | 3,300–6,500 (BEG)      | 2,082               |

*Note:* Biological escapement goal (BEG) and sustainable escapement goal (SEG).

Table 8.—Summary of 2021 summer chum salmon escapement estimates in U.S. (Alaska) compared to existing escapement goals.

| Location             | Assessment method | Escapement goal (type)  | 2021 Summer chum salmon escapement |
|----------------------|-------------------|-------------------------|------------------------------------|
| Yukon (drainagewide) | Sonar             | 500,000–1,200,000 (BEG) | 153,000 <sup>a</sup>               |
| East Fork Andreafsky | Weir              | >40,000 (SEG)           | 2,531                              |
| Anvik                | Sonar             | 350,000–700,000 (BEG)   | 18,819                             |
| Gisasa               | Weir              | none                    | Not operated                       |
| Henshaw              | Weir              | none                    | 3,729                              |
| Chena                | Tower/sonar       | none                    | 578 <sup>b</sup>                   |
| Salcha               | Tower/sonar       | none                    | 2,193 <sup>b</sup>                 |

*Note:* Biological escapement goal (BEG) and sustainable escapement goal (SEG).

<sup>a</sup> A drainagewide summer chum run reconstruction model was developed in 2016 (Hamazaki and Conitz 2015), and the resulting model estimate of escapement for 2021 is presented here.

<sup>b</sup> Incomplete count due to high water events and considered a minimum estimate.

Table 9.—Summary of 2021 fall chum salmon escapement estimates in U.S. (Alaska) compared to existing escapement goals.

| Location                     | Assessment method | Escapement goal (type) | 2021 Fall chum salmon escapement a |
|------------------------------|-------------------|------------------------|------------------------------------|
| Drainagewide                 | Sonar and harvest | 300,000–600,000 (SEG)  | 94,500                             |
| Chandalar River <sup>b</sup> | Sonar             | 85,000–234,000 (SEG)   | 21,162                             |
| Delta River                  | Ground surveys    | 7,000–20,000 (SEG)     | 1,600                              |

*Note:* Sustainable escapement goal (SEG).

<sup>a</sup> Numbers are rounded.

<sup>b</sup> The Chandalar River and North Fork collectively were renamed the Teedriinjik and the Middle Fork was renamed Ch'idriinjik in September of 2015.

Table 10.—Summary of 2021 preliminary fall chum salmon escapement counts to Canada in comparison with existing international interim management escapement goals (IMEG).

| Location                           | Assessment method        | Escapement goal (type) | 2021 Fall chum salmon escapement |
|------------------------------------|--------------------------|------------------------|----------------------------------|
| Fishing Branch River               | Weir & video/sonar count | 22,000–49,000 (IMEG)   | 2,413                            |
| Yukon River Mainstem               | Sonar and harvest        | 70,000–104,000 (IMEG)  | 23,170                           |
| Porcupine River (Canadian portion) | Sonar and harvest        | none                   | 3,486                            |

Table 11.–Pilot Station sonar Chinook salmon passage and Canadian-origin proportion by strata, 2005–2021.

| Year | Strata    | Dates         | Pilot Station passage | Proportion of run | Canadian proportion <sup>a</sup> | Estimated number of Canadian fish |
|------|-----------|---------------|-----------------------|-------------------|----------------------------------|-----------------------------------|
| 2005 | Stratum 1 | 06/04 - 06/17 | 91,136                | 0.35              | 0.60                             | 54,335                            |
|      | Stratum 2 | 06/18 - 07/03 | 119,627               | 0.46              | 0.45                             | 53,533                            |
|      | Stratum 3 | 07/04 - 08/20 | 48,451                | 0.19              | 0.29                             | 14,002                            |
|      | Total     |               | 259,214               | 1.00              | 0.47                             | 121,871                           |
| 2006 | Stratum 1 | 06/07 - 06/24 | 63,374                | 0.28              | 0.44                             | 28,106                            |
|      | Stratum 2 | 06/25 - 07/26 | 165,389               | 0.72              | 0.39                             | 64,312                            |
|      | Total     |               | 228,763               | 1.00              | 0.40                             | 92,417                            |
| 2007 | Stratum 1 | 06/06 - 06/19 | 50,083                | 0.29              | 0.53                             | 26,629                            |
|      | Stratum 2 | 06/20 - 06/30 | 62,907                | 0.37              | 0.37                             | 23,502                            |
|      | Stratum 3 | 07/01 - 08/16 | 57,256                | 0.34              | 0.21                             | 11,772                            |
|      | Total     |               | 170,246               | 1.00              | 0.37                             | 61,903                            |
| 2008 | Stratum 1 | 06/07 - 06/23 | 41,294                | 0.24              | 0.47                             | 19,532                            |
|      | Stratum 2 | 06/24 - 06/29 | 42,554                | 0.24              | 0.33                             | 13,958                            |
|      | Stratum 3 | 06/30 - 08/02 | 90,559                | 0.52              | 0.31                             | 27,711                            |
|      | Total     |               | 174,407               | 1.00              | 0.35                             | 61,201                            |
| 2009 | Stratum 1 | 06/09 - 06/16 | 7,000                 | 0.04              | 0.68                             | 4,750                             |
|      | Stratum 2 | 06/17 - 06/22 | 27,229                | 0.15              | 0.53                             | 14,347                            |
|      | Stratum 3 | 06/23 - 06/29 | 83,866                | 0.47              | 0.41                             | 34,509                            |
|      | Stratum 4 | 06/30 - 07/19 | 59,701                | 0.34              | 0.17                             | 10,265                            |
|      | Total     |               | 177,796               | 1.00              | 0.36                             | 63,871                            |
| 2010 | Stratum 1 | 06/12 - 06/21 | 28,885                | 0.21              | 0.49                             | 14,110                            |
|      | Stratum 2 | 06/22 - 06/27 | 45,306                | 0.33              | 0.50                             | 22,860                            |
|      | Stratum 3 | 06/28 - 09/05 | 63,708                | 0.46              | 0.28                             | 17,891                            |
|      | Total     |               | 137,899               | 1.00              | 0.40                             | 54,861                            |
| 2011 | Stratum 1 | 06/01 - 06/18 | 31,273                | 0.21              | 0.58                             | 18,148                            |
|      | Stratum 2 | 06/19 - 06/27 | 67,686                | 0.45              | 0.36                             | 24,611                            |
|      | Stratum 3 | 06/28 - 08/07 | 49,838                | 0.33              | 0.16                             | 8,034                             |
|      | Total     |               | 148,797               | 1.00              | 0.34                             | 50,792                            |
| 2012 | Stratum 1 | 06/10 - 06/24 | 31,998                | 0.25              | 0.45                             | 14,463                            |
|      | Stratum 2 | 06/25 - 07/02 | 63,648                | 0.50              | 0.47                             | 30,042                            |
|      | Stratum 3 | 07/03 - 07/30 | 31,909                | 0.25              | 0.34                             | 10,753                            |
|      | Total     |               | 127,555               | 1.00              | 0.43                             | 55,258                            |
| 2013 | Stratum 1 | 06/14 - 06/27 | 78,133                | 0.57              | 0.72                             | 56,568                            |
|      | Stratum 2 | 06/28 - 08/02 | 58,672                | 0.43              | 0.26                             | 15,137                            |
|      | Total     |               | 136,805               | 1.00              | 0.52                             | 71,706                            |
| 2014 | Stratum 1 | 06/01 - 06/14 | 45,236                | 0.28              | 0.49                             | 22,347                            |
|      | Stratum 2 | 06/15 - 06/24 | 82,146                | 0.50              | 0.42                             | 34,255                            |
|      | Stratum 3 | 06/25 - 08/04 | 36,513                | 0.22              | 0.18                             | 6,718                             |
|      | Total     |               | 163,895               | 1.00              | 0.39                             | 63,320                            |
| 2015 | Stratum 1 | 05/30 - 06/17 | 30,600                | 0.21              | 0.50                             | 15,178                            |
|      | Stratum 2 | 06/18 - 06/26 | 51,172                | 0.35              | 0.37                             | 18,780                            |
|      | Stratum 3 | 06/27 - 08/17 | 65,087                | 0.44              | 0.33                             | 21,218                            |
|      | Total     |               | 146,859               | 1.00              | 0.38                             | 55,176                            |

-continued-

Table 11.–Page 2 of 2.

| Year  | Strata    | Dates         | Pilot Station<br>passage | Proportion<br>of run | Canadian<br>proportion <sup>a</sup> | Estimated number of<br>Canadian fish |
|---|-----------|---------------|--------------------------|----------------------|-------------------------------------|--------------------------------------|
| 2016  | Stratum 1 | 05/30 - 06/14 | 37,511                   | 0.21                 | 0.52                                | 19,136                               |
|   | Stratum 2 | 06/15 - 06/25 | 86,622                   | 0.49                 | 0.34                                | 29,114                               |
|   | Stratum 3 | 06/26 - 08/24 | 52,765                   | 0.30                 | 0.54                                | 28,282                               |
|   | Total     |               | 176,898                  | 1.00                 | 0.43                                | 76,532                               |
| 2017  | Stratum 1 | 05/31 - 06/13 | 30,088                   | 0.11                 | 0.43                                | 12,857                               |
|   | Stratum 2 | 06/14 - 06/20 | 79,913                   | 0.30                 | 0.49                                | 38,929                               |
|   | Stratum 3 | 06/21 - 06/25 | 69,392                   | 0.26                 | 0.43                                | 30,121                               |
|   | Stratum 4 | 06/26 - 08/11 | 83,621                   | 0.32                 | 0.41                                | 34,008                               |
|   | Total     |               | 263,014                  | 1.00                 | 0.44                                | 115,915                              |
| 2018  | Stratum 1 | 06/02 - 06/13 | 16,275                   | 0.10                 | 0.53                                | 8,621                                |
|   | Stratum 2 | 06/14 - 06/24 | 56,344                   | 0.35                 | 0.47                                | 26,357                               |
|   | Stratum 3 | 06/25 - 07/03 | 57,070                   | 0.35                 | 0.41                                | 23,227                               |
|   | Stratum 4 | 07/04 - 08/05 | 32,209                   | 0.20                 | 0.29                                | 9,402                                |
|   | Total     |               | 161,831                  | 1.00                 | 0.42                                | 67,609                               |
| 2019  | Stratum 1 | 06/02 - 06/23 | 82,035                   | 0.37                 | 0.56                                | 45,637                               |
|   | Stratum 2 | 06/24 - 06/30 | 73,551                   | 0.33                 | 0.42                                | 30,563                               |
|   | Stratum 3 | 07/01 - 08/24 | 64,038                   | 0.29                 | 0.36                                | 22,910                               |
|   | Total     |               | 219,624                  | 1.00                 | 0.45                                | 99,110                               |
| 2020  | Stratum 1 | 06/07-06/22   | 34,551                   | 0.21                 | 0.63                                | 21,891                               |
|   | Stratum 2 | 06/23-06/29   | 64,298                   | 0.40                 | 0.48                                | 30,873                               |
|   | Stratum 3 | 06/30-07/06   | 35,047                   | 0.22                 | 0.44                                | 15,453                               |
|   | Stratum 4 | 07/07-08/17   | 28,356                   | 0.17                 | 0.37                                | 10,468                               |
|   | Total     |               | 162,252                  | 1.00                 | 0.48                                | 78,685                               |
| 2021  | Stratum 1 | 05/31-06/22   | 44,751                   | 0.36                 | 0.62                                | 27,527                               |
|   | Stratum 2 | 06/23-07/06   | 59,173                   | 0.47                 | 0.54                                | 32,065                               |
|   | Stratum 3 | 07/07-08/06   | 20,921                   | 0.17                 | 0.35                                | 7,409                                |
|   | Total     |               | 124,845                  | 1.00                 | 0.54                                | 67,001                               |
| Average annual proportion of Canadian stock |           |               |                          |                      | 0.41                                |                                      |
| Minimum annual proportion of Canadian stock |           |               |                          |                      | 0.34                                |                                      |
| Maximum annual proportion of Canadian stock |           |               |                          |                      | 0.52                                |                                      |

Note: Average, minimum, and maximum values exclude the most recent year data.

<sup>a</sup> Total Canadian proportion is weighted with "Proportion of run".



Table 12.–Genetic baseline (microsatellite, 37 populations) used for stock separation of chum salmon sampled in the Pilot Station sonar drift gillnet test fishery, 2021.

| Stock aggregate name | Populations in baseline   |
|----------------------|---|
| Lower                | Andreafsky, Anvik, California, Chulinak, Clear, Dakli, Kaltag, Nulato, Gisasa, Melozitna, Rodo, Tolstoi |
| Upper Koyukuk+Main   | Henshaw, Jim, Middle Fork Koyukuk, South Fork Koyukuk (early and late run), Tozitna                     |
| Tanana Summer        | Chena, Salcha   |
| Tanana Fall          | Bluff Cabin, Delta, Nenana, Kantishna, Tanana Mainstem, Toklat  |
| Border U.S.          | Big Salt, Black, Chandalar <sup>a</sup> , Sheenjek  |
| Porcupine            | Fishing Branch  |
| Mainstem             | Big Creek, Minto, Pelly, Tatchun  |
| White                | Donjek, Kluane  |
| Teslin               | Teslin  |
| Aggregate name       | Aggregate within aggregate  |
| Summer               | Lower, Middle   |
| Middle               | Upper Koyukuk+Main, Tanana Summer   |
| Fall                 | Tanana Fall, Border U.S., Border Canada, Upper Canada   |
| Fall U.S.            | Tanana Fall, Border U.S.  |
| U.S.                 | Lower, Middle, Tanana Fall, Border U.S.   |
| Border Canada        | Porcupine, Mainstem   |
| Upper Canada         | White, Teslin   |
| Canada               | Border Canada, Upper Canada   |

<sup>a</sup> The Chandalar River and North Fork collectively were renamed the Teedriinjik and the Middle Fork was renamed the Ch'idriinjik in September of 2015.

Table 13.—Preseason Canadian-origin Yukon River Chinook salmon outlooks for 2000–2022 and the observed run sizes for 2000–2021.

| Year | Outlook range <sup>a</sup> |          | Postseason estimate             |
|------|----------------------------|----------|---------------------------------|
|      | Low end                    | High end | Estimated run size <sup>b</sup> |
| 2000 | 91,000                     | 128,000  | 53,000                          |
| 2001 | 37,000                     | 37,000   | 86,000                          |
| 2002 | 49,000                     | 49,000   | 81,000                          |
| 2003 | 62,000                     | 62,000   | 150,000                         |
| 2004 | 69,700                     | 107,200  | 117,000                         |
| 2005 | 117,000                    | 117,000  | 124,000                         |
| 2006 | 93,000                     | 93,000   | 119,000                         |
| 2007 | 74,500                     | 112,900  | 88,000                          |
| 2008 | 80,000                     | 111,000  | 63,000                          |
| 2009 | 60,700                     | 99,800   | 88,000                          |
| 2010 | 77,800                     | 113,000  | 60,000                          |
| 2011 | 65,000                     | 89,000   | 72,000                          |
| 2012 | 54,000                     | 73,000   | 48,000                          |
| 2013 | 49,000                     | 72,000   | 37,000                          |
| 2014 | 32,000                     | 61,000   | 65,000                          |
| 2015 | 59,000                     | 70,000   | 87,000                          |
| 2016 | 65,000                     | 88,000   | 83,000                          |
| 2017 | 73,000                     | 97,000   | 93,000                          |
| 2018 | 71,000                     | 103,000  | 76,000                          |
| 2019 | 69,000                     | 99,000   | 73,000                          |
| 2020 | 59,000                     | 90,000   | 46,000                          |
| 2021 | 42,000                     | 77,000   | 33,000                          |
| 2022 | 41,000                     | 62,000   |                                 |

*Note:* Run sizes are rounded to the nearest 1,000 fish.

<sup>a</sup> The outlook range has been calculated using a variety of different methods. Refer to previous published JTC reports for a full description for a particular year.

<sup>b</sup> Estimated run size is the border passage estimate plus the U.S. and Canada harvest of Canadian-origin Chinook salmon. U.S. harvest estimates are determined using Canadian stock genetic proportion estimates applied to U.S. harvest.

Table 14.–Forecasted 2022 total run size of fall chum salmon based on parent year escapement for each brood year and predicted return per spawner (R/S) rates, Yukon River, 2016–2019.

| Brood year   | Escapement | Estimated production (R/S) | Estimated production | Age | Contribution based on age | Ricker return | Forecasted return |
|--|------------|----------------------------|----------------------|-----|---------------------------|---------------|-------------------|
| 2016   | 832,200    | 0.13                       | 108,186              | 6   | 0.2%                      | 1,473         | 253               |
| 2017   | 1,706,000  | 0.07                       | 119,420              | 5   | 4.4%                      | 28,198        | 4,836             |
| 2018   | 654,300    | 1.34                       | 876,762              | 4   | 91.5%                     | 588,961       | 101,002           |
| 2019   | 527,950    | 1.58                       | 834,161              | 3   | 3.8%                      | 24,763        | 4,247             |
| Total forecasted run size point estimate (unadjusted and adjusted) |            |                            |                      |     |                           | 643,395       | <b>110,337</b>    |
| Total unadjusted forecasted run size, 80% CI:                      |            |                            |                      |     | 534,000                   | to            | 752,000           |
| Total adjusted forecasted run size:                                |            |                            |                      |     | 78,100                    | to            | 148,000           |

*Note:* The 2022 unadjusted forecast was based on previously-established JTC methods. The 2022 forecast point estimate and 80% CI were adjusted down to account for unprecedented poor run sizes in 2020 and 2021. Adjustment to the point estimate was based on the average (17.1%) of observed return to forecast for 2020 and 2021. Adjustment to the 80% CI range is based on the minimum (14.6%) and maximum (19.7%) differences observed in 2020 and 2021. Escapements and forecast range are rounded.

Table 15.–Preseason Yukon River drainagewide fall chum salmon outlooks 1998–2022 and estimated run sizes for 1998–2021.

| Year | Expected run size (preseason) | Estimated run size (postseason) <sup>a</sup> | Performance of preseason outlook (preseason/postseason) |
|------|-------------------------------|--|---|
| 1998 | 880,000                       | 352,000                                      | 2.50  |
| 1999 | 1,197,000                     | 420,000                                      | 2.85  |
| 2000 | 1,137,000                     | 253,000                                      | 4.49  |
| 2001 | 962,000                       | 375,000                                      | 2.57  |
| 2002 | 646,000                       | 428,000                                      | 1.51  |
| 2003 | 647,000                       | 792,000                                      | 0.82  |
| 2004 | 672,000                       | 653,000                                      | 1.03  |
| 2005 | 776,000                       | 2,181,000                                    | 0.36  |
| 2006 | 1,211,000                     | 1,212,000                                    | 1.00  |
| 2007 | 1,106,000                     | 1,161,000                                    | 0.95  |
| 2008 | 1,057,000                     | 857,000                                      | 1.23  |
| 2009 | 791,000                       | 598,000                                      | 1.32  |
| 2010 | 690,000                       | 587,000                                      | 1.18  |
| 2011 | 740,000                       | 1,239,000                                    | 0.60  |
| 2012 | 1,114,000                     | 1,086,000                                    | 1.03  |
| 2013 | 1,029,000                     | 1,212,000                                    | 0.85  |
| 2014 | 932,000                       | 955,000                                      | 0.98  |
| 2015 | 1,060,000                     | 824,000                                      | 1.29  |
| 2016 | 666,000                       | 1,389,000                                    | 0.48  |
| 2017 | 1,560,000                     | 2,288,000                                    | 0.68  |
| 2018 | 1,700,000                     | 1,113,000                                    | 1.53  |
| 2019 | 1,045,000                     | 802,000                                      | 1.30  |
| 2020 | 936,000                       | 194,000                                      | 4.82  |
| 2021 | 652,000                       | 95,000                                       | 6.86  |
| 2022 | 110,000                       |  |   |

*Note:* Run sizes are rounded to the nearest 1,000 fish. The expected run sizes are point estimates (rounded). Ranges were used since 1999 but until 2006 were not always distributed around the point estimate. Starting in 2006, expected run sizes are the midpoint of the outlook range. Refer to previous published JTC reports for a full method description for a particular year.

<sup>a</sup> Postseason estimates are updated annually based on the Bayesian space-state modeling of the drainagewide estimates and may include refined harvest estimates.

Table 16.—Preseason Canadian-origin mainstem Yukon River fall chum salmon outlooks for 1998–2022 and observed run sizes for 1998–2021.

| Year | Expected run size<br>(preseason) | Estimated run size<br>(postseason) | Performance of preseason outlook<br>(preseason/postseason) |
|------|----------------------------------|------------------------------------|--|
| 1998 | 198,000                          | 70,000                             | 2.83   |
| 1999 | 336,000                          | 116,000                            | 2.90   |
| 2000 | 334,000                          | 66,000                             | 5.06   |
| 2001 | 245,000                          | 49,000                             | 5.00   |
| 2002 | 144,000                          | 113,000                            | 1.27   |
| 2003 | 145,000                          | 182,000                            | 0.80   |
| 2004 | 147,000                          | 193,000                            | 0.76   |
| 2005 | 126,000                          | 558,000                            | 0.23   |
| 2006 | 126,000                          | 330,000                            | 0.38   |
| 2007 | 147,000                          | 347,000                            | 0.42   |
| 2008 | 229,000                          | 269,000                            | 0.85   |
| 2009 | 195,000                          | 128,000                            | 1.52   |
| 2010 | 172,000                          | 143,000                            | 1.20   |
| 2011 | 184,000                          | 326,000                            | 0.56   |
| 2012 | 273,000                          | 238,000                            | 1.15   |
| 2013 | 257,000                          | 303,000                            | 0.85   |
| 2014 | 230,000                          | 223,000                            | 1.03   |
| 2015 | 265,000                          | 205,000                            | 1.29   |
| 2016 | 166,000                          | 298,000                            | 0.56   |
| 2017 | 388,000                          | 563,000                            | 0.69   |
| 2018 | 425,000                          | 279,000                            | 1.52   |
| 2019 | 262,000                          | 178,000                            | 1.47   |
| 2020 | 234,000                          | 25,000                             | 9.36   |
| 2021 | 163,000                          | 23,000                             | 7.09   |
| 2022 | 28,000                           |                                    |  |

*Note:* Run sizes are rounded to the nearest 1,000 fish. The 2009 through 2020 preseason expected run sizes are the midpoint of the outlook range. Estimated run sizes are calculated by adding estimated U.S. harvest of Canadian-origin fall chum salmon to the mainstem Yukon River Eagle sonar passage estimate. The proportion of Canadian mainstem fall chum salmon in the total U.S. harvest is assumed to be equal to the proportion of Canadian-origin fall chum salmon in the drainagewide escapement (i.e. 25%).

Table 17.—Preseason Fishing Branch River fall chum salmon outlooks for 1998–2022 and observed run sizes for 1998–2021.

| Year | Expected run size<br>(preseason) | Estimated run size<br>(postseason) <sup>a</sup> | Performance of preseason<br>outlook<br>(preseason/postseason) |
|------|----------------------------------|---|---|
| 1998 | 112,000                          | 25,000  | 4.48  |
| 1999 | 124,000                          | 24,000  | 5.17  |
| 2000 | 150,000                          | 13,000  | 11.54   |
| 2001 | 101,000                          | 33,000  | 3.06  |
| 2002 | 41,000                           | 19,000  | 2.16  |
| 2003 | 29,000                           | 46,000  | 0.63  |
| 2004 | 22,000                           | 32,000  | 0.69  |
| 2005 | 48,000                           | 186,000   | 0.26  |
| 2006 | 54,000                           | 48,000  | 1.13  |
| 2007 | 80,000                           | 50,000  | 1.60  |
| 2008 | 78,000                           | 30,000  | 2.60  |
| 2009 | 49,000                           | 40,000  | 1.23  |
| 2010 | 43,000                           | 20,000  | 2.15  |
| 2011 | 37,000                           | 28,000  | 1.32  |
| 2012 | 55,000                           | 50,000  | 1.10  |
| 2013 | 52,000                           | 39,000 (52,000) <sup>b</sup>                    | —   |
| 2014 | 46,000                           | 13,000 (24,000) <sup>b</sup>                    | —   |
| 2015 | 17,000                           | 13,000  | 1.31  |
| 2016 | 27,000                           | 54,000  | 0.50  |
| 2017 | 62,000                           | 73,000  | 0.85  |
| 2018 | 68,000                           | 29,000  | 2.34  |
| 2019 | 42,000                           | 29,000  | 1.45  |
| 2020 | 37,000                           | 5,000   | 7.40  |
| 2021 | 26,000                           | 2,500   | 10.40   |
| 2022 | 4,000                            |   |   |

Note: Run sizes are rounded to nearest 1,000. The 2009 through 2021 preseason forecasted run sizes are the midpoint of an outlook range. The Fishing Branch River weir monitors the dominant spawning stock within the Porcupine River drainage.

<sup>a</sup> Total run size from 1998 to 2014 is for the Canadian Porcupine River. The total run size is estimated by adding the estimated Canadian harvest of Fishing Branch River fall chum and estimated U.S. harvest of Fishing Branch River fall chum salmon to the Fishing Branch River weir escapement estimate, unless otherwise noted. In 2003, total run size was calculated using the equation;  $((\text{Fishing Branch River escapement}/0.88) + \text{Canadian Porcupine River harvest}) \times 1.15$ . From 2004 to 2009, total run size was calculated using the equation;  $\text{Fishing Branch River escapement}/0.8/0.8$ . In 2010, total run size was calculated using the equation;  $\text{Fishing Branch River escapement}/0.8$ . In 2011, total run size was calculated using the equation;  $(\text{Fishing Branch River escapement} + 75\% \text{ of Canadian Porcupine harvest})/0.68/0.75$ . In 2012, total run size was calculated using the equation;  $\text{Fishing Branch River escapement} \times 1.25 + \text{Canadian Porcupine River harvest} + 5\% \text{ U.S. harvest of fall chum}$ . From 2013 to 2015, the proportion of Fishing Branch River fall chum salmon in the total U.S. harvest is assumed to be equal to the proportion of Fishing Branch River fall chum salmon in the drainagewide escapement. From 2016–2021, the proportion of Fishing Branch River Fall chum in the total U.S. harvest is assumed to be 4%. In 2020, proportion of Fishing Branch-origin fall chum salmon in the total Canadian-origin Porcupine River fall chum salmon harvest was calculated as 63%, estimated by regression of Porcupine sonar to Fishing Branch River weir passage estimates from 2015 to 2019 (excluding an incomplete Porcupine sonar estimate in 2018). For 2016 to 2018, Fishing Branch River proportion within Porcupine River fall chum was considered 80%, based on historical telemetry work. From 2012–2015, 100% of Canadian fall chum salmon harvest in the Porcupine River was included in the Fishing Branch River estimated run size. From 2003 to 2010, 80% of Canadian fall chum salmon harvest in the Porcupine River was included in the total run estimate.

<sup>b</sup> Run size was based on Old Crow sonar counts and proportion of tag recoveries. Numbers in parentheses are the corresponding Canadian-origin Porcupine River sonar-based estimates. Outlook performances are not included due to uncertainty in the assessment methods compared with previous years.




Figure 1.—Map of the Alaska (U.S.) portion of the Yukon River drainage showing communities and fishing districts.




Figure 2.—Primary assessment projects operated in the U.S. and Canada used to assess Chinook and fall chum salmon run strength or escapement.




Figure 3.—Daily (top) and cumulative (bottom) catch per unit effort (CPUE) for Chinook salmon in the Lower Yukon set gillnet test fishery at Big Eddy in 2021.

*Note:* Middle Mouth set gillnet was not operated during 2021. Therefore, 2021 CPUE is not directly comparable to other years with combined south mouth and middle mouth CPUE for set gillnet.






Figure 4.—U.S. (Alaska) harvest of Chinook salmon, Yukon River, 1961–2021.

*Note:* The 2017–2021 harvest estimates are preliminary. Commercial harvests through 2007 were Chinook salmon-directed commercial fishing. Commercial harvests 2008 to present include Chinook salmon incidentally harvested and sold from the chum salmon-directed fisheries. 'Commercial related' refers to the estimated harvest of female Chinook salmon to produce roe sold between 1990 and 2002.




Figure 5.—U.S. (Alaska) harvest of fall chum salmon, Yukon River, 1961–2021.

*Note:* Subsistence harvest estimates of fall chum salmon are minimal prior to 1979 because of timing of harvest surveys. The commercial fishery was closed in 1963, 1987, 1993, 1998, 2000–2002, 2020, and 2021. 'Commercial related' refers to the estimated harvest of female salmon to produce roe sold. The 2017–2021 harvest estimates are preliminary.




Figure 6.—U.S. (Alaska) harvest of coho salmon, Yukon River, 1961–2021.

*Note:* Subsistence harvest estimates of coho salmon are minimal prior to 1979 because of timing of harvest surveys. The commercial fishery was closed 1987, 1993, 1998, 2000–2002, 2020, and 2021. 'Commercial related' refers to the estimated harvest of female salmon to produce roe sold. The 2017–2021 harvest estimates are preliminary.




Figure 7.—Commercial fishing boundaries, tributaries, and major towns within the Yukon Territory, Canada.




Figure 8.—Canadian harvest of Chinook salmon, Yukon River, 1961–2021.




Figure 9.—Estimated spawning escapement estimates and escapement goals (minimum or range) for Canadian-origin Yukon River mainstem Chinook salmon, 1982–2021.




Figure 10.—Chinook salmon ground-based escapement estimates for selected tributaries in the U.S. (Alaska) portion of the Yukon River drainage, 1986–2021.

-continued-




Figure 10.—Page 2 of 2.

*Note:* Escapement goal range relative to years when the goal was in effect. There are no escapement goals at the Henshaw Creek and Gisasa River weirs. Vertical scale is variable.






Figure 11.—Estimated drainagewide escapement of fall chum salmon, Yukon River, 1974–2021.




Figure 12.—Fall chum salmon escapement estimates for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1972–2021.

*Note:* Horizontal lines represent escapement goals or ranges. The vertical scale is variable. Escapement goal is relative to years applied as either minimums or ranges. Sheenjek escapement project was not funded after 2012 and the goal was discontinued in 2016.




Figure 13.—Canadian harvest of fall chum salmon, Yukon River, 1961–2021.




Figure 14.—Estimated spawning escapement and escapement goals (minimum or range) for Canadian-origin fall chum salmon for the mainstem Yukon River and Fishing Branch River, 1972–2021.




Figure 15.—Daily passage estimates of Chinook salmon at the Pilot Station sonar in 2021 (top) and cumulative passage estimate, including 90% confidence intervals (bottom), 2021 compared to historical average.

*Note:* Historical average includes 1995, 1997, 2000, 2002–2008, and 2010–2019.




Figure 16.—Daily passage estimates of chum salmon at the Pilot Station sonar in the fall season in 2021 (top), cumulative passage estimates, including 90% confidence intervals (bottom), compared to median passages.

*Note:* Historical median includes 1995–2020, excluding 1996 and 2009.




Figure 17.—Daily passage estimates of coho salmon at the Pilot Station sonar in 2021 (top), cumulative passage estimates, including 90% confidence intervals (bottom), compared to median passages.

*Note:* Historical median includes 1995–2020, excluding 1996 and 2009.




Figure 18.—Lower Yukon daily water temperatures, comparing 2021 to historical minimum, maximum, and average temperatures.

*Note:* Temperatures were collected in the Yukon River near Emmonak using handheld thermometers (1984–present) and data loggers (2004–present). The years the data types overlap are averaged together.






Figure 19.—Expected versus observed number of Canadian-origin Chinook salmon returning to spawn, 2000–2021.

*Note:* Forecast methods have changed over time and the "expected" value is the published JTC forecast range midpoint. Forecast range error bars are included for years with a published range. The "observed" is estimated Total Canadian-origin run size. This is calculated as the spawning escapement plus estimated U.S. and Canada harvest.




Figure 20.—Yukon River Canadian-origin Chinook salmon recruits versus spawners, Ricker curve (thick line), and 1:1 replacement line (thin line). Brood years 1982–2015 are included.




Figure 21.—Juvenile abundance estimates of Canadian-origin Chinook salmon from the Yukon River based on pelagic trawl research surveys in the northern Bering Sea (2003–2021).

*Note:* Error bars ranges are one deviation above and below the abundance estimates. No survey occurred in 2008 or 2020.




Figure 22.—The relationship between juvenile abundance estimated from surface trawl surveys and adult returns for Canadian-origin Chinook salmon from the Yukon River. Data labels indicate juvenile year, gray shaded area indicates the 80% confidence interval, and black dashed lines indicate the 80% prediction interval. Data include 2003–2017, excluding 2005, 2008, and 2014.




Figure 23.—Historic run size estimates of Canadian-origin Chinook salmon in the Yukon River (gray bars) 1982–2021 and preliminary projected run size for 2022 based on juvenile abundance (gray hashed bar).

*Note:* Error bar range reflects the 80% prediction interval around the 2022 juvenile abundance-based forecast.



## **APPENDIX A: TABLES**

Appendix A1.—Passage estimates from the Pilot Station sonar, 1995 and 1997–2021.

| Year <sup>a</sup> | Chinook            |        |         | Chum      |                   |           | Coho <sup>c</sup> | Pink      | Other <sup>d</sup> | Total     |
|-------------------|--------------------|--------|---------|-----------|-------------------|-----------|-------------------|-----------|--------------------|-----------|
|                   | Large <sup>b</sup> | Small  | Total   | Summer    | Fall <sup>c</sup> | Total     |                   |           |                    |           |
| 1995              | 173,437            | 47,920 | 221,357 | 3,620,102 | 1,148,916         | 4,769,018 | 115,569           | 53,165    | 721,739            | 5,880,848 |
| 1997 <sup>e</sup> | 114,519            | 85,244 | 199,763 | 1,359,117 | 579,767           | 1,938,884 | 118,065           | 3,872     | 376,841            | 2,637,425 |
| 1998              | 88,129             | 19,909 | 108,038 | 824,901   | 375,222           | 1,200,123 | 146,365           | 103,416   | 210,677            | 1,768,619 |
| 1999              | 159,805            | 24,413 | 184,218 | 969,459   | 451,505           | 1,420,964 | 76,174            | 3,947     | 337,701            | 2,023,004 |
| 2000              | 48,321             | 6,239  | 54,560  | 448,665   | 273,206           | 721,871   | 206,365           | 61,389    | 262,627            | 1,306,812 |
| 2001 <sup>f</sup> | 104,060            | 17,029 | 121,089 | 442,546   | 408,961           | 851,507   | 160,272           | 2,846     | 265,749            | 1,401,463 |
| 2002              | 111,290            | 40,423 | 151,713 | 1,097,769 | 367,886           | 1,465,655 | 137,077           | 123,698   | 405,534            | 2,283,677 |
| 2003              | 287,729            | 30,359 | 318,088 | 1,183,009 | 923,540           | 2,106,549 | 280,552           | 11,370    | 379,651            | 3,096,210 |
| 2004              | 138,317            | 62,444 | 200,761 | 1,344,213 | 633,368           | 1,977,581 | 207,844           | 399,339   | 391,939            | 3,177,464 |
| 2005 <sup>g</sup> | 227,154            | 31,861 | 259,015 | 2,570,696 | 1,893,688         | 4,464,384 | 194,372           | 61,091    | 364,250            | 5,343,112 |
| 2006              | 192,296            | 36,467 | 228,763 | 3,780,760 | 964,238           | 4,744,998 | 163,889           | 183,006   | 531,047            | 5,851,703 |
| 2007              | 119,622            | 50,624 | 170,246 | 1,875,491 | 740,195           | 2,615,686 | 192,406           | 126,282   | 761,657            | 3,866,277 |
| 2008              | 138,220            | 36,826 | 175,046 | 1,849,553 | 636,525           | 2,486,078 | 145,378           | 580,127   | 306,225            | 3,692,854 |
| 2009 <sup>h</sup> | 128,154            | 49,642 | 177,796 | 1,477,186 | 274,227           | 1,751,413 | 240,779           | 34,529    | 589,916            | 2,794,433 |
| 2010              | 118,335            | 26,753 | 145,088 | 1,415,027 | 458,103           | 1,873,130 | 177,724           | 917,731   | 567,454            | 3,681,127 |
| 2011              | 117,213            | 31,584 | 148,797 | 2,051,501 | 873,877           | 2,925,378 | 149,533           | 9,754     | 453,537            | 3,686,999 |
| 2012              | 106,529            | 21,026 | 127,555 | 2,136,476 | 778,158           | 2,914,634 | 130,734           | 420,344   | 464,058            | 4,057,325 |
| 2013              | 120,536            | 16,269 | 136,805 | 2,849,683 | 865,295           | 3,714,978 | 110,515           | 6,126     | 732,009            | 4,700,433 |
| 2014              | 120,060            | 43,835 | 163,895 | 2,020,309 | 706,630           | 2,726,939 | 283,421           | 679,126   | 584,831            | 4,438,212 |
| 2015              | 105,063            | 41,796 | 146,859 | 1,591,505 | 669,483           | 2,260,988 | 121,193           | 39,690    | 853,989            | 3,422,719 |
| 2016              | 135,013            | 41,885 | 176,898 | 1,921,748 | 994,760           | 2,916,508 | 168,297           | 1,364,849 | 355,365            | 4,981,917 |
| 2017              | 217,821            | 45,193 | 263,014 | 3,093,735 | 1,829,931         | 4,923,666 | 166,320           | 166,529   | 796,199            | 6,315,728 |
| 2018              | 122,394            | 39,437 | 161,831 | 1,612,688 | 928,664           | 2,541,352 | 136,347           | 689,607   | 547,959            | 4,077,096 |
| 2019              | 172,242            | 47,382 | 219,624 | 1,402,925 | 842,041           | 2,244,966 | 86,401            | 42,353    | 568,576            | 3,161,920 |
| 2020              | 124,905            | 37,347 | 162,252 | 692,602   | 262,439           | 955,041   | 107,680           | 207,942   | 388,287            | 1,821,202 |
| 2021              | 104,267            | 20,578 | 124,845 | 153,718   | 146,197           | 299,915   | 37,255            | 22,181    | 556,464            | 1,040,660 |

-continued-



## Appendix A1.–Page 2 of 2.

---

*Note:* Historical passage estimates at the Pilot Station sonar were adjusted in 2016 after the adoption of a new species apportionment model.

- <sup>a</sup> Estimates for all years were generated with the most current apportionment model and may differ from earlier estimates.
- <sup>b</sup> Chinook salmon >655 mm measured mid eye to tail fork length.
- <sup>c</sup> This estimate may not include the entire run. Most years operated through August 31, except 1995 (September 3), 1998 (September 9), 2000 (September 14), 2008–2014, 2017–2018, and 2021 (September 7).
- <sup>d</sup> Includes sockeye salmon, cisco, whitefish, sheefish, burbot, suckers, Dolly Varden, and northern pike.
- <sup>e</sup> The Yukon River sonar project did not operate at full capacity in 1996 and there are no passage estimates for this year.
- <sup>f</sup> High water levels were experienced on site at Pilot Station in 2001 throughout the season, and passage estimates are considered conservative.
- <sup>g</sup> Estimates include extrapolations for the dates June 10–June 18, 2005 to account for the time before the DIDSON was deployed.
- <sup>h</sup> High water levels were experienced at Pilot Station in 2009 during the summer season and extreme low water occurred during the fall season, and therefore passage estimates are considered conservative.

Appendix A2.—Alaska commercial salmon sales (number of fish) by district and subdistrict, 2021.

| District/Subdistrict             | Number of<br>fishermen <sup>a</sup> | Chinook | Summer<br>chum | Fall chum <sup>b</sup> | Coho <sup>b</sup> | Pink |
|----------------------------------|-------------------------------------|---------|----------------|------------------------|-------------------|------|
| 1                                | —                                   | —       | —              | —                      | —                 | —    |
| 2                                | —                                   | —       | —              | —                      | —                 | —    |
| 3 <sup>b</sup>                   | —                                   | —       | —              | —                      | —                 | —    |
| Total Lower Yukon                | 0                                   | 0       | 0              | 0                      | 0                 | 0    |
| Anvik River                      | —                                   | —       | —              | —                      | —                 | —    |
| 4-A                              | —                                   | —       | —              | —                      | —                 | —    |
| 4-BC                             | —                                   | —       | —              | —                      | —                 | —    |
| Subtotal District 4 <sup>b</sup> | 0                                   | 0       | 0              | 0                      | 0                 | 0    |
| 5-ABC                            | —                                   | —       | —              | —                      | —                 | —    |
| 5-D                              | —                                   | —       | —              | —                      | —                 | —    |
| Subtotal District 5 <sup>b</sup> | 0                                   | 0       | 0              | 0                      | 0                 | 0    |
| 6-ABC <sup>b</sup>               | —                                   | —       | —              | —                      | —                 | —    |
| Total Upper Yukon                | 0                                   | 0       | 0              | 0                      | 0                 | 0    |
| Total Alaska                     | 0                                   | 0       | 0              | 0                      | 0                 | 0    |

*Note:* En dash indicates no commercial fishing activity occurred. Does not include ADF&G test fishery sales.

<sup>a</sup> Number of unique permits fished by district, subdistrict, or area. Totals by area may not add up due to transfers between districts or subdistricts.

<sup>b</sup> Fishery did not operate in 2021.

Appendix A3.—Yukon River Canadian-origin Chinook salmon total run by brood year and escapement by year.

| Brood<br>year     | Age   |        |        |         |        |       | Return  | Spawners | R/S  |
|-------------------|-------|--------|--------|---------|--------|-------|---------|----------|------|
|                   | 3     | 4      | 5      | 6       | 7      | 8     |         |          |      |
| 1974              |       |        |        |         |        | 4,388 |         |          |      |
| 1975              |       |        |        |         | 34,696 | 278   |         |          |      |
| 1976              |       |        |        | 82,801  | 20,859 | 47    |         |          |      |
| 1977              |       |        | 18,964 | 107,561 | 20,000 | 547   | 147,071 |          |      |
| 1978              |       | 5,204  | 28,339 | 63,387  | 32,684 | 793   | 130,406 |          |      |
| 1979              | 1,534 | 3,168  | 21,293 | 99,647  | 44,935 | 1,202 | 171,780 |          |      |
| 1980              | 15    | 6,308  | 10,976 | 78,443  | 30,605 | 4,332 | 130,679 |          |      |
| 1981              | 0     | 1,505  | 29,105 | 124,142 | 65,576 | 1,076 | 221,404 |          |      |
| 1982              | 0     | 5,246  | 13,141 | 32,404  | 27,166 | 171   | 78,128  | 43,538   | 1.79 |
| 1983              | 560   | 4,970  | 32,100 | 86,220  | 13,707 | 108   | 137,665 | 44,475   | 3.10 |
| 1984              | 69    | 11,041 | 37,824 | 81,832  | 20,060 | 192   | 151,018 | 50,005   | 3.02 |
| 1985              | 223   | 11,873 | 36,643 | 59,757  | 4,771  | 64    | 113,331 | 40,435   | 2.80 |
| 1986              | 356   | 18,829 | 42,293 | 114,716 | 16,137 | 138   | 192,470 | 41,425   | 4.65 |
| 1987              | 7     | 2,142  | 27,309 | 69,477  | 15,988 | 18    | 114,941 | 41,307   | 2.78 |
| 1988              | 21    | 6,760  | 35,595 | 83,506  | 12,893 | 68    | 138,844 | 39,699   | 3.50 |
| 1989              | 471   | 10,480 | 68,225 | 126,578 | 31,814 | 0     | 237,568 | 60,299   | 3.94 |
| 1990              | 125   | 4,665  | 22,520 | 56,724  | 4,836  | 9     | 88,880  | 59,212   | 1.50 |
| 1991              | 363   | 7,470  | 89,841 | 126,660 | 11,207 | 0     | 235,540 | 42,728   | 5.51 |
| 1992              | 309   | 4,035  | 24,212 | 39,924  | 2,295  | 0     | 70,775  | 39,155   | 1.81 |
| 1993              | 21    | 5,860  | 34,834 | 84,973  | 7,450  | 477   | 133,615 | 36,244   | 3.69 |
| 1994              | 132   | 2,189  | 20,831 | 27,856  | 8,334  | 0     | 59,341  | 56,449   | 1.05 |
| 1995              | 119   | 2,330  | 15,468 | 48,952  | 10,113 | 10    | 76,991  | 50,673   | 1.52 |
| 1996              | 19    | 2,069  | 23,375 | 43,760  | 11,789 | 2     | 81,013  | 74,060   | 1.09 |
| 1997              | 0     | 4,526  | 22,321 | 94,778  | 6,426  | 14    | 128,065 | 53,821   | 2.38 |
| 1998              | 0     | 5,237  | 41,060 | 80,818  | 6,271  | 0     | 133,386 | 35,497   | 3.76 |
| 1999              | 56    | 2,330  | 25,048 | 73,931  | 1,411  | 0     | 102,775 | 37,184   | 2.76 |
| 2000              | 12    | 4,954  | 40,562 | 49,713  | 1,202  | 0     | 96,443  | 25,870   | 3.73 |
| 2001              | 0     | 2,813  | 63,400 | 51,278  | 2,223  | 0     | 119,713 | 52,564   | 2.28 |
| 2002              | 21    | 4,962  | 29,302 | 20,646  | 227    | 9     | 55,166  | 42,359   | 1.30 |
| 2003              | 0     | 6,118  | 37,202 | 52,067  | 2,261  | 1     | 97,649  | 80,594   | 1.21 |
| 2004              | 0     | 2,531  | 26,680 | 21,938  | 4,763  | 1     | 55,913  | 48,469   | 1.15 |
| 2005              | 9     | 8,232  | 29,477 | 38,855  | 1,755  | 0     | 78,327  | 67,985   | 1.15 |
| 2006              | 15    | 6,009  | 25,248 | 25,697  | 1,567  | 0     | 58,536  | 62,630   | 0.93 |
| 2007              | 47    | 2,858  | 17,737 | 22,193  | 1,694  | 0     | 44,529  | 34,904   | 1.28 |
| 2008              | 1     | 3,131  | 11,091 | 25,750  | 1,853  | 1     | 41,828  | 33,883   | 1.23 |
| 2009              | 173   | 2,325  | 32,868 | 44,942  | 454    | 0     | 80,762  | 65,278   | 1.24 |
| 2010              | 1     | 4,379  | 29,627 | 19,751  | 876    | 0     | 54,634  | 32,014   | 1.71 |
| 2011              | 194   | 10,645 | 52,818 | 42,322  | 1,209  | 1     | 107,188 | 46,307   | 2.31 |
| 2012              | 255   | 9,650  | 44,760 | 31,923  | 858    | 1     | 87,448  | 32,656   | 2.68 |
| 2013              | 92    | 5,116  | 33,631 | 29,713  | 1,453  | 1     | 70,005  | 28,669   | 2.44 |
| 2014              | 115   | 9,566  | 35,089 | 22,475  | 1,315  |       | 68,560  | 63,331   | 1.08 |
| 2015              | 28    | 6,954  | 18,310 | 15,989  |        |       | 41,281  | 82,674   | 0.50 |
| 2016              | 5     | 3,160  | 14,939 |         |        |       |         | 68,798   |      |
| 2017              | 102   | 725    |        |         |        |       |         | 68,315   |      |
| 2018              | 0     |        |        |         |        |       |         | 54,474   |      |
| 2019              |       |        |        |         |        |       |         | 42,052   |      |
| 2020              |       |        |        |         |        |       |         | 30,967   |      |
| 2021              |       |        |        |         |        |       |         | 31,479   |      |
| Average 1982–2014 |       |        |        |         |        |       | 103,518 | 47,506   | 2.31 |
| Contrast          |       |        |        |         |        |       |         | 3.12     |      |

-continued-

*Note:* Spawner data are derived from a 3-area aerial survey index of combined counts from Big Salmon, Little Salmon, and Nisutlin rivers (1982–2001), radiotelemetry (2002–2004), and the mainstem Yukon River sonar at Eagle (2005–2020). Shaded values are preliminary estimates by brood year. Average includes the years with complete brood information through age-7. Ages used were from samples collected at the mainstem sonar test fishery (2007–present) and converted fish wheel data based on a length selectivity method for years 1982–2006 (Hamazaki 2018).

Appendix A4.—Chinook salmon age and sex percentages from selected Yukon River monitoring projects operated in U.S. (Alaska), 2021.

| Location                                     | Sample size |        | Percent by age class |       |       |       |       |       |       | Mean length |
|--|-------------|--------|----------------------|-------|-------|-------|-------|-------|-------|-------------|
|  |             |        | Age-3                | Age-4 | Age-5 | Age-6 | Age-7 | Age-8 | Total |             |
| East Fork Andreafsky River weir <sup>b</sup> | 73          | Male   | 0.0                  | 43.8  | 19.2  | 0.0   | 0.0   | 0.0   | 63.0  | 487         |
|  |             | Female | 0.0                  | 2.7   | 17.8  | 16.4  | 0.0   | 0.0   | 37.0  | 677         |
|  |             | Total  | 0.0                  | 46.6  | 37.0  | 16.4  | 0.0   | 0.0   | 100.0 | 557         |
| Pilot Station test fishery <sup>a</sup>      | 646         | Male   | 0.0                  | 1.7   | 30.8  | 17.2  | 1.1   | 0.2   | 50.9  | 714         |
|  |             | Female | 0.0                  | 0.3   | 17.2  | 28.0  | 3.6   | 0.0   | 49.1  | 769         |
|  |             | Total  | 0.0                  | 2.0   | 48.0  | 45.2  | 4.6   | 0.2   | 100.0 | 741         |
| Henshaw Creek weir                           | 86          | Male   | 0.0                  | 3.5   | 57.0  | 4.7   | 0.0   | 0.0   | 65.1  | 657         |
|  |             | Female | 0.0                  | 0.0   | 20.9  | 14.0  | 0.0   | 0.0   | 34.9  | 759         |
|  |             | Total  | 0.0                  | 3.5   | 77.9  | 18.6  | 0.0   | 0.0   | 100.0 | 694         |
| Eagle test fishery <sup>a</sup>              | 327         | Male   | 0.0                  | 2.1   | 37.9  | 14.1  | 0.9   | 0.0   | 55.0  | 728         |
|  |             | Female | 0.0                  | 0.0   | 7.3   | 34.6  | 3.1   | 0.0   | 45.0  | 806         |
|  |             | Total  | 0.0                  | 2.1   | 45.3  | 48.6  | 4.0   | 0.0   | 100.0 | 763         |
| Salcha River tower <sup>b,c</sup>            | 108         | Male   | 0.0                  | 9.3   | 36.1  | 8.3   | 0.0   | 0.0   | 53.7  | 684         |
|  |             | Female | 0.0                  | 0.0   | 20.4  | 25.9  | 0.0   | 0.0   | 46.3  | 781         |
|  |             | Total  | 0.0                  | 9.3   | 56.5  | 34.3  | 0.0   | 0.0   | 100.0 | 731         |
| Chena River tower <sup>b,c</sup>             | 32          | Male   | 0.0                  | 15.6  | 43.8  | 0.0   | 0.0   | 0.0   | 59.4  | 693         |
|  |             | Female | 0.0                  | 0.0   | 25.0  | 15.6  | 0.0   | 0.0   | 40.6  | 769         |
|  |             | Total  | 0.0                  | 15.6  | 68.8  | 15.6  | 0.0   | 0.0   | 100.0 | 728         |

*Note:* Length is measured mid eye to the fork of tail to the nearest millimeter. Male and female percentages are based on the subset of aged samples and may differ from estimates based on all samples.

<sup>a</sup> Samples were from test fishing with drift gillnets.

<sup>b</sup> Sample size was below established sample size goal.

<sup>c</sup> Carcass samples collected throughout the spawning grounds upriver from the tower project.

Appendix A5.—Yukon River Chinook salmon age, female percentage, and mean length from Eagle sonar project, 2005–2021.

| Year                        | Sample size | Percent by age class |       |       |       |       | Percent female | Mean length |
|-----------------------------|-------------|----------------------|-------|-------|-------|-------|----------------|-------------|
|                             |             | Age-3                | Age-4 | Age-5 | Age-6 | Age-7 |                |             |
| 2005                        | 171         | 0.0                  | 8.2   | 50.3  | 38.0  | 3.5   | 33.9           | 779         |
| 2006                        | 256         | 0.0                  | 16.8  | 60.2  | 22.7  | 0.4   | 37.9           | 737         |
| 2007                        | 389         | 0.0                  | 5.7   | 40.1  | 53.7  | 0.5   | 43.4           | 787         |
| 2008                        | 375         | 0.0                  | 2.7   | 56.3  | 36.5  | 4.5   | 36.8           | 780         |
| 2009                        | 647         | 0.0                  | 7.7   | 33.2  | 59.0  | 0.0   | 39.6           | 791         |
| 2010                        | 336         | 0.0                  | 7.4   | 46.4  | 42.0  | 4.2   | 40.5           | 770         |
| 2011                        | 419         | 0.0                  | 2.1   | 29.6  | 60.4  | 7.9   | 51.3           | 809         |
| 2012                        | 246         | 0.4                  | 6.1   | 29.7  | 59.3  | 4.5   | 49.6           | 780         |
| 2013                        | 265         | 0.0                  | 4.2   | 27.5  | 63.4  | 4.9   | 51.7           | 807         |
| 2014                        | 606         | 0.2                  | 6.6   | 50.5  | 40.1  | 2.6   | 35.1           | 763         |
| 2015                        | 926         | 0.3                  | 10.8  | 34.3  | 52.4  | 2.2   | 42.1           | 776         |
| 2016                        | 666         | 0.0                  | 9.2   | 65.0  | 25.2  | 0.6   | 32.4           | 759         |
| 2017                        | 719         | 0.1                  | 4.2   | 46.5  | 48.1  | 1.1   | 50.9           | 797         |
| 2018                        | 700         | 0.0                  | 10.3  | 43.0  | 45.0  | 1.7   | 43.4           | 769         |
| 2019                        | 554         | 0.0                  | 8.5   | 48.4  | 41.9  | 1.3   | 47.8           | 772         |
| 2020                        | 513         | 0.2                  | 5.2   | 38.4  | 52.9  | 3.3   | 56.0           | 777         |
| 2021                        | 327         | 0.0                  | 2.1   | 45.3  | 48.6  | 4.0   | 45.0           | 763         |
| Average<br>(2005–2020)      | 487         | 0                    | 7     | 44    | 46    | 3     | 43             | 778         |
| 5-yr Average<br>(2016–2020) | 630         | 0                    | 7     | 48    | 43    | 2     | 46             | 775         |

*Note:* Length is measured mid eye to the fork of tail to the nearest millimeter. Age nomenclature (years in freshwater “.” years at sea). Slight modifications have been made to the drift gillnet mesh sizes used at the Eagle sonar during the first three years of operation (2005–2007); however, mesh sizes measuring 5.25, 6.5, 7.5, and 8.5-inch have been used consistently since 2007. Small fish may be underrepresented in the samples due to not fishing gillnets smaller than 5.25-inch mesh.

Appendix A6.—Yukon River Chinook salmon harvest percentage by stock group for U.S. harvest, U.S. and Canada harvest combined, and the percentage of the upper stock group harvest by each country, 1981–2021.

| Year              | Stock groups (U.S. harvest) |        |       | Stock groups (U.S. and Canada harvest) |        |       | Upper stock group |        |
|-------------------|-----------------------------|--------|-------|--|--------|-------|-------------------|--------|
|                   | Lower                       | Middle | Upper | Lower                                  | Middle | Upper | U.S.              | Canada |
| 1981              | 5.9                         | 59.8   | 34.3  | 5.4                                    | 54.5   | 40.1  | 78.1              | 21.9   |
| 1982              | 15.4                        | 27.5   | 57.1  | 13.9                                   | 24.7   | 61.4  | 83.5              | 16.5   |
| 1983              | 14.2                        | 37.0   | 48.9  | 12.9                                   | 33.7   | 53.3  | 83.7              | 16.3   |
| 1984              | 28.0                        | 44.3   | 27.7  | 25.3                                   | 40.2   | 34.5  | 72.7              | 27.3   |
| 1985              | 30.4                        | 24.6   | 45.1  | 27.6                                   | 22.3   | 50.1  | 81.6              | 18.4   |
| 1986              | 22.3                        | 10.9   | 66.8  | 19.5                                   | 9.6    | 70.9  | 82.7              | 17.3   |
| 1987              | 17.4                        | 21.4   | 61.2  | 15.9                                   | 19.6   | 64.5  | 86.7              | 13.3   |
| 1988              | 24.9                        | 18.1   | 57.0  | 21.8                                   | 15.8   | 62.5  | 79.8              | 20.2   |
| 1989              | 27.2                        | 17.7   | 55.1  | 24.4                                   | 15.9   | 59.7  | 82.9              | 17.1   |
| 1990              | 22.8                        | 28.4   | 48.8  | 20.2                                   | 25.2   | 54.7  | 79.2              | 20.8   |
| 1991              | 31.8                        | 28.7   | 39.6  | 28.0                                   | 25.3   | 46.7  | 74.8              | 25.2   |
| 1992              | 18.0                        | 24.1   | 57.8  | 16.3                                   | 21.8   | 61.9  | 84.5              | 15.5   |
| 1993              | 23.7                        | 28.0   | 48.3  | 21.5                                   | 25.4   | 53.1  | 82.6              | 17.4   |
| 1994              | 20.4                        | 24.1   | 55.5  | 18.2                                   | 21.4   | 60.4  | 81.8              | 18.2   |
| 1995              | 20.0                        | 25.0   | 55.0  | 17.9                                   | 22.4   | 59.7  | 82.4              | 17.6   |
| 1996              | 24.0                        | 11.8   | 64.2  | 21.0                                   | 10.4   | 68.6  | 81.9              | 18.1   |
| 1997              | 28.9                        | 18.3   | 52.8  | 26.4                                   | 16.8   | 56.9  | 84.8              | 15.2   |
| 1998              | 34.7                        | 18.5   | 46.8  | 32.7                                   | 17.4   | 49.8  | 88.8              | 11.2   |
| 1999              | 44.1                        | 6.9    | 49.0  | 40.1                                   | 6.3    | 53.6  | 83.0              | 17.0   |
| 2000              | 37.5                        | 13.6   | 48.9  | 33.9                                   | 12.3   | 53.8  | 81.9              | 18.1   |
| 2001              | 37.5                        | 19.0   | 43.5  | 31.6                                   | 16.0   | 52.4  | 69.8              | 30.3   |
| 2002              | 22.1                        | 33.3   | 44.6  | 19.4                                   | 29.2   | 51.4  | 76.3              | 23.5   |
| 2003              | 7.5                         | 31.7   | 60.8  | 6.8                                    | 28.9   | 64.3  | 86.2              | 13.8   |
| 2004              | 16.9                        | 31.6   | 51.5  | 15.3                                   | 28.8   | 55.9  | 83.7              | 16.3   |
| 2005              | 23.4                        | 24.2   | 52.4  | 20.7                                   | 21.4   | 57.9  | 80.1              | 19.9   |
| 2006              | 19.2                        | 30.2   | 50.5  | 17.6                                   | 27.6   | 54.9  | 84.1              | 15.9   |
| 2007              | 13.7                        | 32.3   | 54.0  | 13.0                                   | 30.6   | 56.4  | 90.5              | 9.5    |
| 2008              | 18.2                        | 30.0   | 51.8  | 17.0                                   | 28.0   | 55.0  | 88.1              | 11.9   |
| 2009              | 12.7                        | 35.8   | 51.6  | 11.1                                   | 31.4   | 57.5  | 78.8              | 21.2   |
| 2010              | 18.7                        | 34.3   | 47.0  | 17.8                                   | 32.7   | 49.5  | 90.5              | 9.5    |
| 2011              | 15.6                        | 33.3   | 51.1  | 13.9                                   | 29.8   | 56.3  | 81.0              | 19.0   |
| 2012              | 14.4                        | 37.5   | 48.2  | 13.3                                   | 34.8   | 51.9  | 86.3              | 13.7   |
| 2013              | 16.0                        | 25.0   | 59.0  | 13.4                                   | 21.0   | 65.6  | 75.5              | 24.5   |
| 2014              | 29.8                        | 26.0   | 44.3  | 25.4                                   | 27.8   | 46.8  | 93.4              | 6.6    |
| 2015              | 15.6                        | 36.3   | 48.1  | 13.5                                   | 31.3   | 55.2  | 75.2              | 24.8   |
| 2016              | 15.1                        | 33.5   | 51.5  | 13.3                                   | 29.5   | 57.2  | 80.4              | 19.6   |
| 2017              | 9.3                         | 35.0   | 55.6  | 8.5                                    | 32.1   | 59.3  | 85.9              | 14.1   |
| 2018              | 8.6                         | 31.8   | 59.6  | 7.9                                    | 29.2   | 62.9  | 87.2              | 12.8   |
| 2019              | 14.0                        | 32.3   | 53.7  | 13.3                                   | 30.6   | 56.1  | 91.0              | 9.0    |
| 2020 <sup>a</sup> | 11.1                        | 35.5   | 53.4  | 10.0                                   | 32.1   | 57.8  | 83.7              | 16.3   |
| 2021 <sup>a</sup> | 14.6                        | 23.0   | 62.4  | 12.6                                   | 19.9   | 67.5  | 79.9              | 20.1   |
| Average           |                             |        |       |  |        |       |                   |        |
| 2011–2020         | 14.9                        | 32.6   | 52.5  | 13.2                                   | 29.8   | 56.9  | 84.0              | 16.0   |
| 2016–2020         | 11.6                        | 33.6   | 54.8  | 10.6                                   | 30.7   | 58.7  | 85.7              | 14.3   |
| Minimum           | 5.9                         | 6.9    | 27.7  | 5.4                                    | 6.3    | 34.5  | 69.8              | 6.6    |
| Maximum           | 44.1                        | 59.8   | 66.8  | 40.1                                   | 54.5   | 70.9  | 93.4              | 30.3   |

*Note:* The Lower and Middle stock groups are composed of tributary populations in the Alaska portion of the Yukon River drainage. The Upper stock group is composed of tributary populations in Canada. U.S. fisheries harvest all stock groups, while Canadian fisheries only harvest the Upper (Canadian) stock. Stock composition of U.S. harvest has been estimated annually from dedicated harvest sampling programs. Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Data are preliminary.



Appendix A7.—Stock group percentage by major stock and by country, from chum salmon beginning July 19 at the Pilot Station sonar, Yukon River, 1999–2021.

| Year <sup>a</sup> | Season stock groups |      | U.S. stock groups |                          | Fall stock country groups |        |
|-------------------|---------------------|------|-------------------|--------------------------|---------------------------|--------|
|                   | Summer              | Fall | Tanana fall       | Border U.S. <sup>b</sup> | Fall U.S.                 | Canada |
| 1999              | 16.2                | 83.8 | —                 | —                        | —                         | —      |
| 2000              | 12.0                | 88.0 | —                 | —                        | —                         | —      |
| 2001              | 13.3                | 86.7 | —                 | —                        | —                         | —      |
| 2002              | 19.2                | 80.8 | —                 | —                        | —                         | —      |
| 2003              | —                   | —    | —                 | —                        | —                         | —      |
| 2004              | 13.6                | 86.4 | 31.5              | 27.4                     | 58.8                      | 27.6   |
| 2005              | 11.2                | 88.8 | 20.6              | 42.7                     | 63.3                      | 25.5   |
| 2006              | 18.2                | 81.8 | 16.8              | 36.1                     | 52.9                      | 28.9   |
| 2007              | 21.2                | 78.8 | 22.9              | 25.7                     | 48.6                      | 30.2   |
| 2008              | 16.2                | 83.8 | 21.8              | 31.2                     | 53.1                      | 30.8   |
| 2009              | 24.4                | 75.6 | 19.4              | 30.0                     | 49.4                      | 26.2   |
| 2010              | 24.9                | 75.1 | 24.2              | 19.6                     | 43.8                      | 31.3   |
| 2011              | 13.7                | 86.3 | 13.3              | 38.4                     | 51.7                      | 34.5   |
| 2012              | 20.0                | 80.0 | 25.9              | 31.8                     | 57.8                      | 22.2   |
| 2013              | 11.2                | 88.8 | 33.1              | 23.7                     | 56.7                      | 32.1   |
| 2014              | 9.7                 | 90.3 | 28.7              | 32.2                     | 60.9                      | 29.4   |
| 2015              | 22.7                | 77.3 | 22.0              | 28.8                     | 50.8                      | 26.4   |
| 2016              | 20.1                | 79.9 | 23.5              | 28.9                     | 52.5                      | 27.4   |
| 2017              | 11.9                | 88.1 | 32.5              | 33.2                     | 65.6                      | 22.4   |
| 2018              | 17.3                | 82.7 | 35.1              | 22.9                     | 58.0                      | 24.7   |
| 2019              | 34.8                | 65.2 | 24.3              | 19.8                     | 44.2                      | 21.0   |
| 2020              | 30.0                | 70.0 | 30.8              | 22.9                     | 53.7                      | 16.4   |
| 2021              | 31.0                | 69.0 | 29.1              | 27.8                     | 56.9                      | 12.1   |
| Average           |                     |      |                   |                          |                           |        |
| 2006–2020         | 19.8                | 80.2 | 25.0              | 28.3                     | 53.3                      | 26.9   |
| 2016–2020         | 22.8                | 77.2 | 29.3              | 25.5                     | 54.8                      | 22.4   |
| Minimum           | 9.7                 | 65.2 | 13.3              | 19.6                     | 43.8                      | 16.4   |
| Maximum           | 34.8                | 90.3 | 35.1              | 42.7                     | 65.6                      | 34.5   |

*Note:* July 19 is the date when U.S. management switches from a focus on summer chum to fall chum salmon in this section of the river. Minimum and maximum values exclude the most recent year data. En dash indicates no analysis is available.

<sup>a</sup> Stock identification methods from 1999 through 2002 were based on allozyme analysis. No samples were collected in 2003. Beginning in 2004, analysis was based on microsatellite baseline. Min Max calculations also do not include 1999–2002.

<sup>b</sup> Border U.S. stocks include Big Salt, Teedriinjik (Chandalar), Sheenjek and Draanjik (Black) Rivers.

Appendix A8.—Drainagewide Yukon River fall chum salmon estimated brood year production and return per spawner estimates 1974–2021.

| Brood<br>year | Number of salmon by age <sup>a</sup> |                      |                      |                     | Return    | Spawners <sup>b</sup> | Return/<br>spawner |
|---------------|--------------------------------------|----------------------|----------------------|---------------------|-----------|-----------------------|--------------------|
|               | Age-3                                | Age-4                | Age-5                | Age-6               |           |                       |                    |
| 1974          | 112,017                              | 654,046              | 96,746               | 0                   | 862,809   | 685,200               | 1.26               |
| 1975          | 197,691                              | 1,725,889            | 67,333               | 0                   | 1,990,914 | 2,220,000             | 0.90               |
| 1976          | 143,742                              | 644,242              | 138,736              | 4,889               | 931,609   | 557,600               | 1.67               |
| 1977          | 112,580                              | 1,082,886            | 196,160              | 6,351               | 1,397,976 | 727,500               | 1.92               |
| 1978          | 22,321                               | 374,987              | 106,866              | 0                   | 504,173   | 557,400               | 0.90               |
| 1979          | 45,040                               | 906,515              | 310,715              | 4,233               | 1,266,504 | 1,351,000             | 0.94               |
| 1980          | 13,634                               | 411,169              | 200,180              | 2,852               | 627,834   | 335,850               | 1.87               |
| 1981          | 51,788                               | 997,034              | 339,584              | 8,934               | 1,397,340 | 560,450               | 2.49               |
| 1982          | 12,434                               | 495,669              | 173,136              | 782                 | 682,021   | 247,900               | 2.75               |
| 1983          | 15,223                               | 935,414              | 233,352              | 4,040               | 1,188,029 | 508,350               | 2.34               |
| 1984          | 6,581                                | 427,316              | 162,759              | 9,142               | 605,797   | 361,350               | 1.68               |
| 1985          | 47,598                               | 917,968              | 305,462              | 2,604               | 1,273,632 | 698,400               | 1.82               |
| 1986          | 1,454                                | 524,145              | 340,461              | 5,702               | 871,763   | 535,300               | 1.63               |
| 1987          | 12,165                               | 677,093              | 347,344              | 7,733               | 1,044,335 | 717,700               | 1.46               |
| 1988          | 12,138                               | 212,320              | 161,775              | 33,287 <sup>c</sup> | 419,520   | 353,100               | 1.19               |
| 1989          | 3,286                                | 303,344              | 410,542 <sup>c</sup> | 20,898              | 738,069   | 540,900               | 1.36               |
| 1990          | 683                                  | 665,743 <sup>c</sup> | 455,593              | 33,287              | 1,155,306 | 498,650               | 2.32               |
| 1991          | 0 <sup>c</sup>                       | 1,127,210            | 398,358              | 13,019              | 1,538,588 | 593,200               | 2.59               |
| 1992          | 7,834                                | 699,580              | 207,567              | 4,124               | 919,104   | 419,600               | 2.19               |
| 1993          | 9,889                                | 482,144              | 107,945              | 3,258               | 603,236   | 382,400               | 1.58               |
| 1994          | 4,550                                | 237,392              | 149,212              | 2,529 <sup>c</sup>  | 393,684   | 940,000               | 0.42               |
| 1995          | 2,496                                | 266,589              | 73,353 <sup>c</sup>  | 420                 | 342,859   | 1,150,000             | 0.30               |
| 1996          | 420                                  | 174,530 <sup>c</sup> | 130,130              | 8,369               | 313,449   | 879,800               | 0.36               |
| 1997          | 2,529 <sup>c</sup>                   | 243,894              | 119,474              | 3,632               | 369,530   | 537,200               | 0.69               |
| 1998          | 440                                  | 270,880              | 59,802               | 6,308               | 337,430   | 281,100               | 1.20               |
| 1999          | 29,245                               | 719,543              | 195,655              | 17,176              | 961,620   | 288,100               | 3.34               |
| 2000          | 9,048                                | 320,241              | 114,194              | 0                   | 443,483   | 224,300               | 1.98               |
| 2001          | 131,012                              | 2,049,118            | 718,937              | 34,751              | 2,933,817 | 329,300               | 8.91               |
| 2002          | 0                                    | 464,740              | 250,284              | 15,218              | 730,242   | 400,200               | 1.82               |
| 2003          | 27,597                               | 875,066              | 477,379              | 17,995              | 1,398,037 | 712,800               | 1.96               |
| 2004          | 0                                    | 362,236              | 155,305              | 2,524               | 520,066   | 576,600               | 0.90               |
| 2005          | 2,435                                | 398,145              | 92,321               | 3,893               | 496,794   | 1,890,000             | 0.26               |
| 2006          | 26,832                               | 397,089              | 359,551              | 30,530 <sup>d</sup> | 814,002   | 940,600               | 0.87               |

-continued-

Appendix A8.–Page 2 of 2.

| Brood<br>year     | Number of salmon by age <sup>a</sup> |                      |                      |                  | Return    | Spawners <sup>b</sup> | Return/<br>spawner |
|-------------------|--------------------------------------|----------------------|----------------------|------------------|-----------|-----------------------|--------------------|
|                   | 3                                    | 4                    | 5                    | 6                |           |                       |                    |
| 2007              | 95,157                               | 862,242              | 188,603 <sup>d</sup> | 9,065            | 1,155,067 | 954,200               | 1.21               |
| 2008              | 12,406                               | 854,621 <sup>d</sup> | 414,560              | 9,476            | 1,291,064 | 638,900               | 2.02               |
| 2009              | 11,945 <sup>d</sup>                  | 785,988              | 426,012              | 22,616           | 1,246,561 | 504,800               | 2.47               |
| 2010              | 2,296                                | 496,329              | 245,677              | 9,166            | 753,467   | 506,900               | 1.49               |
| 2011              | 22,952                               | 486,301              | 181,968              | 1,775            | 692,997   | 910,400               | 0.76               |
| 2012              | 69,059                               | 1,168,828            | 328,388              | 5,644            | 1,571,918 | 689,100               | 2.28               |
| 2013              | 29,099                               | 1,901,133            | 318,686              | 3,232            | 2,252,151 | 853,800               | 2.64               |
| 2014              | 57,087                               | 758,789              | 126,619              | 2,483            | 944,977   | 741,300               | 1.27               |
| 2015              | 29,716                               | 663,718              | 86,658               | 430 <sup>d</sup> | 780,522   | 541,000               | 1.44               |
| 2016              | 8,045                                | 89,521               | 6,437 <sup>d</sup>   | 1,436            | 105,439   | 832,200               | 0.13 <sup>e</sup>  |
| 2017              | 5,571                                | 87,443 <sup>d</sup>  | 29,989               |                  | 123,003   | 1,706,000             | 0.07 <sup>e</sup>  |
| 2018              | 940 <sup>d</sup>                     |                      |                      |                  |           | 654,300               |                    |
| 2019              |                                      |                      |                      |                  |           | 528,000               |                    |
| 2020              |                                      |                      |                      |                  |           | 178,400               |                    |
| 2021              |                                      |                      |                      |                  |           | 94,500                |                    |
| Average 1974–2015 |                                      |                      |                      |                  | 970,531   | 674,815               | 1.75               |
| Minimum           |                                      |                      |                      |                  | 313,449   | 224,300               | 0.26               |
| Maximum           |                                      |                      |                      |                  | 2,933,817 | 2,220,000             | 8.91               |

*Note:* Spawner data are derived from Bayesian spawner-recruit model 1974–2021. Average includes the years with complete brood information through age-6. Minimums and maximum indicate the lowest and highest values for each year presented through 2015.

<sup>a</sup> Age composition is based on samples from the Lower Yukon test fishery gillnets, weighted by test fish catch per unit effort. Prior to 1983 commercial sampling was used to supplement test fishery age samples.

<sup>b</sup> Contrast in escapement data is 9.90. Values are rounded to the nearest 100.

<sup>c</sup> Based upon expanded test fish age composition estimates for years in which the test fishery terminated early both in 1994 and 2000.

<sup>d</sup> Combination of Mt. Village test fishery weighted ages with Lower Yukon test fishery to bolster sample sizes.

<sup>e</sup> Return per spawner includes preliminary estimates from incomplete brood year (shaded value).

Appendix A9.–Escapement, rebuilding and interim goals for Canadian-origin Chinook and fall chum salmon stocks, 1985–2022.

| Year              | Canadian origin stock targets |                             |                            |
|-------------------|-------------------------------|-----------------------------|----------------------------|
|                   | Chinook salmon                | Fall chum salmon            |                            |
|                   | Mainstem                      | Mainstem                    | Fishing Branch River       |
| 1985              | 33,000-43,000                 |                             |                            |
| 1986              | 33,000-43,000                 |                             |                            |
| 1987              | 33,000-43,000                 | 90,000-135,000              | 50,000-120,000             |
| 1988              | 33,000-43,000                 | 90,000-135,000              | 50,000-120,000             |
| 1989              | 33,000-43,000                 | 90,000-135,000              | 50,000-120,000             |
| 1990              | 18,000                        | 80,000                      | 50,000-120,000             |
| 1991              | 18,000                        | 80,000                      | 50,000-120,000             |
| 1992              | 18,000                        | 51,000                      | 50,000-120,000             |
| 1993              | 18,000                        | 51,000                      | 50,000-120,000             |
| 1994              | 18,000                        | 61,000                      | 50,000-120,000             |
| 1995              | 18,000                        | 80,000                      | 50,000-120,000             |
| 1996              | 28,000                        | 65,000                      | 50,000-120,000             |
| 1997              | 28,000                        | 49,000                      | 50,000-120,000             |
| 1998              | 28,000                        | 80,000                      | 50,000-120,000             |
| 1999              | 28,000                        | 80,000                      | 50,000-120,000             |
| 2000              | 28,000                        | 80,000                      | 50,000-120,000             |
| 2001              | 28,000                        | 80,000                      | 50,000-120,000             |
| 2002              | 28,000                        | 60,000                      | 50,000-120,000             |
| 2003 <sup>a</sup> | 28,000 <sup>b</sup>           | 65,000                      | 15,000                     |
| 2004              | 28,000                        | 65,000                      | 13,000                     |
| 2005              | 28,000                        | 65,000                      | 24,000                     |
| 2006              | 28,000                        | 80,000                      | 28,000                     |
| 2007              | 33,000-43,000                 | 80,000                      | 34,000                     |
| 2008              | 45,000 <sup>c</sup>           | 80,000                      | 22,000-49,000 <sup>d</sup> |
| 2009              | 45,000                        | 80,000                      | 22,000-49,000              |
| 2010              | 42,500-55,000 <sup>e</sup>    | 70,000-104,000 <sup>f</sup> | 22,000-49,000              |
| 2011              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2012              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2013              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2014              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2015              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2016              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2017              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2018              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2019              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2020              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2021              | 42,500-55,000                 | 70,000-104,000              | 22,000-49,000              |
| 2022              | —                             | 70,000-104,000              | 22,000-49,000              |

-continued-

*Note:* As per the Yukon River Salmon Agreement (YRSA), the Yukon River Panel (YRP) may recommend that both parties manage the current year salmon run to achieve annual stabilization/rebuilding/interim spawning escapement goals that differ from the escapement goals outlined in Appendix 1 and 2 of the YRSA. The goals shown in this table document what both parties managed to achieve in each year, based on recommendations by the YRP. All single numbers are considered minimums. The Chinook salmon mainstem escapement objective for the 2022 season has yet to be confirmed.

- <sup>a</sup> Treaty was signed by governments in December 2002.
- <sup>b</sup> In 2003, the Chinook salmon goal was set at 25,000 fish. However, if the U.S. conducted a commercial fishery the goal would be increased to 28,000 fish.
- <sup>c</sup> Interim management escapement goal (IMEG) assessed using sonar near Eagle (previous years were measured by mark–recapture abundance estimates).
- <sup>d</sup> Interim Management Escapement Goal (IMEG) established for 2008–2010, based on percentile method.
- <sup>e</sup> IMEG of 42,500 to 55,000 fish recommended in 2010, based on levels selected from several unpublished analyses.
- <sup>f</sup> IMEG established in 2010 based on brood table of Canadian-origin mainstem stocks (1982–2003).

Appendix A10.–Fall chum salmon age and sex percentages with average lengths from selected Yukon River monitoring projects, 2021.

| Location                                     | Sample size |         | Age |      |      |     |     | Total | Mean length |
|--|-------------|---------|-----|------|------|-----|-----|-------|-------------|
|  |             |         | 3   | 4    | 5    | 6   | 7   |       |             |
| Emmonak, Alaska <sup>a</sup>                 | 134         | Males   | 1.5 | 38.1 | 5.2  | 0.7 | 0.0 | 45.5  | 566         |
|  |             | Females | 0.7 | 49.3 | 4.5  | 0.0 | 0.0 | 54.5  | 561         |
|  |             | Total   | 2.2 | 87.4 | 9.7  | 0.7 | 0.0 | 100.0 | 563         |
| Mt. Village, Alaska <sup>a</sup>             | 123         | Males   | 0.8 | 45.5 | 1.6  | 0.0 | 0.0 | 47.9  | 565         |
|  |             | Females | 0.0 | 48.8 | 3.3  | 0.0 | 0.0 | 52.1  | 564         |
|  |             | Total   | 0.8 | 94.3 | 4.9  | 0.0 | 0.0 | 100.0 | 564         |
| Delta River, Alaska <sup>b</sup>             | 160         | Males   | 6.9 | 52.5 | 9.4  | 0.0 | 0.0 | 68.8  | 563         |
|  |             | Females | 1.2 | 23.8 | 6.2  | 0.0 | 0.0 | 31.2  | 547         |
|  |             | Total   | 8.1 | 76.3 | 15.6 | 0.0 | 0.0 | 100.0 | 558         |
| Yukon mainstem at Eagle, Alaska <sup>a</sup> | 78          | Males   | 0.0 | 56.4 | 3.9  | 0.0 | 0.0 | 60.3  | 575         |
|  |             | Females | 0.0 | 34.6 | 5.1  | 0.0 | 0.0 | 39.7  | 555         |
|  |             | Total   | 0.0 | 91.0 | 9.0  | 0.0 | 0.0 | 100.0 | 567         |
| Fishing Branch River, Canada <sup>c</sup>    | 184         | Males   | 1.6 | 42.4 | 4.9  | 0.5 | 0.0 | 49.5  | 575         |
|  |             | Females | 0.0 | 41.3 | 9.2  | 0.0 | 0.0 | 50.5  | 567         |
|  |             | Total   | 1.6 | 83.7 | 14.1 | 0.5 | 0.0 | 100.0 | 571         |

*Note:* Length is measured mid eye to the fork of tail to the nearest millimeter. Data is unweighted.

<sup>a</sup> Samples were from test fishing with drift gillnets, structure is scales.

<sup>b</sup> Samples were handpicked carcasses from east and middle channels, structure is vertebrae.

<sup>c</sup> Samples were collected from live fish passing the Fishing Branch River weir, supplemented by opportunistic carcass sampling, structure is scales.

## **APPENDIX B: TABLES**

Appendix B1.—Alaskan and Canadian total utilization of Yukon River Chinook, chum, and coho salmon, 1961–2021.

| Year | Alaska/U.S. <sup>a, b</sup> |              |           | Yukon/Canada <sup>c</sup> |                           |        | Total   |              |           |
|------|-----------------------------|--------------|-----------|---------------------------|---------------------------|--------|---------|--------------|-----------|
|      | Chinook                     | Other salmon | Total     | Chinook                   | Other salmon <sup>d</sup> | Total  | Chinook | Other salmon | Total     |
| 1961 | 141,152                     | 461,597      | 602,749   | 13,246                    | 9,076                     | 22,322 | 154,398 | 470,673      | 625,071   |
| 1962 | 105,844                     | 434,663      | 540,507   | 13,937                    | 9,436                     | 23,373 | 119,781 | 444,099      | 563,880   |
| 1963 | 141,910                     | 429,396      | 571,306   | 10,077                    | 27,696                    | 37,773 | 151,987 | 457,092      | 609,079   |
| 1964 | 109,818                     | 504,420      | 614,238   | 7,408                     | 12,221                    | 19,629 | 117,226 | 516,641      | 633,867   |
| 1965 | 134,706                     | 484,587      | 619,293   | 5,380                     | 11,789                    | 17,169 | 140,086 | 496,376      | 636,462   |
| 1966 | 104,822                     | 309,502      | 414,324   | 4,452                     | 13,324                    | 17,776 | 109,274 | 322,826      | 432,100   |
| 1967 | 146,104                     | 352,397      | 498,501   | 5,150                     | 16,961                    | 22,111 | 151,254 | 369,358      | 520,612   |
| 1968 | 118,530                     | 270,818      | 389,348   | 5,042                     | 11,633                    | 16,675 | 123,572 | 282,451      | 406,023   |
| 1969 | 104,999                     | 424,399      | 529,398   | 2,624                     | 7,776                     | 10,400 | 107,623 | 432,175      | 539,798   |
| 1970 | 93,019                      | 585,760      | 678,779   | 4,663                     | 3,711                     | 8,374  | 97,682  | 589,471      | 687,153   |
| 1971 | 136,091                     | 547,448      | 683,539   | 6,447                     | 17,471                    | 23,918 | 142,538 | 564,919      | 707,457   |
| 1972 | 113,098                     | 461,617      | 574,715   | 5,729                     | 7,532                     | 13,261 | 118,827 | 469,149      | 587,976   |
| 1973 | 99,696                      | 779,158      | 878,854   | 4,522                     | 10,182                    | 14,704 | 104,218 | 789,340      | 893,558   |
| 1974 | 117,847                     | 1,229,678    | 1,347,525 | 5,631                     | 11,646                    | 17,277 | 123,478 | 1,241,324    | 1,364,802 |
| 1975 | 76,959                      | 1,307,037    | 1,383,996 | 6,000                     | 20,600                    | 26,600 | 82,959  | 1,327,637    | 1,410,596 |
| 1976 | 105,950                     | 1,026,908    | 1,132,858 | 5,025                     | 5,200                     | 10,225 | 110,975 | 1,032,108    | 1,143,083 |
| 1977 | 117,014                     | 1,090,758    | 1,207,772 | 7,527                     | 12,479                    | 20,006 | 124,541 | 1,103,237    | 1,227,778 |
| 1978 | 130,476                     | 1,615,312    | 1,745,788 | 5,881                     | 9,566                     | 15,447 | 136,357 | 1,624,878    | 1,761,235 |
| 1979 | 159,232                     | 1,596,133    | 1,755,365 | 10,375                    | 22,084                    | 32,459 | 169,607 | 1,618,217    | 1,787,824 |
| 1980 | 197,665                     | 1,730,960    | 1,928,625 | 22,846                    | 23,718                    | 46,564 | 220,511 | 1,754,678    | 1,975,189 |
| 1981 | 188,477                     | 2,097,871    | 2,286,348 | 18,109                    | 22,781                    | 40,890 | 206,586 | 2,120,652    | 2,327,238 |
| 1982 | 152,808                     | 1,265,457    | 1,418,265 | 17,208                    | 16,091                    | 33,299 | 170,016 | 1,281,548    | 1,451,564 |

-continued-



Appendix B1.—Page 2 of 3.

| Year | Alaska/U.S. <sup>a, b</sup> |              |           | Yukon/Canada <sup>c</sup> |                           |        | Total   |              |           |
|------|-----------------------------|--------------|-----------|---------------------------|---------------------------|--------|---------|--------------|-----------|
|      | Chinook                     | Other salmon | Total     | Chinook                   | Other salmon <sup>d</sup> | Total  | Chinook | Other salmon | Total     |
| 1983 | 198,436                     | 1,678,597    | 1,877,033 | 18,952                    | 29,490                    | 48,442 | 217,388 | 1,708,087    | 1,925,475 |
| 1984 | 162,683                     | 1,548,101    | 1,710,784 | 16,795                    | 29,767                    | 46,562 | 179,478 | 1,577,868    | 1,757,346 |
| 1985 | 187,327                     | 1,657,984    | 1,845,311 | 19,301                    | 41,515                    | 60,816 | 206,628 | 1,699,499    | 1,906,127 |
| 1986 | 146,004                     | 1,758,825    | 1,904,829 | 20,364                    | 14,843                    | 35,207 | 166,368 | 1,773,668    | 1,940,036 |
| 1987 | 192,007                     | 1,276,066    | 1,468,073 | 17,614                    | 44,786                    | 62,400 | 209,621 | 1,320,852    | 1,530,473 |
| 1988 | 150,009                     | 2,360,718    | 2,510,727 | 21,427                    | 33,915                    | 55,342 | 171,436 | 2,394,633    | 2,566,069 |
| 1989 | 157,632                     | 2,292,211    | 2,449,843 | 17,944                    | 23,490                    | 41,434 | 175,576 | 2,315,701    | 2,491,277 |
| 1990 | 149,433                     | 1,055,515    | 1,204,948 | 19,227                    | 34,304                    | 53,531 | 168,660 | 1,089,819    | 1,258,479 |
| 1991 | 154,651                     | 1,335,111    | 1,489,762 | 20,607                    | 35,653                    | 56,260 | 175,258 | 1,370,764    | 1,546,022 |
| 1992 | 169,642                     | 880,535      | 1,050,177 | 17,903                    | 21,312                    | 39,215 | 187,545 | 901,847      | 1,089,392 |
| 1993 | 161,718                     | 362,551      | 524,269   | 16,611                    | 14,150                    | 30,761 | 178,329 | 376,701      | 555,030   |
| 1994 | 171,654                     | 567,074      | 738,728   | 21,198                    | 38,342                    | 59,540 | 192,852 | 605,416      | 798,268   |
| 1995 | 179,748                     | 1,455,736    | 1,635,484 | 20,884                    | 46,109                    | 66,993 | 200,632 | 1,501,845    | 1,702,477 |
| 1996 | 141,649                     | 1,143,992    | 1,285,641 | 19,612                    | 24,395                    | 44,007 | 161,261 | 1,168,387    | 1,329,648 |
| 1997 | 176,025                     | 560,777      | 736,802   | 16,528                    | 15,900                    | 32,428 | 192,553 | 576,677      | 769,230   |
| 1998 | 99,760                      | 201,480      | 301,240   | 5,937                     | 8,168                     | 14,105 | 105,697 | 209,648      | 315,345   |
| 1999 | 125,427                     | 250,198      | 375,625   | 12,468                    | 19,736                    | 32,204 | 137,895 | 269,934      | 407,829   |
| 2000 | 45,867                      | 120,424      | 166,291   | 4,879                     | 9,283                     | 14,162 | 50,746  | 129,707      | 180,453   |
| 2001 | 56,620                      | 131,500      | 188,120   | 10,144                    | 9,872                     | 20,016 | 66,764  | 141,372      | 208,136   |
| 2002 | 69,240                      | 137,688      | 206,928   | 9,258                     | 8,567                     | 17,825 | 78,498  | 146,255      | 224,753   |
| 2003 | 101,000                     | 214,323      | 315,323   | 9,619                     | 11,435                    | 21,054 | 110,619 | 225,758      | 336,377   |
| 2004 | 114,370                     | 214,744      | 329,114   | 11,238                    | 9,930                     | 21,168 | 125,608 | 224,674      | 350,282   |
| 2005 | 86,369                      | 493,542      | 579,911   | 11,371                    | 18,583                    | 29,954 | 97,740  | 512,125      | 609,865   |

-continued-

Appendix B1.—Page 3 of 3.

| Year              | Alaska/U.S. <sup>a, b</sup> |              |           | Yukon/Canada <sup>c</sup> |                           |        | Total   |              |           |
|-------------------|-----------------------------|--------------|-----------|---------------------------|---------------------------|--------|---------|--------------|-----------|
|                   | Chinook                     | Other salmon | Total     | Chinook                   | Other salmon <sup>d</sup> | Total  | Chinook | Other salmon | Total     |
| 2006              | 96,067                      | 553,299      | 649,366   | 9,072                     | 11,908                    | 20,980 | 105,139 | 565,207      | 670,346   |
| 2007              | 90,753                      | 548,568      | 639,321   | 5,094                     | 14,332                    | 19,426 | 95,847  | 562,900      | 658,747   |
| 2008              | 50,362                      | 500,029      | 550,391   | 3,713                     | 9,566                     | 13,279 | 54,075  | 509,595      | 563,670   |
| 2009              | 35,111                      | 368,717      | 403,828   | 4,758                     | 2,011                     | 6,769  | 39,869  | 370,728      | 410,597   |
| 2010              | 55,092                      | 415,968      | 471,060   | 2,706                     | 5,891                     | 8,597  | 57,798  | 421,859      | 479,657   |
| 2011              | 41,625                      | 780,784      | 822,409   | 4,884                     | 8,226                     | 13,110 | 46,509  | 789,010      | 835,519   |
| 2012              | 30,831                      | 935,740      | 966,571   | 2,200                     | 7,033                     | 9,233  | 33,031  | 942,773      | 975,804   |
| 2013              | 12,741                      | 1,037,537    | 1,050,278 | 2,146                     | 6,170                     | 8,316  | 14,887  | 1,043,707    | 1,058,594 |
| 2014              | 3,287                       | 950,408      | 953,695   | 103                       | 5,166                     | 5,269  | 3,390   | 955,574      | 958,964   |
| 2015              | 7,595                       | 872,084      | 879,679   | 1,204                     | 4,453                     | 5,657  | 8,799   | 876,537      | 885,336   |
| 2016              | 21,689                      | 1,376,854    | 1,398,543 | 2,946                     | 5,750                     | 8,696  | 24,635  | 1,382,604    | 1,407,239 |
| 2017 <sup>e</sup> | 38,347                      | 1,370,813    | 1,409,160 | 3,631                     | 5,787                     | 9,418  | 41,978  | 1,376,600    | 1,418,578 |
| 2018 <sup>e</sup> | 32,213                      | 1,225,903    | 1,258,116 | 3,098                     | 4,856                     | 7,954  | 35,311  | 1,230,759    | 1,266,070 |
| 2019 <sup>e</sup> | 51,782                      | 687,642      | 739,424   | 3,104                     | 3,759                     | 6,863  | 54,886  | 691,401      | 746,287   |
| 2020 <sup>e</sup> | 22,780                      | 68,885       | 91,665    | 2,543                     | 100                       | 2,643  | 25,323  | 68,985       | 94,308    |
| 2021 <sup>e</sup> | 1,945                       | 2,249        | 4,194     | 322                       | 21                        | 343    | 2,267   | 2,270        | 4,537     |
| Average           |                             |              |           |                           |                           |        |         |              |           |
| 1961–2020         | 109,729                     | 873,280      | 983,009   | 9,973                     | 15,692                    | 25,665 | 119,702 | 888,972      | 1,008,674 |
| 2011–2020         | 26,289                      | 930,665      | 956,954   | 2,586                     | 5,130                     | 7,716  | 28,875  | 935,795      | 964,670   |
| 2016–2020         | 33,362                      | 946,019      | 979,382   | 3,064                     | 4,050                     | 7,115  | 36,427  | 950,070      | 986,496   |
| Minimum           | 3,287                       | 68,885       | 91,665    | 103                       | 100                       | 2,643  | 3,390   | 68,985       | 94,308    |
| Maximum           | 198,436                     | 2,360,718    | 2,510,727 | 22,846                    | 46,109                    | 66,993 | 220,511 | 2,394,633    | 2,566,069 |

Note: Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Catch in number of salmon. Includes estimated number of salmon harvested for the commercial production of salmon roe.

<sup>b</sup> Commercial, subsistence, personal use, test fish retained for subsistence, and sport catches combined. Beginning in 2017 report includes harvest from the Coastal District communities of Scammon Bay and Hooper Bay even though not all stocks are bound for the Yukon River. Coastal District harvest information is included in the following years: 1978, 1987–1989 and 1992 to present.

<sup>c</sup> Catch in number of salmon. Commercial, Aboriginal, domestic, and sport catches combined.

<sup>d</sup> Includes coho salmon harvests in First Nations public angling and commercial fisheries, most of which was harvested in the Old Crow Aboriginal fishery (99.8%).

<sup>e</sup> Data are preliminary; particularly not yet published Alaska subsistence harvest data from 2017–2021.

Appendix B2.—Alaska harvest of Yukon River Chinook salmon, 1961–2021.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial<br>related <sup>c</sup> | Personal<br>use <sup>d</sup> | Test<br>fish sales | Sport<br>fish | Yukon Area<br>total |
|------|--------------------------|-------------------------|------------------------------------|------------------------------|--------------------|---------------|---------------------|
| 1961 | 21,488                   | 119,664                 |                                    |                              |                    |               | 141,152             |
| 1962 | 11,110                   | 94,734                  |                                    |                              |                    |               | 105,844             |
| 1963 | 24,862                   | 117,048                 |                                    |                              |                    |               | 141,910             |
| 1964 | 16,231                   | 93,587                  |                                    |                              |                    |               | 109,818             |
| 1965 | 16,608                   | 118,098                 |                                    |                              |                    |               | 134,706             |
| 1966 | 11,507                   | 93,315                  |                                    |                              |                    |               | 104,822             |
| 1967 | 16,448                   | 129,656                 |                                    |                              |                    |               | 146,104             |
| 1968 | 12,004                   | 106,526                 |                                    |                              |                    |               | 118,530             |
| 1969 | 13,972                   | 91,027                  |                                    |                              |                    |               | 104,999             |
| 1970 | 13,874                   | 79,145                  |                                    |                              |                    |               | 93,019              |
| 1971 | 25,584                   | 110,507                 |                                    |                              |                    |               | 136,091             |
| 1972 | 20,258                   | 92,840                  |                                    |                              |                    |               | 113,098             |
| 1973 | 24,343                   | 75,353                  |                                    |                              |                    |               | 99,696              |
| 1974 | 19,758                   | 98,089                  |                                    |                              |                    |               | 117,847             |
| 1975 | 13,121                   | 63,838                  |                                    |                              |                    |               | 76,959              |
| 1976 | 18,174                   | 87,776                  |                                    |                              |                    |               | 105,950             |
| 1977 | 20,101                   | 96,757                  |                                    |                              |                    | 156           | 117,014             |
| 1978 | 30,785                   | 99,168                  |                                    |                              |                    | 523           | 130,476             |
| 1979 | 31,005                   | 127,673                 |                                    |                              |                    | 554           | 159,232             |
| 1980 | 42,724                   | 153,985                 |                                    |                              |                    | 956           | 197,665             |
| 1981 | 29,690                   | 158,018                 |                                    |                              |                    | 769           | 188,477             |
| 1982 | 28,158                   | 123,644                 |                                    |                              |                    | 1,006         | 152,808             |
| 1983 | 49,478                   | 147,910                 |                                    |                              |                    | 1,048         | 198,436             |
| 1984 | 42,428                   | 119,904                 |                                    |                              |                    | 351           | 162,683             |
| 1985 | 39,771                   | 146,188                 |                                    |                              |                    | 1,368         | 187,327             |
| 1986 | 45,238                   | 99,970                  |                                    |                              |                    | 796           | 146,004             |
| 1987 | 55,039                   | 134,760 <sup>e</sup>    |                                    | 1,706                        |                    | 502           | 192,007             |

-continued-

Appendix B2.–Page 2 of 3.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial<br>related <sup>c</sup> | Personal<br>use <sup>d</sup> | Test<br>fish sales | Sport<br>fish | Yukon Area<br>total |
|------|--------------------------|-------------------------|------------------------------------|------------------------------|--------------------|---------------|---------------------|
| 1988 | 45,495                   | 100,364                 |                                    | 2,125                        | 1,081              | 944           | 150,009             |
| 1989 | 48,462                   | 104,198                 |                                    | 2,616                        | 1,293              | 1,063         | 157,632             |
| 1990 | 48,587                   | 95,247 <sup>e</sup>     | 413                                | 2,594                        | 2,048              | 544           | 149,433             |
| 1991 | 46,773                   | 104,878 <sup>e</sup>    | 1,538                              |                              | 689                | 773           | 154,651             |
| 1992 | 47,077                   | 120,245 <sup>e</sup>    | 927                                |                              | 962                | 431           | 169,642             |
| 1993 | 63,915                   | 93,550                  | 560                                | 426                          | 1,572              | 1,695         | 161,718             |
| 1994 | 53,902                   | 113,137                 | 703                                |                              | 1,631              | 2,281         | 171,654             |
| 1995 | 50,620                   | 122,728                 | 1,324                              | 399                          | 2,152              | 2,525         | 179,748             |
| 1996 | 45,671                   | 89,671                  | 521                                | 215                          | 1,698              | 3,873         | 141,649             |
| 1997 | 57,117                   | 112,841                 | 769                                | 313                          | 2,811              | 2,174         | 176,025             |
| 1998 | 54,124                   | 43,618                  | 81                                 | 357                          | 926                | 654           | 99,760              |
| 1999 | 53,305                   | 69,275                  | 288                                | 331                          | 1,205              | 1,023         | 125,427             |
| 2000 | 36,404                   | 8,515                   | -                                  | 75                           | 597                | 276           | 45,867              |
| 2001 | 55,819                   | -                       | -                                  | 122                          | -                  | 679           | 56,620              |
| 2002 | 43,742                   | 24,128                  | 230                                | 126                          | 528                | 486           | 69,240              |
| 2003 | 56,959                   | 40,438                  | -                                  | 204                          | 680                | 2,719         | 101,000             |
| 2004 | 55,713                   | 56,151                  | -                                  | 201                          | 792                | 1,513         | 114,370             |
| 2005 | 53,409                   | 32,029                  | -                                  | 138                          | 310                | 483           | 86,369              |
| 2006 | 48,593                   | 45,829                  | -                                  | 89                           | 817                | 739           | 96,067              |
| 2007 | 55,174                   | 33,634                  | -                                  | 136                          | 849                | 960           | 90,753              |
| 2008 | 45,186                   | 4,641                   | -                                  | 126                          | -                  | 409           | 50,362              |
| 2009 | 33,805                   | 316                     | -                                  | 127                          | -                  | 863           | 35,111              |
| 2010 | 44,559                   | 9,897                   | -                                  | 162                          | -                  | 474           | 55,092              |
| 2011 | 40,980                   | 82 <sup>f</sup>         | -                                  | 89                           | -                  | 474           | 41,625              |
| 2012 | 30,415                   | -                       | -                                  | 71                           | -                  | 345           | 30,831              |
| 2013 | 12,533                   | -                       | -                                  | 42                           | -                  | 166           | 12,741              |
| 2014 | 3,286                    | -                       | -                                  | 1                            | -                  | 0             | 3,287               |

-continued-

Appendix B2.–Page 3 of 3.

| Year      | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales | Sport fish   | Yukon Area total |
|-----------|--------------------------|-------------------------|---------------------------------|---------------------------|-----------------|--------------|------------------|
| 2015      | 7,577                    | -                       | -                               | 5                         | -               | 13           | 7,595            |
| 2016      | 21,612                   | -                       | -                               | 57                        | -               | 20           | 21,689           |
| 2017      | 38,036 <sup>g</sup>      | 168 <sup>f</sup>        | -                               | 125 <sup>g</sup>          | -               | 18           | 38,347           |
| 2018      | 31,812 <sup>g</sup>      | -                       | -                               | 201 <sup>g</sup>          | -               | 200          | 32,213           |
| 2019      | 48,379 <sup>g</sup>      | 3,110 <sup>h</sup>      | -                               | 244 <sup>g</sup>          | -               | 38           | 51,782           |
| 2020      | 22,668 <sup>g</sup>      | -                       | -                               | 112 <sup>g</sup>          | -               | 49           | 22,780           |
| 2021      | 1,945 <sup>g</sup>       | -                       | -                               | 0 <sup>g</sup>            | -               | <sup>i</sup> | 1,945            |
| Averages  |                          |                         |                                 |                           |                 |              |                  |
| 1961–2020 | 34,924                   | 84,767                  | 669                             | 437                       | 1,192           | 839          | 109,729          |
| 2011–2020 | 25,730                   | 1,120                   | -                               | 95                        | -               | 132          | 26,289           |
| 2016–2020 | 32,501                   | 1,639                   | -                               | 148                       | -               | 65           | 33,362           |
| Minimum   | 3,286                    | 82                      | 81                              | 1                         | 310             | 0            | 3,287            |
| Maximum   | 63,915                   | 158,018                 | 1,538                           | 2,616                     | 2,811           | 3,873        | 198,436          |

*Note:* Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Includes test fish harvest and commercial retained fish (not sold) that were utilized for subsistence. Coastal District harvest information is included in the following years: 1975–1978, 1987–1989 and 1992–present even though not all stocks harvested in the Coastal District are bound for the Yukon River.

<sup>b</sup> Includes ADF&G test fish sales prior to 1988.

<sup>c</sup> Includes an estimate of the number of salmon harvested for the commercial production of salmon roe; including carcasses from subsistence caught fish. These data are only available since 1990.

<sup>d</sup> Regulations did not provide for personal use fisheries in the Yukon River drainage prior to 1987 and in 1990, 1991, and 1994 therefore fishing occurred under subsistence regulations.

<sup>e</sup> Includes Chinook salmon sold illegally.

<sup>f</sup> No Chinook salmon were sold in the summer season. A total of 82 and 168 Chinook salmon were sold in District 1 and 2 in the fall season in 2011 and 2017 respectively.

<sup>g</sup> Data are not yet published and are considered preliminary.

<sup>h</sup> Incidental harvest to chum salmon directed fishery in the summer season and allowed sales in the fall season.

<sup>i</sup> Data are unavailable at this time.

Appendix B3.—Alaska harvest of Yukon River summer chum salmon, 1970–2021.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales | Sport fish | Yukon Area total |
|------|--------------------------|-------------------------|---------------------------------|---------------------------|-----------------|------------|------------------|
| 1970 | 166,504                  | 137,006                 |                                 |                           |                 |            | 303,510          |
| 1971 | 171,487                  | 100,090                 |                                 |                           |                 |            | 271,577          |
| 1972 | 108,006                  | 135,668                 |                                 |                           |                 |            | 243,674          |
| 1973 | 161,012                  | 285,509                 |                                 |                           |                 |            | 446,521          |
| 1974 | 227,811                  | 589,892                 |                                 |                           |                 |            | 817,703          |
| 1975 | 211,888                  | 710,295                 |                                 |                           |                 |            | 922,183          |
| 1976 | 186,872                  | 600,894                 |                                 |                           |                 |            | 787,766          |
| 1977 | 159,502                  | 534,875                 |                                 |                           |                 | 316        | 694,693          |
| 1978 | 171,383                  | 1,052,226               | 25,761                          |                           |                 | 451        | 1,249,821        |
| 1979 | 155,970                  | 779,316                 | 40,217                          |                           |                 | 328        | 975,831          |
| 1980 | 167,705                  | 928,609                 | 139,106                         |                           |                 | 483        | 1,235,903        |
| 1981 | 117,629                  | 1,006,938               | 272,763                         |                           |                 | 612        | 1,397,942        |
| 1982 | 117,413                  | 461,403                 | 255,610                         |                           |                 | 780        | 835,206          |
| 1983 | 149,180                  | 744,879                 | 250,590                         |                           |                 | 998        | 1,145,647        |
| 1984 | 166,630                  | 588,597                 | 277,443                         |                           |                 | 585        | 1,033,255        |
| 1985 | 157,744                  | 516,997                 | 417,016                         |                           |                 | 1,267      | 1,093,024        |
| 1986 | 182,337                  | 721,469                 | 467,381                         |                           |                 | 895        | 1,372,082        |
| 1987 | 200,346                  | 442,238                 | 180,303                         | 4,262                     |                 | 846        | 827,995          |
| 1988 | 227,829                  | 1,148,650               | 468,032                         | 2,225                     | 3,587           | 1,037      | 1,851,360        |
| 1989 | 169,496                  | 955,806 <sup>e</sup>    | 496,934                         | 1,891                     | 10,605          | 2,132      | 1,636,864        |
| 1990 | 115,609                  | 302,625                 | 214,552                         | 1,827                     | 8,263           | 472        | 643,348          |
| 1991 | 118,540                  | 349,113 <sup>e</sup>    | 308,989                         |                           | 3,934           | 1,037      | 781,613          |
| 1992 | 142,192                  | 332,313 <sup>e</sup>    | 211,264                         |                           | 1,967           | 1,308      | 689,044          |
| 1993 | 125,574                  | 96,522                  | 43,594                          | 674                       | 1,869           | 564        | 268,797          |
| 1994 | 124,807                  | 80,284                  | 178,457                         |                           | 3,212           | 350        | 387,110          |
| 1995 | 136,083                  | 259,774                 | 558,640                         | 780                       | 6,073           | 1,174      | 962,524          |
| 1996 | 124,738                  | 147,127                 | 535,106                         | 905                       | 7,309           | 1,946      | 817,131          |

-continued-

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales    | Sport fish   | Yukon Area total |
|------|--------------------------|-------------------------|---------------------------------|---------------------------|--------------------|--------------|------------------|
| 1997 | 112,820                  | 95,242                  | 133,010                         | 391                       | 2,590              | 662          | 344,715          |
| 1998 | 87,366                   | 28,611                  | 187                             | 84                        | 3,019              | 421          | 119,688          |
| 1999 | 83,784                   | 29,389                  | 24                              | 382                       | 836                | 555          | 114,970          |
| 2000 | 78,072                   | 6,624                   | 0                               | 30                        | 648                | 161          | 85,535           |
| 2001 | 72,155                   | <sup>f</sup>            | 0                               | 146                       | 0                  | 82           | 72,383           |
| 2002 | 87,056                   | 13,558                  | 19                              | 175                       | 218                | 384          | 101,410          |
| 2003 | 82,272                   | 10,685                  | 0                               | 148                       | 119                | 1,638        | 94,862           |
| 2004 | 77,934                   | 26,410                  | 0                               | 231                       | 217                | 203          | 104,995          |
| 2005 | 93,259                   | 41,264                  | 0                               | 152                       | 134                | 435          | 135,244          |
| 2006 | 115,078                  | 92,116                  | 0                               | 262                       | 456                | 583          | 208,495          |
| 2007 | 92,926                   | 198,201                 | 0                               | 184                       | 10                 | 245          | 291,566          |
| 2008 | 86,514                   | 151,186                 | 0                               | 138                       | 80                 | 371          | 238,289          |
| 2009 | 80,539                   | 170,272                 | 0                               | 308                       | 0                  | 174          | 251,293          |
| 2010 | 88,373                   | 232,888                 | 0                               | 319                       | 0                  | 1,183        | 322,763          |
| 2011 | 96,020                   | 275,161                 | 0                               | 439                       | 0                  | 294          | 371,914          |
| 2012 | 126,992                  | 319,575                 | 0                               | 321                       | 2,412              | 271          | 449,571          |
| 2013 | 115,114                  | 485,587                 | 0                               | 138                       | 2,304              | 1,423        | 604,566          |
| 2014 | 86,900                   | 530,644                 | 0                               | 235                       | 0                  | 374          | 618,153          |
| 2015 | 83,567                   | 358,856                 | 0                               | 220                       | 2,494 <sup>g</sup> | 194          | 445,331          |
| 2016 | 87,902                   | 525,809                 | 0                               | 176                       | 380                | 264          | 614,531          |
| 2017 | 87,437 <sup>h</sup>      | 556,516                 | 0                               | 438 <sup>h</sup>          | 1,819              | 186          | 646,396          |
| 2018 | 76,926 <sup>h</sup>      | 576,700                 | 0                               | 509 <sup>h</sup>          | 1,028              | 200          | 655,363          |
| 2019 | 63,303 <sup>h</sup>      | 227,089                 | 0                               | 294 <sup>h</sup>          | 230                | 36           | 290,952          |
| 2020 | 42,597 <sup>h</sup>      | 13,955                  | 0                               | 67 <sup>h</sup>           | 0                  | 1,684        | 58,303           |
| 2021 | 1,253 <sup>h</sup>       | <sup>f</sup>            | 0                               | 0 <sup>h</sup>            | 0                  | <sup>i</sup> | 1,253            |

-continued-

Appendix B3.–Page 3 of 3.

| Year      | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales | Sport fish | Yukon Area total |
|-----------|--------------------------|-------------------------|---------------------------------|---------------------------|-----------------|------------|------------------|
| Averages  |                          |                         |                                 |                           |                 |            |                  |
| 1961–2020 | 126,532                  | 386,969                 | 130,357                         | 609                       | 2,057           | 649        | 617,502          |
| 2011–2020 | 86,676                   | 386,989                 | 0                               | 284                       | 1,067           | 493        | 475,508          |
| 2016–2020 | 71,633                   | 380,014                 | 0                               | 297                       | 691             | 474        | 453,109          |
| Minimum   | 42,597                   | 6,624                   | 0                               | 30                        | 0               | 36         | 58,303           |
| Maximum   | 227,829                  | 1,148,650               | 558,640                         | 4,262                     | 10,605          | 2,132      | 1,851,360        |

*Note:* Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Includes test fish giveaways and commercial retained fish (not sold) that were utilized for subsistence. Coastal District harvest information is included in the following years: 1987–1989 and 1992–present even though not all stocks harvested in the Coastal District are bound for the Yukon River.

<sup>b</sup> Includes ADF&G test fish sales prior to 1988.

<sup>c</sup> Includes an estimate of the number of salmon harvested for the commercial production of salmon roe; including carcasses from subsistence caught fish.

<sup>d</sup> Regulations did not provide for personal use fisheries in the Yukon River drainage prior to 1987 and in 1990, 1991, and 1994 therefore fishing occurred under subsistence regulations.

<sup>e</sup> Includes illegal sales of summer chum salmon.

<sup>f</sup> Summer season commercial fishery was not conducted.

<sup>g</sup> Test fish sales includes both the Lower Yukon Test Fishery sales and Purse Seine Test Fishery sales.

<sup>h</sup> Data are not yet published and are considered preliminary.

<sup>i</sup> Data are unavailable at this time.



Appendix B4.—Alaska harvest of Yukon River fall chum salmon, 1961–2021.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial<br>related <sup>c</sup> | Personal<br>use <sup>d</sup> | Test<br>fish sales <sup>e</sup> | Yukon Area<br>total |
|------|--------------------------|-------------------------|------------------------------------|------------------------------|---------------------------------|---------------------|
| 1961 | 101,772 <sup>f, g</sup>  | 42,461                  | 0                                  |                              |                                 | 144,233             |
| 1962 | 87,285 <sup>f, g</sup>   | 53,116                  | 0                                  |                              |                                 | 140,401             |
| 1963 | 99,031 <sup>f, g</sup>   |                         |                                    |                              |                                 | 99,031              |
| 1964 | 120,360 <sup>f, g</sup>  | 8,347                   | 0                                  |                              |                                 | 128,707             |
| 1965 | 112,283 <sup>f, g</sup>  | 23,317                  | 0                                  |                              |                                 | 135,600             |
| 1966 | 51,503 <sup>f, g</sup>   | 71,045                  | 0                                  |                              |                                 | 122,548             |
| 1967 | 68,744 <sup>f, g</sup>   | 38,274                  | 0                                  |                              |                                 | 107,018             |
| 1968 | 44,627 <sup>f, g</sup>   | 52,925                  | 0                                  |                              |                                 | 97,552              |
| 1969 | 52,063 <sup>f, g</sup>   | 131,310                 | 0                                  |                              |                                 | 183,373             |
| 1970 | 55,501 <sup>f, g</sup>   | 209,595                 | 0                                  |                              |                                 | 265,096             |
| 1971 | 57,162 <sup>f, g</sup>   | 189,594                 | 0                                  |                              |                                 | 246,756             |
| 1972 | 36,002 <sup>f, g</sup>   | 152,176                 | 0                                  |                              |                                 | 188,178             |
| 1973 | 53,670 <sup>f, g</sup>   | 232,090                 | 0                                  |                              |                                 | 285,760             |
| 1974 | 93,776 <sup>f, g</sup>   | 289,776                 | 0                                  |                              |                                 | 383,552             |
| 1975 | 86,591 <sup>f, g</sup>   | 275,009                 | 0                                  |                              |                                 | 361,600             |
| 1976 | 72,327 <sup>f, g</sup>   | 156,390                 | 0                                  |                              |                                 | 228,717             |
| 1977 | 82,771 <sup>g</sup>      | 257,986                 | 0                                  |                              |                                 | 340,757             |
| 1978 | 84,239 <sup>g</sup>      | 236,383                 | 10,628                             |                              |                                 | 331,250             |
| 1979 | 214,881                  | 359,946                 | 18,466                             |                              |                                 | 593,293             |
| 1980 | 167,637                  | 293,430                 | 5,020                              |                              |                                 | 466,087             |
| 1981 | 177,240                  | 466,451                 | 11,285                             |                              |                                 | 654,976             |
| 1982 | 132,092                  | 224,187                 | 805                                |                              |                                 | 357,084             |
| 1983 | 187,864                  | 302,598                 | 5,064                              |                              |                                 | 495,526             |
| 1984 | 172,495                  | 208,232                 | 2,328                              |                              |                                 | 383,055             |
| 1985 | 203,947                  | 267,744                 | 2,525                              |                              |                                 | 474,216             |
| 1986 | 163,466                  | 139,442                 | 577                                |                              |                                 | 303,485             |
| 1987 | 342,819 <sup>h</sup>     |                         |                                    | 19,066                       |                                 | 361,885             |
| 1988 | 153,848                  | 133,763                 | 3,227                              | 3,881                        | 27,663                          | 322,382             |
| 1989 | 211,303                  | 270,195                 | 14,749                             | 5,082                        | 20,973                          | 522,302             |

-continued-

## Appendix B4.–Page 2 of 3.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial<br>related <sup>c</sup> | Personal<br>use <sup>d</sup> | Test<br>fish sales <sup>e</sup> | Yukon Area<br>total |
|------|--------------------------|-------------------------|------------------------------------|------------------------------|---------------------------------|---------------------|
| 1990 | 167,900                  | 124,174                 | 12,168                             | 5,176                        | 9,224                           | 318,642             |
| 1991 | 145,524                  | 230,852                 | 23,366                             | 0                            | 3,936                           | 403,678             |
| 1992 | 107,808                  | 15,721 <sup>j</sup>     | 3,301                              | 0                            | 1,407                           | 128,237             |
| 1993 | 76,882                   | <sup>i</sup>            |                                    | 163                          | 0                               | 77,045              |
| 1994 | 123,565                  | 3,631                   | 4,368                              | 0                            | 0                               | 131,564             |
| 1995 | 130,860                  | 250,766                 | 32,324                             | 863                          | 1,121                           | 415,934             |
| 1996 | 129,258                  | 88,342                  | 17,288                             | 356                          | 1,717                           | 236,961             |
| 1997 | 95,141                   | 56,713                  | 1,474                              | 284                          | 867                             | 154,479             |
| 1998 | 62,901                   | <sup>i</sup>            |                                    | 2                            | 0                               | 62,903              |
| 1999 | 89,940                   | 20,371                  | 0                                  | 262                          | 1,171                           | 111,744             |
| 2000 | 19,395                   | <sup>i</sup>            |                                    | 1                            | 0                               | 19,396              |
| 2001 | 35,703                   | <sup>i</sup>            |                                    | 10                           | 0                               | 35,713              |
| 2002 | 19,674                   | <sup>i</sup>            |                                    | 3                            | 0                               | 19,677              |
| 2003 | 56,930                   | 10,996                  | 0                                  | 394                          | 0                               | 68,320              |
| 2004 | 62,526                   | 4,110                   | 0                                  | 230                          | 0                               | 66,866              |
| 2005 | 91,534                   | 180,249                 | 0                                  | 133                          | 87                              | 272,003             |
| 2006 | 84,002                   | 174,542                 | 0                                  | 333                          | 0                               | 258,877             |
| 2007 | 101,221                  | 90,677                  | 0                                  | 173                          | 0                               | 192,071             |
| 2008 | 89,357                   | 119,265                 | 0                                  | 181                          | 0                               | 208,803             |
| 2009 | 66,119                   | 25,876                  | 0                                  | 78                           | 0                               | 92,073              |
| 2010 | 68,645                   | 2,550                   | 0                                  | 3,209                        | 0                               | 74,404              |
| 2011 | 80,202                   | 238,979                 | 0                                  | 347                          | 0                               | 319,528             |
| 2012 | 99,309                   | 289,692                 | 0                                  | 410                          | 166                             | 389,577             |
| 2013 | 113,384                  | 238,051                 | 0                                  | 383                          | 121                             | 351,939             |
| 2014 | 92,529                   | 115,599                 | 0                                  | 278                          | 30                              | 208,436             |
| 2015 | 86,600                   | 191,470                 | 0                                  | 80                           | 50                              | 278,200             |
| 2016 | 84,617                   | 465,511                 | 0                                  | 283                          | 668                             | 551,079             |
| 2017 | 85,093 <sup>k</sup>      | 489,702                 | 0                                  | 626 <sup>k</sup>             | 1,246                           | 576,667             |
| 2018 | 64,494 <sup>k</sup>      | 387,788                 | 0                                  | 514 <sup>k</sup>             | 907                             | 453,703             |

-continued-

Appendix B4.–Page 3 of 3.

| Year      | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales <sup>e</sup> | Yukon Area total |
|-----------|--------------------------|-------------------------|---------------------------------|---------------------------|------------------------------|------------------|
| 2019      | 63,862 <sup>k</sup>      | 268,360 <sup>l</sup>    | 0                               | 408 <sup>k</sup>          | 275                          | 332,905          |
| 2020      | 6,207 <sup>k</sup>       | <sup>i</sup>            | 0                               | 37 <sup>k</sup>           | 0                            | 6,244            |
| 2021      | 703 <sup>k</sup>         | <sup>i</sup>            | 0                               | 0 <sup>k</sup>            | 0                            | 703              |
| Averages  |                          |                         |                                 |                           |                              |                  |
| 1961–2020 | 101,475                  | 176,342                 | 3,188                           | 1,272                     | 2,171                        | 259,035          |
| 2011–2020 | 77,630                   | 298,350                 | 0                               | 337                       | 346                          | 346,828          |
| 2016–2020 | 60,855                   | 402,840                 | 0                               | 374                       | 619                          | 384,120          |
| Minimum   | 6,207                    | 2,550                   | 0                               | 0                         | 0                            | 6,244            |
| Maximum   | 342,819                  | 489,702                 | 32,324                          | 19,066                    | 27,663                       | 654,976          |

*Note:* Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Includes test fish harvest and commercial retained fish (not sold) that were utilized for subsistence. Coastal District harvest information is included in the following years: 1978, 1987–1989 and 1992–present even though not all stocks harvested in the Coastal District are bound for the Yukon River.

<sup>b</sup> Includes fish sold in the round and estimated numbers of female salmon commercially harvested for production of salmon roe (see Bergstrom et al. 1992). Includes ADF&G test fish prior to 1988. Beginning in 1999, commercial harvest may include some commercial related harvest.

<sup>c</sup> Includes an estimate of number of salmon harvested for the commercial production of salmon roe and the carcasses used for subsistence. In prior JTC reports, subsistence plus commercial related harvests are noted as subsistence "use".

<sup>d</sup> Regulations did not provide for personal use fisheries in the Yukon River drainage prior to 1987 and in 1990, 1991, and 1994 therefore fishing occurred under subsistence regulations.

<sup>e</sup> Test fish sales is the number of salmon sold by ADF&G test fisheries.

<sup>f</sup> Catches estimated because harvests of species other than Chinook salmon were not differentiated.

<sup>g</sup> Minimum estimates from 1961–1978 because subsistence surveys were conducted prior to the end of the fishing season.

<sup>h</sup> Includes an estimated 95,768 and 119,168 fall chum salmon illegally sold in Districts 5 (Yukon River) and 6 (Tanana River), respectively.

<sup>i</sup> Commercial fishery was not conducted.

<sup>j</sup> Commercial fishery operated only in District 6, the Tanana River.

<sup>k</sup> Data are not yet published and are considered preliminary.

<sup>l</sup> Commercial harvest includes an estimated 63,000 summer chum salmon that is removed for the total run size estimate.

Appendix B5.—Alaska harvest of Yukon River coho salmon, 1961–2021.

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales <sup>e</sup> | Sport fish <sup>f</sup> | Yukon Area total |
|------|--------------------------|-------------------------|---------------------------------|---------------------------|------------------------------|-------------------------|------------------|
| 1961 | 9,192 <sup>g, h</sup>    | 2,855                   | 0                               |                           |                              |                         | 12,047           |
| 1962 | 9,480 <sup>g, h</sup>    | 22,926                  | 0                               |                           |                              |                         | 32,406           |
| 1963 | 27,699 <sup>g, h</sup>   | 5,572                   | 0                               |                           |                              |                         | 33,271           |
| 1964 | 12,187 <sup>g, h</sup>   | 2,446                   | 0                               |                           |                              |                         | 14,633           |
| 1965 | 11,789 <sup>g, h</sup>   | 350                     | 0                               |                           |                              |                         | 12,139           |
| 1966 | 13,192 <sup>g, h</sup>   | 19,254                  | 0                               |                           |                              |                         | 32,446           |
| 1967 | 17,164 <sup>g, h</sup>   | 11,047                  | 0                               |                           |                              |                         | 28,211           |
| 1968 | 11,613 <sup>g, h</sup>   | 13,303                  | 0                               |                           |                              |                         | 24,916           |
| 1969 | 7,776 <sup>g, h</sup>    | 15,093                  | 0                               |                           |                              |                         | 22,869           |
| 1970 | 3,966 <sup>g, h</sup>    | 13,188                  | 0                               |                           |                              |                         | 17,154           |
| 1971 | 16,912 <sup>g, h</sup>   | 12,203                  | 0                               |                           |                              |                         | 29,115           |
| 1972 | 7,532 <sup>g, h</sup>    | 22,233                  | 0                               |                           |                              |                         | 29,765           |
| 1973 | 10,236 <sup>g, h</sup>   | 36,641                  | 0                               |                           |                              |                         | 46,877           |
| 1974 | 11,646 <sup>g, h</sup>   | 16,777                  | 0                               |                           |                              |                         | 28,423           |
| 1975 | 20,708 <sup>g, h</sup>   | 2,546                   | 0                               |                           |                              |                         | 23,254           |
| 1976 | 5,241 <sup>g, h</sup>    | 5,184                   | 0                               |                           |                              |                         | 10,425           |
| 1977 | 16,333 <sup>h</sup>      | 38,863                  | 0                               |                           |                              | 112                     | 55,308           |
| 1978 | 7,787 <sup>h</sup>       | 26,152                  | 0                               |                           |                              | 302                     | 34,241           |
| 1979 | 9,794                    | 17,165                  | 0                               |                           |                              | 50                      | 27,009           |
| 1980 | 20,158                   | 8,745                   | 0                               |                           |                              | 67                      | 28,970           |
| 1981 | 21,228                   | 23,680                  | 0                               |                           |                              | 45                      | 44,953           |
| 1982 | 35,894                   | 37,176                  | 0                               |                           |                              | 97                      | 73,167           |
| 1983 | 23,905                   | 13,320                  | 0                               |                           |                              | 199                     | 37,424           |
| 1984 | 49,020                   | 81,940                  | 0                               |                           |                              | 831                     | 131,791          |
| 1985 | 32,264                   | 57,672                  | 0                               |                           |                              | 808                     | 90,744           |
| 1986 | 34,468                   | 47,255                  | 0                               |                           |                              | 1,535                   | 83,258           |
| 1987 | 82,371 <sup>i</sup>      | <sup>j</sup>            |                                 | 2,523                     |                              | 1,292                   | 86,186           |
| 1988 | 69,679                   | 99,907                  | 0                               | 1,250                     | 13,720                       | 2,420                   | 186,976          |
| 1989 | 40,924                   | 85,493                  | 0                               | 872                       | 3,945                        | 1,811                   | 133,045          |

-continued-

| Year | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial<br>related <sup>c</sup> | Personal<br>use <sup>d</sup> | Test<br>fish sales <sup>e</sup> | Sport<br>fish <sup>f</sup> | Yukon Area<br>total |
|------|--------------------------|-------------------------|------------------------------------|------------------------------|---------------------------------|----------------------------|---------------------|
| 1990 | 43,460                   | 41,032                  | 3,255                              | 1,181                        | 2,650                           | 1,947                      | 93,525              |
| 1991 | 37,388                   | 103,180                 | 3,506                              | 0                            | 2,971                           | 2,775                      | 149,820             |
| 1992 | 51,980                   | 6,556 <sup>k</sup>      | 1,423                              | 0                            | 1,629                           | 1,666                      | 63,254              |
| 1993 | 15,812                   | <sup>j</sup>            |                                    | 0                            | 0                               | 897                        | 16,709              |
| 1994 | 41,775                   | 120                     | 4,331                              | 0                            | 0                               | 2,174                      | 48,400              |
| 1995 | 28,377                   | 45,939                  | 1,074                              | 417                          | 193                             | 1,278                      | 77,278              |
| 1996 | 30,404                   | 52,643                  | 3,339                              | 198                          | 1,728                           | 1,588                      | 89,900              |
| 1997 | 23,945                   | 35,320                  | 0                                  | 350                          | 498                             | 1,470                      | 61,583              |
| 1998 | 18,121                   | 1                       | 0                                  | 9                            | 0                               | 758                        | 18,889              |
| 1999 | 20,891                   | 1,601                   | 0                                  | 147                          | 236                             | 609                        | 23,484              |
| 2000 | 14,939                   | <sup>j</sup>            |                                    | 0                            | 0                               | 554                        | 15,493              |
| 2001 | 22,122                   | <sup>j</sup>            |                                    | 34                           | 0                               | 1,248                      | 23,404              |
| 2002 | 15,489                   | <sup>j</sup>            |                                    | 20                           | 0                               | 1,092                      | 16,601              |
| 2003 | 23,872                   | 25,243                  | 0                                  | 549                          | 0                               | 1,477                      | 51,141              |
| 2004 | 20,795                   | 20,232                  | 0                                  | 233                          | 0                               | 1,623                      | 42,883              |
| 2005 | 27,250                   | 58,311                  | 0                                  | 107                          | 0                               | 627                        | 86,295              |
| 2006 | 19,706                   | 64,942                  | 0                                  | 279                          | 0                               | 1,000                      | 85,927              |
| 2007 | 19,624                   | 44,575                  | 0                                  | 135                          | 0                               | 597                        | 64,931              |
| 2008 | 16,855                   | 35,691                  | 0                                  | 50                           | 0                               | 341                        | 52,937              |
| 2009 | 16,006                   | 8,311                   | 0                                  | 70                           | 0                               | 964                        | 25,351              |
| 2010 | 13,045                   | 3,750                   | 0                                  | 1,062                        | 0                               | 944                        | 18,801              |
| 2011 | 12,344                   | 76,303                  | 0                                  | 232                          | 0                               | 463                        | 89,342              |
| 2012 | 21,533                   | 74,789                  | 0                                  | 100                          | 39                              | 131                        | 96,592              |
| 2013 | 14,457                   | 66,199                  | 0                                  | 109                          | 1                               | 266                        | 81,032              |
| 2014 | 17,098                   | 104,692                 | 0                                  | 174                          | 0                               | 1,855                      | 123,819             |
| 2015 | 18,107                   | 129,700                 | 0                                  | 145                          | 8                               | 593                        | 148,553             |
| 2016 | 8,815                    | 201,482                 | 0                                  | 266                          | 11                              | 670                        | 211,244             |
| 2017 | 7,281 <sup>1</sup>       | 139,915                 | 0                                  | 200 <sup>1</sup>             | 63                              | 291                        | 147,750             |
| 2018 | 5,527 <sup>1</sup>       | 110,587                 | 0                                  | 131 <sup>1</sup>             | 48                              | 544                        | 116,837             |

-continued-

| Year      | Subsistence <sup>a</sup> | Commercial <sup>b</sup> | Commercial related <sup>c</sup> | Personal use <sup>d</sup> | Test fish sales <sup>e</sup> | Sport fish <sup>f</sup> | Yukon Area total |
|-----------|--------------------------|-------------------------|---------------------------------|---------------------------|------------------------------|-------------------------|------------------|
| 2019      | 5,014 <sup>1</sup>       | 58,591                  | 0                               | 68 <sup>1</sup>           | 40                           | 72                      | 63,785           |
| 2020      | 2,922 <sup>1</sup>       | j                       | 0                               | 79 <sup>1</sup>           | 0                            | 1,337                   | 4,338            |
| 2021      | 293 <sup>1</sup>         | j                       | 0                               | 0 <sup>1</sup>            | 0                            | - <sup>m</sup>          | 293              |
| Averages  |                          |                         |                                 |                           |                              |                         |                  |
| 1961–2020 | 21,252                   | 41,642                  | 308                             | 323                       | 842                          | 941                     | 60,349           |
| 2011–2020 | 11,310                   | 106,918                 | 0                               | 150                       | 21                           | 622                     | 108,329          |
| 2016–2020 | 5,912                    | 127,644                 | 0                               | 149                       | 32                           | 583                     | 108,791          |
| Minimum   | 2,922                    | 1                       | 0                               | 0                         | 0                            | 45                      | 4,338            |
| Maximum   | 82,371                   | 201,482                 | 4,331                           | 2,523                     | 13,720                       | 2,775                   | 211,244          |

*Note:* Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Includes test fish harvest and commercial retained fish (not sold) that were utilized for subsistence. Coastal District harvest information is included in the following years: 1978, 1988, 1989, and 1992–present even though not all stocks harvested in the Coastal District are bound for the Yukon River.

<sup>b</sup> Includes fish sold in the round and estimated numbers of female salmon commercially harvested for production of salmon roe (see Bergstrom et al. 1992: 1990 Yukon Area Annual Management Report). Includes ADF&G test fish prior to 1988. Beginning in 1999, commercial harvest may include some commercial related harvest.

<sup>c</sup> Includes an estimate of number of salmon harvested for the commercial production of salmon roe and the carcasses used for subsistence.

<sup>d</sup> Regulations did not provide for personal use fisheries in the Yukon River drainage prior to 1987 and in 1990, 1991, and 1994 therefore fishing occurred under subsistence regulations.

<sup>e</sup> Test fish sales is the number of salmon sold by ADF&G test fisheries.

<sup>f</sup> The majority of the sport-fish harvest is taken in the Tanana River drainage.

<sup>g</sup> Catches estimated because harvests of species other than Chinook salmon were not differentiated.

<sup>h</sup> Minimum estimates from 1961–1978 because subsistence surveys were conducted prior to the end of the fishing season.

<sup>i</sup> Includes an estimated 5,015 and 31,276 coho salmon illegally sold in Districts 5 (Yukon River) and 6 (Tanana River), respectively.

<sup>j</sup> Commercial fishery was not conducted.

<sup>k</sup> Commercial fishery operated only in District 6, the Tanana River.

<sup>l</sup> Data are not yet published and are considered preliminary.

<sup>m</sup> Data are unavailable at this time.

Appendix B6.—Alaska (U.S.) and Canada total utilization of Yukon River Chinook and fall chum salmon, 1961–2021.

| Year | Chinook salmon      |                        |         | Fall chum salmon    |                        |         |
|------|---------------------|------------------------|---------|---------------------|------------------------|---------|
|      | Canada <sup>a</sup> | Alaska <sup>b, c</sup> | Total   | Canada <sup>a</sup> | Alaska <sup>b, c</sup> | Total   |
| 1961 | 13,246              | 141,152                | 154,398 | 9,076               | 144,233                | 153,309 |
| 1962 | 13,937              | 105,844                | 119,781 | 9,436               | 140,401                | 149,837 |
| 1963 | 10,077              | 141,910                | 151,987 | 27,696              | 99,031 <sup>d</sup>    | 126,727 |
| 1964 | 7,408               | 109,818                | 117,226 | 12,187              | 128,707                | 140,894 |
| 1965 | 5,380               | 134,706                | 140,086 | 11,789              | 135,600                | 147,389 |
| 1966 | 4,452               | 104,822                | 109,274 | 13,192              | 122,548                | 135,740 |
| 1967 | 5,150               | 146,104                | 151,254 | 16,961              | 107,018                | 123,979 |
| 1968 | 5,042               | 118,530                | 123,572 | 11,633              | 97,552                 | 109,185 |
| 1969 | 2,624               | 104,999                | 107,623 | 7,776               | 183,373                | 191,149 |
| 1970 | 4,663               | 93,019                 | 97,682  | 3,711               | 265,096                | 268,807 |
| 1971 | 6,447               | 136,091                | 142,538 | 16,911              | 246,756                | 263,667 |
| 1972 | 5,729               | 113,098                | 118,827 | 7,532               | 188,178                | 195,710 |
| 1973 | 4,522               | 99,696                 | 104,218 | 10,135              | 285,760                | 295,895 |
| 1974 | 5,631               | 117,847                | 123,478 | 11,646              | 383,552                | 395,198 |
| 1975 | 6,000               | 76,959                 | 82,959  | 20,600              | 361,600                | 382,200 |
| 1976 | 5,025               | 105,950                | 110,975 | 5,200               | 228,717                | 233,917 |
| 1977 | 7,527               | 117,014                | 124,541 | 12,479              | 340,757                | 353,236 |
| 1978 | 5,881               | 130,476                | 136,357 | 9,566               | 331,250                | 340,816 |
| 1979 | 10,375              | 159,232                | 169,607 | 22,084              | 593,293                | 615,377 |
| 1980 | 22,846              | 197,665                | 220,511 | 22,218              | 466,087                | 488,305 |
| 1981 | 18,109              | 188,477                | 206,586 | 22,281              | 654,976                | 677,257 |
| 1982 | 17,208              | 152,808                | 170,016 | 16,091              | 357,084                | 373,175 |
| 1983 | 18,952              | 198,436                | 217,388 | 29,490              | 495,526                | 525,016 |
| 1984 | 16,795              | 162,683                | 179,478 | 29,267              | 383,055                | 412,322 |
| 1985 | 19,301              | 187,327                | 206,628 | 41,265              | 474,216                | 515,481 |
| 1986 | 20,364              | 146,004                | 166,368 | 14,543              | 303,485                | 318,028 |
| 1987 | 17,614              | 192,007                | 209,621 | 44,480              | 361,885 <sup>d</sup>   | 406,365 |
| 1988 | 21,427              | 150,009                | 171,436 | 33,565              | 322,382                | 355,947 |
| 1989 | 17,944              | 157,632                | 175,576 | 23,020              | 522,302                | 545,322 |
| 1990 | 19,227              | 149,433                | 168,660 | 33,622              | 318,642                | 352,264 |
| 1991 | 20,607              | 154,651                | 175,258 | 35,418              | 403,678                | 439,096 |
| 1992 | 17,903              | 169,642                | 187,545 | 20,815              | 128,237 <sup>e</sup>   | 149,052 |
| 1993 | 16,611              | 161,718                | 178,329 | 14,090              | 77,045 <sup>d</sup>    | 91,135  |
| 1994 | 21,198              | 171,654                | 192,852 | 38,008              | 131,564                | 169,572 |
| 1995 | 20,884              | 179,748                | 200,632 | 45,600              | 415,934                | 461,534 |
| 1996 | 19,612              | 141,649                | 161,261 | 24,354              | 236,961                | 261,315 |
| 1997 | 16,528              | 176,025                | 192,553 | 15,600              | 154,479                | 170,079 |
| 1998 | 5,937               | 99,760                 | 105,697 | 7,954               | 62,903                 | 70,857  |
| 1999 | 12,468              | 125,427                | 137,895 | 19,636              | 111,744                | 131,380 |

-continued-

## Appendix B6.–Page 2 of 2.

| Year              | Chinook salmon      |                        |         | Fall chum salmon    |                        |         |
|-------------------|---------------------|------------------------|---------|---------------------|------------------------|---------|
|                   | Canada <sup>a</sup> | Alaska <sup>b, c</sup> | Total   | Canada <sup>a</sup> | Alaska <sup>b, c</sup> | Total   |
| 2000              | 4,879               | 45,867                 | 50,746  | 9,246               | 19,396 <sup>d</sup>    | 28,642  |
| 2001              | 10,144              | 56,620 <sup>f</sup>    | 66,764  | 9,872               | 35,713 <sup>d</sup>    | 45,585  |
| 2002              | 9,258               | 69,240                 | 78,498  | 8,092               | 19,677 <sup>d</sup>    | 27,769  |
| 2003              | 9,619               | 101,000                | 110,619 | 10,905              | 68,320                 | 79,225  |
| 2004              | 11,238              | 114,370                | 125,608 | 9,750               | 66,866                 | 76,616  |
| 2005              | 11,371              | 86,369                 | 97,740  | 18,572              | 272,003                | 290,575 |
| 2006              | 9,072               | 96,067                 | 105,139 | 11,796              | 258,877                | 270,673 |
| 2007              | 5,094               | 90,753                 | 95,847  | 13,830              | 192,071                | 205,901 |
| 2008              | 3,713               | 50,362                 | 54,075  | 9,566               | 208,803                | 218,369 |
| 2009              | 4,758               | 35,111                 | 39,869  | 2,011               | 92,073                 | 94,084  |
| 2010              | 2,706               | 55,092                 | 57,798  | 5,787               | 74,404                 | 80,191  |
| 2011              | 4,884               | 41,625 <sup>f</sup>    | 46,509  | 8,163               | 319,528                | 327,691 |
| 2012              | 2,200               | 30,831 <sup>f</sup>    | 33,031  | 7,023               | 389,577                | 396,600 |
| 2013              | 2,146               | 12,741 <sup>f</sup>    | 14,887  | 6,170               | 351,939                | 358,109 |
| 2014              | 103                 | 3,287 <sup>f</sup>     | 3,390   | 5,033               | 208,436                | 213,469 |
| 2015              | 1,204               | 7,595 <sup>f</sup>     | 8,799   | 4,453               | 278,200                | 282,653 |
| 2016              | 2,946               | 21,689 <sup>f</sup>    | 24,635  | 5,750               | 551,079                | 556,829 |
| 2017 <sup>g</sup> | 3,631               | 38,347 <sup>f</sup>    | 41,978  | 5,716               | 576,667                | 582,383 |
| 2018 <sup>g</sup> | 3,098               | 32,213 <sup>f</sup>    | 35,311  | 4,831               | 453,703                | 458,534 |
| 2019 <sup>g</sup> | 3,104               | 51,782 <sup>f</sup>    | 54,886  | 3,759               | 332,905                | 336,664 |
| 2020 <sup>g</sup> | 2,543               | 22,780 <sup>f</sup>    | 25,323  | 100                 | 6,244                  | 6,344   |
| 2021 <sup>g</sup> | 322                 | 1,945 <sup>f</sup>     | 2,267   | 21                  | 703                    | 724     |
| Averages          |                     |                        |         |                     |                        |         |
| 1961–2020         | 9,973               | 109,729                | 119,702 | 15,523              | 259,035                | 274,557 |
| 2011–2020         | 2,586               | 26,289                 | 28,875  | 5,100               | 346,828                | 351,928 |
| 2016–2020         | 3,064               | 33,362                 | 36,427  | 4,031               | 384,120                | 388,151 |
| Minimum           | 103                 | 3,287                  | 3,390   | 100                 | 6,244                  | 6,344   |
| Maximum           | 22,846              | 198,436                | 220,511 | 45,600              | 654,976                | 677,257 |

*Note:* Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Catches in number of salmon. Includes commercial, Aboriginal, domestic, and sport catches combined.

<sup>b</sup> Catch in number of salmon. Includes estimated number of salmon harvested for the commercial production of salmon roe (see Bergstrom et al. 1992: 1990 Yukon Area Annual Management Report).

<sup>c</sup> Commercial, subsistence, personal-use, test fish, and sport catches combined. Coastal District harvest information is included in the following years: 1975–1978, 1987–1989 and 1992–present even though not all stocks harvested in the Coastal District are bound for the Yukon River.

<sup>d</sup> Commercial fishery did not operate within the Alaskan portion of the drainage.

<sup>e</sup> Commercial fishery operated only in District 6, the Tanana River.

<sup>f</sup> No Chinook salmon directed commercial fishery was conducted during the summer season.

<sup>g</sup> Data are preliminary, particularly not yet published Alaska subsistence and personal use harvest data from 2017–2021.



Appendix B7.—Canadian harvest of Yukon River Chinook salmon, 1961–2021.

| Year | Mainstem Yukon River harvest |          |                         |                             |                 |                            | Porcupine River | Total<br>Canadian<br>harvest |                                    |
|------|------------------------------|----------|-------------------------|-----------------------------|-----------------|----------------------------|-----------------|------------------------------|------------------------------------|
|      | Commercial                   | Domestic | First Nation<br>fishery | Public Angling <sup>a</sup> | Test<br>fishery | Combined<br>non-commercial | Total           |                              | First Nation<br>fishery<br>harvest |
| 1961 | 3,446                        |          | 9,300                   |                             |                 | 9,300                      | 12,746          | 500                          | 13,246                             |
| 1962 | 4,037                        |          | 9,300                   |                             |                 | 9,300                      | 13,337          | 600                          | 13,937                             |
| 1963 | 2,283                        |          | 7,750                   |                             |                 | 7,750                      | 10,033          | 44                           | 10,077                             |
| 1964 | 3,208                        |          | 4,124                   |                             |                 | 4,124                      | 7,332           | 76                           | 7,408                              |
| 1965 | 2,265                        |          | 3,021                   |                             |                 | 3,021                      | 5,286           | 94                           | 5,380                              |
| 1966 | 1,942                        |          | 2,445                   |                             |                 | 2,445                      | 4,387           | 65                           | 4,452                              |
| 1967 | 2,187                        |          | 2,920                   |                             |                 | 2,920                      | 5,107           | 43                           | 5,150                              |
| 1968 | 2,212                        |          | 2,800                   |                             |                 | 2,800                      | 5,012           | 30                           | 5,042                              |
| 1969 | 1,640                        |          | 957                     |                             |                 | 957                        | 2,597           | 27                           | 2,624                              |
| 1970 | 2,611                        |          | 2,044                   |                             |                 | 2,044                      | 4,655           | 8                            | 4,663                              |
| 1971 | 3,178                        |          | 3,260                   |                             |                 | 3,260                      | 6,438           | 9                            | 6,447                              |
| 1972 | 1,769                        |          | 3,960                   |                             |                 | 3,960                      | 5,729           |                              | 5,729                              |
| 1973 | 2,199                        |          | 2,319                   |                             |                 | 2,319                      | 4,518           | 4                            | 4,522                              |
| 1974 | 1,808                        | 406      | 3,342                   |                             |                 | 3,748                      | 5,556           | 75                           | 5,631                              |
| 1975 | 3,000                        | 400      | 2,500                   |                             |                 | 2,900                      | 5,900           | 100                          | 6,000                              |
| 1976 | 3,500                        | 500      | 1,000                   |                             |                 | 1,500                      | 5,000           | 25                           | 5,025                              |
| 1977 | 4,720                        | 531      | 2,247                   |                             |                 | 2,778                      | 7,498           | 29                           | 7,527                              |
| 1978 | 2,975                        | 421      | 2,485                   |                             |                 | 2,906                      | 5,881           |                              | 5,881                              |
| 1979 | 6,175                        | 1,200    | 3,000                   |                             |                 | 4,200                      | 10,375          |                              | 10,375                             |
| 1980 | 9,500                        | 3,500    | 7,546                   | 300                         |                 | 11,346                     | 20,846          | 2,000                        | 22,846                             |
| 1981 | 8,593                        | 237      | 8,879                   | 300                         |                 | 9,416                      | 18,009          | 100                          | 18,109                             |
| 1982 | 8,640                        | 435      | 7,433                   | 300                         |                 | 8,168                      | 16,808          | 400                          | 17,208                             |
| 1983 | 13,027                       | 400      | 5,025                   | 300                         |                 | 5,725                      | 18,752          | 200                          | 18,952                             |
| 1984 | 9,885                        | 260      | 5,850                   | 300                         |                 | 6,410                      | 16,295          | 500                          | 16,795                             |
| 1985 | 12,573                       | 478      | 5,800                   | 300                         |                 | 6,578                      | 19,151          | 150                          | 19,301                             |
| 1986 | 10,797                       | 342      | 8,625                   | 300                         |                 | 9,267                      | 20,064          | 300                          | 20,364                             |
| 1987 | 10,864                       | 330      | 6,069                   | 300                         |                 | 6,699                      | 17,563          | 51                           | 17,614                             |
| 1988 | 13,217                       | 282      | 7,178                   | 650                         |                 | 8,110                      | 21,327          | 100                          | 21,427                             |
| 1989 | 9,789                        | 400      | 6,930                   | 300                         |                 | 7,630                      | 17,419          | 525                          | 17,944                             |
| 1990 | 11,324                       | 247      | 7,109                   | 300                         |                 | 7,656                      | 18,980          | 247                          | 19,227                             |
| 1991 | 10,906                       | 227      | 9,011                   | 300                         |                 | 9,538                      | 20,444          | 163                          | 20,607                             |
| 1992 | 10,877                       | 277      | 6,349                   | 300                         |                 | 6,926                      | 17,803          | 100                          | 17,903                             |

-continued-

| Year | Mainstem Yukon River harvest |          |                         |                             |                 |                            |        | Porcupine River                    | Total<br>Canadian<br>harvest |
|------|------------------------------|----------|-------------------------|-----------------------------|-----------------|----------------------------|--------|------------------------------------|------------------------------|
|      | Commercial                   | Domestic | First Nation<br>fishery | Public Angling <sup>a</sup> | Test<br>fishery | Combined<br>non-commercial | Total  | First Nation<br>fishery<br>harvest |                              |
| 1993 | 10,350                       | 243      | 5,576                   | 300                         |                 | 6,119                      | 16,469 | 142                                | 16,611                       |
| 1994 | 12,028                       | 373      | 8,069                   | 300                         |                 | 8,742                      | 20,770 | 428                                | 21,198                       |
| 1995 | 11,146                       | 300      | 7,942                   | 700                         |                 | 8,942                      | 20,088 | 796                                | 20,884                       |
| 1996 | 10,164                       | 141      | 8,451                   | 790                         |                 | 9,382                      | 19,546 | 66                                 | 19,612                       |
| 1997 | 5,311                        | 288      | 8,888                   | 1,230                       |                 | 10,406                     | 15,717 | 811                                | 16,528                       |
| 1998 | 390                          | 24       | 4,687                   | -                           | 737             | 5,448                      | 5,838  | 99                                 | 5,937                        |
| 1999 | 3,160                        | 213      | 8,804                   | 177                         |                 | 9,194                      | 12,354 | 114                                | 12,468                       |
| 2000 | -                            | -        | 4,068                   | -                           | 761             | 4,829                      | 4,829  | 50                                 | 4,879                        |
| 2001 | 1,351                        | 89       | 7,421                   | 146                         | 767             | 8,423                      | 9,774  | 370                                | 10,144                       |
| 2002 | 708                          | 59       | 7,139                   | 128                         | 1,036           | 8,362                      | 9,070  | 188                                | 9,258                        |
| 2003 | 2,672                        | 115      | 6,121                   | 275                         | 263             | 6,774                      | 9,446  | 173                                | 9,619                        |
| 2004 | 3,785                        | 88       | 6,483                   | 423                         | 167             | 7,161                      | 10,946 | 292                                | 11,238                       |
| 2005 | 4,066                        | 99       | 6,376                   | 436                         |                 | 6,911                      | 10,977 | 394                                | 11,371                       |
| 2006 | 2,332                        | 63       | 5,757                   | 606                         |                 | 6,426                      | 8,758  | 314                                | 9,072                        |
| 2007 | -                            | -        | 4,175                   | 2 <sup>b</sup>              | 617             | 4,794                      | 4,794  | 300                                | 5,094                        |
| 2008 | 1 <sup>c</sup>               | -        | 2,885                   | -                           | 513             | 3,398                      | 3,399  | 314                                | 3,713                        |
| 2009 | 364                          | 17       | 3,791                   | 125                         | -               | 3,933                      | 4,297  | 461                                | 4,758                        |
| 2010 | -                            | -        | 2,455 <sup>d</sup>      | 1 <sup>e</sup>              | -               | 2,456                      | 2,456  | 250                                | 2,706                        |
| 2011 | 4 <sup>c</sup>               | -        | 4,550 <sup>d</sup>      | 40                          | -               | 4,590                      | 4,594  | 290                                | 4,884                        |
| 2012 | -                            | -        | 2,000 <sup>d</sup>      | -                           | -               | 2,000                      | 2,000  | 200                                | 2,200                        |
| 2013 | 2 <sup>c</sup>               | -        | 1,902 <sup>d</sup>      | -                           | -               | 1,902                      | 1,904  | 242                                | 2,146                        |
| 2014 | -                            | -        | 100                     | -                           | -               | 100                        | 100    | 3                                  | 103                          |
| 2015 | -                            | -        | 1,000                   | -                           | -               | 1,000                      | 1,000  | 204                                | 1,204                        |
| 2016 | 1 <sup>c</sup>               | -        | 2,768                   | -                           | -               | 2,768                      | 2,769  | 177                                | 2,946                        |
| 2017 | -                            | -        | 3,500                   | -                           | -               | 3,500                      | 3,500  | 131                                | 3,631                        |
| 2018 | 1 <sup>c</sup>               | -        | 2,789                   | -                           | -               | 2,789                      | 2,790  | 308                                | 3,098                        |
| 2019 | -                            | -        | 2,764                   | -                           | -               | 2,764                      | 2,764  | 340                                | 3,104                        |
| 2020 | -                            | -        | 2,363                   | -                           | -               | 2,363                      | 2,363  | 180                                | 2,543                        |
| 2021 | -                            | -        | 306                     | -                           | -               | 306                        | 306    | 16                                 | 322                          |

-continued-

Appendix B7.–Page 3 of 3.

| Year      | Mainstem Yukon River harvest |          |                         |                             |                 |                            |        | Porcupine River                    | Total<br>Canadian<br>harvest |
|-----------|------------------------------|----------|-------------------------|-----------------------------|-----------------|----------------------------|--------|------------------------------------|------------------------------|
|           | Commercial                   | Domestic | First Nation<br>fishery | Public Angling <sup>a</sup> | Test<br>fishery | Combined<br>non-commercial | Total  | First Nation<br>fishery<br>harvest |                              |
| Averages  |                              |          |                         |                             |                 |                            |        |                                    |                              |
| 1961–2020 | 5,717 <sup>f</sup>           | 393      | 4,890                   | 342 <sup>f</sup>            | 608             | 5,353                      | 9,736  | 249                                | 9,973                        |
| 2011–2020 | 2 <sup>f</sup>               | -        | 2,374                   | 21                          | -               | 2,378                      | 2,378  | 208                                | 2,586                        |
| 2016–2020 | 1                            | -        | 2,837                   | -                           | -               | 2,837                      | 2,837  | 227                                | 3,064                        |
| Minimum   | 1                            | 17       | 100                     | 1                           | 167             | 100                        | 100    | 3                                  | 103                          |
| Maximum   | 13,217                       | 3,500    | 9,300                   | 1,230                       | 1,036           | 11,346                     | 21,327 | 2,000                              | 22,846                       |

*Note:* Minimum and maximum values exclude the most recent year data. Dash indicates fishery did not occur.

<sup>a</sup> Public angling harvest unknown before 1980.

<sup>b</sup> Public angling fishery involved non-retention of Chinook salmon for most of the season thus effectively closed.

<sup>c</sup> Closed during Chinook salmon season, harvested in chum salmon fishery.

<sup>d</sup> Adjusted to account for underreporting.

<sup>e</sup> Fishery was closed, 1 fish mistakenly caught and retained.

<sup>f</sup> Excluding years when no directed fishery occurred.

Appendix B8.—Canadian harvest of Yukon River fall chum salmon, 1961–2021.

| Year | Mainstem Yukon River Harvest |          |                      |              |                                      | Total <sup>a</sup> | Porcupine River              | Total Canadian harvest |
|------|------------------------------|----------|----------------------|--------------|--------------------------------------|--------------------|------------------------------|------------------------|
|      | Commercial                   | Domestic | First Nation fishery | Test fishery | Combined non-commercial <sup>a</sup> |                    | First Nation fishery harvest |                        |
| 1961 | 3,276                        |          | 3,800                |              | 3,800                                | 7,076              | 2,000                        | 9,076                  |
| 1962 | 936                          |          | 6,500                |              | 6,500                                | 7,436              | 2,000                        | 9,436                  |
| 1963 | 2,196                        |          | 5,500                |              | 5,500                                | 7,696              | 20,000                       | 27,696                 |
| 1964 | 1,929                        |          | 4,200                |              | 4,200                                | 6,129              | 6,058                        | 12,187                 |
| 1965 | 2,071                        |          | 2,183                |              | 2,183                                | 4,254              | 7,535                        | 11,789                 |
| 1966 | 3,157                        |          | 1,430                |              | 1,430                                | 4,587              | 8,605                        | 13,192                 |
| 1967 | 3,343                        |          | 1,850                |              | 1,850                                | 5,193              | 11,768                       | 16,961                 |
| 1968 | 453                          |          | 1,180                |              | 1,180                                | 1,633              | 10,000                       | 11,633                 |
| 1969 | 2,279                        |          | 2,120                |              | 2,120                                | 4,399              | 3,377                        | 7,776                  |
| 1970 | 2,479                        |          | 612                  |              | 612                                  | 3,091              | 620                          | 3,711                  |
| 1971 | 1,761                        |          | 150                  |              | 150                                  | 1,911              | 15,000                       | 16,911                 |
| 1972 | 2,532                        |          |                      |              | 0                                    | 2,532              | 5,000                        | 7,532                  |
| 1973 | 2,806                        |          | 1,129                |              | 1,129                                | 3,935              | 6,200                        | 10,135                 |
| 1974 | 2,544                        | 466      | 1,636                |              | 2,102                                | 4,646              | 7,000                        | 11,646                 |
| 1975 | 2,500                        | 4,600    | 2,500                |              | 7,100                                | 9,600              | 11,000                       | 20,600                 |
| 1976 | 1,000                        | 1,000    | 100                  |              | 1,100                                | 2,100              | 3,100                        | 5,200                  |
| 1977 | 3,990                        | 1,499    | 1,430                |              | 2,929                                | 6,919              | 5,560                        | 12,479                 |
| 1978 | 3,356                        | 728      | 482                  |              | 1,210                                | 4,566              | 5,000                        | 9,566                  |
| 1979 | 9,084                        | 2,000    | 11,000               |              | 13,000                               | 22,084             |                              | 22,084                 |
| 1980 | 9,000                        | 4,000    | 3,218                |              | 7,218                                | 16,218             | 6,000                        | 22,218                 |
| 1981 | 15,260                       | 1,611    | 2,410                |              | 4,021                                | 19,281             | 3,000                        | 22,281                 |
| 1982 | 11,312                       | 683      | 3,096                |              | 3,779                                | 15,091             | 1,000                        | 16,091                 |
| 1983 | 25,990                       | 300      | 1,200                |              | 1,500                                | 27,490             | 2,000                        | 29,490                 |
| 1984 | 22,932                       | 535      | 1,800                |              | 2,335                                | 25,267             | 4,000                        | 29,267                 |
| 1985 | 35,746                       | 279      | 1,740                |              | 2,019                                | 37,765             | 3,500                        | 41,265                 |
| 1986 | 11,464                       | 222      | 2,200                |              | 2,422                                | 13,886             | 657                          | 14,543                 |
| 1987 | 40,591                       | 132      | 3,622                |              | 3,754                                | 44,345             | 135                          | 44,480                 |

-continued-

| Year              | Mainstem Yukon River Harvest |          |                      |                    |                                      |                    | Porcupine River              | Total Canadian harvest |
|-------------------|------------------------------|----------|----------------------|--------------------|--------------------------------------|--------------------|------------------------------|------------------------|
|                   | Commercial                   | Domestic | First Nation fishery | Test fishery       | Combined non-commercial <sup>a</sup> | Total <sup>a</sup> | First Nation fishery harvest |                        |
| 1988              | 30,263                       | 349      | 1,882                |                    | 2,231                                | 32,494             | 1,071                        | 33,565                 |
| 1989              | 17,549                       | 100      | 2,462                | 300                | 2,562                                | 20,111             | 2,909                        | 23,020                 |
| 1990              | 27,537                       | 0        | 3,675                |                    | 3,675                                | 31,212             | 2,410                        | 33,622                 |
| 1991              | 31,404                       | 0        | 2,438                |                    | 2,438                                | 33,842             | 1,576                        | 35,418                 |
| 1992              | 18,576                       | 0        | 304                  |                    | 304                                  | 18,880             | 1,935                        | 20,815                 |
| 1993              | 7,762                        | 0        | 4,660                |                    | 4,660                                | 12,422             | 1,668                        | 14,090                 |
| 1994              | 30,035                       | 0        | 5,319                |                    | 5,319                                | 35,354             | 2,654                        | 38,008                 |
| 1995              | 39,012                       | 0        | 1,099                |                    | 1,099                                | 40,111             | 5,489                        | 45,600                 |
| 1996              | 20,069                       | 0        | 1,260                |                    | 1,260                                | 21,329             | 3,025                        | 24,354                 |
| 1997              | 8,068                        | 0        | 1,238                |                    | 1,238                                | 9,306              | 6,294                        | 15,600                 |
| 1998 <sup>b</sup> | -                            |          | 1,795                |                    | 1,795                                | 1,795              | 6,159                        | 7,954                  |
| 1999              | 10,402                       | 0        | 3,234                |                    | 3,234                                | 13,636             | 6,000                        | 19,636                 |
| 2000              | 1,319                        | 0        | 2,927                |                    | 2,927                                | 4,246              | 5,000                        | 9,246                  |
| 2001              | 2,198                        | 3        | 3,077                | 1 <sup>b</sup>     | 3,080                                | 5,278              | 4,594                        | 9,872                  |
| 2002              | 3,065                        | 0        | 3,167                | 2,756 <sup>b</sup> | 3,167                                | 6,232              | 1,860                        | 8,092                  |
| 2003              | 9,030                        | 0        | 1,493                | 990 <sup>b</sup>   | 1,493                                | 10,523             | 382                          | 10,905                 |
| 2004              | 7,365                        | 0        | 2,180                | 995 <sup>b</sup>   | 2,180                                | 9,545              | 205                          | 9,750                  |
| 2005              | 11,931                       | 13       | 2,035                |                    | 2,048                                | 13,979             | 4,593                        | 18,572                 |
| 2006              | 4,096                        | 0        | 2,521                |                    | 2,521                                | 6,617              | 5,179                        | 11,796                 |
| 2007              | 7,109                        | 0        | 2,221                | 3,765 <sup>b</sup> | 2,221                                | 9,330              | 4,500                        | 13,830                 |
| 2008              | 4,062                        | 0        | 2,068                |                    | 2,068                                | 6,130              | 3,436                        | 9,566                  |
| 2009              | 293                          | 0        | 820                  |                    | 820                                  | 1,113              | 898                          | 2,011                  |
| 2010              | 2,186                        | 0        | 1,523 <sup>c</sup>   |                    | 1,523                                | 3,709              | 2,078                        | 5,787                  |
| 2011              | 5,312                        | 0        | 1,000 <sup>c</sup>   |                    | 1,000                                | 6,312              | 1,851                        | 8,163                  |
| 2012              | 3,205                        | 0        | 700 <sup>c</sup>     |                    | 700                                  | 3,905              | 3,118                        | 7,023                  |
| 2013              | 3,369                        | 18       | 500 <sup>c</sup>     |                    | 518                                  | 3,887              | 2,283                        | 6,170                  |

-continued-

| Year      | Mainstem Yukon River Harvest |          |                      |              |                                      | Porcupine River    |                              | Total Canadian harvest |
|-----------|------------------------------|----------|----------------------|--------------|--------------------------------------|--------------------|------------------------------|------------------------|
|           | Commercial                   | Domestic | First Nation fishery | Test fishery | Combined non-commercial <sup>a</sup> | Total <sup>a</sup> | First Nation fishery harvest |                        |
| 2014      | 2,485                        | 19       | 546                  |              | 565                                  | 3,050              | 1,983                        | 5,033                  |
| 2015      | 2,862                        | 35       | 1,000 <sup>c</sup>   |              | 1,035                                | 3,897              | 556                          | 4,453                  |
| 2016      | 1,745                        | 0        | 1,000 <sup>c</sup>   |              | 1,000                                | 2,745              | 3,005                        | 5,750                  |
| 2017      | 2,404                        | 0        | 1,000 <sup>c</sup>   |              | 1,000                                | 3,404              | 2,312                        | 5,716                  |
| 2018      | 1,957                        | 0        | 1,000 <sup>c</sup>   |              | 1,000                                | 2,957              | 1,874                        | 4,831                  |
| 2019      | 1,728                        | 31       | 1,000 <sup>c</sup>   |              | 1,031                                | 2,759              | 1,000                        | 3,759                  |
| 2020      | -                            | -        | 0                    |              | 0                                    | 0                  | 100                          | 100                    |
| 2021      | -                            | -        | 0                    |              | 0                                    | 0                  | 21                           | 21                     |
| Averages  |                              |          |                      |              |                                      |                    |                              |                        |
| 1961–2020 | 9,351                        | 414      | 2,173                | 1,468        | 2,448                                | 11,487             | 4,104                        | 15,523                 |
| 2011–2020 | 2,785                        | 11       | 775                  | -            | 785                                  | 3,292              | 1,808                        | 5,100                  |
| 2016–2020 | 1,959                        | 8        | 800                  | -            | 806                                  | 2,373              | 1,658                        | 4,031                  |
| Minimum   | 0                            | 0        | 0                    | 0            | 0                                    | 0                  | 100                          | 100                    |
| Maximum   | 40,591                       | 4,600    | 11,000               | 3,765        | 13,000                               | 44,345             | 20,000                       | 45,600                 |

*Note:* Minimum and maximum values exclude the most recent year data. Dash indicates fishery did not occur or the value is not able to be calculated.

<sup>a</sup> Test fishery was not included in totals as it was live release.

<sup>b</sup> The chum salmon test fishery practiced live-release therefore not included in the annual harvest totals.

<sup>c</sup> Adjusted to account for underreporting.

Appendix B9.—Chinook salmon aerial survey indices for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1961–2021.

| Year | Andreafsky River   |                    | Anvik River        |                         | Nulato River            |                  |                    | Gisasa River     |
|------|--------------------|--------------------|--------------------|-------------------------|-------------------------|------------------|--------------------|------------------|
|      | East Fork          | West Fork          | Drainagewide total | Index area <sup>a</sup> | North Fork <sup>b</sup> | South Fork       | Both forks         |                  |
| 1961 | 1,003              | -                  | 1,226              |                         | 376 <sup>c</sup>        | 167              | 543                | 266 <sup>c</sup> |
| 1962 | 675 <sup>c</sup>   | 762 <sup>c</sup>   | -                  | -                       | -                       | -                | -                  | -                |
| 1963 | -                  | -                  | -                  | -                       | -                       | -                | -                  | -                |
| 1964 | 867                | 705                | -                  | -                       | -                       | -                | -                  | -                |
| 1965 | -                  | 344 <sup>c</sup>   | 650 <sup>c</sup>   | -                       | -                       | -                | -                  | -                |
| 1966 | 361                | 303                | 638                | -                       | -                       | -                | -                  | -                |
| 1967 | -                  | 276 <sup>c</sup>   | 336 <sup>c</sup>   | -                       | -                       | -                | -                  | -                |
| 1968 | 383                | 383                | 310 <sup>c</sup>   | -                       | -                       | -                | -                  | -                |
| 1969 | 274 <sup>c</sup>   | 231 <sup>c</sup>   | 296 <sup>c</sup>   | -                       | -                       | -                | -                  | -                |
| 1970 | 665                | 574 <sup>c</sup>   | 368                | -                       | -                       | -                | -                  | -                |
| 1971 | 1,904              | 1,682              | -                  | -                       | -                       | -                | -                  | -                |
| 1972 | 798                | 582 <sup>c</sup>   | 418                | -                       | -                       | -                | -                  | -                |
| 1973 | 825                | 788                | 222                | -                       | -                       | -                | -                  | -                |
| 1974 | -                  | 285 <sup>c</sup>   | -                  | -                       | 55 <sup>c</sup>         | 23 <sup>c</sup>  | 78 <sup>c</sup>    | 161              |
| 1975 | 993                | 301 <sup>c</sup>   | 730                | -                       | 123                     | 81               | 204                | 385              |
| 1976 | 818                | 643                | 1,053              | -                       | 471                     | 177              | 648                | 332              |
| 1977 | 2,008              | 1,499              | 1,371              | -                       | 286                     | 201              | 487                | 255              |
| 1978 | 2,487              | 1,062              | 1,324              | -                       | 498                     | 422              | 920                | 45 <sup>c</sup>  |
| 1979 | 1,180              | 1,134              | 1,484              | -                       | 1,093                   | 414              | 1,507              | 484              |
| 1980 | 958                | 1,500              | 1,330              | 1,192                   | 954 <sup>c</sup>        | 369 <sup>c</sup> | 1,323 <sup>c</sup> | 951              |
| 1981 | 2,146 <sup>c</sup> | 231 <sup>c</sup>   | 807 <sup>c</sup>   | 577 <sup>c</sup>        | -                       | 791 <sup>c</sup> | 791 <sup>c</sup>   |                  |
| 1982 | 1,274              | 851                | -                  |                         | -                       | -                | -                  | 421              |
| 1983 | -                  | -                  | 653 <sup>c</sup>   | 376 <sup>c</sup>        | 526                     | 480              | 1,006              | 572              |
| 1984 | 1,573              | 1,993              | 641 <sup>c</sup>   | 574 <sup>c</sup>        | -                       | -                | -                  | -                |
| 1985 | 1,617              | 2,248              | 1,051              | 720                     | 1,600                   | 1,180            | 2,780              | 735              |
| 1986 | 1,954              | 3,158              | 1,118              | 918                     | 1,452                   | 1,522            | 2,974              | 1,346            |
| 1987 | 1,608              | 3,281              | 1,174              | 879                     | 1,145                   | 493              | 1,638              | 731              |
| 1988 | 1,020              | 1,448              | 1,805              | 1,449                   | 1,061                   | 714              | 1,775              | 797              |
| 1989 | 1,399              | 1,089              | 442 <sup>c</sup>   | 212 <sup>c</sup>        | -                       | -                | -                  | -                |
| 1990 | 2,503              | 1,545              | 2,347              | 1,595                   | 568 <sup>c</sup>        | 430 <sup>c</sup> | 998 <sup>c</sup>   | 884 <sup>c</sup> |
| 1991 | 1,938              | 2,544              | 875 <sup>c</sup>   | 625 <sup>c</sup>        | 767                     | 1,253            | 2,020              | 1,690            |
| 1992 | 1,030 <sup>c</sup> | 2,052 <sup>c</sup> | 1,536              | 931                     | 348                     | 231              | 579                | 910              |
| 1993 | 5,855              | 2,765              | 1,720              | 1,526                   | 1,844                   | 1,181            | 3,025              | 1,385            |
| 1994 | 300 <sup>c</sup>   | 213 <sup>c</sup>   | 913 <sup>c</sup>   | 913 <sup>c</sup>        | -                       | -                | -                  | 2,775            |
| 1995 | 1,635              | 1,108              | 1,996              | 1,147                   | 968                     | 681              | 1,649              | 410              |
| 1996 |                    | 624                | 839                | 709                     | -                       | 100              | 100 <sup>c</sup>   | -                |
| 1997 | 1,140              | 1,510              | 3,979              | 2,690                   | -                       | -                | -                  | 144 <sup>c</sup> |
| 1998 | 1,027              | 1,249 <sup>c</sup> | 709 <sup>c</sup>   | 648 <sup>c</sup>        | 507                     | 546              | 1,053              | 889 <sup>c</sup> |

-continued-

## Appendix B9.—Page 2 of 2.

| Year             | Andreafsky River   |                  | Anvik River        |                         | Nulato River            |              |                    | Gisasa River       |
|------------------|--------------------|------------------|--------------------|-------------------------|-------------------------|--------------|--------------------|--------------------|
|                  | East Fork          | West Fork        | Drainagewide total | Index area <sup>a</sup> | North Fork <sup>b</sup> | South Fork   | Both forks         |                    |
| 1999             | -                  | 870 <sup>c</sup> | 950 <sup>c</sup>   | 950 <sup>c</sup>        | -                       | -            | -                  | -                  |
| 2000             | 1,018              | 427              | 1,721              | 1,394                   | -                       | -            | -                  | -                  |
| 2001             | 1,059              | 565              | 1,420              | 1,177                   | 1,116                   | 768          | 1,884 <sup>d</sup> | 1,298 <sup>c</sup> |
| 2002             | 1,447              | 917              | 1,713              | 1,329                   | 687                     | 897          | 1,584              | 506                |
| 2003             | 1,116 <sup>c</sup> | 1,578            | 973 <sup>c</sup>   | 973 <sup>c</sup>        | -                       | -            | -                  | -                  |
| 2004             | 2,879              | 1,317            | 3,679              | 3,304                   | 856                     | 465          | 1,321              | 731                |
| 2005             | 1,715              | 1,492            | 2,421              | 1,922                   | 323                     | 230          | 553                | 958                |
| 2006             | 591 <sup>c</sup>   | 824              | 1,886              | 1,776 <sup>e</sup>      | 620                     | 672          | 1,292              | 843                |
| 2007             | 1,758              | 976              | 1,650              | 1,497                   | 1,684                   | 899          | 2,583              | 593                |
| 2008             | 278 <sup>c</sup>   | 262 <sup>c</sup> | 992 <sup>c</sup>   | 827 <sup>c</sup>        | 415                     | 507          | 922                | 487                |
| 2009             | 84 <sup>c</sup>    | 1,678            | 832                | 590                     | 1,418                   | 842          | 2,260              | 515                |
| 2010             | 537 <sup>c</sup>   | 858              | 974                | 721                     | 356                     | 355          | 711                | 264                |
| 2011             | 620                | 1,173            | 642                | 501                     | 788                     | 613          | 1,401              | 906                |
| 2012             | -                  | 227 <sup>c</sup> | 722                | 451                     | 682                     | 692          | 1,374              | <sup>c</sup>       |
| 2013             | 1,441              | 1,090            | 940                | 656                     | 586                     | 532          | 1,118              | 201 <sup>c</sup>   |
| 2014             | -                  | 1,695            | 1,584              | 800                     | <sup>c</sup>            | <sup>c</sup> | <sup>c</sup>       | <sup>c</sup>       |
| 2015             | 2,167              | 1,356            | 2,616              | 1,726                   | 999                     | 565          | 1,564              | 558                |
| 2016             | -                  | -                | -                  | -                       | -                       | -            | -                  | -                  |
| 2017             | -                  | 942              | 1,101              | 894                     | 500                     | 443          | 943                |                    |
| 2018             | 746                | 455              | 1,109 <sup>c</sup> | 800                     | 438                     | 432          | 870                | 452                |
| 2019             | 1,547              | 904              | 1,432              | 1,043                   | 656                     | 485          | 1,141              | -                  |
| 2020             | 335                | 508              | 675                | 506                     | 459                     | 403          | 862                | 419                |
| 2021             | -                  | -                | -                  | -                       | -                       | -            | -                  | -                  |
| SEG <sup>f</sup> | <sup>g</sup>       | 640–<br>1,600    | 1,100–1,700        |                         | 940–<br>1,900           |              |                    | <sup>h</sup>       |
| Averages         |                    |                  |                    |                         |                         |              |                    |                    |
| 1961–2020        | 1,297              | 1,091            | 1,215              | 1,065                   | 756                     | 559          | 1,276              | 703                |
| 2011–2020        | 1,143              | 928              | 1,202              | 820                     | 639                     | 521          | 1,159              | 507                |
| 2016–2020        | 876                | 702              | 1,079              | 811                     | 513                     | 441          | 954                | 436                |
| Minimum          | 84                 | 213              | 222                | 212                     | 55                      | 23           | 78                 | 45                 |
| Maximum          | 5,855              | 3,281            | 3,979              | 3,304                   | 1,844                   | 1,522        | 3,025              | 2,775              |

Note: Aerial survey counts are peak counts only. Survey rating was fair or good unless otherwise noted. Minimum and maximum values exclude the most recent year data. Dash indicates no survey.

<sup>a</sup> Anvik River Index Area includes mainstem counts between Beaver Creek and McDonald Creek.

<sup>b</sup> Nulato River mainstem aerial survey counts below the forks are included with the North Fork.

<sup>c</sup> Incomplete, poor timing and/or poor survey conditions resulting in minimal, inaccurate, or no counts.

<sup>d</sup> In 2001, the Nulato River escapement goal was established for both forks combined.

<sup>e</sup> The count represents the index area and an additional 8 river miles downstream of Yellow River confluence.

<sup>f</sup> Sustainable Escapement Goal.

<sup>g</sup> Aerial escapement goal for Andreafsky River was discontinued in 2010. Note: weir-based goal replaced East Fork Andreafsky River aerial survey goal.

<sup>h</sup> Gisasa River aerial escapement goal was discontinued in 2010.



Appendix B10.—Chinook salmon escapement counts and percentage females counted for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1986–2021.

| Year | East Fork Andreafsky River weir |        | Nulato River tower | Henshaw Creek weir |        | Gisasa River weir |        | Chena River tower/sonar |                     | Salcha River tower/sonar |                     |
|------|---------------------------------|--------|--------------------|--------------------|--------|-------------------|--------|-------------------------|---------------------|--------------------------|---------------------|
|      | No. fish                        | % Fem. | No. fish           | No. fish           | % Fem. | No. fish          | % Fem. | No. fish                | % Fem. <sup>a</sup> | No. fish                 | % Fem. <sup>a</sup> |
| 1986 | 1,530 <sup>b</sup>              | 29     | -                  | -                  | -      | -                 | -      | 9,065 <sup>c</sup>      | 25                  | -                        | 35                  |
| 1987 | 2,011 <sup>b</sup>              | 53     | -                  | -                  | -      | -                 | -      | 6,404 <sup>c</sup>      | 58                  | 4,771 <sup>c</sup>       | 63                  |
| 1988 | 1,341 <sup>b</sup>              | 42     | -                  | -                  | -      | -                 | -      | 3,346 <sup>c</sup>      | 61                  | 4,322 <sup>c</sup>       | 40                  |
| 1989 | -                               | 5      | -                  | -                  | -      | -                 | -      | 2,730 <sup>c</sup>      | 65                  | 3,294 <sup>c</sup>       | 62                  |
| 1990 | -                               | 38     | -                  | -                  | -      | -                 | -      | 5,603 <sup>c</sup>      | 47                  | 10,728 <sup>c</sup>      | 47                  |
| 1991 | -                               | 28     | -                  | -                  | -      | -                 | -      | 3,172 <sup>c</sup>      | 32                  | 5,608 <sup>c</sup>       | 47                  |
| 1992 | -                               | 26     | -                  | -                  | -      | -                 | -      | 5,580 <sup>c</sup>      | 38                  | 7,862 <sup>c</sup>       | 34                  |
| 1993 | -                               | 29     | -                  | -                  | -      | -                 | -      | 12,241                  | 17                  | 10,008                   | 28                  |
| 1994 | 7,801                           | 35     | 1,795              | -                  | -      | 2,888             | -      | 11,877                  | 45                  | 18,404                   | 45                  |
| 1995 | 5,841                           | 42     | 1,412              | -                  | -      | 4,023             | 46     | 11,394 <sup>c</sup>     | 66                  | 13,643                   | 56                  |
| 1996 | 2,955                           | 42     | 756                | -                  | -      | 1,991             | 20     | 7,153 <sup>c</sup>      | 44                  | 7,570 <sup>c</sup>       | 51                  |
| 1997 | 3,186                           | 37     | 4,766              | -                  | -      | 3,764             | 26     | 13,390                  | 40                  | 18,514                   | 50                  |
| 1998 | 4,034                           | 29     | 1,536              | -                  | -      | 2,414             | 16     | 4,745                   | 41                  | 5,027                    | 30                  |
| 1999 | 3,444                           | 29     | 1,932              | -                  | -      | 2,644             | 26     | 6,485                   | 66                  | 9,198                    | 55                  |
| 2000 | 1,609                           | 32     | 908                | 193                | 30     | 2,089             | 34     | 4,694 <sup>c</sup>      | 26                  | 4,595                    | 44                  |
| 2001 | 1,148                           | 64     | -                  | 1,091              | 36     | 3,052             | 49     | 9,696                   | 43                  | 13,328                   | 38                  |
| 2002 | 4,123 <sup>d</sup>              | 21     | 2,696              | 649                | 31     | 2,025             | 21     | 6,967 <sup>c</sup>      | 32                  | 9,000 <sup>e</sup>       | 35                  |
| 2003 | 4,336                           | 48     | 1,716 <sup>f</sup> | 748                | 39     | 1,901             | 38     | 11,100                  | 45                  | 15,500 <sup>e</sup>      | 42                  |
| 2004 | 8,045                           | 35     | -                  | 1,248              | 23     | 1,774             | 34     | 9,645                   | 63                  | 15,761                   | 63                  |
| 2005 | 2,239                           | 50     | -                  | 1,059              | 42     | 3,111             | 36     | - <sup>d</sup>          | 42                  | 5,988                    | 54                  |
| 2006 | 6,463                           | 44     | -                  | - <sup>d</sup>     | -      | 3,031             | 29     | 2,936                   | 46                  | 10,679                   | 43                  |
| 2007 | 4,504                           | 45     | -                  | 740                | 43     | 1,427             | 41     | 3,806                   | 40                  | 6,425                    | 36                  |
| 2008 | 4,242                           | 39     | -                  | 766                | 27     | 1,738             | 15     | 3,208                   | 44                  | 5,415 <sup>e</sup>       | 39                  |
| 2009 | 3,004                           | 47     | -                  | 1,637              | 54     | 1,955             | 28     | 5,253                   | 55                  | 12,774                   | 39                  |
| 2010 | 2,413                           | 49     | -                  | 857                | 49     | 1,516             | 30     | 2,382                   | 31                  | 6,135                    | 33                  |
| 2011 | 5,213                           | 20     | -                  | 1,796              | 34     | 2,692             | 19     | - <sup>d</sup>          | 32                  | 7,200 <sup>e</sup>       | 42                  |
| 2012 | 2,517                           | 27     | -                  | 922                | 43     | 1,323             | 39     | 2,220 <sup>g</sup>      | 56                  | 7,165                    | 60                  |
| 2013 | 1,998                           | 39     | -                  | 772                | 47     | 1,126             | 34     | 1,859 <sup>d</sup>      | 40                  | 5,465                    | 50                  |
| 2014 | 5,949                           | 48     | -                  | - <sup>d</sup>     | -      | 1,589             | 19     | 7,192 <sup>h</sup>      | 33                  | - <sup>d</sup>           | 32                  |
| 2015 | 5,474                           | 40     | -                  | 2,391              | 41     | 1,319             | 30     | 6,294                   | 55                  | 6,288 <sup>i</sup>       | 43                  |
| 2016 | 2,676                           | 49     | -                  | 1,354              | 48     | 1,395             | 27     | 6,665 <sup>i</sup>      | 23                  | 2,675 <sup>i</sup>       | 39                  |
| 2017 | 2,970                           | 26     | -                  | 677                | 42     | 1,083             | 28     | 4,201 <sup>i</sup>      | 45                  | 4,195 <sup>i</sup>       | 41                  |

-continued-

Appendix B10.–Page 2 of 2.

| Year                | East Fork Andreafsky River weir |        | Nulato River tower | Henshaw Creek weir |                | Gisasa River weir |                | Chena River tower |                     | Salcha River tower |                     |
|---------------------|---------------------------------|--------|--------------------|--------------------|----------------|-------------------|----------------|-------------------|---------------------|--------------------|---------------------|
|                     | No. fish                        | % Fem. | No. fish           | No. fish           | % Fem.         | No. fish          | % Fem.         | No. fish          | % Fem. <sup>a</sup> | No. fish           | % Fem. <sup>a</sup> |
| 2018 <sup>j</sup>   | 4,114                           | 25     | -                  | -                  | - <sup>d</sup> | -                 | -              | 4,227             | 55                  | 4,053              | 56                  |
| 2019 <sup>j</sup>   | 5,111                           | 34     | -                  | 438                | 61             | 1,328             | 24             | 2,018             | - <sup>k</sup>      | 4,678              | 44                  |
| 2020 <sup>j,1</sup> | -                               | -      | -                  | -                  | -              | -                 | -              | - <sup>m</sup>    | -                   | -                  | -                   |
| 2021 <sup>j</sup>   | 1,418                           | 37     | -                  | 130                | 35             | -                 | - <sup>l</sup> | 1,417             | 41                  | 2,082              | 46                  |
| SEG <sup>n</sup>    | 2,100–4,900                     |        |                    |                    |                |                   |                |                   |                     |                    |                     |
| BEG <sup>o</sup>    |                                 |        |                    |                    |                |                   |                | 2,800–5,700       |                     | 3,300–6,500        |                     |
| Averages            |                                 |        |                    |                    |                |                   |                |                   |                     |                    |                     |
| 1986–2020           | 3,803                           | 37     | 1,946              | 1,020              | 40             | 2,128             | 30             | 6,173             | 44                  | 8,321              | 45                  |
| 2011–2020           | 4,002                           | 34     |                    | 1,193              | 45             | 1,482             | 27             | 4,335             | 42                  | 5,215              | 45                  |
| 2016–2020           | 3,718                           | 34     |                    | 823                | 50             | 1,269             | 26             | 4,278             | 41                  | 3,900              | 45                  |
| Minimum             | 1,148                           | 5      | 756                | 193                | 23             | 1,083             | 15             | 1,859             | 17                  | 2,675              | 28                  |
| Maximum             | 8,045                           | 64     | 4,766              | 2,391              | 61             | 4,023             | 49             | 13,390            | 66                  | 18,514             | 63                  |

*Note:* Minimum and maximum values exclude the most recent year data. No. = number; Fem. = female. Dashes indicate no survey or a value cannot be calculated.

<sup>a</sup> Adjustment factor was applied.

<sup>b</sup> Tower counts.

<sup>c</sup> Mark–recapture population estimate.

<sup>d</sup> Project operations were hindered by high water most of the season.

<sup>e</sup> Estimate includes an expansion for missed counting days based on average run timing.

<sup>f</sup> Weir count.

<sup>g</sup> Estimate includes an expansion for missed counting days based on using 2 DIDSON sonars to assess Chinook salmon passage.

<sup>h</sup> Due to high water, DIDSON sonar was used and preliminary species apportionment was estimated using average run timing.

<sup>i</sup> Final estimate uses a binomial mixed-effects model to create passage estimates for periods of missed counts.

<sup>j</sup> Preliminary.

<sup>k</sup> Only 8 fish were sampled for sex; value not presented due to low sample size.

<sup>l</sup> Projects did not operate due to COVID-19 or funding.

<sup>m</sup> Total escapement could not be determined. Sonar only operated 17 days due to flooding and debris.

<sup>n</sup> Sustainable Escapement Goal (SEG).

<sup>o</sup> Biological Escapement Goal (BEG).

Appendix B11.—Estimated run size and spawning escapement of Canadian-origin Yukon River mainstem Chinook salmon, 1982–2021.

| Year | Historic mark-recapture border passage estimate <sup>a</sup> | Eagle sonar estimate | U.S. harvest above Eagle sonar <sup>b</sup> | Canadian mainstem border passage estimate | Canadian mainstem harvest | Spawning escapement estimate <sup>c</sup> | Canadian origin total run size estimate <sup>d</sup> |
|------|--|----------------------|---|---|---------------------------|---|--|
| 1982 | 36,598   |                      |   | 60,346 <sup>e</sup>                       | 16,808                    | 43,538                                    | 147,587  |
| 1983 | 47,741   |                      |   | 63,227 <sup>e</sup>                       | 18,752                    | 44,475                                    | 160,221  |
| 1984 | 43,911 <sup>f</sup>  |                      |   | 66,300 <sup>e</sup>                       | 16,295                    | 50,005                                    | 111,035  |
| 1985 | 29,881   |                      |   | 59,586 <sup>e</sup>                       | 19,151                    | 40,435                                    | 145,359  |
| 1986 | 36,479   |                      |   | 61,489 <sup>e</sup>                       | 20,064                    | 41,425                                    | 159,082  |
| 1987 | 30,823   |                      |   | 58,870 <sup>e</sup>                       | 17,563                    | 41,307                                    | 174,128  |
| 1988 | 44,445   |                      |   | 61,026 <sup>e</sup>                       | 21,327                    | 39,699                                    | 145,675  |
| 1989 | 42,620   |                      |   | 77,718 <sup>e</sup>                       | 17,419                    | 60,299                                    | 164,516  |
| 1990 | 56,679   |                      |   | 78,192 <sup>e</sup>                       | 18,980                    | 59,212                                    | 151,188  |
| 1991 | 41,187   |                      |   | 63,172 <sup>e</sup>                       | 20,444                    | 42,728                                    | 124,382  |
| 1992 | 43,185   |                      |   | 56,958 <sup>e</sup>                       | 17,803                    | 39,155                                    | 154,219  |
| 1993 | 45,027   |                      |   | 52,713 <sup>e</sup>                       | 16,469                    | 36,244                                    | 131,528  |
| 1994 | 46,680   |                      |   | 77,219 <sup>e</sup>                       | 20,770                    | 56,449                                    | 172,885  |
| 1995 | 52,353   |                      |   | 70,761 <sup>e</sup>                       | 20,088                    | 50,673                                    | 168,502  |
| 1996 | 47,955   |                      |   | 93,606 <sup>e</sup>                       | 19,546                    | 74,060                                    | 182,564  |
| 1997 | 53,400   |                      |   | 69,538 <sup>e</sup>                       | 15,717                    | 53,821                                    | 161,700  |
| 1998 | 22,588   |                      |   | 41,335 <sup>e</sup>                       | 5,838                     | 35,497                                    | 88,282   |
| 1999 | 23,716   |                      |   | 49,538 <sup>e</sup>                       | 12,354                    | 37,184                                    | 110,446  |
| 2000 | 16,173   |                      |   | 30,699 <sup>e</sup>                       | 4,829                     | 25,870                                    | 52,842   |
| 2001 | 52,207   |                      |   | 62,338 <sup>e</sup>                       | 9,774                     | 52,564                                    | 85,663   |
| 2002 | 49,214   |                      |   | 51,428 <sup>g</sup>                       | 9,070                     | 42,358                                    | 81,486   |
| 2003 | 56,929   |                      |   | 90,040 <sup>g</sup>                       | 9,446                     | 80,594                                    | 149,980  |
| 2004 | 48,111   |                      |   | 59,415 <sup>g</sup>                       | 10,946                    | 48,469                                    | 117,246  |
| 2005 | 42,245   | 81,528               | 2,566                                       | 78,962 <sup>h</sup>                       | 10,977                    | 67,985                                    | 123,612  |
| 2006 | 36,748   | 73,691               | 2,303                                       | 71,388 <sup>h</sup>                       | 8,758                     | 62,630                                    | 119,485  |
| 2007 | 22,120   | 41,697               | 1,999                                       | 39,698 <sup>h</sup>                       | 4,794                     | 34,904                                    | 88,018   |
| 2008 | 14,666   | 38,097               | 815   | 37,282 <sup>h</sup>                       | 3,399                     | 33,883                                    | 62,611   |
| 2009 | -  | 69,957               | 382   | 69,575 <sup>h</sup>                       | 4,297                     | 65,278                                    | 87,221   |
| 2010 | -  | 35,074               | 604   | 34,470 <sup>h</sup>                       | 2,456                     | 32,014                                    | 59,741   |
| 2011 | -  | 51,271               | 370   | 50,901 <sup>h</sup>                       | 4,594                     | 46,307                                    | 71,725   |
| 2012 | -  | 34,747               | 91  | 34,656 <sup>h</sup>                       | 2,000                     | 32,656                                    | 48,498   |
| 2013 | -  | 30,725               | 152   | 30,573 <sup>h</sup>                       | 1,904                     | 28,669                                    | 37,177   |
| 2014 | -  | 63,482               | 51  | 63,431 <sup>h</sup>                       | 100                       | 63,331                                    | 64,886   |
| 2015 | -  | 84,015               | 341   | 83,674 <sup>h</sup>                       | 1,000                     | 82,674                                    | 87,323   |
| 2016 | -  | 72,329               | 762   | 71,567 <sup>h</sup>                       | 2,769                     | 68,798                                    | 83,043   |

-continued-

| Year      | Historic mark-recapture border passage estimate <sup>a</sup> | Eagle sonar estimate | U.S. harvest above Eagle sonar <sup>b</sup> | Canadian mainstem border passage estimate | Canadian mainstem harvest | Spawning escapement estimate <sup>c</sup> | Canadian origin total run size estimate <sup>d</sup> |
|-----------|--|----------------------|---|---|---------------------------|---|--|
| 2017      | -  | 73,313               | 1,498                                       | 71,815 <sup>h</sup>                       | 3,500                     | 68,315                                    | 92,622   |
| 2018      | -  | 57,893               | 629   | 57,264 <sup>h</sup>                       | 2,790                     | 54,474                                    | 76,530   |
| 2019      | -  | 45,560               | 744   | 44,816 <sup>h</sup>                       | 2,764                     | 42,052                                    | 72,620   |
| 2020      | -  | 33,550               | 220   | 33,330 <sup>h</sup>                       | 2,363                     | 30,967                                    | 45,501   |
| 2021      | -  | 31,796               | 38  | 31,758 <sup>h</sup>                       | 306                       | 31,452                                    | 32,970   |
| Averages  |  |                      |   |   |                           |   |  |
| 1982–2020 | 40,136   | 55,433               | 845   | 59,716                                    | 10,716                    | 49,000                                    | 111,824  |
| 2009–2020 | -  | 54,326               | 487   | 53,839                                    | 2,545                     | 51,295                                    | 68,907   |
| 2014–2020 | -  | 61,449               | 606   | 60,842                                    | 2,184                     | 58,659                                    | 74,646   |
| Minimum   | 14,666   | 30,725               | 51  | 30,573                                    | 100                       | 25,870                                    | 37,177   |
| Maximum   | 56,929   | 84,015               | 2,566                                       | 93,606                                    | 21,327                    | 82,674                                    | 182,564  |

Note: Minimum and maximum values exclude the most recent year data. Dashes indicate no survey or a value cannot be calculated.

<sup>a</sup> From 1982–2008, a mark–recapture program was used to determine border passage; fish were sampled and tagged near the border using fish wheels and sampled for marks/tags in upstream fisheries. The Eagle sonar project replaced the mark–recapture program in 2005.

<sup>b</sup> U.S. harvests between the sonar site and border prior to 2008 is unknown because subsistence harvest in the Eagle area extended above and below the sonar site but were most likely in the hundreds for Chinook salmon. Starting in 2008, subsistence harvests between the sonar site and the U.S./Canada border were recorded specifically for the purpose of estimating border passage.

<sup>c</sup> Canadian spawning escapement estimated as border passage minus Canadian harvest.

<sup>d</sup> Canadian total origin run size is estimated as the border passage plus the U.S. harvest of Canadian origin fish. In 1984, border passage was estimated using harvest and escapement estimate based on proportion of aerial surveys.

<sup>e</sup> Chinook salmon passage for Yukon mainstem at U.S./Canada border from 1982–2001 was reconstructed using a linear relationship with 3-area index (aerial surveys of Little Salmon, Big Salmon, and Nisutlin rivers in 2002–2007) plus Canadian harvests.

<sup>f</sup> In 1984, border passage was estimated using harvest and escapement estimates based on proportion of aerial surveys.

<sup>g</sup> Border passage estimated in 2002–2004 using escapement estimate from a radio tagging proportion study, plus Canadian harvest.

<sup>h</sup> Since 2005, border passage was estimated as fish counted by the Eagle sonar minus the U.S. harvest upriver from the sonar project.

Appendix B12.—Chinook salmon escapement counts for selected spawning areas in the Canadian (Yukon) portion of the Yukon River drainage, 1961–2021.

| Year | Tatchun<br>Creek <sup>a</sup> | Weirs          |                    | Sonars        |                   |                 |                |                    |                  | Whitehorse Fishway |                            |
|------|-------------------------------|----------------|--------------------|---------------|-------------------|-----------------|----------------|--------------------|------------------|--------------------|----------------------------|
|      |                               | Blind<br>Creek | Chandindu<br>River | Big<br>Salmon | Klondike<br>River | Teslin<br>River | Pelly<br>River | Porcupine<br>River | Takhini<br>River | Count              | % hatchery<br>contribution |
| 1961 |                               |                |                    |               |                   |                 |                |                    |                  | 1,068              | 0                          |
| 1962 |                               |                |                    |               |                   |                 |                |                    |                  | 1,500              | 0                          |
| 1963 |                               |                |                    |               |                   |                 |                |                    |                  | 483                | 0                          |
| 1964 |                               |                |                    |               |                   |                 |                |                    |                  | 595                | 0                          |
| 1965 |                               |                |                    |               |                   |                 |                |                    |                  | 903                | 0                          |
| 1966 |                               | <sup>b</sup>   |                    |               |                   |                 |                |                    |                  | 563                | 0                          |
| 1967 |                               |                |                    |               |                   |                 |                |                    |                  | 533                | 0                          |
| 1968 |                               |                |                    |               |                   |                 |                |                    |                  | 414                | 0                          |
| 1969 |                               |                |                    |               |                   |                 |                |                    |                  | 334                | 0                          |
| 1970 | 100                           |                |                    |               |                   |                 |                |                    |                  | 625                | 0                          |
| 1971 | 130                           |                |                    |               |                   |                 |                |                    |                  | 856                | 0                          |
| 1972 | 80                            |                |                    |               |                   |                 |                |                    |                  | 391                | 0                          |
| 1973 | 99                            |                |                    |               |                   |                 |                |                    |                  | 224                | 0                          |
| 1974 | 192                           |                |                    |               |                   |                 |                |                    |                  | 273                | 0                          |
| 1975 | 175                           |                |                    |               |                   |                 |                |                    |                  | 313                | 0                          |
| 1976 | 52                            |                |                    |               |                   |                 |                |                    |                  | 121                | 0                          |
| 1977 | 150                           |                |                    |               |                   |                 |                |                    |                  | 277                | 0                          |
| 1978 | 200                           |                |                    |               |                   |                 |                |                    |                  | 725                | 0                          |
| 1979 | 150                           |                |                    |               |                   |                 |                |                    |                  | 1,184              | 0                          |
| 1980 | 222                           |                |                    |               |                   |                 |                |                    |                  | 1,383              | 0                          |
| 1981 | 133                           |                |                    |               |                   |                 |                |                    |                  | 1,555              | 0                          |
| 1982 | 73                            |                |                    |               |                   |                 |                |                    |                  | 473                | 0                          |
| 1983 | 264                           |                |                    |               |                   |                 |                |                    |                  | 905                | 0                          |
| 1984 | 153                           |                |                    |               |                   |                 |                |                    |                  | 1,042              | 0                          |
| 1985 | 190                           |                |                    |               |                   |                 |                |                    |                  | 508                | 0                          |
| 1986 | 155                           |                |                    |               |                   |                 |                |                    |                  | 557                | 0                          |
| 1987 | 159                           |                |                    |               |                   |                 |                |                    |                  | 327                | 0                          |
| 1988 | 152                           |                |                    |               |                   |                 |                |                    |                  | 405                | 16                         |
| 1989 | 100                           |                |                    |               |                   |                 |                |                    |                  | 549                | 19                         |
| 1990 | 643                           |                |                    |               |                   |                 |                |                    |                  | 1,407              | 24                         |
| 1991 |                               |                |                    |               |                   |                 |                |                    |                  | 1,266 <sup>c</sup> | 51 <sup>c</sup>            |
| 1992 | 106                           |                |                    |               |                   |                 |                |                    |                  | 758 <sup>c</sup>   | 84 <sup>c</sup>            |

-continued-

## Appendix B12.–Page 2 of 3.

| Year              | Tatchun<br>Creek <sup>a</sup> | Weirs            |                    | Sonars        |                   |                    |                    |                    |                    | Whitehorse Fishway |                            |
|-------------------|-------------------------------|------------------|--------------------|---------------|-------------------|--------------------|--------------------|--------------------|--------------------|--------------------|----------------------------|
|                   |                               | Blind<br>Creek   | Chandindu<br>River | Big<br>Salmon | Klondike<br>River | Teslin<br>River    | Pelly<br>River     | Porcupine<br>River | Takhini<br>River   | Count              | % hatchery<br>contribution |
| 1993              | 183                           |                  |                    |               |                   |                    |                    |                    |                    | 668 <sup>c</sup>   | 73 <sup>c</sup>            |
| 1994              | 477                           |                  |                    |               |                   |                    |                    |                    |                    | 1,577 <sup>c</sup> | 54 <sup>c</sup>            |
| 1995              | 397                           | 826 <sup>d</sup> |                    |               |                   |                    |                    |                    |                    | 2,103              | 57                         |
| 1996              | 423                           |                  |                    |               |                   |                    |                    |                    |                    | 2,958              | 35                         |
| 1997              | 1,198                         | 957              |                    |               |                   |                    |                    |                    |                    | 2,084              | 24                         |
| 1998              | 405                           | 373              | 132                |               |                   |                    |                    |                    |                    | 777                | 95                         |
| 1999              | 252                           | 892              | 239                |               |                   |                    |                    |                    |                    | 1,118              | 74                         |
| 2000              | 276 <sup>e</sup>              |                  | 4 <sup>f</sup>     |               |                   |                    |                    |                    |                    | 677                | 69                         |
| 2001              |                               |                  | 129 <sup>g</sup>   |               |                   |                    |                    |                    |                    | 988                | 36                         |
| 2002              |                               |                  | <sup>h</sup>       |               |                   |                    |                    |                    |                    | 605                | 39                         |
| 2003              |                               | 1,155            | 185 <sup>i</sup>   |               |                   |                    |                    |                    |                    | 1,443              | 70                         |
| 2004              |                               | 792              |                    |               |                   |                    |                    |                    |                    | 1,989              | 76                         |
| 2005              |                               | 525              |                    | 5,618         |                   |                    |                    |                    |                    | 2,632              | 57                         |
| 2006              |                               | 677              |                    | 7,308         |                   |                    |                    |                    |                    | 1,720              | 47                         |
| 2007              |                               | 304              |                    | 4,506         |                   |                    |                    |                    |                    | 427                | 56                         |
| 2008              |                               | 276              |                    | 1,431         |                   |                    |                    |                    |                    | 399                | 54                         |
| 2009              |                               | 716              |                    | 9,261         | 5,147             |                    |                    |                    |                    | 828                | 47                         |
| 2010              |                               | 270              |                    | 3,817         | 803               |                    |                    |                    |                    | 672                | 49                         |
| 2011              |                               | 360              |                    | 5,156         | 1,181             |                    |                    |                    |                    | 1,534              | 48                         |
| 2012              |                               | 157              |                    | 2,584         |                   | 3,454 <sup>j</sup> |                    |                    |                    | 1,030              | 59                         |
| 2013              |                               | 312              |                    | 3,242         |                   | 9,916              |                    |                    |                    | 1,139              | 67                         |
| 2014              |                               | 602              |                    | 6,321         |                   | 17,507             |                    | 3,066              |                    | 1,601              | 78                         |
| 2015              |                               | 964              |                    | 10,078        |                   | 20,463             |                    | 4,851              |                    | 1,465              | 60                         |
| 2016              |                               | 664              |                    | 6,761         |                   |                    | 5,807 <sup>k</sup> | 6,665              |                    | 1,556              | 42                         |
| 2017              |                               | <sup>l</sup>     |                    | 5,672         |                   |                    | 9,081              | 1,191              | 1,872 <sup>k</sup> | 1,226              | 39                         |
| 2018              |                               | 612              |                    | 5,159         |                   |                    | 9,751              | 3,414              | 1,554              | 691                | 37                         |
| 2019              |                               |                  |                    | 3,874         |                   |                    | 6,927              | 4,740              |                    | 282                | 13                         |
| 2020              |                               |                  |                    | 1,635         | 470               |                    | 5,678              | <sup>m</sup>       |                    | 216                | 24                         |
| 2021 <sup>n</sup> | 17                            |                  |                    | 1,958         | 855               |                    | 4,980              | 409                | 247                | 274                | 36                         |

-continued-

Appendix B12.–Page 3 of 3.

| Year      | Tatchun<br>Creek <sup>a</sup> | Weirs          |                    | Sonars        |                   |                 |                |                    |                  | Whitehorse Fishway |                            |
|-----------|-------------------------------|----------------|--------------------|---------------|-------------------|-----------------|----------------|--------------------|------------------|--------------------|----------------------------|
|           |                               | Blind<br>Creek | Chandindu<br>River | Big<br>Salmon | Klondike<br>River | Teslin<br>River | Pelly<br>River | Porcupine<br>River | Takhini<br>River | Count              | % hatchery<br>contribution |
| Averages  |                               |                |                    |               |                   |                 |                |                    |                  |                    |                            |
| 1961–2020 | 243                           | 589            | 138                | 5,151         | 1,900             | 15,962          | 7,449          | -                  | -                | 949                | 28                         |
| 2011–2020 | -                             | 524            | -                  | 5,048         | 826               | 15,962          | 7,449          | 3,988              | -                | 1,074              | 47                         |
| 2016–2020 | -                             | 638            | -                  | 4,620         | -                 | -               | 7,449          | 4,003              | 1,713            | 794                | 31                         |
| Minimum   | 52                            | 157            | 4                  | 1,431         | 470               | 9,916           | 5,678          | 1,191              | 1,554            | 121                | 0                          |
| Maximum   | 1,198                         | 1,155          | 239                | 10,078        | 5,147             | 20,463          | 9,751          | 6,665              | 1,872            | 2,958              | 95                         |

*Note:* Minimum and maximum values exclude the most recent year data. Dashes indicate a value cannot be calculated. Unless otherwise noted, blank cells indicate years when the project was not operated.

<sup>a</sup> All aerial surveys prior to 1980, subsequently foot surveys except 1982 and 1986 (aerial), and weir counts from 1997–2000 and 2021.

<sup>b</sup> Incomplete and/or poor survey conditions resulting in minimal or inaccurate counts (Tatchun 1966 only 7 fish observed).

<sup>c</sup> Counts and estimated percentages may be slightly exaggerated. In some or all of these years, a number of adipose-clipped fish ascended the Fishway and were counted more than once. These fish would have been released into the Fishway as fry between 1989 and 1994, inclusive.

<sup>d</sup> Details lacking; no reported data beyond annual passage estimate

<sup>e</sup> Flood conditions caused early termination of this program.

<sup>f</sup> High water delayed project installation, therefore counts are incomplete.

<sup>g</sup> Weir was breached from July 31–August 7 due to high water.

<sup>h</sup> Resistance board weir (RBW) tested for 3 weeks.

<sup>i</sup> Combination RBW and conduit weir tested and operational from July 10–30.

<sup>j</sup> Chinook counted on the left bank due to high water; estimate should be considered a minimum

<sup>k</sup> Sonar feasibility year.

<sup>l</sup> High water conditions prevented weir operation.

<sup>m</sup> Project cancelled due to COVID-19.

<sup>n</sup> Data are preliminary.

Appendix B13.—Summer chum salmon escapement counts for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1973–2021.

| Year | Andreafsky River                          |                      |                     | Anvik River                   |           | Rodo River          | Kaltag River | Nulato River          |                         |                      |
|------|---|----------------------|---------------------|-------------------------------|-----------|---------------------|--------------|-----------------------|-------------------------|----------------------|
|      | East Fork                                 |                      | West Fork           |                               |           |                     |              | South Fork            | North Fork <sup>a</sup> | Mainstem             |
|      | Sonar, tower, or weir counts <sup>c</sup> |                      |                     | Tower and aerial <sup>d</sup> |           |                     |              |                       |                         |                      |
|      | Aerial <sup>b</sup>                       |                      | Aerial <sup>b</sup> |                               | Sonar     | Aerial <sup>b</sup> | Tower        | Aerial <sup>b</sup>   | Aerial <sup>b</sup>     | Tower                |
| 1973 | 10,149 <sup>e</sup>                       |                      | 51,835              | 249,015                       | -         |                     |              |                       |                         |                      |
| 1974 | 3,215 <sup>e</sup>                        |                      | 33,578              | 411,133                       | -         | 16,137              |              | 29,016                | 29,334                  |                      |
| 1975 | 223,485                                   |                      | 235,954             | 900,967                       | -         | 25,335              |              | 51,215                | 87,280                  |                      |
| 1976 | 105,347                                   |                      | 118,420             | 511,475                       | -         | 38,258              |              | 9,230 <sup>e</sup>    | 30,771                  |                      |
| 1977 | 112,722                                   |                      | 63,120              | 358,771                       | -         | 16,118              |              | 11,385                | 58,275                  |                      |
| 1978 | 127,050                                   |                      | 57,321              | 307,270                       | -         | 17,845              |              | 12,821                | 41,659                  |                      |
| 1979 | 66,471                                    |                      | 43,391              | -                             | 277,712   | -                   |              | 1,506                 | 35,598                  |                      |
| 1980 | 36,823 <sup>e</sup>                       |                      | 114,759             | -                             | 482,181   | -                   |              | 3,702 <sup>e</sup>    | 11,244 <sup>e</sup>     |                      |
| 1981 | 81,555                                    | 152,665              | -                   | -                             | 1,479,582 | -                   |              | 14,348                | -                       |                      |
| 1982 | 7,501 <sup>e</sup>                        | 181,352              | 7,267 <sup>e</sup>  | -                             | 444,581   | -                   |              | -                     | -                       |                      |
| 1983 | -   | 113,328              | -                   | -                             | 362,912   | -                   |              | 1,263 <sup>e</sup>    | 19,749                  |                      |
| 1984 | 95,200 <sup>e</sup>                       | 72,598               | 238,565             | -                             | 891,028   | -                   |              | -                     | -                       |                      |
| 1985 | 66,146                                    | -                    | 52,750              | -                             | 1,080,243 | 24,576              |              | 10,494                | 19,344                  |                      |
| 1986 | 83,931                                    | 152,730              | 99,373              | -                             | 1,085,750 | -                   |              | 16,848                | 47,417                  |                      |
| 1987 | 6,687 <sup>e</sup>                        | 45,221               | 35,535              | -                             | 455,876   | -                   |              | 4,094                 | 7,163                   |                      |
| 1988 | 43,056                                    | 68,937               | 45,432              | -                             | 1,125,449 | 13,872              |              | 15,132                | 26,951                  |                      |
| 1989 | 21,460 <sup>e</sup>                       | -                    | -                   | -                             | 636,906   | -                   |              | -                     | -                       |                      |
| 1990 | 11,519 <sup>e</sup>                       | -                    | 20,426 <sup>e</sup> | -                             | 403,627   | 1,941 <sup>e</sup>  |              | 3,196 <sup>e, f</sup> | 1,419 <sup>e</sup>      |                      |
| 1991 | 31,886                                    | -                    | 46,657              | -                             | 847,772   | 3,977               |              | 13,150                | 12,491                  |                      |
| 1992 | 11,308 <sup>e</sup>                       | -                    | 37,808 <sup>e</sup> | -                             | 775,626   | 4,465               |              | 5,322                 | 12,358                  |                      |
| 1993 | 10,935 <sup>e</sup>                       | -                    | 9,111 <sup>e</sup>  | -                             | 517,409   | 7,867               |              | 5,486                 | 7,698                   |                      |
| 1994 | -   | 200,981 <sup>g</sup> | -                   | -                             | 1,124,689 | -                   | 47,295       | -                     | -                       | 148,762 <sup>g</sup> |
| 1995 | -   | 172,148              | -                   | -                             | 1,339,418 | 12,849              | 77,193       | 10,875                | 29,949                  | 236,890              |
| 1996 | -   | 108,450              | -                   | -                             | 933,240   | 4,380               | 51,269       | 8,490 <sup>e</sup>    | -                       | 129,694              |
| 1997 | -   | 51,139               | -                   | -                             | 605,751   | 2,775 <sup>e</sup>  | 48,018       | -                     | -                       | 157,975              |
| 1998 | -   | 67,720               | -                   | -                             | 487,300   | -                   | 8,113        | -                     | -                       | 49,140               |
| 1999 | -   | 32,587               | -                   | -                             | 437,355   | -                   | 5,339        | -                     | -                       | 30,076               |
| 2000 | 2,094 <sup>e</sup>                        | 24,785               | 18,989 <sup>e</sup> | -                             | 196,350   | -                   | 6,727        | -                     | -                       | 24,308               |

-continued-



## Appendix B13.—Page 2 of 5.

| Year              | Andreafsky River    |   |                     | Anvik River                   |                 | Rodo River          | Kaltag River | Nulato River        |                         |                     |
|-------------------|---------------------|---|---------------------|-------------------------------|-----------------|---------------------|--------------|---------------------|-------------------------|---------------------|
|                   | East Fork           |   | West Fork           | Tower and aerial <sup>d</sup> | Sonar           | Aerial <sup>b</sup> | Tower        | South Fork          | North Fork <sup>a</sup> | Mainstem            |
|                   | Aerial <sup>b</sup> | Sonar, tower, or weir counts <sup>c</sup> | Aerial <sup>b</sup> |                               |                 |                     |              | Aerial <sup>b</sup> | Aerial <sup>b</sup>     | Tower               |
| 2001              | -                   | 2,134 <sup>g</sup>                        | -                   | -                             | 224,059         | -                   | -            | -                   | -                       | -                   |
| 2002              | -                   | 44,194                                    | -                   | -                             | 459,058         | -                   | 13,583       | -                   | -                       | 72,232              |
| 2003              | -                   | 22,461                                    | -                   | -                             | 256,920         | -                   | 3,056        | -                   | -                       | 19,590 <sup>g</sup> |
| 2004              | -                   | 64,883                                    | -                   | -                             | 365,354         | -                   | 5,247        | -                   | -                       | -                   |
| 2005              | -                   | 20,127                                    | -                   | -                             | 525,392         | -                   | 22,093       | -                   | -                       | -                   |
| 2006              | 3,100 <sup>e</sup>  | 102,260                                   | 617                 | -                             | 605,487         | -                   | -            | 7,772               | 11,658                  | -                   |
| 2007              | -                   | 69,642                                    | -                   | -                             | 459,038         | -                   | -            | 21,825              | 15,277                  | -                   |
| 2008              | 9,300               | 57,259                                    | 25,850              | -                             | 374,933         | -                   | -            | 12,070              | 10,715                  | -                   |
| 2009              | 736                 | 8,770                                     | 3,877               | -                             | 193,098         | 621                 | -            | 2,120               | 567                     | -                   |
| 2010              | 1,982               | 72,893                                    | 24,380              | -                             | 396,174         | -                   | -            | 1,891               | 1,038                   | -                   |
| 2011              | 12,889              | 100,473                                   | 10,020              | -                             | 642,529         | 6,011               | -            | 9,454               | 8,493                   | -                   |
| 2012              | -                   | 56,680                                    | -                   | -                             | 484,091         | 15,606              | -            | 20,600              | 14,948                  | -                   |
| 2013              | 10,965              | 61,234                                    | 9,685               | 38,915                        | 577,876         | -                   | -            | 13,695              | 13,230                  | -                   |
| 2014              | -                   | 37,793                                    | 9,650               | 54,061                        | 399,796         | -                   | -            | -                   | -                       | -                   |
| 2015              | 6,004 <sup>e</sup>  | 48,809                                    | 2,837 <sup>e</sup>  | 36,871                        | 374,968         | 3,685               | -            | 4,102               | 9,525                   | -                   |
| 2016              | -                   | 50,362                                    | -                   | -                             | 337,821         | -                   | -            | -                   | -                       | -                   |
| 2017              | -                   | 55,532                                    | 11,655              | 38,191                        | 415,139         | -                   | -            | 4,890               | 7,882                   | -                   |
| 2018              | 16,206              | 36,330                                    | 13,837              | 30,309                        | 305,098         | -                   | -            | 3,930               | 1,164                   | -                   |
| 2019              | 26,048              | 49,881                                    | 17,198              | 15,499                        | 249,014         | -                   | -            | 2,612               | 4,898                   | -                   |
| 2020              | 10,628              | -   | 9,932               | 8,461                         | -               | -                   | -            | 861                 | 722                     | -                   |
| 2021 <sup>1</sup> | -                   | 2,531                                     | -                   | -                             | 18,819          | -                   | -            | -                   | -                       | -                   |
| Goal <sup>h</sup> |                     | >40,000                                   |                     |                               | 350,000–700,000 |                     |              |                     |                         |                     |
| Average           |                     |   |                     |                               |                 |                     |              |                     |                         |                     |
| 1973–2020         | 40,529              | 72,920                                    | 47,414              | 227,764                       | 576,502         | 12,018              | 26,176       | 10,419              | 19,294                  | 96,519              |
| 2011–2020         | 13,790              | 55,233                                    | 10,602              | 31,758                        | 420,704         | 8,434               | -            | 7,518               | 7,608                   | -                   |
| 2016–2020         | 17,627              | 48,026                                    | 13,156              | 23,115                        | 326,768         | -                   | -            | 3,073               | 3,667                   | -                   |
| Minimum           | 736                 | 2,134                                     | 617                 | 8,461                         | 193,098         | 621                 | 3,056        | 861                 | 567                     | 19,590              |
| Maximum           | 223,485             | 200,981                                   | 238,565             | 900,967                       | 1,479,582       | 38,258              | 77,193       | 51,215              | 87,280                  | 236,890             |

-continued-

## Appendix B13.—Page 3 of 5.

| Year | Henshaw<br>Creek | Gisasa River        |                     | Hogatza River          |                  | Tozitna<br>River                | Chena River         |                     | Salcha River        |        |
|------|------------------|---------------------|---------------------|------------------------|------------------|---------------------------------|---------------------|---------------------|---------------------|--------|
|      | Weir             | Aerial <sup>b</sup> | Weir                | Clear &<br>Caribou Cr. | Clear<br>Creek   | Weir and<br>Aerial <sup>b</sup> | Aerial <sup>b</sup> | Tower               | Aerial <sup>b</sup> | Tower  |
| 1973 |                  |                     |                     |                        |                  |                                 | 79 <sup>e</sup>     |                     | 290                 |        |
| 1974 |                  | 22,022              |                     |                        |                  | 1,823                           | 4,349               |                     | 3,510               |        |
| 1975 |                  | 56,904              |                     | 22,355                 |                  | 3,512                           | 1,670               |                     | 7,573               |        |
| 1976 |                  | 21,342              |                     | 20,744                 |                  | 725 <sup>e</sup>                | 685                 |                     | 6,484               |        |
| 1977 |                  | 2,204 <sup>e</sup>  |                     | 10,734                 |                  | 761 <sup>e</sup>                | 610                 |                     | 677 <sup>e</sup>    |        |
| 1978 |                  | 9,280 <sup>e</sup>  |                     | 5,102                  |                  | 2,262                           | 1,609               |                     | 5,405               |        |
| 1979 |                  | 10,962              |                     | 14,221                 |                  | -                               | 1,025 <sup>e</sup>  |                     | 3,060               |        |
| 1980 |                  | 10,388              |                     | 19,786                 |                  | 580                             | 338                 |                     | 4,140               |        |
| 1981 |                  | -                   |                     | -                      |                  | -                               | 3,500               |                     | 8,500               |        |
| 1982 |                  | 334 <sup>e</sup>    |                     | 4,984 <sup>e</sup>     |                  | 874                             | 1,509               |                     | 3,756               |        |
| 1983 |                  | 2,356 <sup>e</sup>  |                     | 28,141                 |                  | 1,604                           | 1,097               |                     | 716 <sup>e</sup>    |        |
| 1984 |                  | -                   |                     | 184 <sup>e</sup>       |                  | -                               | 1,861               |                     | 9,810               |        |
| 1985 |                  | 13,232              |                     | 22,566                 |                  | 1,030                           | 1,005               |                     | 3,178               |        |
| 1986 |                  | 12,114              |                     | -                      |                  | 1,778                           | 1,509               |                     | 8,028               |        |
| 1987 |                  | 2,123               |                     | 5,669 <sup>e</sup>     |                  | -                               | 333                 |                     | 3,657               |        |
| 1988 |                  | 9,284               |                     | 6,890                  |                  | 2,983                           | 432                 |                     | 2,889 <sup>e</sup>  |        |
| 1989 |                  | -                   |                     | -                      |                  | -                               | 714 <sup>e</sup>    |                     | 1,574 <sup>e</sup>  |        |
| 1990 |                  | 450 <sup>e</sup>    |                     | 2,177 <sup>e</sup>     |                  | 36                              | 245 <sup>e</sup>    |                     | 450 <sup>e</sup>    |        |
| 1991 |                  | 7,003               |                     | 9,947                  |                  | 93                              | 115 <sup>e</sup>    |                     | 154 <sup>e</sup>    |        |
| 1992 |                  | 9,300               |                     | 2,986                  |                  | 794                             | 848 <sup>e</sup>    |                     | 3,222               |        |
| 1993 |                  | 1,581               |                     | -                      |                  | 970                             | 168                 | 5,483               | 212                 | 5,809  |
| 1994 |                  | 6,827               | 51,116 <sup>g</sup> | 8,247 <sup>i</sup>     |                  | -                               | 1,137               | 9,984               | 4,916               | 39,450 |
| 1995 |                  | 6,458               | 136,886             | -                      | 116,735          | 4,985                           | 185 <sup>e</sup>    | 3,519 <sup>g</sup>  | 934 <sup>e</sup>    | 30,784 |
| 1996 |                  | -                   | 158,752             | 27,090 <sup>i</sup>    | 100,912          | 2,310                           | 2,061               | 12,810 <sup>g</sup> | 9,722               | 74,827 |
| 1997 |                  | 686 <sup>e</sup>    | 31,800              | 1,821 <sup>e</sup>     | 76,454           | 428 <sup>e</sup>                | 594 <sup>e</sup>    | 9,439 <sup>g</sup>  | 3,968 <sup>e</sup>  | 35,741 |
| 1998 |                  | -                   | 21,142              | 120 <sup>e</sup>       | 212 <sup>g</sup> | 7 <sup>e</sup>                  | 24 <sup>e</sup>     | 5,901               | 370 <sup>e</sup>    | 17,289 |
| 1999 |                  | -                   | 10,155              | -                      | 11,283           | -                               | 520                 | 9,165               | 150                 | 23,221 |
| 2000 | 24,457           | -                   | 11,410              | -                      | 19,376           | 480                             | 105                 | 3,515               | 228                 | 20,516 |
| 2001 | 34,777           | -                   | 17,946              | -                      | 3,674            | 12,527                          | 2                   | 4,773               | -                   | 14,900 |

-continued-

## Appendix B13.–Page 4 of 5.

|                   | Henshaw<br>Creek | Gisasa River        |         | Hogatza River          |                     | Tozitna<br>River                | Chena River         |                     | Salcha River        |                     |
|-------------------|------------------|---------------------|---------|------------------------|---------------------|---------------------------------|---------------------|---------------------|---------------------|---------------------|
|                   |                  |                     |         | Clear &<br>Caribou Cr. | Clear<br>Creek      |                                 |                     |                     |                     |                     |
| Year              | Weir             | Aerial <sup>b</sup> | Weir    | Aerial <sup>b</sup>    | Tower               | Weir and<br>Aerial <sup>b</sup> | Aerial <sup>b</sup> | Tower               | Aerial <sup>b</sup> | Tower               |
| 2002              | 25,249           | -                   | 33,481  | -                      | 13,150              | 18,789                          | -                   | 1,021 <sup>g</sup>  | 78                  | 27,012 <sup>j</sup> |
| 2003              | 21,400           | -                   | 25,999  | -                      | 6,159               | 8,487                           | -                   | 573 <sup>g</sup>    | -                   | -                   |
| 2004              | 86,474           | -                   | 37,851  | -                      | 15,661              | 25,003                          | -                   | 15,163 <sup>g</sup> | -                   | 47,861              |
| 2005              | 237,481          | -                   | 172,259 | -                      | 26,420              | 39,700                          | 219                 | 16,873 <sup>g</sup> | 4,320               | 194,933             |
| 2006              | -                | 1,000               | 261,306 | -                      | 29,166 <sup>j</sup> | 22,629                          | 469                 | 35,109 <sup>g</sup> | 152                 | 113,960             |
| 2007              | 44,425           | -                   | 46,257  | -                      | 6,029 <sup>j</sup>  | 8,470                           | -                   | 4,999               | 4 <sup>e</sup>      | 13,069              |
| 2008              | 96,731           | 20,470              | 36,938  | -                      | -                   | 9,133                           | 37                  | 1,300 <sup>g</sup>  | 0 <sup>e</sup>      | 2,213 <sup>g</sup>  |
| 2009              | 156,933          | 1,060               | 25,904  | 3,981                  | -                   | 8,434                           | -                   | 16,516              | -                   | 31,035              |
| 2010              | 105,398          | 1,096               | 47,669  | 840                    | -                   | -                               | -                   | 7,561               | -                   | 22,185              |
| 2011              | 248,247          | 13,228              | 95,796  | 3,665                  | -                   | 11,351                          | 4,600               | -                   | 1,154               | 66,564 <sup>k</sup> |
| 2012              | 292,082          | -                   | 83,423  | 23,022                 | -                   | 11,045                          | 1,180               | 6,882               | -                   | 46,252              |
| 2013              | 285,008          | 9,300 <sup>e</sup>  | 80,055  | -                      | -                   | -                               | 135 <sup>e</sup>    | 21,372              | -                   | 60,981              |
| 2014              | -                | -                   | 32,523  | -                      | -                   | -                               | 1,317               | 13,303 <sup>g</sup> | 1993 <sup>e</sup>   | - <sup>g</sup>      |
| 2015              | 238,529          | 5,601               | 42,747  | 6,080                  | -                   | -                               | -                   | 8,620               | 0 <sup>e</sup>      | 12,812              |
| 2016              | 286,780          | -                   | 66,670  | -                      | -                   | -                               | -                   | 6,493 <sup>g</sup>  | -                   | 2,897 <sup>g</sup>  |
| 2017              | 360,687          | -                   | 73,584  | -                      | -                   | -                               | -                   | 21,156 <sup>g</sup> | -                   | 29,093 <sup>g</sup> |
| 2018              | -                | 8,058               | -       | 3,307                  | -                   | -                               | -                   | 13,084 <sup>g</sup> | -                   | 22,782 <sup>g</sup> |
| 2019              | 34,342           | -                   | 19,099  | -                      | -                   | -                               | -                   | 2,704               | -                   | 2,117               |
| 2020              | -                | 754                 | -       | -                      | -                   | -                               | -                   | 357 <sup>g</sup>    | -                   | -                   |
| 2021 <sup>i</sup> | 3,729            | -                   | -       | -                      | -                   | -                               | -                   | 578                 | -                   | 2,193               |
| Goal <sup>h</sup> |                  |                     |         |                        |                     |                                 |                     |                     |                     |                     |
| Average           |                  |                     |         |                        |                     |                                 |                     |                     |                     |                     |
| 1973–2020         | 151,706          | 9,152               | 64,831  | 10,186                 | 32,710              | 6,568                           | 1,008               | 9,544               | 2,924               | 38,332              |
| 2011–2020         | 249,382          | 7,388               | 61,737  | 9,019                  | -                   | 11,198                          | 1,808               | 10,441              | 1,049               | 30,437              |
| 2016–2020         | 227,270          | 4,406               | 53,118  | 3,307                  | -                   | -                               | -                   | 8,759               | -                   | 14,222              |
| Minimum           | 21,400           | 334                 | 10,155  | 120                    | 212                 | 7                               | 2                   | 357                 | 0                   | 2,117               |
| Maximum           | 360,687          | 56,904              | 261,306 | 28,141                 | 116,735             | 39,700                          | 4,600               | 35,109              | 9,810               | 194,933             |

-continued-

*Note:* Unless otherwise noted blank cells indicate years prior to the project being operational. Dashes indicate years in which no information was collected. Minimum and maximum values exclude the most recent year data.

- <sup>a</sup> Includes mainstem counts below the confluence of the North and South Forks, unless otherwise noted.
- <sup>b</sup> Aerial survey counts are peak counts only, survey rating is fair or good unless otherwise noted.
- <sup>c</sup> East Fork Andreafsky passage estimated with sonar 1981–1984, tower counts 1986–1988; weir counts from 1994–present. The project did not operate in 1985, 1989–1993 and 2020.
- <sup>d</sup> From 1972–1979, counting tower operated; escapement estimate listed is the tower counts plus expanded aerial survey counts below the tower.
- <sup>e</sup> Incomplete survey and/or poor survey timing or conditions resulted in minimal or inaccurate count.
- <sup>f</sup> Mainstem counts below the confluence of the North and South Forks of the Nulato River included in the South Fork counts.
- <sup>g</sup> Incomplete count due to late installation and/or early removal of project or high water events.
- <sup>h</sup> Biological escapement goal (Andreafsky) or sustainable escapement goal (Anvik).
- <sup>i</sup> Bureau of Land management helicopter survey.
- <sup>j</sup> Project operated as a video monitoring system.
- <sup>k</sup> Estimate includes an expansion for missed counting days based on average run timing. Minimum documented abundance from successful counting days was 30,411 (standard error not reported).
- <sup>l</sup> Data are preliminary.

Appendix B14.—Fall chum salmon abundance estimates or escapement estimates for selected spawning areas in the U.S. (Alaska) portions of the Yukon River drainage, 1971–2021.

| Year | Yukon<br>River<br>mainstem<br>sonar<br>estimate <sup>a</sup> | Tanana River drainage        |  |                             |                                       | Upper Yukon River drainage                                  |   |                                |
|------|--|------------------------------|--|-----------------------------|---------------------------------------|---|---|--------------------------------|
|      |  | Toklat<br>River <sup>b</sup> | Kantishna<br>River<br>abundance<br>estimate <sup>c</sup> | Delta<br>River <sup>d</sup> | Bluff<br>Cabin<br>Slough <sup>e</sup> | Upper Tanana<br>River<br>abundance<br>estimate <sup>f</sup> | Teedriinjik-<br>Chandalar<br>River <sup>g</sup> | Sheenjek<br>River <sup>h</sup> |
| 1971 |  |                              |  |                             |                                       |   |   |                                |
| 1972 |  |                              |  | 5,384 <sup>i</sup>          |                                       |   |   |                                |
| 1973 |  |                              |  | 10,469 <sup>i</sup>         |                                       |   |   |                                |
| 1974 |  | 41,798                       |  | 5,915 <sup>i</sup>          |                                       |   |   | 117,921 <sup>j</sup>           |
| 1975 |  | 92,265                       |  | 3,734                       |                                       |   |   | 227,935 <sup>j</sup>           |
| 1976 |  | 52,891                       |  | 6,312                       |                                       |   |   | 34,649 <sup>j</sup>            |
| 1977 |  | 34,887                       |  | 16,876                      |                                       |   |   | 59,878 <sup>j</sup>            |
| 1978 |  | 37,001                       |  | 11,136 <sup>i</sup>         |                                       |   |   | 42,661 <sup>j</sup>            |
| 1979 |  | 158,336                      |  | 8,355 <sup>i</sup>          |                                       |   |   | 120,129 <sup>j</sup>           |
| 1980 |  | 26,346 <sup>k</sup>          |  | 5,137 <sup>i</sup>          | 3,190 <sup>l</sup>                    |   |   | 38,093 <sup>j</sup>            |
| 1981 |  | 15,623                       |  | 23,508 <sup>i</sup>         | 6,120 <sup>l</sup>                    |   |   | 102,137 <sup>m</sup>           |
| 1982 |  | 3,624                        |  | 4,235 <sup>i</sup>          | 1,156                                 |   |   | 43,042 <sup>m</sup>            |
| 1983 |  | 21,869                       |  | 7,705 <sup>i</sup>          | 12,715                                |   |   | 64,989 <sup>m</sup>            |
| 1984 |  | 16,758                       |  | 12,411 <sup>i</sup>         | 4,017                                 |   |   | 36,173 <sup>m</sup>            |
| 1985 |  | 22,750                       |  | 17,276                      | 2,655 <sup>l</sup>                    |   |   | 179,727 <sup>m, n</sup>        |
| 1986 |  | 17,976                       |  | 6,703                       | 3,458                                 |   | 59,313  | 84,207 <sup>n, o</sup>         |
| 1987 |  | 22,117                       |  | 21,180 <sup>i</sup>         | 9,395                                 |   | 52,416  | 153,267 <sup>n, o</sup>        |
| 1988 |  | 13,436                       |  | 18,024 <sup>i</sup>         | 4,481 <sup>l</sup>                    |   | 33,619  | 45,206 <sup>o</sup>            |
| 1989 |  | 30,421                       |  | 21,342                      | 5,386 <sup>l</sup>                    |   | 69,161  | 99,116 <sup>o</sup>            |
| 1990 |  | 34,739                       |  | 8,992                       | 1,632                                 |   | 78,631  | 77,750 <sup>o</sup>            |
| 1991 |  | 13,347                       |  | 32,905                      | 7,198                                 |   |   | 86,496 <sup>p</sup>            |
| 1992 |  | 14,070                       |  | 8,893                       | 3,615 <sup>l</sup>                    |   |   | 78,808                         |
| 1993 |  | 27,838                       |  | 19,857 <sup>i</sup>         | 5,550 <sup>l</sup>                    |   |   | 42,922                         |
| 1994 |  | 76,057                       |  | 23,777                      | 2,277 <sup>l</sup>                    |   |   | 150,565                        |
| 1995 | 1,148,196  | 54,513 <sup>k</sup>          |  | 20,587 <sup>i</sup>         | 19,460                                | 268,173   | 323,586   | 241,855                        |
| 1996 | <sup>q</sup>   | 18,264                       |  | 19,758                      | 7,074 <sup>d</sup>                    | 134,563   | 230,450   | 246,889                        |
| 1997 | 579,767  | 14,511                       |  | 7,705                       | 5,707 <sup>d</sup>                    | 71,661  | 211,914   | 80,423 <sup>r</sup>            |
| 1998 | 375,222  | 15,605                       |  | 7,804                       | 3,549 <sup>d</sup>                    | 62,014  | 83,899  | 33,058                         |
| 1999 | 451,505  | 4,551                        | 27,199   | 16,534                      | 7,559 <sup>d</sup>                    | 97,843  | 92,685  | 14,229                         |

-continued-

## Appendix B14.—Page 2 of 4.

| Year                    | Yukon<br>River<br>mainstem<br>sonar<br>estimate <sup>a</sup> | Tanana River drainage        |  |                             |                                       | Upper Yukon River drainage                                  |   |                                |
|-------------------------|--|------------------------------|--|-----------------------------|---------------------------------------|---|---|--------------------------------|
|                         |  | Toklat<br>River <sup>b</sup> | Kantishna<br>River<br>abundance<br>estimate <sup>c</sup> | Delta<br>River <sup>d</sup> | Bluff<br>Cabin<br>Slough <sup>e</sup> | Upper Tanana<br>River<br>abundance<br>estimate <sup>f</sup> | Teedriinjik-<br>Chandalar<br>River <sup>g</sup> | Sheenjek<br>River <sup>h</sup> |
| 2000                    | 273,206  | 8,911                        | 21,450   | 3,001                       | 1,595                                 | 34,844  | 71,048  | 30,084 <sup>s</sup>            |
| 2001                    | 408,961  | 6,007 <sup>t</sup>           | 22,992   | 8,103                       | 1,808 <sup>l</sup>                    | 96,556 <sup>u</sup>   | 112,664   | 53,932                         |
| 2002                    | 367,886  | 28,519                       | 56,665   | 11,992                      | 3,116                                 | 109,961   | 94,472  | 31,642                         |
| 2003                    | 923,540  | 21,492                       | 87,359   | 22,582                      | 10,600 <sup>l</sup>                   | 193,418   | 221,343   | 44,047 <sup>v</sup>            |
| 2004                    | 633,368  | 35,480                       | 76,163   | 25,073                      | 10,270 <sup>l</sup>                   | 123,879   | 169,848   | 37,878                         |
| 2005                    | 1,894,078  | 17,779 <sup>j</sup>          | 107,719  | 28,132                      | 11,964 <sup>l</sup>                   | 337,755   | 526,838   | 561,863 <sup>n</sup>           |
| 2006                    | 964,238  |                              | 71,135   | 14,055                      |                                       | 202,669   | 254,778   | 160,178 <sup>n</sup>           |
| 2007                    | 740,195  |                              | 81,843   | 18,610                      |                                       | 320,811   | 243,805   | 65,435 <sup>n</sup>            |
| 2008                    | 636,525  |                              |  | 23,055                      | 1,198 <sup>l</sup>                    |   | 178,278   | 50,353 <sup>n</sup>            |
| 2009                    | <sup>q</sup>   |                              |  | 13,492                      | 2,900 <sup>l</sup>                    |   | <sup>q</sup>                                    | 54,126 <sup>n</sup>            |
| 2010                    | 458,103  |                              |  | 17,993                      | 1,610 <sup>l</sup>                    |   | 167,532   | 22,053                         |
| 2011                    | 873,877  |                              |  | 23,639                      | 2,655 <sup>l</sup>                    |   | 298,223   | 97,976 <sup>n</sup>            |
| 2012                    | 778,158  |                              |  | 9,377 <sup>e</sup>          |                                       |   | 205,791   | 104,701 <sup>n</sup>           |
| 2013                    | 865,295  | 9,161 <sup>l</sup>           |  | 31,955                      | 5,554 <sup>l</sup>                    |   | 252,710   |                                |
| 2014                    | 706,630  |                              |  | 32,480 <sup>e</sup>         | 4,095 <sup>l</sup>                    |   | 226,489   |                                |
| 2015                    | 669,483  | 8,422 <sup>l</sup>           |  | 33,401 <sup>e</sup>         | 6,020 <sup>l</sup>                    |   | 164,486   |                                |
| 2016                    | 994,760  | 16,885 <sup>l</sup>          |  | 21,913 <sup>e</sup>         | 4,936 <sup>l</sup>                    |   | 295,023   |                                |
| 2017                    | 1,829,931  |                              |  | 48,783 <sup>e</sup>         |                                       |   | 509,115   |                                |
| 2018                    | 928,664  | 19,141 <sup>l</sup>          |  | 39,641 <sup>e</sup>         | 5,554 <sup>l</sup>                    |   | 170,356   |                                |
| 2019                    | 842,041  |                              |  | 51,748 <sup>e</sup>         | 4,664 <sup>l</sup>                    |   | 116,323   |                                |
| 2020                    | 262,439  | 1,330 <sup>l</sup>           |  | 9,854 <sup>e</sup>          | 1,124 <sup>l</sup>                    |   |   |                                |
| 2021                    | <sup>w</sup> 146,172   |                              |  | 1,613                       | 1,085 <sup>l</sup>                    |   | 21,162  |                                |
| Escapement <sup>x</sup> | 300,000 <sup>y</sup>   |                              |  | 7,000 <sup>z</sup>          |                                       |   | 85,000 <sup>z</sup>                             |                                |
| Goal Ranges             | 600,000  |                              |  | 20,000                      |                                       |   | 234,000   |                                |
| Averages                |  |                              |  |                             |                                       |   |   |                                |
| 1971–2020               | 775,253  | 28,506                       | 61,392   | 17,579                      | 5,293                                 | 158,011   | 190,165   | 97,856                         |
| 2011–2020               | 875,128  | 10,988                       | -  | 30,279                      | 4,325                                 | -   | 248,724   | 101,339                        |
| 2016–2020               | 971,567  | 12,452                       | -  | 34,388                      | 4,070                                 | -   | 272,704   | -                              |
| Minimum                 | 262,439  | 1,330                        | 21,450   | 3,001                       | 1,124                                 | 34,844  | 33,619  | 14,229                         |
| Maximum                 | 1,894,078  | 158,336                      | 107,719  | 51,748                      | 19,460                                | 337,755   | 526,838   | 561,863                        |

-continued-

*Note:* Minimum and maximum values exclude the most recent year data. Dashes indicate a value cannot be calculated.

- <sup>a</sup> New model estimates generated in 2015 and applied to dataset back to 1995 and used since.
- <sup>b</sup> Expanded total abundance estimates for upper Toklat River index area using stream life curve (SLC) developed with 1987–1993 data. Index area includes Geiger Creek, Sushana River, and mainstem floodplain sloughs from approximately 0.25 mile upstream of roadhouse.
- <sup>c</sup> Fall chum salmon abundance estimate for the Kantishna and Toklat River drainages is based on a mark–recapture program. Number of tagging and recovery wheels changed over the years.
- <sup>d</sup> Population estimate generated from replicate foot surveys and stream life data (area under the curve method), unless otherwise indicated.
- <sup>e</sup> Peak foot survey, unless otherwise indicated.
- <sup>f</sup> Fall chum salmon abundance estimate for the upper Tanana River drainage is based on a mark–recapture program. Upper Tanana River consists of that portion upstream of the confluence with the Kantishna River. Number of tagging and recovery wheels changed over the years.
- <sup>g</sup> Single-beam sonar estimate for 1986–1990 (not used in run reconstruction), split-beam sonar estimate 1995–2006, DIDSON in use since 2007, project was aborted in 2009 and not operated in 2020. Sonar counts on the Teedriinjik are extrapolated after conclusion of the project through October 9 from 1995–present, with 2018 expanded to October 14 due to late run timing.
- <sup>h</sup> Single-beam sonar estimate beginning in 1981, split-beam sonar estimate 2003–2004, and DIDSON 2005–2012. Sonar counts on the Sheenjek River are extrapolated after conclusion of the project through October 9 from 2005–2012.
- <sup>i</sup> Estimates are a total spawner abundance, using migratory time density curves and stream life data.
- <sup>j</sup> Total escapement estimate using sonar to aerial survey expansion factor of 2.22.
- <sup>k</sup> Minimal estimate because of late timing of ground surveys with respect to peak of spawning.
- <sup>l</sup> Aerial survey count, unless otherwise indicated.
- <sup>m</sup> Project started late, estimated escapements expanded for portion missed using average run timing curves based on Teedriinjik (1986–1990) and Sheenjek (1991–1993) rivers.
- <sup>n</sup> Sonar counts include both banks in 1985–1987, 2005–2009, and 2011–2012.
- <sup>o</sup> Expanded estimates for period approximating second week of August through fourth week of September, using annual Chandalar River run timing data (1986–1990).
- <sup>p</sup> Total abundance estimates are for the period approximating second week of August through fourth week of September (1991–2012). Comparative escapement estimates before 1986 are considered more conservative; approximating the period end of August through September.
- <sup>q</sup> Project operated all or partial season, estimate was not useable.
- <sup>r</sup> Data interpolated due to high water from August 29–September 3, 1997 during buildup to peak passage on the Sheenjek River.
- <sup>s</sup> Sheenjek sonar project ended early (September 12) because of low water therefore estimate was expanded based on average run timing (62%).
- <sup>t</sup> Minimal estimate because Sushana River was breached by the main channel and uncountable.
- <sup>u</sup> Low numbers of tags deployed and recovered resulted in an estimate with an extremely large confidence interval (95% CI +/- 41,072).
- <sup>v</sup> Sheenjek sonar project ended on peak daily passages due to late run timing, estimate was expanded based on run timing (87%) at Rampart.
- <sup>w</sup> Data are preliminary.
- <sup>x</sup> Escapement Goals (EG) expressed as ranges.
- <sup>y</sup> Drainagewide escapement goal is related to mainstem passage estimate based on the sonar near Pilot Station minus upriver harvests.
- <sup>z</sup> Escapement goal revised to a sustainable escapement goal range in 2019 based on percentile method.

Appendix B15.–Fall chum salmon escapement estimates for selected spawning areas in Canadian (Yukon) portions of the Yukon River drainage, 1971–2021.

| Year | Porcupine Drainage                |                       | Mainstem                         |                            |                              |                              |
|------|-----------------------------------|-----------------------|----------------------------------|----------------------------|------------------------------|------------------------------|
|      | Fishing Branch River <sup>a</sup> | Porcupine River Sonar | Yukon River Index <sup>b,c</sup> | Koidern River <sup>b</sup> | Kluane River <sup>b, d</sup> | Teslin River <sup>b, e</sup> |
| 1971 | 312,800 <sup>f</sup>              |                       |                                  |                            |                              |                              |
| 1972 | 35,230 <sup>g</sup>               |                       |                                  |                            | 198 <sup>h,l</sup>           |                              |
| 1973 | 15,991                            |                       | 383                              |                            | 2,500                        |                              |
| 1974 | 31,841                            |                       |                                  |                            | 400                          |                              |
| 1975 | 353,282                           |                       | 7,671                            |                            | 362 <sup>h</sup>             |                              |
| 1976 | 36,584 <sup>f</sup>               |                       |                                  |                            | 20                           |                              |
| 1977 | 88,400 <sup>f</sup>               |                       |                                  |                            | 3,555                        |                              |
| 1978 | 40,800 <sup>f</sup>               |                       |                                  |                            | 0 <sup>h</sup>               |                              |
| 1979 | 119,898 <sup>f</sup>              |                       |                                  |                            | 4,640 <sup>h</sup>           |                              |
| 1980 | 55,268 <sup>f</sup>               |                       |                                  |                            | 3,150                        |                              |
| 1981 | 57,386 <sup>i</sup>               |                       |                                  |                            | 25,806                       |                              |
| 1982 | 15,901 <sup>f</sup>               |                       | 1,020 <sup>j</sup>               |                            | 5,378                        |                              |
| 1983 | 27,200 <sup>f</sup>               |                       | 7,560                            |                            | 8,578 <sup>h</sup>           |                              |
| 1984 | 15,150 <sup>f</sup>               |                       | 2,800 <sup>k</sup>               | 1,300                      | 7,200                        | 200                          |
| 1985 | 56,223                            |                       | 10,760                           | 1,195                      | 7,538                        | 356                          |
| 1986 | 31,811                            |                       | 825                              | 14                         | 16,686                       | 213                          |
| 1987 | 49,038                            |                       | 6,115                            | 50                         | 12,000                       |                              |
| 1988 | 23,645                            |                       | 1,550                            | 0                          | 6,950                        | 140                          |
| 1989 | 44,042                            |                       | 5,320                            | 40                         | 3,050                        | 210 <sup>l</sup>             |
| 1990 | 35,000 <sup>m</sup>               |                       | 3,651                            | 1                          | 4,683                        | 739                          |
| 1991 | 37,870                            |                       | 2,426                            | 53                         | 11,675                       | 468                          |
| 1992 | 22,539                            |                       | 4,438                            | 4                          | 3,339                        | 450                          |
| 1993 | 28,707                            |                       | 2,620                            | 0                          | 4,610                        | 555                          |
| 1994 | 65,247                            |                       | 1,429 <sup>j</sup>               | 20 <sup>j</sup>            | 10,734                       | 209 <sup>l</sup>             |
| 1995 | 51,971 <sup>n</sup>               |                       | 4,701                            | 0                          | 16,456                       | 633                          |
| 1996 | 77,302                            |                       | 4,977                            |                            | 14,431                       | 315                          |
| 1997 | 27,031                            |                       | 2,189                            |                            | 3,350                        | 207                          |
| 1998 | 13,687                            |                       | 7,292                            |                            | 7,337                        | 235                          |
| 1999 | 12,958                            |                       |                                  |                            | 5,136                        | 19 <sup>i</sup>              |
| 2000 | 5,057                             |                       | 933 <sup>l</sup>                 |                            | 1,442                        | 204                          |
| 2001 | 21,737                            |                       | 2,453                            |                            | 4,884                        | 5                            |
| 2002 | 13,636                            |                       | 973                              |                            | 7,147                        | 64                           |
| 2003 | 29,713                            |                       | 7,982                            |                            | 39,347                       | 390                          |
| 2004 | 20,417                            |                       | 3,440                            |                            | 18,982                       | 167                          |
| 2005 | 119,058                           |                       | 16,425                           |                            | 34,600                       | 585                          |
| 2006 | 30,954                            |                       | 6,553                            |                            | 18,208                       | 620                          |
| 2007 | 32,150                            |                       |                                  |                            |                              |                              |
| 2008 | 19,086 <sup>n</sup>               |                       |                                  |                            |                              |                              |
| 2009 | 25,828 <sup>o</sup>               |                       |                                  |                            |                              |                              |
| 2010 | 15,413 <sup>o</sup>               |                       |                                  |                            |                              |                              |
| 2011 | 13,085 <sup>n,o</sup>             | 14,640 <sup>q</sup>   |                                  |                            |                              |                              |
| 2012 | 22,399 <sup>o</sup>               | 33,496 <sup>q</sup>   |                                  |                            |                              |                              |
| 2013 | 25,376 <sup>p</sup>               | 35,615                |                                  |                            |                              |                              |
| 2014 | 7,304 <sup>p</sup>                | 17,244 <sup>r</sup>   |                                  |                            |                              |                              |
| 2015 | 8,351                             | 21,397                |                                  |                            |                              |                              |

-continued-



## Appendix B15.–Page 2 of 2.

| Year              | Porcupine Drainage                |                       | Mainstem                         |                            |                              |                              |
|-------------------|-----------------------------------|-----------------------|----------------------------------|----------------------------|------------------------------|------------------------------|
|                   | Fishing Branch River <sup>a</sup> | Porcupine River Sonar | Yukon River Index <sup>b,c</sup> | Koidern River <sup>b</sup> | Kluane River <sup>b, d</sup> | Teslin River <sup>b, e</sup> |
| 2016              | 29,397                            | 54,395                |                                  |                            |                              |                              |
| 2017              | 48,524                            | 67,818                |                                  |                            | 16,265 <sup>t</sup>          |                              |
| 2018              | 10,151                            |                       |                                  |                            | 1,734                        |                              |
| 2019              | 18,171                            | 27,447                |                                  |                            | 928                          |                              |
| 2020              | 4,795                             |                       | 323                              |                            | 120                          |                              |
| 2021 <sup>v</sup> | 2,413                             | 3,486                 | 1,131                            |                            | 64                           |                              |
| Goal <sup>w</sup> | 50,000-120,000                    |                       |                                  |                            |                              |                              |
| IMEG <sup>x</sup> | 22,000-49,000                     |                       |                                  |                            |                              |                              |
| Averages          |                                   |                       |                                  |                            |                              |                              |
| 1971–2020         | 46,068                            | 34,007                | 4,326                            | 223                        | 8,337                        | 317                          |
| 2011–2020         | 18,755                            | 34,007                | -                                | -                          | -                            | -                            |
| 2016–2020         | 22,208                            | 49,887                | -                                | -                          | 3,822                        | -                            |
| Minimum           | 4,795                             | 14,640                | 323                              | 0                          | 0                            | 5                            |
| Maximum           | 353,282                           | 67,818                | 16,425                           | 1,300                      | 39,347                       | 739                          |

Note: Minimum and maximum values exclude the most recent year data. Dashes indicate a value cannot be calculated.

<sup>a</sup> Weir count, unless otherwise indicated. Weir counts from 1972–1975, 1985–1989, 1991–1992, 1996–2012 were expanded to represent the remainder of the run after the project was terminated for the season through October 25.

<sup>b</sup> Aerial survey, unless otherwise indicated.

<sup>c</sup> Index area includes Tatchun Creek to Fort Selkirk.

<sup>d</sup> Index area includes Duke River to end of spawning sloughs below Swede Johnson Creek.

<sup>e</sup> Index area includes Boswell Creek area (5 km below to 5 km above confluence).

<sup>f</sup> Total escapement estimated using weir to aerial survey expansion factor of 2.72, unless otherwise indicated.

<sup>g</sup> Weir installed September 22. Estimate consists of weir count of 17,190 after September 22, and tagging passage estimate of 17,935 before weir installation.

<sup>h</sup> Foot survey, unless otherwise indicated.

<sup>i</sup> Initial aerial survey count doubled before applying the weir/aerial expansion factor of 2.72 because only half of the spawning area was surveyed.

<sup>j</sup> Boat survey.

<sup>k</sup> Total index area not surveyed. Survey included the mainstem Yukon River between Yukon Crossing to 30 km below Fort Selkirk.

<sup>l</sup> Incomplete and/or poor survey conditions resulting in minimal or inaccurate counts

<sup>m</sup> Weir not operated. Although only 7,541 chum salmon were counted on a single survey flown October 26, a population estimate of approximately 27,000 fish was made through date of survey, based upon historic average aerial-to-weir expansion of 28%. Actual population of spawners was reported by DFO as between 30,000–40,000 fish considering aerial survey timing.

<sup>n</sup> Incomplete count caused by late installation and/or early removal of project or high water events.

<sup>o</sup> Run timing was late and counts were expanded to represent the remainder of the run after the project was terminated for the season.

<sup>p</sup> Fishing Branch River weir did not operate, and escapement was estimated from a sonar operated on the upper Porcupine River minus Old Crow harvest and the proportion of radio tags to Fishing Branch River.

<sup>q</sup> Counts taken from corresponding R&E reports. Polynomial expansion calculated from last day of counts to Oct 14.

<sup>r</sup> Left bank estimate (15,363) was re-calculated post 2014 season after extensive review of 2014 sonar file data. The 2014 in season right bank estimate was deemed substandard and discarded. The 2014 post season estimate (1881) was calculated using the average proportion of right bank passage from 2015 and 2016.

<sup>s</sup> High water in August and early ice up prevented a complete passage estimate for Porcupine River fall chum salmon.

<sup>t</sup> Aerial surveys resumed following permanent diversion of Kluane Lake headwaters in 2016 by glacial retreat.

<sup>u</sup> Project cancelled due to COVID-19

<sup>v</sup> Data are preliminary

<sup>w</sup> Escapement goal in Pacific Salmon Treaty for Fishing Branch River fall chum salmon.

<sup>x</sup> Interim Management Escapement Goal (IMEG) established for 2010–2018 based on brood table of Canadian origin mainstem stocks (1982–2003).

Appendix B16.—Estimated spawning escapement of Canadian-origin mainstem Yukon River fall chum salmon, 1980–2021.

| Date | Eagle sonar estimate | Eagle sonar expanded estimate <sup>a</sup> | U.S. harvest above Eagle sonar <sup>b</sup> | U.S./Canada mainstem border passage estimate <sup>b</sup> | Canadian mainstem harvest | Spawning escapement estimate <sup>c</sup> |
|------|----------------------|--|---|---|---------------------------|---|
| 1980 |                      |  |   | 39,130  | 16,218                    | 22,912                                    |
| 1981 |                      |  |   | 66,347  | 19,281                    | 47,066 <sup>d</sup>                       |
| 1982 |                      |  |   | 47,049  | 15,091                    | 31,958                                    |
| 1983 |                      |  |   | 118,365   | 27,490                    | 90,875                                    |
| 1984 |                      |  |   | 81,900  | 25,267                    | 56,633 <sup>d</sup>                       |
| 1985 |                      |  |   | 99,775  | 37,765                    | 62,010                                    |
| 1986 |                      |  |   | 101,826   | 13,886                    | 87,940                                    |
| 1987 |                      |  |   | 125,121   | 44,345                    | 80,776                                    |
| 1988 |                      |  |   | 69,280  | 32,494                    | 36,786                                    |
| 1989 |                      |  |   | 55,861  | 20,111                    | 35,750                                    |
| 1990 |                      |  |   | 82,947  | 31,212                    | 51,735                                    |
| 1991 |                      |  |   | 112,303   | 33,842                    | 78,461                                    |
| 1992 |                      |  |   | 67,962  | 18,880                    | 49,082                                    |
| 1993 |                      |  |   | 42,165  | 12,422                    | 29,743                                    |
| 1994 |                      |  |   | 133,712   | 35,354                    | 98,358                                    |
| 1995 |                      |  |   | 198,203   | 40,111                    | 158,092                                   |
| 1996 |                      |  |   | 143,758   | 21,329                    | 122,429                                   |
| 1997 |                      |  |   | 94,725  | 9,306                     | 85,419                                    |
| 1998 |                      |  |   | 48,047  | 1,795                     | 46,252                                    |
| 1999 |                      |  |   | 72,188 <sup>e</sup>                                       | 13,636                    | 58,552                                    |
| 2000 |                      |  |   | 57,978 <sup>e</sup>                                       | 4,246                     | 53,732                                    |
| 2001 |                      |  |   | 38,769 <sup>e</sup>                                       | 5,278                     | 33,491                                    |
| 2002 |                      |  |   | 104,853 <sup>e</sup>                                      | 6,232                     | 98,621                                    |
| 2003 |                      |  |   | 153,656 <sup>e</sup>                                      | 10,523                    | 143,133                                   |
| 2004 |                      |  |   | 163,625 <sup>e</sup>                                      | 9,545                     | 154,080                                   |
| 2005 |                      |  |   | 451,477   | 13,979                    | 437,498                                   |
| 2006 | 236,386              | 245,290                                    | 17,775                                      | 227,515 <sup>f,g</sup>                                    | 6,617                     | 220,898                                   |
| 2007 | 235,871              | 265,008                                    | 18,691                                      | 246,317 <sup>f,g</sup>                                    | 9,330                     | 236,987                                   |
| 2008 | 171,347              | 185,409                                    | 11,381                                      | 174,028 <sup>f,g</sup>                                    | 6,130                     | 167,898                                   |
| 2009 | 95,462               | 101,734                                    | 6,995                                       | 94,739 <sup>f</sup>                                       | 1,113                     | 93,626                                    |
| 2010 | 125,547              | 132,930                                    | 11,432                                      | 121,498 <sup>f</sup>                                      | 3,709                     | 117,789                                   |
| 2011 | 212,162              | 224,355                                    | 12,477                                      | 211,878 <sup>f</sup>                                      | 6,312                     | 205,566                                   |
| 2012 | 147,710              | 153,248                                    | 11,681                                      | 141,567 <sup>f</sup>                                      | 3,905                     | 137,662                                   |
| 2013 | 200,754              | 216,791                                    | 12,642                                      | 204,149 <sup>f</sup>                                      | 3,887                     | 200,262                                   |
| 2014 | 167,715              | 172,887                                    | 13,041                                      | 159,846 <sup>f</sup>                                      | 3,050                     | 156,796                                   |
| 2015 | 112,136              | 125,095                                    | 12,540                                      | 112,555 <sup>f</sup>                                      | 3,897                     | 108,658                                   |

-continued-

## Appendix B16.—Page 2 of 2.

| Date              | Eagle sonar estimate | Eagle sonar expanded estimate <sup>a</sup> | U.S. harvest above Eagle sonar | U.S./Canada mainstem border passage estimate <sup>b</sup> | Canadian mainstem harvest | Spawning escapement estimate <sup>c</sup> |
|-------------------|----------------------|--|--------------------------------|---|---------------------------|---|
| 2016              | 144,035              | 161,027                                    | 13,015                         | 148,012 <sup>f</sup>                                      | 2,745                     | 145,267                                   |
| 2017 <sup>h</sup> | 407,166              | 419,099                                    | 14,110                         | 404,989 <sup>f</sup>                                      | 3,404                     | 401,585                                   |
| 2018 <sup>h</sup> | 136,732              | 168,798                                    | 11,715                         | 157,083 <sup>f</sup>                                      | 2,957                     | 154,126                                   |
| 2019 <sup>h</sup> | 101,678              | 113,256                                    | 10,759                         | 102,497 <sup>f</sup>                                      | 2,759                     | 99,738                                    |
| 2020 <sup>h</sup> | 20,766               | 23,512                                     | 0                              | 23,512 <sup>f</sup>                                       | 0                         | 23,512                                    |
| 2021 <sup>h</sup> | 19,668               | 23,170                                     | 0                              | 23,170 <sup>f</sup>                                       | 0                         | 23,170                                    |
| Goal <sup>i</sup> |                      |  |                                |   |                           | >80,000                                   |
| IMEG <sup>c</sup> |                      |  |                                |   |                           | 70,000–104,000                            |
| Averages          |                      |  |                                |   |                           |   |
| 1980–2020         | 167,698              | 180,563                                    | 11,884                         | 129,298   | 14,133                    | 115,165                                   |
| 2011–2020         | 165,085              | 177,807                                    | 11,198                         | 166,609   | 3,292                     | 163,317                                   |
| 2016–2020         | 162,075              | 177,138                                    | 9,920                          | 167,219   | 2,373                     | 164,846                                   |
| Minimum           | 20,766               | 23,512                                     | 0                              | 23,512  | 0                         | 22,912                                    |
| Maximum           | 407,166              | 419,099                                    | 18,691                         | 451,477   | 44,345                    | 437,498                                   |

*Note:* Table includes information on U.S./Canada border passage estimates, Eagle area subsistence harvest between the sonar and the border (where applicable), and Canadian mainstem harvest. Estimates for subsistence caught salmon between the sonar site and border (Eagle area) prior to 2008 include an unknown portion caught below the sonar site. This number is most likely in the thousands for chum salmon. Starting in 2008, the estimates for subsistence-caught salmon only include salmon harvested between the sonar site and the U.S./Canada border. Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Sonar estimates include an expansion for fish that may have passed after operations ceased through October 18. In 2018, expanded to October 23 due to late run timing.

<sup>b</sup> Border passage estimate is based on a mark–recapture estimate unless otherwise indicated.

<sup>c</sup> Estimated mainstem border passage minus Canadian mainstem harvest (excludes Fishing Branch River). Current interim management escapement goal (IMEG) is 70,000 to 104,000 fall chum salmon. IMEG was established in 2010 based on brood table of Canadian-origin mainstem stocks (1982–2003).

<sup>d</sup> Escapement estimate based on mark–recapture program unavailable. Estimate based on assumed average exploitation rate.

<sup>e</sup> From 1999–2004, border passage estimates were revised using a Stratified Population Analysis System (Arnason et. al 1995).

<sup>f</sup> From 2006–present, border passage estimate is based on sonar minus harvest from U.S. residents upstream of deployment.

<sup>g</sup> Mark–recapture border passage estimates include 217,810; 235,956; and 132,048 fish from 2006–2008 respectively, during transition to sonar.

<sup>h</sup> Data are preliminary as harvest information is not published yet.

<sup>i</sup> Escapement goal in Pacific Salmon Treaty for mainstem Yukon River Canadian-origin fall chum salmon.

Appendix B17.—Coho salmon passage estimates or escapement estimates for selected spawning areas in the U.S. (Alaska) portion of the Yukon River drainage, 1972–2021.

| Year | Yukon<br>River<br>mainstem<br>sonar<br>estimate <sup>a</sup> | Nenana River drainage  |                                 |                        |                          | Upper Tanana River drainage               |                                  |                                   |
|------|--|------------------------|---------------------------------|------------------------|--------------------------|---|----------------------------------|-----------------------------------|
|      |  | Lost<br>Slough         | Nenana<br>mainstem <sup>b</sup> | Wood<br>Creek          | Seventeen<br>Mile Slough | Delta<br>Clearwater<br>River <sup>c</sup> | Clearwater<br>Lake and<br>outlet | Richardson<br>Clearwater<br>River |
| 1972 |  |                        |                                 |                        |                          | 632 (b)                                   | 417 (f)                          | 454 (f) <sup>d</sup>              |
| 1973 |  |                        |                                 |                        |                          | 3,322 (u)                                 | 551 (u)                          | 375 (u)                           |
| 1974 |  | 1,388 (f)              |                                 |                        | 27 (f)                   | 3,954 (h) <sup>d</sup>                    | 560 (f)                          | 652 (h)                           |
| 1975 |  | 827 (f)                |                                 |                        | 956 (f)                  | 5,100 (b)                                 | 1,575 (b)                        |                                   |
| 1976 |  | 118 (f)                |                                 |                        | 281 (f)                  | 1,920 (b)                                 | 1,500 (b)                        | 80 (f) <sup>d</sup>               |
| 1977 |  | 524 (f) <sup>d</sup>   |                                 | 310 (g)                | 1,167 (f)                | 4,793 (b)                                 | 730 (b)                          | 327 (f)                           |
| 1978 |  | 350 (f)                |                                 | 300 (g)                | 466 (f)                  | 4,798 (b)                                 | 570 (b)                          |                                   |
| 1979 |  | 227 (f)                |                                 |                        | 1,987 (f)                | 8,970 (b)                                 | 1,015 (b)                        | 372 (f)                           |
| 1980 |  | 499 (f) <sup>d</sup>   |                                 | 1,603 (g)              | 592 (f)                  | 3,946 (b)                                 | 1,545 (b)                        | 611 (f)                           |
| 1981 |  | 274 (f)                |                                 | 849 (w) <sup>e</sup>   | 1,005 (f)                | 8,563 (u) <sup>f</sup>                    | 459 (f)                          | 550 (f)                           |
| 1982 |  |                        |                                 | 1,436 (w) <sup>e</sup> | (f)                      | 8,365 (g) <sup>f</sup>                    |                                  |                                   |
| 1983 |  | 766 (f)                |                                 | 1,042 (w)              | 103 (f)                  | 8,019 (b) <sup>f</sup>                    | 253 (f)                          | 88 (f)                            |
| 1984 |  | 2,677 (f)              |                                 | 8,826 (w)              | (f)                      | 11,061 (b)                                | 1,368 (f)                        | 428 (f)                           |
| 1985 |  | 1,584 (f)              |                                 | 4,470 (w)              | 2,081 (f)                | 5,358 (b)                                 | 750 (f)                          |                                   |
| 1986 |  | 794 (f)                |                                 | 1,664 (w)              | 218 (b)                  | 10,857 (b)                                | 3,577 (f)                        | 146 (f) <sup>d</sup>              |
| 1987 |  | 2,511 (f)              |                                 | 2,387 (w)              | 3,802 (f)                | 22,300 (b)                                | 4,225 (b)                        |                                   |
| 1988 |  | 348 (f)                |                                 | 2,046 (w)              |                          | 21,600 (b)                                | 825 (b)                          |                                   |
| 1989 |  |                        |                                 | 412 (w)                | 824 (f) <sup>d</sup>     | 11,000 (b)                                | 1,600 (b)                        | 483 (f)                           |
| 1990 |  | 688 (f)                | 1,308 (f)                       |                        | (h) <sup>d</sup>         | 8,325 (b)                                 | 2,375 (b)                        |                                   |
| 1991 |  | 564 (f)                | 447 (f)                         |                        | 52 (f)                   | 23,900 (b)                                | 3,150 (b)                        |                                   |
| 1992 |  | 372 (f)                |                                 |                        | 490 (f)                  | 3,963 (b)                                 | 229 (b)                          | 500 (f)                           |
| 1993 |  | 350 (f)                | 419 (f)                         | 666 (w) <sup>g</sup>   | 581 (h)                  | 10,875 (b)                                | 3,525 (b)                        |                                   |
| 1994 |  | 944 (h)                | 1,648 (h)                       | 1,317 (w) <sup>h</sup> | 2,909 (h)                | 62,675 (b)                                | 3,425 (b)                        | 5,800 (f)                         |
| 1995 | 115,569  | 4,169 (f)              | 2,218 (h)                       | 500 (w)                | 1,512 (h)                | 20,100 (b)                                | 3,625 (b)                        |                                   |
| 1996 | <sup>i</sup>   | 2,040 (h)              | 2,171 (h)                       | 201 (u) <sup>d</sup>   | 3,668 (g/b)              | 14,075 (b)                                | 1,125 (h) <sup>d</sup>           |                                   |
| 1997 | 118,065  | 1,524 (h)              | 1,446 (h)                       | <sup>j</sup>           | 1,996 (h)                | 11,525 (b)                                | 2,775 (b)                        |                                   |
| 1998 | 146,365  | 1,360 (h) <sup>d</sup> | 2,771 (h) <sup>d</sup>          | <sup>j</sup>           | 1,413 (g/b)              | 11,100 (b)                                | 2,775 (b)                        |                                   |
| 1999 | 76,174   | 1,002 (h) <sup>d</sup> | 745 (h) <sup>d</sup>            | 370 (h)                | 662 (h) <sup>d</sup>     | 10,975 (b)                                |                                  |                                   |

-continued-

| Year | Yukon<br>River<br>mainstem<br>sonar<br>estimate <sup>a</sup> | Nenana River drainage |                                 |                    |                          | Upper Tanana River drainage               |                                  |                                   |  |
|------|--|-----------------------|---------------------------------|--------------------|--------------------------|---|----------------------------------|-----------------------------------|--|
|      |  | Lost<br>Slough        | Nenana<br>mainstem <sup>b</sup> | Wood<br>Creek      | Seventeen<br>Mile Slough | Delta<br>Clearwater<br>River <sup>c</sup> | Clearwater<br>Lake and<br>outlet | Richardson<br>Clearwater<br>River |  |
|      |  |                       |                                 |                    |                          |   |                                  |                                   |  |
| 2000 | 206,365  | 55 (h) <sup>d</sup>   | 68 (h) <sup>d</sup>             | j                  | 879 (h) <sup>d</sup>     | 9,225 (b)                                 | 1,025 (b)                        | 2,175 (h)                         |  |
| 2001 | 160,272  | 242 (h)               | 859 (h)                         | 699 (h)            | 3,753 (h)                | 46,985 (b)                                | 4,425 (b)                        | 1,531 (f)                         |  |
| 2002 | 137,077  | 0 (h)                 | 328 (h)                         | 935 (h)            | 1,910 (h)                | 38,625 (b)                                | 5,900 (b)                        | 874 (f)                           |  |
| 2003 | 280,552  | 85 (h)                | 658 (h)                         | 3,055 (h)          | 4,535 (h)                | 102,800 (b)                               | 8,800 (b)                        | 6,232 (h)                         |  |
| 2004 | 207,844  | 220 (h)               | 450 (h)                         | 840 (h)            | 3,370 (h)                | 37,550 (b)                                | 2,925 (b)                        | 8,626 (h)                         |  |
| 2005 | 194,622  | 430 (h)               | 325 (h)                         | 1,030 (h)          | 3,890 (h)                | 34,293 (b)                                | 2,100 (b)                        | 2,024 (h)                         |  |
| 2006 | 163,889  | 194 (h)               | 160 (h)                         | 634 (h)            | 1,916 (h)                | 16,748 (b)                                | 4,375 (b)                        | 271 (h)                           |  |
| 2007 | 192,406  | 63 (h)                | 520 (h)                         | 605 (h)            | 1,733 (h)                | 14,650 (b)                                | 2,075 (b)                        | 553 (h)                           |  |
| 2008 | 145,378  | 1,342 (h)             | 1,539 (h)                       | 578 (h)            | 1,652 (h)                | 7,500 (b)                                 | 1,275 (b)                        | 265 (h)                           |  |
| 2009 | i  | 410 (h)               |                                 | 470 (h)            | 680 (h)                  | 16,850 (b)                                | 5,450 (b)                        | 155 (h)                           |  |
| 2010 | 177,724  | 1,110 (h)             | 280 (h)                         | 340 (h)            | 720 (h)                  | 5,867 (b)                                 | 813 (b)                          | 1,002 (h)                         |  |
| 2011 | 149,533  | 369 (h)               |                                 | 0 (h) <sup>j</sup> | 912 (h)                  | 6,180 (b)                                 | 2,092 (b)                        | 575 (h)                           |  |
| 2012 | 130,734  |                       | 106 (h)                         | 0 (h) <sup>j</sup> | 405 (h)                  | 5,230 (b)                                 | 396 (h)                          | 515 (h)                           |  |
| 2013 | 110,515  | 721 (h)               |                                 | 55 (h)             | 425 (h)                  | 6,222 (b)                                 | 2,221 (h)                        | 647 (h)                           |  |
| 2014 | 283,421  | 333 (h)               | 378 (h)                         | 649 (h)            | 886 (h)                  | 4,285 (b)                                 | 434 (h)                          | 1,941 (h)                         |  |
| 2015 | 121,193  | 242 (h)               | 1,789 (h)                       | 1,419 (h)          | 3,890 (h)                | 19,533 (b)                                | 1,621 (h)                        | 3,742 (h)                         |  |
| 2016 | 168,297  | 334 (h)               | 1,680 (h)                       | 1,327 (h)          | 2,746 (h)                | 6,767 (b)                                 | 1,421 (h)                        | 1,350 (h)                         |  |
| 2017 | 166,320  | 1,278 (h)             | 862 (h)                         | 2,025 (h)          | 1,942 (h)                | 9,617 (b)                                 |                                  |                                   |  |
| 2018 | 136,347  | 1,822 (h)             | 241 (h)                         | 361 (h)            | 347 (h)                  | 2,884 (b)                                 | 2,465 (h)                        | 976 (h)                           |  |
| 2019 | 86,401   |                       | 749 (h)                         | 184 (h)            | 424 (h)                  | 2,043 (b)                                 | 258 (h)                          | 300 (h)                           |  |
| 2020 | 107,680  | 28 (h)                | 206 (h)                         | 231 (h)            | 507 (h)                  | 2,557 (b)                                 | 210 (h)                          | 475 (h)                           |  |
| 2021 | 37,257 <sup>k</sup>  | 126 (h)               | 104 (h)                         | 226 (h)            | 213 (h)                  | 913 (b)                                   | 130 (h)                          | 17 (h)                            |  |

-continued-

Appendix B17.—Page 3 of 3.

| Year             | Yukon<br>River<br>mainstem<br>sonar<br>estimate <sup>a</sup> | Nenana River drainage |                                 |               |                          | Upper Tanana River drainage               |                                  |                                   |
|------------------|--|-----------------------|---------------------------------|---------------|--------------------------|---|----------------------------------|-----------------------------------|
|                  |  | Lost<br>Slough        | Nenana<br>mainstem <sup>b</sup> | Wood<br>Creek | Seventeen<br>Mile Slough | Delta<br>Clearwater<br>River <sup>c</sup> | Clearwater<br>Lake and<br>outlet | Richardson<br>Clearwater<br>River |
|                  |  |                       |                                 |               |                          |   |                                  |                                   |
| SEG <sup>1</sup> |  |                       |                                 |               |                          | 5,200–17,000                              |                                  |                                   |
| Averages         |  |                       |                                 |               |                          |   |                                  |                                   |
| 1972–2020        | 157,614  | 818                   | 903                             | 1,185         | 1,496                    | 14,663                                    | 2,061                            | 1,326                             |
| 2011–2020        | 146,044  | 641                   | 751                             | 625           | 1,248                    | 6,532                                     | 1,235                            | 1,169                             |
| 2016–2020        | 133,009  | 866                   | 748                             | 826           | 1,193                    | 4,774                                     | 1,089                            | 775                               |
| Minimum          | 76,174   | 0                     | 68                              | 0             | 27                       | 632                                       | 210                              | 80                                |
| Maximum          | 283,421  | 4,169                 | 2,771                           | 8,826         | 4,535                    | 102,800                                   | 8,800                            | 8,626                             |

*Note:* Only peak counts presented. Survey rating is fair to good, unless otherwise noted. Denotations of survey methods include: (b)=boat, (f)=fixed wing, (g)=ground/foot, (h)=helicopter, (u)=undocumented, and (w)=weir. Minimum and maximum values exclude the most recent year data.

<sup>a</sup> Passage estimates for coho salmon are incomplete. The sonar project is terminated prior to the end of the coho salmon run. New model estimates generated in 2015 and applied to dataset back to 1995 and used since.

<sup>b</sup> Index area includes mainstem Nenana River between confluences of Lost Slough and Teklanika River.

<sup>c</sup> Index area is lower 28km (17.5 mi) of system.

<sup>d</sup> Poor survey resulted in minimal count.

<sup>e</sup> Weir was operated at the mouth of Clear Creek (Shores Landing).

<sup>f</sup> Expanded estimate based on partial survey counts and historic distribution of spawners from 1977–1980.

<sup>g</sup> Weir project terminated on October 4, 1993. Weir normally operated until mid- to late October.

<sup>h</sup> Weir project terminated September 27, 1994. Weir normally operated until mid- to late October.

<sup>i</sup> Project operated all or partial season, estimate was not useable.

<sup>j</sup> No survey of Wood Creek due to obstructions in creek or surveyed with zero fish observed.

<sup>k</sup> Data are preliminary.

<sup>1</sup> Sustainable escapement goal (SEG) established January 2004 (replaces BEG of greater than 9,000 fish established March 1993), based on boat survey counts of coho salmon in the lower 17.5 river miles during the period October 21–27.

Appendix B18.–Yukon River Salmon Agreement specified obligations for harvest shares, border passage and spawning escapement for mainstem Canadian-origin Yukon River Chinook salmon, 2001–2021

| Year              | Total  |                          |        |                |        |   | Yukon River                  |                                |                               |                      |        |        |                                     |        |
|-------------------|--|--------------------------|--------|----------------|--------|---|------------------------------|--------------------------------|-------------------------------|----------------------|--------|--------|-------------------------------------|--------|
|                   | Total<br>estimated<br>Canadian-<br>origin<br>run size <sup>a</sup> | allowable                |        | U.S. share (%) |        | Border<br>passage<br>objective <sup>d</sup> | Canada share                 |                                |                               | Panel goal           |        |        | Spawning<br>escapement <sup>g</sup> |        |
|                   |  | catch (TAC) <sup>b</sup> |        | of TAC         |        |   | U.S.<br>harvest <sup>c</sup> | Border<br>passage <sup>e</sup> | Canada<br>mainstem<br>harvest | or IMEG <sup>f</sup> |        |        |                                     |        |
|                   |  | From                     | To     | 0.74           | 0.8    |   |                              |                                |                               | (%) of TAC           | 0.20   | 0.26   |                                     | From   |
| 2001              | 77,354   | 49,354                   |        | 36,522         | 39,483 | 23,325                                      | 39,351                       | 54,029                         | 9,871                         | 12,832               | 9,774  | 28,000 |                                     | 44,255 |
| 2002              | 73,417   | 45,417                   |        | 33,609         | 36,334 | 30,058                                      | 38,446                       | 43,359                         | 9,083                         | 11,808               | 9,070  | 28,000 |                                     | 34,289 |
| 2003              | 118,022  | 90,022                   |        | 66,616         | 72,018 | 59,940                                      | 48,705                       | 58,082                         | 18,004                        | 23,406               | 9,446  | 28,000 |                                     | 48,636 |
| 2004              | 105,942  | 77,942                   |        | 57,677         | 62,354 | 57,831                                      | 45,927                       | 48,111                         | 15,588                        | 20,265               | 10,946 | 28,000 |                                     | 37,165 |
| 2005              | 86,895   | 58,895                   |        | 43,582         | 47,116 | 44,650                                      | 41,546                       | 42,245                         | 11,779                        | 15,313               | 10,977 | 28,000 |                                     | 31,268 |
| 2006              | 84,845   | 56,845                   |        | 42,065         | 45,476 | 48,097                                      | 41,074                       | 36,748                         | 11,369                        | 14,780               | 8,758  | 28,000 |                                     | 27,990 |
| 2007              | 70,440   | 27,440                   | 37,440 | 20,306         | 29,952 | 48,320                                      | 40,611                       | 22,120                         | 5,488                         | 9,734                | 4,794  | 33,000 | 43,000                              | 17,326 |
| 2008 <sup>e</sup> | 62,358   | 17,358                   |        | 12,845         | 13,886 | 25,329                                      | 48,992                       | 37,029                         | 3,472                         | 4,513                | 3,399  | 45,000 |                                     | 33,630 |
| 2009              | 87,221   | 42,221                   |        | 31,244         | 33,777 | 17,646                                      | 54,711                       | 69,575                         | 8,444                         | 10,977               | 4,297  | 45,000 |                                     | 65,278 |
| 2010              | 59,736   | 4,736                    | 17,236 | 3,505          | 13,789 | 25,271                                      | 45,214                       | 34,465                         | 947                           | 4,481                | 2,456  | 42,500 | 55,000                              | 32,009 |
| 2011              | 71,725   | 16,725                   | 29,225 | 12,377         | 23,380 | 20,824                                      | 47,972                       | 50,901                         | 3,345                         | 7,599                | 4,594  | 42,500 | 55,000                              | 46,307 |
| 2012              | 48,498   | 0                        | 5,998  | 0              | 4,798  | 13,842                                      | 43,280                       | 34,656                         | 0                             | 1,559                | 2,000  | 42,500 | 55,000                              | 32,656 |
| 2013              | 37,177   | 0                        | 0      | 0              | 0      | 6,604                                       | 42,500                       | 30,573                         | 0                             | 0                    | 1,904  | 42,500 | 55,000                              | 28,669 |
| 2014              | 64,886   | 9,886                    | 22,386 | 7,316          | 17,909 | 1,455                                       | 46,399                       | 63,431                         | 1,977                         | 5,820                | 100    | 42,500 | 55,000                              | 63,331 |
| 2015              | 87,323   | 32,323                   | 44,823 | 23,919         | 35,858 | 3,649                                       | 51,559                       | 83,674                         | 6,465                         | 11,654               | 1,000  | 42,500 | 55,000                              | 82,674 |
| 2016              | 82,765   | 27,765                   | 40,265 | 20,546         | 32,212 | 11,198                                      | 50,511                       | 71,567                         | 5,553                         | 10,469               | 2,769  | 42,500 | 55,000                              | 68,798 |
| 2017              | 93,188   | 38,188                   | 50,688 | 28,259         | 40,551 | 21,373                                      | 52,908                       | 71,815                         | 7,638                         | 13,179               | 3,500  | 42,500 | 55,000                              | 68,315 |
| 2018              | 76,356   | 21,356                   | 33,856 | 15,804         | 27,085 | 19,092                                      | 49,037                       | 57,264                         | 4,271                         | 8,803                | 2,790  | 42,500 | 55,000                              | 54,474 |
| 2019              | 72,620   | 17,620                   | 30,120 | 13,039         | 24,096 | 27,804                                      | 48,178                       | 44,816                         | 3,524                         | 7,831                | 2,764  | 42,500 | 55,000                              | 42,052 |
| 2020              | 45,501   | 0                        | 3,001  | 0              | 2,401  | 12,171                                      | 42,890                       | 33,330                         | 0                             | 780                  | 2,363  | 42,500 | 55,000                              | 30,967 |
| 2021              | 32,972   | 0                        | 0      | 0              | 0      | 1,214                                       | 42,500                       | 31,758                         | 0                             | 0                    | 306    | 42,500 | 55,000                              | 31,452 |

-continued-

## Appendix B18.—Page 2 of 2

*Note:* The table does not represent a dataset, its intent is to represent the information at the time. Data presented for each year is from the assessment methods of that year, and represents final values (may not be the same as preliminary values published in that years annual JTC report, or as retroactively finalized values using revised calculation techniques). Gray shaded boxes indicate Yukon River Salmon Agreement performance obligations that were not met.

- <sup>a</sup> Total estimated Canadian-origin run size is calculated as border passage plus Alaskan harvest of Canadian-origin Chinook salmon. From 2001 to 2012, these values were not specifically presented in annual JTC reports, and have been retroactively calculated based on best available historical information, from the assessment methods used in that year.
- <sup>b</sup> Total run size, total allowable catch (TAC) and harvest share calculations are finalized post-season. TAC is calculated by subtracting the IMEG from the total run size. Delivering the IMEG plus the midpoint of Canada's harvest share to the Alaska-Yukon border is part of the U.S. obligation as per the Pacific Salmon Treaty's Yukon River Salmon Agreement.
- <sup>c</sup> Scale pattern analysis was used to determine the U.S. Harvest stock proportions prior to 2004. Since 2004 U.S. Harvest estimates of the Canadian-origin stock were estimated by applying the stock proportions collected from harvest sampling to number of fish harvested in Alaska. Beginning in 2014, the U.S. harvest includes harvest from the Coastal District. Values from 2001-2012 were obtained from the annual ADF&G report "Origins of Chinook Salmon in Yukon Area Fisheries", and values from 2013 onwards have been reported in the annual JTC Report.
- <sup>d</sup> Border passage objective is calculated post season as the agreed spawning escapement goal plus the mid-point of the Canadian harvest share. For years where the escapement goal is a range, this is represented as the average of the Canadian Harvest Share, plus the lower end of the escapement goal.
- <sup>e</sup> From 2001 to 2007 the border passage was estimated from a mark recapture project. Beginning in 2008 border passage was estimated from the Eagle sonar, minus any Alaskan harvest upstream of the sonar. The bold horizontal line between 2007 and 2008 indicates the JTC's recommendation to use the Eagle sonar as the primary assessment tool for the border passage estimate. Values from this year forward are sonar based.
- <sup>f</sup> Yukon River Panel goals have changed over time, and have been both points and ranges. IMEGs are not biologically based escapement goals.
- <sup>g</sup> Spawning escapement is calculated as the border passage estimate minus the harvest in Canada using the assessment methods of that year.
- <sup>h</sup> In the 2001 JTC report, there are some references to a lower goal of 18,000 although further reports state the goal of 28,000 was the only goal for this year.
- <sup>i</sup> In 2002 and 2003, the Chinook salmon goal was set at 25,000 fish. However, if the U.S. conducted a commercial fishery the goal would be increased to 28,000 fish.
- <sup>j</sup> In 2004, the escapement target for Canadian-origin Upper Yukon Chinook salmon was >28,000 Chinook salmon. If the run was gauged to be sufficiently strong, the escapement target could range up to 38,000 Chinook salmon, although the Panel did not describe what constituted a "strong" run.



Appendix B19.—Summary of management and conservation measures implemented in the U.S. (Alaska) and Canada for Chinook salmon, 2001–2021.

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)  | Canadian management actions (First Nation fishery) | Canadian management actions (commercial, domestic, public angling)   |
|------|--|---|--|--|
| 2001 | Subsistence fishing schedule implemented (and continued in following years). | No commercial fishing for Chinook or summer chum salmon.  | Unrestricted                                       | Test fishery implemented in early season; commercial/domestic openings determined by weekly estimates of abundance, public angling open. |
| 2002 |  | Chinook commercial fishing shifted to midpoint of run and later.  | Unrestricted                                       | Test fishery implemented in early season; commercial/domestic openings determined by weekly estimates of abundance, public angling open. |
| 2003 |  | Chinook commercial fishing shifted to midpoint of run and later.  | Unrestricted                                       | Test fishery implemented in early season; commercial/domestic openings determined by weekly estimates of abundance, public angling open. |
| 2004 |  | Chinook commercial fishing shifted to midpoint of run and later.  | Unrestricted                                       | Test fishery implemented in early season; commercial/domestic openings determined by weekly estimates of abundance, public angling open. |
| 2005 |  | Chinook commercial fishing shifted to midpoint of run and later.  | Unrestricted                                       | Commercial/domestic openings determined by weekly estimates of abundance, public angling open.   |
| 2006 |  | Chinook commercial fishing delayed until start of second pulse.   | Unrestricted                                       | Commercial/domestic openings determined by weekly estimates of abundance, public angling open.   |
| 2007 |  | Short fishing period on historic first quarter point date. Majority of harvest spread over middle 50% of the run. | Unrestricted                                       | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River.     |
| 2008 | Protection on 2nd and 3rd pulses.  | Chinook commercial fishing closed.  | Voluntary reduction in harvest.                    | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River.     |
| 2009 | 1st and 2nd pulse closure.   | Chinook commercial fishing closed and no sale of incidental catch; summer chum fishing delayed.                   | Voluntary reduction in harvest in early season.    | Commercial/domestic openings determined by weekly estimates of abundance, public angling open.   |

-continued-

Appendix B19.–Page 2 of 10.

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)  | Canadian management actions (First Nation fishery) | Canadian management actions (commercial, domestic, public angling)   |
|------|--|---|--|--|
| 2010 |  | Chinook commercial fishing closed; summer chum fishing delayed.   | Voluntary reduction in harvest.                    | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery.   |
| 2011 | 1st and 2nd pulse closure; additional fishing time reductions in upper districts; 7.5 inch mesh size restriction all season.   | Chinook commercial fishing closed and no sale of incidental catch; summer chum fishing delayed; summer chum fishing restricted to certain areas of low Chinook abundance.   | Voluntary reduction in harvest in early season.    | Chinook commercial/domestic fishing closed; recreational fishing varied to non-retention in the public angling fishery, angling closure at Tatchun River, public angling restrictions lifted late in the season. |
| 2012 | 1st and 2nd pulse closure; additional fishing time reductions in upper districts; 6 inch mesh size restriction after closures.   | Chinook commercial fishing closed and no sale of incidental catch; summer chum fishing delayed and restricted to areas of low Chinook abundance; chum fish wheels attended at all times and Chinook released alive.   | Voluntary reduction in harvest.                    | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River.   |
| 2013 | 1st, 2nd and 3rd pulse closures - limited opportunity in between pulses; additional fishing time reductions in upper districts; 6 inch mesh size restriction all season. | Chinook commercial fishing closed and no sale of incidental catch. Summer chum fishing with beach seines and dip nets, all Chinook released alive. Gillnet summer chum fishing restricted to 5.5 inch and 30 meshes; delayed and restricted to areas of low Chinook abundance; chum fish wheels attended at all times and Chinook released alive. | Voluntary reduction in harvest.                    | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River and Teslin River.  |

-continued-

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|--|--|---|--|
| 2014 | Entire mainstem river closed to Chinook-directed fishing; no gillnets allowed greater than 4 inch mesh size to harvest non-salmon species; opportunity to harvest summer chum salmon in Districts 1–4 using elective gear that allows immediate and live release of Chinook allowed (dip nets, beach seines, and fish wheels); short openings with 6 inch or smaller gillnets allowed in each districts after greater than 90% of Chinook salmon run had passed through; greater than 99% in District 5. | Chinook commercial fishing closed; liberal opportunity for summer chum fishing with beach seines and dip nets - all Chinook released immediately and alive; 6 inch or smaller gillnet summer chum fishing delayed until majority of Chinook run complete; no sale of incidental Chinook; chum fish wheels had to be attended at all times and all Chinook released immediately to the water; concurrent subsistence and commercial openings. | Regulatory removal of TAC until 3rd quartile, voluntary reduction or closure maintained by majority of First Nations. | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River and Teslin River |

-continued-

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|--|--|---|--|
| 2015 | Entire river closed to Chinook-directed fishing; no gillnets allowed greater than 4 inch mesh size to harvest non-salmon species; opportunity to harvest summer chum salmon in Districts 1–4 using selective gear that allows immediate and live release of Chinook (dipnets, beach seines, and fish wheels); short openings with 6 inch or smaller gillnets allowed in each district between pulses of Chinook salmon when summer chum abundance was high. Subsistence fishing was allowed in Subdistrict 5-D on the early trickle of Chinook salmon. Subsistence schedules liberalized in Districts 4 and 5 once Chinook salmon border escapement was surpassed. | Chinook commercial fishing closed; liberal opportunity for summer chum fishing with beach seines and dipnets - all Chinook released immediately and alive; 6 inch or smaller gillnet summer chum fishing delayed until majority of Chinook run complete; no sale of incidental Chinook; fish wheels had to be attended at all times and all Chinook released immediately to the water; concurrent subsistence and commercial openings. | Regulatory removal of TAC until 2nd quartile, voluntary reduction or closure maintained by majority of First Nations. | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River. |

-continued-

| Year | U.S. management actions (subsistence)   | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|---|--|---|--|
| 2016 | Early season only: Districts 1–5 using selective gear requiring live release of Chinook (dipnets, beach seines, and fish wheels); Subdistrict 5-D had open fishing on the early trickle with 6 inch gillnets. Reduced regulatory schedule fishing with gillnets restricted to 6” in most districts. Followed by surgical openings with 7.5 inch gillnets late in the run. Subsistence schedules liberalized in Districts 4 and 5 once Chinook salmon border escapement was surpassed. | Chinook commercial fishing closed; liberal opportunity for summer chum fishing with selective gear - all Chinook released immediately and alive; 6 inch or smaller gillnet summer chum fishing delayed until majority of Chinook run complete; no sale of incidental Chinook. No concurrent subsistence and commercial openings. | Open with recommendation for reduced harvest (30%), voluntary reduction or closure maintained by majority of First Nations. | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River. |

-continued-

Appendix B19.–Page 6 of 10.

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|--|--|---|--|
| 2017 | Early season only: Districts 1–5 placed on regulatory schedule fishing with gillnets restricted to 6 inch prior to the first pulse. Fishing restricted to selective gear requiring live release of Chinook (dipnets, beach seines, and fish wheels), then reopened to regulatory schedule with 7.5 inch of smaller mesh. Coastal District, Koyukuk and Innoko Rivers, and Subdistrict 5-D remained open with 7.5 inch or smaller mesh size all season. | Chinook commercial fishing closed; liberal opportunity for summer chum fishing with selective gear - all Chinook released immediately and alive; 6 inch or smaller gillnet summer chum fishing delayed until majority of Chinook salmon run had entered the river. No sale of incidental Chinook salmon in summer season; one commercial period occurred in District 1 where Chinook salmon caught during fall chum directed commercial fishing were allowed to be sold. No concurrent commercial and subsistence openings in Districts 1 and 2. | Open with recommendation for reduced harvest, voluntary reduction or closure maintained by majority of First Nations. | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River. |

-continued-

Appendix B19.–Page 7 of 10.

| Year | U.S. management actions (subsistence)   | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|---|--|---|--|
| 2018 | Early season: Districts 1–5 placed on half regulatory schedule fishing with gillnets restricted to 6 inch. Two subsistence periods (one per week) were cancelled in Districts 1–4A. Later in the season, limited opportunity (one reduced time opening per week) was provided with 7.5 inch mesh in Districts 1-4. District 5 remained restricted to 6 inch mesh through the third pulse of the Chinook salmon run. Coastal District, Koyukuk and Innoko Rivers remained open with 7.5 inch or smaller mesh all season. | Chinook commercial fishing closed; liberal opportunity for summer chum fishing with selective gear - all Chinook released immediately and alive; 6 inch or smaller gillnet summer chum fishing delayed until majority of Chinook salmon run had entered the river. No sale of incidental Chinook salmon. No concurrent commercial and subsistence openings in Districts 1 and 2. | Open with recommendation for reduced harvest; voluntary reduction or closure maintained by majority of First Nations. | Chinook commercial/domestic fishing closed; varied to non-retention in the public angling fishery, angling closure at Tatchun River. |

-continued-

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)  | Canadian management actions (First Nation fishery)   | Canadian management actions (commercial, domestic, public angling)   |
|------|--|---|--|--|
| 2019 | Most of season: Districts 1-5 placed on half regulatory schedule fishing. 6 inch or smaller mesh restrictions added for at least 2 periods in Districts 1-6. One subsistence period was cancelled in Districts 1-4. Fishing was closed for 10 days in Subdistrict 5-D. Coastal District, Koyukuk and Innoko Rivers remained open with 7.5 inch or smaller mesh all season. | Summer chum commercial fishing delayed due to late run timing; 6 inch or smaller gillnet summer chum commercial fishing occurred after the majority of Chinook run complete. Sale of incidental Chinook salmon allowed in the summer season after over 200,000 Chinook salmon had been counted at Pilot Station sonar. Sale of incidental Chinook salmon allowed during fall chum-directed commercial fishing. No concurrent commercial and subsistence openings. | Season commenced on July 1 with an opening and full allocation available for First Nation Chinook Fishery. Voluntary reduction or closure maintained by majority of First Nations. First Nation Governments were notified in early August advised to implement additional precautionary measures due to lower than expected passage at Eagle sonar and unlikelihood of achieving the midpoint of the IMEG. | Commercial and domestic fishery conditions of licence limited harvesters to gillnets with a 6 inch or smaller mesh size; Chinook commercial/domestic fisheries were closed. In advance of the Chinook return, retention varied to zero in the public angling fishery. A complete angling closure was enacted on the Yukon River and its tributaries as a Chinook conservation measure. Similarly, chum commercial/domestic fishery opening delayed to mid-September due to Chinook late run timing and low returns. Salmon angling fishery reopened in late September. |

-continued-



| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial)   | Canadian management actions (First Nation fishery)   | Canadian management actions (commercial, domestic, public angling)   |
|------|--|--|--|--|
| 2020 | Start of season; Districts 1-4 on half time and 6 inch or smaller mesh gillnets. Fishing in most districts closed or restricted to selective gear types in late June in response to late run timing. Fishing re-opened in most districts on reduced schedule with 6 inch mesh. Eagle sonar midpoint projections were poor; District 5 closed in late July for the rest of the summer season. Additional closures of 4 inch mesh were implemented throughout the drainage to avoid any harvest of Chinook salmon. | Summer chum commercial fishing delayed due to late run timing; 6 inch or smaller gillnet summer chum commercial fishing occurred after the majority of Chinook run was complete. Only 5 commercial periods were fished in the Lower Yukon due to low summer chum salmon run. No Chinook commercial fishing; less than 350 fish retained for subsistence from gillnet openings. No commercial fishing occurred in Upper Yukon Area. | Season commenced July 1 with an opening available for limited First Nation (FN) Chinook Fishery. FN Governments manage FN Fisheries as per Yukon First Nation Self-Governing Agreements. FNs initiate harvest in conservative manner. Late July, FN Governments advised to implement additional precautionary measures due to lower than expected passage at Eagle sonar and unlikelihood of achieving the IMEG. Early August FN Governments implement voluntary Chinook harvest restriction followed by harvest restriction for chum. | Conditions of licence in the commercial and domestic fisheries obligated harvesters to gillnets with 6 inch or smaller mesh size; Chinook and chum commercial and domestic fisheries closed for duration of the season. Chinook and chum retention prohibited in the public angling fishery from June 26 to November 30 and September 11 to November 30, respectively. Public angling fishery closed from July 29 to November 30. Public angling fishery effectively closed for duration of salmon season. |

-continued-

Appendix B19.–Page 10 of 10.

| Year | U.S. management actions (subsistence)  | U.S. management actions (commercial) | Canadian management actions (First Nation fishery)  | Canadian management actions (commercial, domestic, public angling)   |
|------|--|--------------------------------------|---|--|
| 2021 | Subsistence salmon fishing closed starting on June 2 in District 1 and the Coastal District when the first Chinook salmon began entering the river. All districts, subdistricts, and tributaries closed based on run timing of early Chinook salmon. Salmon fishing remained closed all season. Gillnets with 4 inch or smaller mesh size was allowed for harvest of nonsalmon but were restricted to 60 feet or less in length. | No commercial fishing occurred.      | Season commenced July 1 with no harvest allocation for First Nation (FN) Chinook Fishery. FN Governments manage FN Fisheries as per Yukon First Nation Self-Governing Agreements. Early July, FN Governments advised to not harvest Chinook due to lower than expected passage at Eagle sonar and unlikeliness of achieving the IMEG. | Conditions of licence in the commercial and domestic fisheries restricted harvesters to gillnets with 6" or smaller mesh size; Chinook commercial and domestic fisheries closed for duration of the season. Public angling fishery closed for duration of salmon season. |

*Note:* Personal Use (PU) and Sport Fisheries are not listed. PU fisheries which occur only in the Tanana River drainage and Sport Fisheries which occur primarily in US tributaries are therefore of no concern to Canadian Chinook Salmon stocks.

Appendix B20.—Yukon River Salmon Agreement specified obligations for harvest shares, border passage and spawning escapement for mainstem Canadian-origin fall chum salmon, 2001–2021.

| Year | Total<br>estimated<br>Canadian-<br>origin run<br>size <sup>a</sup> | Total allowable<br>catch (TAC) <sup>b</sup> |         | U.S. share (%) of<br>TAC <sup>c</sup> |         | U.S.<br>harvest of<br>Canadian-<br>origin <sup>d</sup> | Border<br>passage<br>objective <sup>e</sup> | Border<br>passage<br><sup>f</sup> | Canada share<br>(%) of TAC |         | Canada<br>mainstem<br>harvest | Yukon River<br>Treaty goal or<br>Panel Interim<br>Management<br>Escapement |         | Spawning<br>escapement<br><sup>h</sup> |
|------|--|---|---------|---------------------------------------|---------|--|---|-----------------------------------|----------------------------|---------|-------------------------------|--|---------|--|
|      |  |   |         |                                       |         |  |   |                                   |                            |         |                               | Goal <sup>g</sup>  |         |  |
|      |  | From  | To      | 65%                                   | 71%     |  |   |                                   | From                       | To      |                               |  |         |  |
| 2001 | 90,100   | 10,100                                      |         | 6,565                                 | 7,154   | 8,789  | 83,240                                      | 38,908                            | 2,946                      | 3,535   | 4,919                         | 80,000   |         | 33,989                                 |
| 2002 | 89,900   | 29,900                                      |         | 19,435                                | 21,179  | 4,848  | 69,593                                      | 91,808                            | 8,721                      | 10,465  | 6,158                         | 60,000   |         | 85,650                                 |
| 2003 | 170,800  | 105,800                                     |         | 68,770                                | 74,941  | 17,044   | 98,944                                      | 142,591                           | 30,859                     | 37,030  | 10,973                        | 65,000   |         | 131,618                                |
| 2004 | 181,300  | 116,300                                     |         | 75,595                                | 82,379  | 16,637   | 102,313                                     | 125,000                           | 33,921                     | 40,705  | 9,545                         | 65,000   |         | 115,455                                |
| 2005 | 504,500  | 439,500                                     |         | 237,750                               | 244,750 | 67,332   | 263,250                                     | 451,477                           | 194,750                    | 201,750 | 13,744                        | 65,000   |         | 437,733                                |
| 2006 | 284,200  | 204,200                                     |         | 120,100                               | 127,100 | 64,669   | 160,600                                     | 217,810                           | 77,100                     | 84,100  | 6,617                         | 80,000   |         | 211,193                                |
| 2007 | 278,500  | 198,500                                     |         | 117,250                               | 124,250 | 47,449   | 157,750                                     | 235,956                           | 74,250                     | 81,250  | 9,330                         | 80,000   |         | 226,626                                |
| 2008 | 237,000  | 157,000                                     |         | 96,500                                | 103,500 | 49,954   | 137,000                                     | 180,379                           | 53,500                     | 60,500  | 6,130                         | 80,000   |         | 174,249                                |
| 2009 | 128,000  | 48,000                                      |         | 31,200                                | 34,000  | 22,886   | 95,400                                      | 94,739                            | 14,000                     | 16,800  | 1,115                         | 80,000   |         | 93,624                                 |
| 2010 | 143,000  | 39,000                                      | 73,000  | 25,350                                | 51,708  | 18,601   | 88,463                                      | 121,580                           | 11,375                     | 25,550  | 3,709                         | 70,000   | 104,000 | 117,871                                |
| 2011 | 326,000  | 222,000                                     | 256,000 | 129,000                               | 153,000 | 79,882   | 168,000                                     | 211,929                           | 86,000                     | 110,000 | 6,312                         | 70,000   | 104,000 | 205,617                                |
| 2012 | 238,000  | 134,000                                     | 168,000 | 85,000                                | 109,000 | 97,394   | 124,000                                     | 141,648                           | 42,000                     | 66,000  | 3,905                         | 70,000   | 104,000 | 137,743                                |
| 2013 | 303,000  | 199,000                                     | 233,000 | 117,500                               | 141,500 | 87,985   | 156,500                                     | 204,149                           | 74,500                     | 98,500  | 3,887                         | 70,000   | 104,000 | 200,262                                |
| 2014 | 223,000  | 119,000                                     | 153,000 | 77,350                                | 101,500 | 50,098   | 116,604                                     | 159,846                           | 34,709                     | 58,500  | 3,050                         | 70,000   | 104,000 | 156,796                                |
| 2015 | 205,000  | 101,000                                     | 135,000 | 65,650                                | 92,500  | 69,583   | 109,479                                     | 112,555                           | 29,459                     | 49,500  | 3,897                         | 70,000   | 104,000 | 108,658                                |
| 2016 | 298,000  | 194,000                                     | 228,000 | 115,000                               | 139,000 | 137,749  | 154,000                                     | 148,012                           | 72,000                     | 96,000  | 2,745                         | 70,000   | 104,000 | 145,267                                |
| 2017 | 563,000  | 459,000                                     | 493,000 | 247,500                               | 271,500 | 144,167  | 286,500                                     | 404,989                           | 204,500                    | 228,500 | 3,404                         | 70,000   | 104,000 | 401,585                                |
| 2018 | 279,000  | 175,000                                     | 209,000 | 105,500                               | 129,500 | 113,426  | 144,500                                     | 157,083                           | 62,500                     | 86,500  | 2,957                         | 70,000   | 104,000 | 154,126                                |
| 2019 | 178,000  | 74,000                                      | 108,000 | 48,100                                | 76,500  | 83,226   | 99,692                                      | 102,497                           | 21,584                     | 37,800  | 2,759                         | 70,000   | 104,000 | 99,738                                 |
| 2020 | 25,000   | 0   | 0       | 0                                     | 0       | 1,561  | 70,000                                      | 23,512                            | 0                          | 0       | 0                             | 70,000   | 104,000 | 23,512                                 |
| 2021 | 23,000   | 0   | 0       | 0                                     | 0       | 176  | 70,000                                      | 23,170                            | 0                          | 0       | 0                             | 70,000   | 104,000 | 23,170                                 |

-continued-

*Note:* The table does not represent a dataset, its intent is to represent the information at the time. Data presented for each year is from the assessment methods of that year. Harvest numbers were taken from the following year JTC summary report. Gray shaded boxes indicate Yukon River Salmon Agreement performance obligations that were not met.

- <sup>a</sup> For 2001 to 2002 values were not specifically presented in JTC reports, and have been retroactively calculated. 2003 and 2004 values were preliminary and taken from the 2005 JTC summary report.
- <sup>b</sup> Total run size, total allowable catch (TAC) and harvest share calculations are finalized post-season. TAC is calculated by subtracting the IMEG from the total run size. Delivering the IMEG plus the midpoint of Canada's harvest share to the Alaska-Yukon border is part of the U.S. obligation as per the Pacific Salmon Treaty's Yukon River Salmon Agreement.
- <sup>c</sup> Includes 50% of the portion of total allowable catch if greater than 120,000 chum salmon.
- <sup>d</sup> Assumed Canadian portion is 25% for all years.
- <sup>e</sup> Border passage objective is calculated post season as the agreed spawning escapement goal plus the mid-point of the Canadian harvest share. For years where the escapement goal is a range, this is represented as the average of the Canadian harvest shares, plus the lower end of the escapement goal.
- <sup>f</sup> From 2001 to 2007 the border passage was estimated from a mark recapture project. From 2008 on border passage was estimated from the Eagle sonar, minus any Alaskan harvest upstream of the sonar. The bold horizontal line between 2007 and 2008 indicates the JTC's recommendation to use the Eagle sonar as the primary assessment tool for the border passage estimate. Values from this year forward are sonar based.
- <sup>g</sup> Yukon River Panel goals have changed over time, and have been both points and ranges. IMEGs are not biologically based escapement goals.
- <sup>h</sup> Spawning escapement is calculated as the border passage estimate minus the harvest in Canada.

Appendix B21.–Summary of management and conservation measures implemented in the U.S. (Alaska) fall season fisheries and Canada Yukon mainstem for fall chum salmon fisheries, 2001–2021.

| Year | <u>U.S. management actions</u>  |  | <u>Canada management actions</u>  |   |
|------|---|--|---|---|
|      | Subsistence   | Commercial                                     | First Nation fishery  | Commercial, Domestic, and Public Angling  |
| 2001 | Full and partial closures to begin season, followed by full regulatory schedules. | Closed   | Unrestricted  | Commercial closed, limited to one 48 hr opening Sep. 12 to 14   |
| 2002 | Full schedule to begin season, time and gear restrictions later in season.        | Closed   | Early season restrictions due to low escapement projections, restrictions lifted Sep. 25. | Commercial and domestic closed except two, 96 hr openings between Oct. 2-13. Public angling implemented non salmon retention Aug. 20. |
| 2003 | Started season restricted then on regulatory schedules by mid season.             | Only directed at coho salmon at end of season. | Unrestricted  | Commercial fishery opened Sep. 7-9. Commercial fishery opened 5 days/week from Sep. 9-Oct. 24.  |
| 2004 | Started on windows schedule relaxed mid season.                                   | Only directed at coho salmon at end of season. | Unrestricted  | Commercial and domestic opened Sep. 12-14, then open 4-5 days for the following 4 weeks. Public angling open                          |
| 2005 | Relaxed subsistence schedule.   | Delayed opening to first quarter point.        | Unrestricted  | Commercial and Domestic opened Aug. 27 for 5 days, open continuously Sep. 3-Oct. 15. Public angling open                              |
| 2006 | Relaxed subsistence schedule.   | Delayed opening to first quarter point.        | Unrestricted  | Commercial and domestic opened Sep. 3 for 4 days. Open Sep. 10 for 4 days. Open Sep. 17 and 30 for 5 days. Open Oct. 1-14.            |

-continued-

Appendix B21.–Page 2 of 3

| Year | <u>U.S. management actions</u>                  |   | <u>Canada management actions</u> |   |
|------|---|---|----------------------------------|---|
|      | Subsistence                                     | Commercial  | First Nation fishery             | Commercial, Domestic, and Public Angling  |
| 2007 | Open on schedule.                               | Delayed opening to mid-point.   | Unrestricted                     | Commercial and domestic open Sep. 18 for 7 days and Sep. 28 for 14 days (21 days total).  |
| 2008 | Open on schedule.                               | Fished July during summer to fall transition and after three quarter point. | Unrestricted                     | Commercial and domestic open Aug. 31 for 4 days, open Sep. 5-9, Sep. 12-16, and Sep. 19-Oct. 7.   |
| 2009 | Open on schedule-some restrictions were taken.  | Fished during summer to fall transition and after three quarter point.      | Unrestricted                     | Commercial and domestic closed in the early season. Limited 4 day opening Oct. 8-12.  |
| 2010 | Open on schedule-some restrictions were taken.  | Only directed at coho salmon in September.                                  | Unrestricted                     | Commercial and domestic limited 24 hour opening, Sep. 22-23. Public angling open  |
| 2011 | Open on schedule.                               | Open throughout season  | Unrestricted                     | Commercial and domestic opened Aug. 26; two 4 day openings on Sep. 2 and Sep. 9; open Sep. 16-Oct. 16. Public angling open.                             |
| 2012 | Open on schedule then relaxed to 7 days a week. | Open throughout season.   | Unrestricted                     | Commercial and domestic open on Aug. 31 for 4 days, open continuously Sep. 7-Oct. 31. Public angling open.  |
| 2013 | Open on schedule then relaxed to 7 days a week. | Fished during summer to fall transition and after three quarter point.      | Unrestricted                     | Commercial and domestic fishery open Aug. 27 for 3 days; open continuously Sep. 2-Oct. 14. Public angling open.   |
| 2014 | Open on schedule then relaxed to 7 days a week. | Open throughout season.   | Unrestricted                     | Commercial and domestic opened Aug. 26 for 6 days below confluence of Yukon River and Coffee Ck.; open continuously Sep. 3-Oct.31. Public angling open. |

-continued-

Appendix B21.—Page 3 of 3.

| Year | <u>U.S. management actions</u>  |                         |                      | <u>Canada management actions</u>   |
|------|---|-------------------------|----------------------|--|
|      | Subsistence   | Commercial              | First Nation fishery | Commercial, Domestic, and Public Angling   |
| 2015 | Open on schedule then relaxed to 7 days a week. Porcupine River mainstem, some restrictions were implemented.             | Open throughout season. | Unrestricted         | Commercial and domestic open Aug. 28 for 5 days below confluence of Yukon River and Coffee Ck.; open continuously Sep. 4-Oct. 21. Public angling open.     |
| 2016 | Open on schedule then relaxed to 7 days a week. Porcupine River mainstem, some restrictions were implemented.             | Open throughout season. | Unrestricted         | Commercial and domestic open Aug. 30 for 8 days below the confluence of Yukon River and Coffee Ck.; open continuously Sep. 8-Oct. 21. Public angling open. |
| 2017 | Open on schedule then relaxed to 7 days a week. Porcupine River mainstem, some restrictions were implemented.             | Open throughout season. | Unrestricted         | Commercial, domestic and public angling fisheries open Aug. 31-Oct. 19.  |
| 2018 | Open on schedule then relaxed to 7 days a week on mainstem. Porcupine River mainstem, some restrictions were implemented. | Open throughout season. | Unrestricted         | Commercial, domestic and public angling fisheries open Aug. 31-Oct. 19.  |
| 2019 | Open on schedule then relaxed to 7 days a week on mainstem. Porcupine River mainstem closed all season.                   | Open throughout season. | Unrestricted         | Fishery opening delayed to protect late running Chinook salmon. Commercial and domestic opened Sep. 12-Oct. 31. Public angling open.                       |
| 2020 | Open on schedule then restricted followed by full closure. Porcupine River drainage closed all season.                    | Commercial closed.      | Closed               | Commercial, domestic, public angling closed for the season.  |
| 2021 | Closed all season. Porcupine River drainage closed all season.  | Commercial closed.      | Closed               | Commercial, domestic, public angling closed for the season; Chum catch and retention limits varied to zero at beginning of season.                         |

*Note:* Personal Use (PU) and Sport Fisheries are not listed. PU fisheries occur only in the Tanana River drainage and are not bound for Canada and sport fisheries do not occur on fall chum salmon.

Appendix B22.—Yukon River Salmon Agreement specified obligations for spawning escapement for Fishing Branch River fall chum salmon, 2001–2021.

| Year              | Total estimated<br>Fishing Branch<br>River run size <sup>a</sup> | Estimated % of<br>Fishing Branch<br>River stock<br>within Canadian<br>Porcupine River<br>stock <sup>b</sup> | Canada Fishing<br>Branch River<br>harvest <sup>c</sup> | U.S. Fishing<br>Branch River<br>harvest <sup>d</sup> | Yukon River Treaty goal or<br>Panel interim management<br>escapement goal <sup>e</sup> |         | Spawning<br>Escapement |
|-------------------|--|---|--|--|--|---------|------------------------|
|                   |  |   |  |  | From   | To      |                        |
| 2001              |  |   |  |  | 50,000   | 120,000 | 21,669                 |
| 2002              |  |   |  |  | 50,000   | 120,000 | 13,563                 |
| 2003              | 35,112   | 88  |  |  |  | 15,000  | 29,519                 |
| 2004              | 25,600   | 80  |  |  |  | 13,000  | 20,274                 |
| 2005              | 151,760  | 80  |  |  |  | 24,000  | 121,413                |
| 2006              | 38,560   | 80  |  |  |  | 28,000  | 30,849                 |
| 2007              | 42,160   | 80  |  |  |  | 34,000  | 33,750                 |
| 2008              | 24,800   | 80  |  |  | 22,000   | 49,000  | 20,055                 |
| 2009              | 32,000   | 80  |  |  | 22,000   | 49,000  | 25,828                 |
| 2010              | 16,000   | 80  |  |  | 22,000   | 49,000  | 15,773                 |
| 2011              | 21,000   | 75  | 1,388  |  | 22,000   | 49,000  | 13,085                 |
| 2012              | 37,500   | 75  | 2,339  | 19,479   | 22,000   | 49,000  | 22,399                 |
| 2013 <sup>f</sup> | 36,705   | 74  | 1,689  | 10,306   | 22,000   | 49,000  | 25,376                 |
| 2014 <sup>f</sup> | 9,998  | 46  | 912  | 1,830  | 22,000   | 49,000  | 7,304                  |
| 2015              | 13,000   | 73  | 406  | 4,136  | 22,000   | 49,000  | 8,351                  |
| 2016              | 54,000   | 80  | 2,404  | 22,040   | 22,000   | 49,000  | 29,397                 |
| 2017              | 73,000   | 80  | 1,850  | 23,067   | 22,000   | 49,000  | 48,524                 |
| 2018              | 29,000   | 80  | 1,499  | 17,680   | 22,000   | 49,000  | 10,151                 |
| 2019              | 29,000   | 66  | 660  | 10,366   | 22,000   | 49,000  | 18,171                 |
| 2020              | 5,000  | 63  | 63   | 250  | 22,000   | 49,000  | 4,795                  |
| 2021              | 2,500  | 69  | 14   | 28   | 22,000   | 49,000  | 2,413                  |

-continued-



*Note:* The table does not represent a dataset, its intent is to represent the information at the time. Data presented for each year is from the assessment methods of that year. Harvest numbers were taken from the following year JTC summary report. Gray shaded boxes indicate Yukon River Salmon Agreement performance obligations that were not met.

- <sup>a</sup> Total run size is finalized post-season. 2003-2012 values are calculated using reported proportion of Fishing Branch River chum salmon within reported Porcupine River Total Run Size from summary year's JTC report. 2013 and 2014 values are calculated using Porcupine River sonar counts and the proportion of tagged chum salmon that reached Fishing Branch, plus proportion of Fishing Branch chum salmon within U.S. harvest.
- <sup>b</sup> Fishing Branch proportions of Porcupine River stock are presented as published in that year's JTC Report (2003 ,2004, 2011 - 2021), except for 2005-2010, when they were assumed to follow the 80% proportion detailed in the 2004/2005 JTC Report.
- <sup>c</sup> Prior to 2011, annual Canadian harvest of Fishing Branch River chum salmon was not considered in total run size calculation (with the exception of 2003).
- <sup>d</sup> Prior to 2012, annual U.S. harvest was not considered in total run size calculation. For 2012, U.S. harvest of Porcupine River chum salmon is considered 5% of total U.S. harvest. From 2013-2015, the proportion of Fishing Branch River chum salmon within total U.S. harvest was assumed to be equal to the proportion of Fishing Branch River escapement in the drainagewide escapement. From 2016-present, U.S. harvest of Fishing Branch River chum salmon is assumed to be 4% of total U.S. harvest.
- <sup>e</sup> Yukon River Panel goals have changed over time, and have been both points and ranges. Interim management escapement goals (IMEG) are not biologically based escapement goals.
- <sup>f</sup> Fishing Branch River weir did not operate. Escapement was estimated from a sonar operated on the upper Porcupine River minus upstream Old Crow harvest then multiplied by the proportion of tags to Fishing Branch River. Escapement taken from 2015 summary JTC report.

Appendix B23.–Summary of management and conservation measures implemented for fall chum salmon in the U.S. (Alaska) and Canada on the Porcupine River, 2001–2021.

| Year | <u>Subsistence</u>   | <u>First Nation fishery</u>  |
|------|--|--|
|      | U.S. management actions  | Canada management actions  |
| 2001 | Open   | Open   |
| 2002 | Closed to begin fall season, followed by some restrictions, open at end of season.                             | Porcupine River restrictions to 25% of normal allocation. Vuntut Gwitchin restricted to 2 days/week from Sep. 4-Oct. 11. |
| 2003 | Open with some restrictions.   | Closed Aug. 10-Oct. 15.  |
| 2004 | Open   | Voluntary closure Aug. 10-Oct. 15.   |
| 2005 | Open   | Open   |
| 2006 | Open   | Open   |
| 2007 | Open   | Open   |
| 2008 | Open   | Open   |
| 2009 | Open, followed by some restrictions taken, open at end of season.  | Closed from noon Sep. 21-noon Oct. 1.  |
| 2010 | Open   | Open   |
| 2011 | Open   | Open   |
| 2012 | Open   | Open   |
| 2013 | Open   | Open   |
| 2014 | Open   | Conservative harvest suggested.  |
| 2015 | Porcupine River mainstem closed all fall season.   | Recommend no fishery.  |
| 2016 | Porcupine River mainstem closed at start of fall season, followed by some restrictions, open at end of season. | Conservative harvest suggested.  |
| 2017 | Open, then some restrictions on Porcupine River mainstem, open at end of season.                               | Conservative harvest suggested.  |
| 2018 | Open, then some restrictions on Porcupine River mainstem, followed by closure.                                 | Conservative harvest suggested.  |
| 2019 | Porcupine River mainstem closed all fall season.   | Conservative harvest suggested.  |
| 2020 | Porcupine River drainage closed all fall season.   | Closed   |
| 2021 | Porcupine River drainage closed all fall season.   | Closed   |

*Note:* Personal Use (PU) and Sport Fisheries are not listed. PU fisheries occur only in the Tanana River drainage and are not bound for Canada and sport fisheries do not occur on fall chum salmon.

**APPENDIX C: BERING SEA-ALEUTIAN ISLANDS  
BYCATCH SUMMARY AND IMPACT ON YUKON RIVER  
CANADIAN-ORIGIN SALMON**

## **Yukon River Salmon Bycatch Summary**

**January 2022, DRAFT**

The Yukon River Salmon Agreement identifies the need to identify, quantify, and undertake efforts to reduce marine catches and bycatch of Yukon River salmon. This section provides an overview of information on U.S. groundfish fisheries in the Bering Sea-Aleutian Islands (BSAI) management region, bycatch regulations, and bycatch impacts on Yukon River Canadian-origin salmon.

### ***Bycatch impacts on Canadian-origin salmon***

Yukon River Canadian-origin salmon are caught as bycatch in BSAI groundfish fisheries along with other salmon stocks from Alaska, the west coast of Canada and the United States, eastern Asia, and Russia. The total number of salmon captured as bycatch is always much greater than the number of returning adult Canadian-origin salmon that are removed from the Yukon River due to bycatch. For example, the total annual bycatch of Chinook salmon in BSAI pollock fishery has varied from approximately 5,000 to 122,000 (Table 1), but the adult equivalent (AEQ) bycatch of Canadian-origin Chinook salmon varied from approximately 400 to 2,400 fish over the same time period (Table 2). The average bycatch impact rate by the pollock fishery on the Canadian-origin Chinook salmon run is estimated to be 1.0% with an annual impact rate less than 3.1% (Ianelli and Stram, 2018). Average bycatch impact rates to western Alaska chum salmon (not Canadian-origin chum salmon) is estimated to be 0.4% with an annual rate less than 1.3% (Murphy et al. 2017). Ongoing regulatory and management measures implemented by the North Pacific Fisheries Management Council (NPFMC) are a key factor limiting bycatch impact rates on Canadian-origin salmon in BSAI groundfish fisheries.

### ***Current BSAI bycatch information***

- Total bycatch of Chinook salmon in BSAI groundfish fisheries (pelagic trawl, bottom trawl, and hook-and-line fisheries) during 2021 ( $n=15,827$ ) was 52% lower than the recent 5-year average (Table 1). Chinook salmon bycatch in the BSAI pollock fishery accounted for 87% ( $n=13,783$ ) of the bycatch during 2021.
- Total bycatch of non-Chinook salmon (primarily chum salmon) in BSAI groundfish fisheries (pelagic trawl, bottom trawl, and hook-and-line fisheries) during 2021 ( $n=535,282$ ) was a 49% increase in the recent 5-year average (Table 1). Bycatch of non-Chinook salmon in the BSAI pollock fishery accounted for 99% ( $n=530,626$ ) of the bycatch during 2021.
- Bycatch impacts to Canadian-origin Chinook salmon by BSAI Pollock fishery is estimated by run year. The 2017 run is the most recent year for which bycatch impact estimates are available for Canadian-origin Chinook salmon.
  - The total Canadian-origin Chinook salmon run in 2017 was 93,188. Adult equivalent models estimate that an additional 772 Canadian-origin Chinook salmon would have contributed to the 2017 run if they had not been captured as bycatch in the BSAI pollock fishery (Table 2). This represents an impact rate of 0.83% on the Canadian-origin Chinook salmon run during 2017.

## Background Information

### *Bycatch management*

- U.S. groundfish trawl fisheries in the BSAI management area are managed to limit the bycatch of salmon under the Magnuson-Stevens Fisheries Conservation and Management Act by the NPFMC and are regulated by National Marine Fisheries Service (NMFS).
- The pollock fishery is the primary focus of bycatch management as it accounts for an average of 88% of the total Chinook salmon bycatch and 99% of the non-Chinook salmon bycatch in the BSAI management area.
- The pollock fishery is managed according to the Fishery Management Plan (FMP) for Groundfish of the BSAI Management Area.

<https://www.npfmc.org/wp-content/PDFdocuments/fmp/BSAI/BSAIfmp.pdf>

### *Bycatch regulations*

- The BSAI groundfish FMP contains regulatory measures to reduce salmon bycatch.
- The BSAI pollock fishery is one of the most heavily regulated and monitored fisheries in the world and includes 100% observer coverage.
- Notable bycatch reduction measures include amendment 91 and amendment 110.
- Amendment 91 (<https://alaskafisheries.noaa.gov/rules-notices/search>) was implemented in 2011 and, among other things, established bycatch caps.
- Amendment 110 (<https://alaskafisheries.noaa.gov/rules-notices/search>) was implemented in 2016 and, among other things, established abundance-based bycatch caps to further protect western Alaska and Canadian-origin Chinook salmon stocks harvested for subsistence purposes. Bycatch caps are set relative to the combined in-river run size for the Unalakleet, Upper Yukon (Canadian-origin), and Kuskokwim River Chinook salmon stock groups (termed the three-system index).

### *Bycatch impact methods*

- The number of salmon captured as bycatch in a given year is not equivalent to the number of adult salmon that would have returned to the Canadian portion of the Yukon River drainage in that year for two reasons.
  - Salmon stocks throughout the North Pacific are captured as bycatch in the BSAI groundfish fisheries. Information on stock origin is required to evaluate the impact of bycatch to a given stock or stock group.
  - Salmon are predominately captured as bycatch during their immature life-history stage and will spend one or more years in the ocean before returning to freshwater. Bycatch numbers of immature salmon require an adjustment for natural mortality before they can be compared to the number of mature adults returning to freshwater. Bycatch estimates that are adjusted for natural mortality are referred to as Adult Equivalent (AEQ) bycatch.
- Bycatch impacts on Yukon River Canadian-origin salmon require stock-specific Adult Equivalent (AEQ) estimates of bycatch. These estimates rely on the following data inputs: total salmon bycatch, bycatch stock mixtures, bycatch age composition, salmon maturity schedules, and assumptions on the natural mortality of salmon in marine habitats (Ianelli and Stram 2014).

- The bycatch AEQ analysis has not been updated since the last Yukon River Panel bycatch summary. AEQ analysis may not be updated annually depending on the regulatory application and need through the NPFMC. Updated AEQ analysis will be reported in the annual Yukon River Panel bycatch summary as it occurs.

### ***Additional resources***

- Bycatch numbers are reported by the National Marine Fisheries Service, available at: <https://alaskafisheries.noaa.gov/fisheries-catch-landings?tid=286>
- Bycatch updates are reported by the North Pacific Fisheries Management Council, available at: <https://www.npfmc.org/bsai-salmon-bycatch/>

### ***References***

- Ianelli, J. N. and D. L. Stram. 2014. Estimating impacts of the pollock fishery bycatch on western Alaska Chinook salmon. *ICES J. Mar. Sci.* **72**: 1159-1172. doi:10.1093/icesjms/fsu173
- Ianelli, J. N., and D. L. Stram. 2018. Chinook Bycatch Mortality Update. Discussion paper presented to the North Pacific Fishery Management Council, April 2018. Available online at: <http://npfmc.legistar.com/gateway.aspx?M=F&ID=e172520e-fc22-46e8-b5aa-72ba233f129e.pdf>
- Murphy, J.M. E.V. Farley, J.N. Ianelli, and D.L. Stram. 2017. Distribution, diet, and bycatch of chum salmon in the Eastern Bering Sea. *N. Pac. Anadr. Fish. Comm. Bull.* 6:219-234. doi: 10.23849/npafcb6/219.234

Table 1.—Numbers of Chinook and non-Chinook (chum) salmon captured as bycatch in the Bering Sea-Aleutian Islands (BSAI) groundfish fisheries by season (A-season: winter, B-season: summer/fall), 1991–2021.

| Year              | BSAI Chinook Salmon Bycatch |               |                   |               |                   |               | BSAI Non-Chinook Salmon Bycatch |               |                   |               |                   |               |
|-------------------|-----------------------------|---------------|-------------------|---------------|-------------------|---------------|---------------------------------|---------------|-------------------|---------------|-------------------|---------------|
|                   | A-season                    |               | B-season          |               | Annual            |               | A-season                        |               | B-season          |               | Annual            |               |
|                   | Pollock Fisheries           | All Fisheries | Pollock Fisheries | All Fisheries | Pollock Fisheries | All Fisheries | Pollock Fisheries               | All Fisheries | Pollock Fisheries | All Fisheries | Pollock Fisheries | All Fisheries |
| 1991 <sup>a</sup> | 38,791                      | 46,392        | 2,114             | 2,488         | 40,905            | 48,880        | 2,850                           | 3,015         | 26,101            | 27,245        | 28,951            | 30,260        |
| 1992 <sup>a</sup> | 25,691                      | 31,418        | 10,259            | 10,536        | 35,950            | 41,954        | 1,951                           | 2,120         | 38,324            | 39,329        | 40,275            | 41,449        |
| 1993 <sup>a</sup> | 17,264                      | 24,688        | 21,252            | 21,325        | 38,516            | 46,013        | 1,593                           | 1,848         | 240,597           | 241,422       | 242,191           | 243,270       |
| 1994              | 28,451                      | 38,921        | 4,686             | 4,899         | 33,137            | 43,820        | 3,990                           | 5,599         | 88,681            | 88,949        | 92,672            | 94,548        |
| 1995              | 10,579                      | 18,939        | 4,405             | 4,497         | 14,984            | 23,436        | 1,707                           | 3,033         | 17,556            | 18,842        | 19,264            | 21,875        |
| 1996              | 36,068                      | 43,316        | 19,554            | 19,888        | 55,622            | 63,204        | 221                             | 665           | 77,014            | 77,395        | 77,236            | 78,060        |
| 1997              | 10,935                      | 16,401        | 33,973            | 34,128        | 44,908            | 50,529        | 2,083                           | 2,710         | 63,904            | 64,285        | 65,987            | 66,995        |
| 1998              | 16,132                      | 19,869        | 40,308            | 40,679        | 56,440            | 60,548        | 4,090                           | 4,520         | 60,866            | 61,177        | 64,956            | 65,697        |
| 1999              | 6,352                       | 8,793         | 5,627             | 5,805         | 11,979            | 14,598        | 362                             | 393           | 44,909            | 46,739        | 45,271            | 47,132        |
| 2000              | 3,422                       | 6,567         | 1,539             | 1,655         | 4,961             | 8,222         | 212                             | 350           | 58,358            | 58,976        | 58,571            | 59,326        |
| 2001              | 18,484                      | 24,871        | 14,961            | 15,676        | 33,445            | 40,547        | 2,386                           | 2,903         | 54,621            | 57,827        | 57,007            | 60,730        |
| 2002              | 21,794                      | 26,276        | 12,701            | 13,407        | 34,495            | 39,683        | 1,377                           | 1,697         | 79,274            | 80,784        | 80,651            | 82,481        |
| 2003              | 33,478                      | 40,058        | 13,055            | 13,603        | 45,661            | 53,661        | 3,831                           | 3,831         | 184,513           | 184,559       | 188,344           | 188,390       |
| 2004              | 24,925                      | 30,766        | 26,663            | 29,272        | 51,762            | 60,038        | 426                             | 426           | 451,907           | 452,131       | 452,333           | 452,560       |
| 2005              | 27,960                      | 33,622        | 40,861            | 41,462        | 68,184            | 75,084        | 594                             | 594           | 710,196           | 710,926       | 710,790           | 711,520       |
| 2006              | 58,547                      | 62,547        | 24,362            | 24,568        | 82,752            | 87,115        | 1,323                           | 1,323         | 305,674           | 305,852       | 306,997           | 307,175       |
| 2007              | 72,943                      | 78,156        | 51,781            | 51,844        | 122,195           | 130,000       | 8,481                           | 8,489         | 84,387            | 85,152        | 92,868            | 93,641        |
| 2008              | 16,495                      | 18,828        | 4,811             | 5,009         | 21,307            | 23,837        | 247                             | 247           | 14,732            | 14,732        | 14,980            | 14,980        |
| 2009              | 9,882                       | 11,289        | 2,697             | 2,825         | 12,579            | 14,115        | 48                              | 48            | 45,397            | 45,397        | 45,445            | 45,445        |
| 2010              | 7,649                       | 9,480         | 2,069             | 2,921         | 9,737             | 12,399        | 40                              | 40            | 13,238            | 13,237        | 13,278            | 13,278        |
| 2011              | 7,137                       | 7,602         | 18,362            | 19,007        | 25,499            | 26,609        | 297                             | 414           | 191,138           | 194,405       | 191,435           | 194,819       |
| 2012              | 7,765                       | 8,981         | 3,578             | 3,949         | 11,343            | 12,929        | 11                              | 307           | 22,172            | 23,766        | 22,183            | 24,073        |
| 2013              | 8,237                       | 9,186         | 4,797             | 6,821         | 13,016            | 15,989        | 215                             | 447           | 125,101           | 126,554       | 125,316           | 127,001       |
| 2014              | 11,539                      | 13,837        | 3,498             | 4,261         | 15,037            | 18,106        | 577                             | 1,629         | 218,865           | 222,634       | 219,442           | 224,263       |
| 2015              | 12,304                      | 17,502        | 6,025             | 7,752         | 18,329            | 25,254        | 4,756                           | 6,158         | 232,996           | 237,196       | 237,752           | 243,354       |
| 2016              | 16,828                      | 25,721        | 5,098             | 6,840         | 21,926            | 32,568        | 3,903                           | 4,838         | 339,098           | 342,503       | 343,001           | 347,341       |
| 2017              | 21,828                      | 27,008        | 8,248             | 9,272         | 30,076            | 36,277        | 1,906                           | 2,313         | 465,772           | 469,134       | 467,678           | 471,447       |
| 2018              | 8,631                       | 11,251        | 5,095             | 6,130         | 13,740            | 17,394        | 1,201                           | 2,120         | 293,863           | 306,926       | 295,064           | 309,045       |
| 2019              | 15,781                      | 20,088        | 9,203             | 11,323        | 24,984            | 31,412        | 2,239                           | 4,509         | 345,643           | 354,294       | 347,882           | 358,804       |
| 2020              | 18,369                      | 20,436        | 13,925            | 14,531        | 32,294            | 34,967        | 807                             | 1,161         | 319,338           | 321,540       | 320,478           | 323,032       |
| 2021              | 9,502                       | 10,718        | 4,281             | 5,109         | 13,783            | 15,827        | 160                             | 372           | 530,466           | 534,910       | 530,626           | 535,282       |

[https://www.fisheries.noaa.gov/sites/default/files/akro/chinook\\_salmon\\_mortality2021.html](https://www.fisheries.noaa.gov/sites/default/files/akro/chinook_salmon_mortality2021.html);

[https://www.fisheries.noaa.gov/sites/default/files/akro/chum\\_salmon\\_mortality2021.html](https://www.fisheries.noaa.gov/sites/default/files/akro/chum_salmon_mortality2021.html)

<sup>a</sup> Community Development Quota (CDQ) bycatch not included.

Table 2.—Estimated adult equivalent (AEQ) bycatch of Canadian-origin Chinook salmon from the Yukon River in the Bering Sea-Aleutian Islands (BSAI) pollock fisheries by run year, run size of the Canadian-origin Chinook salmon, and bycatch exploitation rates, 1994–2017 (Ianelli and Stram, 2018).

| Run Year | Canadian-Origin AEQ Bycatch | Canadian-Origin Run | Canadian-Origin Impact Rate |
|----------|-----------------------------|---------------------|-----------------------------|
| 1994     | 1,035                       | 172,885             | 0.60%                       |
| 1995     | 817                         | 169,789             | 0.48%                       |
| 1996     | 998                         | 182,504             | 0.55%                       |
| 1997     | 995                         | 161,700             | 0.62%                       |
| 1998     | 760                         | 88,282              | 0.86%                       |
| 1999     | 588                         | 110,446             | 0.53%                       |
| 2000     | 347                         | 52,842              | 0.66%                       |
| 2001     | 508                         | 85,663              | 0.59%                       |
| 2002     | 835                         | 81,487              | 1.02%                       |
| 2003     | 1,044                       | 149,979             | 0.70%                       |
| 2004     | 1,214                       | 117,247             | 1.04%                       |
| 2005     | 1,267                       | 123,612             | 1.02%                       |
| 2006     | 1,843                       | 119,485             | 1.54%                       |
| 2007     | 2,361                       | 87,899              | 2.69%                       |
| 2008     | 1,918                       | 62,610              | 3.06%                       |
| 2009     | 1,127                       | 87,899              | 1.28%                       |
| 2010     | 518                         | 59,741              | 0.87%                       |
| 2011     | 359                         | 71,726              | 0.50%                       |
| 2012     | 351                         | 48,494              | 0.72%                       |
| 2013     | 364                         | 37,177              | 0.98%                       |
| 2014     | 401                         | 64,886              | 0.62%                       |
| 2015     | 455                         | 87,323              | 0.52%                       |
| 2016     | 532                         | 82,765              | 0.64%                       |
| 2017     | 772                         | 93,188              | 0.83%                       |