

(2) Alaska Department of Fish and Game, Division of Wildlife Conservation, 1300 College Rd., Fairbanks, Alaska, 99701 USA


Food limitation is among the hypothesized causes for the declines of various pinniped populations in the North Pacific, with juveniles identified as the age class most vulnerable to changes in prey availability. In PWS, harbor seal (*Phoca vitulina richardsi*) numbers have decreased by two-thirds since the early 1980's, only partly due to the 1989 Exxon Valdez oil spill. To test the food limitation hypothesis we measured body composition in newly weaned pups (n=69), yearling (n=46) and 2-3-year-old (n=14) harbor seals in late June 1997-2000 using isotope dilution. We obtained blubber biopsies and estimated non-pup diets using QFASA and a prey database of 26 species (n=1151). Pups were consistently heavy ( $29.4 \pm 0.50$  kg) and fat ( $42.2 \pm 0.68\%$ ) compared to other harbor seal populations, with only minor interannual variation. Yearlings and 2-3-year olds, respectively, were also comparatively and consistently fat ( $25.0 \pm 0.53\%$  and  $23.0 \pm 0.87\%$ ). Diets of these juveniles were dominated by herring ( $53.7 \pm 2.99\%$  and  $46.2 \pm 3.79\%$ ), pollock ( $15.0 \pm 1.66\%$  and  $21.2 \pm 1.92\%$ ) and yellowfin sole ( $18.3 \pm 1.49\%$  and  $20.1 \pm 2.78\%$ ). Although high-fat (19%) eulachon averaged only 4.5% of diets, it contributed >15% to blubber signatures and thus fat stores. Minor prey species included sandlance, shrimp, various flatfish and cephalopods. Body fat content was not related to any single species level in the diet or to diet energy density, however, it was positively correlated with Levin's measure of diet diversity ( $P < 0.001$ ,  $r = 0.54$ ); diet diversity did not differ with year ( $P = 0.536$ ). Our results suggest that lactating females were not nutritionally compromised, given the robust body condition of weaned pups. Additionally, juveniles with more diverse diets were in better condition independent of prey species eaten, provided that some high-quality prey were obtained to balance low-energy prey consumed. These data suggest that food limitation at this time of year or in these age classes was not a likely source of decline in this population.

**High Body Energy Stores and Condition are Linked with Diet Diversity in Juvenile Harbor Seals in Prince William Sound (PWS), Alaska: New Insights from Quantitative Fatty Acid Signature Analysis (QFASA)**

Iverson, Sara J.<sup>1</sup>; Frost, Kathryn J.<sup>2</sup>; Lang, Shelley L. C.<sup>1</sup>

(1) Department of Biology, Dalhousie University, Halifax, NS B3H 4J1 Canada

# 15<sup>th</sup> Biennial Conference



NORTH CAROLINA 2003

## on the Biology of Marine Mammals

Greensboro, NC, USA  
14 - 19 December 2003



**Society for Marine Mammalogy**

### CONFERENCE COMMITTEE

Edward O. Keith – *Chair*

Lemnuel Aragonés  
Edmund Gerstein  
Carolyn Kurle  
Frances Michaelis  
Alejandro Ortega  
Aldemaro Romero  
Caryn Self Sullivan

### SCIENTIFIC PROGRAM COMMITTEE

Aleta A. Hohn – *Chair*

Robin Baird  
Annalisa Berta  
Bob Bonde  
Frank Fish  
Tero Härkönen  
Denise Herzing  
Sascha Hooker  
Paul Jepson  
Richard Merrick  
Monica Muelbert  
Patricia Rosel  
Michael Scott  
Ursula Siebert  
Garry Stenson  
Rob Stewart  
Jim Sumich  
Fernando Trujillo  
Gordon Waring  
Doug Wartzok  
Ben Wilson  
Graham Worthy

Zoey Zahorodny – *Assistant to the  
Scientific Program Committee*