TROUT & OTHER SPECIES FOUND IN SOUTHEAST ALASKA




Green to bluish back with silvery to yellowish-green sides. Broad pinkish band along side and black spots on back, sides, and tail. Upper jaw usually does not extend past eye on adult.



CUTTHROAT TROUT

Yellowish-green body with no pinkish band along sides. Many black spots on back, sides, and tail. Upper jaw extends well past eye on adults. Red to orange slash on underside of lower jaw.






Waterbody **Community Rustabach Lake Haines Upper & Lower Dewey Lakes** Skagway **Dorothy Lake** Juneau Salmon Creek Reservoir Juneau Sitka Deep Lake Sitka **Green Lake** Sitka **Heart Lake** Long Lake Sitka **Thimbleberry Lake** Sitka **Crystal Lake** Petersburg Ketchikan **Grace Lake** Ketchikan Lake Ketchikan **Shelokum Lake** Ketchikan Ketchikan **Perseverance Lake Emerald (Texas)Lake** Hvder

Eastern Brook Trout are not native to Alaska and are found only in the bodies of water listed above.



Light brown to dark purple back, fading to bluish-gray with black spots on sides. Very large dorsal fin with red and violet spots.



Sea-run rainbow trout with bluish-gray back and bright silvery sides. Slender body profile; 8-10 anal fin rays. Small black spots on back, sides, and tail. Pink/red coloration develops after returning to fresh water to spawn.



The body has a background coloration that can range from emerald green to bluish-gray or silver and is covered with lots of small light red spots (usually smaller than its pupil). The tail is slightly forked and there are no distinct spots on head or tail. Lower fins are red or orange with a white anterior border. Easily confused with Eastern Brook Trout, but Dolly Varden do not have wormlike markings on their back or dorsal fin. Found both in fresh and salt waters.