

2.7-1 Ratio of ESSN: Sport Catch of Kasilof LR Kings

Table 10.—Late-run Kasilof River Chinook salmon harvest and abundance, 1996–2015.

Year	ESSN harvest ^a	Inriver sport harvest	Inriver abundance ^b
1996	3,365	833	ND
1997	3,296	1,101	ND
1998	1,480	637	ND
1999	2,754	658	ND
2000	1,072	1,086	ND
2001	1,749	1,378	ND
2002	2,758	451	ND
2003	4,310	1,144	ND
2004	6,310	1,038	ND
2005	6,285	1,052	12,097
2006	2,897	883	8,611
2007	3,577	1,062	8,522
2008	2,204	793	8,276
2009	1,626	2,164	ND
2010	2,305	1,310	ND
2011	2,538	1,660	ND
2012	205	55	ND
2013	637	762	ND
2014	891	314	ND
2015	1,564	790	ND
Min.	205	55	8,276
Average	→ 2,591	→ 959	9,377
Max.	6,310	2,164	12,097

Source: Alaska Sport Fishing Survey database [Internet]. 1996– . Anchorage, AK: Alaska Department of Fish and Game. Division of Sport Fish (cited December 2016). Available from: <http://www.adfg.alaska.gov/sf/sportfishingsurvey/>; Fleischman and McKinley (2013). Eskelin et al. (2013). Eskelin and Barclay. (2015, 2016), and Shields and Dupuis (2016).

Note: ND means no data collected.

^a Uses 2010, 2011, and 2013–2015 eastside setnet genetic stock allocation estimates to calculate Kasilof River Chinook salmon harvest component.

^b Mark-recapture tagging abundance estimates from Reimer and Fleischman (2012).

Source: ADFG Fishery Management Report
Begich et. al FMR 17-06

Submitted by: Kenai River Sport Fishing
Association