

**Interim Report to the Alaska Board of Game on
Intensive Management for Caribou
with Wolf Predation Control
in Game Management Units 9B, 17B&C, and 19A&B,
the Mulchatna Caribou Herd**

**Prepared by the Division of Wildlife Conservation
August 2014**



Interim annual updates are limited to sections that have changed substantially since the prior annual report in February. For complete information, see the prior annual report.

1) **Description of IM Program¹ and Department recommendation for reporting period**

A) **This report is an annual evaluation for a predation control program authorized by the Alaska Board of Game (Board) under 5 AAC 92.111**

B) **Month this report was submitted by the Department to the Board:**

February ___ (annual report) August X (interim annual update²) Year 2014

2) **Prey data**

Date(s) and method of most recent summer abundance assessment for caribou (if statistical variation available, describe method here and show result in Table 1)

The last successful photo-census of post-calving aggregation was conducted on July 2, 2014. Data are currently under analysis.

Compared to IM area, was a similar trend and magnitude of difference in abundance observed in nearby non-treatment area(s) since program inception (Y/N) N/A and in the last year (Y/N) N/A? Describe comparison if necessary:

Not Applicable: This program was initiated in March, 2012 (RY11). It is too early to determine trends in abundance that resulted from these activities.

Date(s) of most recent age and sex composition survey (if statistical variation available, describe method here and show result in Table 1):

October 23, 2013

Compared to IM area, was a similar composition trend and magnitude of difference in composition observed in nearby non-treatment area(s) since program inception (Y/N) N/A and in the last year (Y/N) N/A? Describe comparison if necessary:

Observed calf ratios in the eastern segment of the MCH with no predator control remain lower than in the western segment (14 and 23 calves:100 cows, respectively; Table 1). This program was initiated in March 2012 (RY2011), and it is too early to determine trends in bull ratios.

¹ For purpose and context of this report format, see *Intensive Management Protocol, section on Tools for Program Implementation and Assessment*

² The interim annual update may be limited only to sections that changed substantially since prior annual report

Table 1. Caribou abundance, age and sex composition in assessment area (L) since program implementation in year 1 (not exclusively limited to inception of predation control) to reauthorization review in year 2017 in Mulchatna Caribou Herd Predation Management Area. Regulatory year is 1 July to 30 June (e.g, RY 2010 is 1 July 2010 to 30 June 2011).

Eastern Segment of the MCH (No Predator Control)

		Composition (number per 100 cows)		
Period	RY	Calves	Bulls	Total <i>n</i>
Year 0	2010	17	13	2,581
Year 1	2011	14	18	2,649
Year 2	2012	22	17	2,217
Year 3	2013	14	27	1,479

Western Segment of the MCH (Active Predator Control)

		Composition (number per 100 cows)		
Period	RY	Calves	Bulls	Total <i>n</i>
Year 0	2010	23	23	2,011
Year 1	2011	28	34	1,995
Year 2	2012	38	29	2,636
Year 3	2013	23	27	1,743

All Areas Combined

		Composition (number per 100 cows)			
Period	RY	Abundance (variation)	Calves	Bulls	Total <i>n</i>
Year 0	2010	-	20	17	4,592
Year 1	2011	-	19	22	5,282 ^a
Year 2	2012	25,000-35,000 ^b	30	23	4,853
Year 3	2013	20,000-30,000 ^b	19	27	3,222

^a Includes caribou not assigned to the Eastern or Western Segment of the MCH.

^b Preliminary estimate of abundance based on the Rigest methodology (Rigest et al. 1998).

Describe trend in abundance or composition:

Not Applicable: This program was initiated in March 2012 (RY2011). It is too early to determine trends in abundance or composition resulting from these activities.

Table 2. Caribou harvest in assessment area (M). Methods for estimating unreported harvest are described in Survey and Inventory reports.

Period	RY	Reported			Estimated		Total harvest	Other mortality ^a	Total
		Male	Female	Unk Sex	Unreported	Illegal			
Year 0	2010 ^b	250	220	4	Unk	Unk	470	Unk	474
Year 1	2011 ^b	240	243	9	Unk	Unk	492	Unk	492
Year 2	2012 ^b	171	182	4	Unk	Unk	357	Unk	357
Year 3	2013 ^b	69	35	1	Unk	Unk	77	Unk	105

^aClarify (vehicle mortality, Defense of Life and Property, Mortuary, etc.).

^bData from ADF&G database, August 6, 2014.

Describe trend in harvest:

There has been a fairly steady decline in the reported harvest since 1999 due to a combination of progressively, more-restrictive hunting regulations, decreasing population size, and changes in caribou distribution.

Describe any other harvest related trend if appropriate:

During RY92-RY02 the reported harvest of bulls changed from greater than 74% bulls to an approximately equal bull:cow harvest. However, in RY13 the % bulls in the harvest increased to 69% although the overall harvest dropped dramatically from the previous year. The overall decrease in harvest in RY13 was partially due to the reduced snowfall, which led to poor winter travel conditions, and caribou distribution.

3) Predator data

Date(s) and method of most recent spring abundance assessment for wolves (if statistical variation available, describe method here and list in Table 3):

A minimum abundance estimate survey was conducted in February, 2012.

Date(s) and method of most recent fall abundance assessment for wolves (if statistical variation available, describe method here and list in Table 3):

Not Applicable: Fall abundance has not been estimated due to logistic and weather constraints.

Other research or evidence of trend or abundance status in wolves:

In RY11 local residents in some villages harvested an increased number of wolves in the wolf assessment area ‘N’. In RY12 and RY13 wolf harvest decreased as did the public control removal from the predation control area. This was at least in part due to very poor winter tracking (snow) conditions, but could also be a sign that wolf numbers have been reduced due to the high harvest in RY11. This is supported by local residents and air taxi pilots reporting fewer wolf sightings in the area.

Table 3. Wolf abundance objectives and removal in wolf assessment area (N) of Mulchatna Caribou Herd Predation Management Area. Removal objective is to annually remove 100 % of the wolves in the wolf predation control area (O), so estimated or confirmed number remaining in the control area (O) by the May calving season each regulatory year is 0.

Subunits 9B and 17B&C

Period	RY	Harvest removal from area N		Dept. control removal from area O	Public control removal from area O	Total removal ^a from area N	Minimum Spring abundance (variation) in area N
		Trap	Hunt				
Year 1	2011	25	69	0	11	104	14
Year 2 ^b	2012	0	18	0	4	18	-
Year 3 ^c	2013	8	2	0	0	10	-

^a Additional removal may be Defense of Life and Property, vehicle kill, etc.

^b ADF&G database, December 9, 2013.

^c ADF&G database, August 5, 2014.

4) Habitat data and nutritional condition of prey species

Where active habitat enhancement is occurring or was recommended in the Operational Plan, describe progress toward objectives:

Objective(s):

Not Applicable: There are no demonstrated methods to improve caribou habitat and no reason to believe that habitat is limiting the caribou population. Nutritional indices indicate the habitat is sustaining caribou at a very high nutritional level at this time.

Area treated and method: Not Applicable

Observation on treatment response: Not Applicable

Evidence of progress toward objective(s) (choose one: Apparent Statistical):

Not Applicable

Similar trend in nearby non-treatment areas? Not Applicable

Describe any substantial change in habitat not caused by active program: Not Applicable

Table 4. Nutritional indicators for caribou in assessment area (L) of the Mulchatna Caribou herd Predation Management Area.

Period	RY	Pregnancy Females >2 yrs age ^a	Female Calf Weights at 10.5 months in lbs. (n)
Year 0	2010	79%	124 (20)
Year 1	2011	78%	119 (13)
Year 2	2012	78%	127 (14)
Year 3	2013	90%	128 (14)

^a Pregnancy rate is based on known-aged animals from a collared sample of adult female caribou. Pregnancy status is determined in May based on observed characteristics of pregnancy (antler retention, udder development, and/or presence of a calf at heel).

Where objectives on nutritional condition were listed in the Operational Plan, describe trend in condition indices since inception of (a) habitat enhancement or (b) enhanced harvest: N/A

Evidence of trend: N/A

Similar trend in nearby non-treatment areas? N/A

5) Costs specific to implementing Intensive Management

Table 5. Cost (\$1000 = 1.0) of agency salary based on estimate of proportional time of field level staff and cost of operations for intensive management activities (e.g., predator control or habitat enhancement beyond normal Survey and Inventory work) performed by personnel in the Department or work by other state agencies (e.g., Division of Forestry) or contractors in Mulchatna Caribou Herd Predation Management Area. Fiscal year (FY) is also 1 July to 30 June but the year is one greater than the comparable RY (e.g, FY 2010 is 1 July 2009 to 30 June 2010).

Period	FY	Predation control ^a		Other IM activities		Total IM cost	Research cost ^d
		Time ^b	Cost ^c	Time ^b	Cost ^c		
Year 1	2012	0.0	0.0	1.0	36.0	36.0	415.0
Year 2	2013	0.0	0.0	0.5	6.0	6.0	421.2
Year 3	2014	0.0	0.0	0.3	3.0	3.0	176.3

^aState or private funds only.

^bPerson-months (22 days per month)

^cSalary plus operations

^dSeparate from implementing IM program but beneficial for understanding of ecological or human response to management treatment (scientific approach that is not unique to IM).